



HAL
open science

Robots, computer algebra and eight connected components

Jose Capco, Mohab Safey El Din, Josef Schicho

► **To cite this version:**

Jose Capco, Mohab Safey El Din, Josef Schicho. Robots, computer algebra and eight connected components. ISSAC '20: International Symposium on Symbolic and Algebraic Computation, Jul 2020, Kalamata / Virtual, Greece. pp.62-69, 10.1145/3373207.3404048 . hal-02925478

HAL Id: hal-02925478

<https://hal.sorbonne-universite.fr/hal-02925478v1>

Submitted on 29 Aug 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Robots, computer algebra and eight connected components

Jose Capco*, Mohab Safey El Din[†], Josef Schicho[‡]

August 29, 2020

Abstract

Answering connectivity queries in semi-algebraic sets is a long-standing and challenging computational issue with applications in robotics, in particular for the analysis of kinematic singularities. One task there is to compute the number of connected components of the complementary of the singularities of the kinematic map. Another task is to design a continuous path joining two given points lying in the same connected component of such a set. In this paper, we push forward the current capabilities of computer algebra to obtain computer-aided proofs of the analysis of the kinematic singularities of various robots used in industry.

We first show how to combine mathematical reasoning with easy symbolic computations to study the kinematic singularities of an infinite family (depending on parameters) modelled by the UR-series produced by the company “Universal Robots”. Next, we compute roadmaps (which are curves used to answer connectivity queries) for this family of robots. We design an algorithm for “solving” positive dimensional polynomial system depending on parameters. The meaning of solving here means partitioning the parameter’s space into semi-algebraic components over which the number of connected components of the semi-algebraic set defined by the input system is invariant. Practical experiments confirm our computer-aided proof and show that such an algorithm can already be used to analyze the kinematic singularities of the UR-series family. The number of connected components of the complementary of the kinematic singularities of generic robots in this family is 8.

1 Introduction

The individual parts of a serial robot, called links, are moved by controlling the angle of each joint connecting two links. The inverse kinematics problem asks for the values of all angles producing a desired position of the end effector, where “position” includes not

*Innsbruck University, Innsbruck, Austria, email:jose.capco@uibk.ac.at.

[†]Sorbonne Université, CNRS, LIP6, Paris, France, email:Mohab.Safey@lip6.fr.

[‡]JKU University, Linz, Austria, email:Josef.Schicho@risc.jku.at

just the location of the end effector in 3-space but also the orientation. In a sense, this means inverting a function which is called the forward kinematics map in robotics, which determines the position of the end effector for given angles by a well-known formula (see Section 2). In robot controlling, the inverse kinematics problem is often solved incrementally: starting from some known initial angle configuration and its corresponding end effector position, we want to compute the change of the angles required to achieve a desired small change in the end effector position.

Kinematic singularities are defined as critical points of the forward map, i.e. angle configurations where the Jacobian matrix of the forward map is rank deficient. There are two known facts that make rather difficulty to control a robot in a singular or near a singular configuration. First, if an end effector velocity or force outside the image of the singular Jacobian is desired, then the necessary joint velocity or torque is either not defined or very large (see [23] §4.3 and [31] §5.9). The second reason is that industrial controllers are based on Newton’s method for the incremental solution of the inverse problem, and this method is not guaranteed to converge if it is used with a starting point close to the singular set. For these reasons, engineers prefer to plan the robot movements avoiding kinematic singularities.

For a general serial robot with six joints, the singular set is a hypersurface defined locally by the Jacobian determinant of the forward map. Its complementary, a real manifold, is not connected. Counting the number of connected components of this manifold and answering connectivity queries in this set is then of crucial importance in this application domain.

Answering connectivity queries in semi-algebraic sets is a classical problem of algorithmic semi-algebraic geometry which has attracted a lot of attention through the development of the so-called ROADMAP algorithms (see e.g. [12, 11, 9, 6, 8, 27, 28, 21]). Up to our knowledge, such algorithms had never been developed enough and implemented efficiently to tackle real-life applications.

In this paper, we push forward the capabilities of computer algebra in this application domain by solving connectivity queries for the non singular configuration sets of industrial robots from the UR series of the company “Universal Robots”. For a particular robot in this series, the UR5, the number of components of the non singular configuration set is 8 (see Section 3). For two points in the same component, we show how to construct a connecting path, in two ways: either by an ad hoc way (which has its own algorithmic interest) taking advantage of the specialty of the geometric parameters of UR5, and by using the ROADMAP algorithm (see Section 5). Next, we go further and extend our analysis of UR5 to the whole UR series and prove that outside a proper Zariski closed set (UR5 is outside this closed set) the number of connected components of the non singular configuration set is constant. These are computer-aided mathematical proofs involving «easy» symbolic computations

The next contribution is based on the fact that the family of UR robots is determined by a finite list of real parameters. Hence, an algorithmic way of tackling the problem of analyzing kinematic singularities of the whole UR family is to «solve» a *positive dimensional* system *depending on parameters* (i.e. after specialization of the parameters, the specialized system

is positive dimensional). We design an algorithm that decomposes the parameter’s space into semi-algebraic subsets, such that the number of connected components of the non singular configurations is constant in each of these subsets. As far as we know, this is the first algorithm of that type which is designed.

We also implemented this algorithm and used it for the analysis of the kinematic singularities of the UR series. Computations are heavy but already doable (on a standard laptop) within 10 hours. This is a computational way to retrieve the same results as our computer-aided mathematical proofs. These computations show that computer algebra today is efficient enough to solve connectivity queries that are of practical interest in industrial robot applications.

2 Robotics problem formulation

We define a *manipulator* or *robot* as follows: we have finite ordered rigid bodies called *links* which are connected by n revolute *joints* that are also ordered. To each joint we associate a coordinate system or a *frame*. The links are connected in a serial manner i.e. if we consider the robot as a graph such that the vertices are joints and the edges are links then this graph is a path (the first and last joint has degree 1 and all other joints have degree 2) and the joints allow rotation about its axes, so that if a joint rotates then all other subsequent links rotate about the axes of this joint. A reference coordinate system is chosen for the final joint which is called the *end-effector*¹.

In theoretical kinematics one may forget that the links are rigid bodies so that *collision* between links are disregarded. In this case we may as well think of a robot as a differentiable map $F : \text{SO}(2)^n \rightarrow \text{SE}(3)$ where $\text{SO}(2)$ is the one-dimensional group of rotations around a fixed line, parameterised by the rotation angle, and $\text{SE}(3)$ is the six-dimensional group of Euclidean congruence transformations. This map is defined in the following way:

- The i -th coordinate of an element in $\text{SO}(2)^n$ is associated to the i -th (revolute) joint parameter.
- For joint values $\vec{\theta} := (\theta_1, \dots, \theta_n)$ in $\text{SO}(2)^n$, the image $F(\vec{\theta})$ is the transformation of the end-effector from the initial position corresponding to all angles being zero to the final position obtained by composing the n rotations.

The map F itself is called the *kinematic map* (of the robot). Its domain is called the *configuration space*, while its image is called the *work-space* or the *kinematic image*.

We use the Denavit-Hartenberg (DH) convention when describing relations between two joint frames. It is standard in robotics ; its advantages are discussed in e.g. [31, §3.2], [1, §4.2]. The transformation between the frames is given by the following rule:

- The z -axis of the reference frame will be the axis of rotation of the joint.

¹this is usually another frame, but this is just an additional fixed transformation in $\text{SE}(3)$ and w.l.o.g. we assume that the final offset, distance and twist is 0

- To obtain the next frame, one starts with a rotation about the z -axis of the reference frame, called the *rotation*, followed by
- a translation along the z -axis of the reference frame, called the *offset*, followed by
- a translation along the x -axis, called the *distance*, followed by
- a rotation about the x -axis, called the *twist*.

The transformation between frame i to frame $i + 1$ is

$$R_z(\theta_i)T_z(d_i)T_x(a_i)R_x(\alpha_i)$$

where R_z, T_z, T_x, R_x are rotations or translations with respect to z - or x -axis parameterised by the angle of rotation θ_i (the i -th joint parameter), the offset d_i , the distance a_i and the angle of twist α_i of the i -th frame. For a given robot with n joints all DH parameters except for the rotation are fixed values. So that image of F for given joint values (the rotations) $(\theta_1, \dots, \theta_n)$ is just the multiplication of these transformations in $\text{SE}(3)$. The parameters d_1, d_n, a_n, α_n are assumed to be 0. This is not a loss of generality, because we can freely choose the frame at the base and at the end-effector. More detailed discussion on these can be seen in [31].

Example 2.1. *The UR5 robot has the following DH parameters:*

distances (m.) $(a_1, \dots, a_6) := (0, -\frac{425}{1000}, -\frac{39225}{100000}, 0, 0, 0)$

offsets (m.) $(d_1, \dots, d_6) := (0, 0, 0, \frac{10915}{100000}, \frac{9465}{100000}, 0)$

twist angles (rad.) $(\alpha_1, \dots, \alpha_6) := (\frac{\pi}{2}, 0, 0, \frac{\pi}{2}, -\frac{\pi}{2}, 0)$

For example, the following joint angles (rotations, in rad.)

$$(\theta_1, \dots, \theta_6) := \left(\frac{1}{10}, \frac{2}{10}, \frac{3}{10}, \frac{4}{10}, \frac{5}{10}, \frac{6}{10} \right)$$

leads to the following transformation in $(R, \vec{t}) \in \text{SE}(3)$ (represented as elements in $\text{SO}(3) \times \mathbb{R}^3$) where:

$$R \simeq \begin{pmatrix} 0.047 & -0.977 & -0.209 \\ -0.393 & 0.174 & -0.903 \\ 0.918 & 0.123 & -0.376 \end{pmatrix}, \quad \vec{t} \simeq (-6.768, -1.7784, -3.336).$$

Definition. *Given the kinematic map of a manipulator $F : \text{SO}(2)^n \rightarrow \text{SE}(3)$, the kinematic singularities in the configuration space are the points $P \in (\mathbb{P}^1)^n$ such that the Jacobian of F at P is rank-deficient.*

In this paper, we will only deal with 6-jointed manipulators. Therefore the kinematic map is a differentiable map from the 6-dimensional configuration space $(\text{SO}(2))^6$ to the group $\text{SE}(3)$, which is also 6-dimensional. For non-singular points of the map, the Jacobian is therefore invertible, and F is a local homeomorphism. Here is a well-known geometric description of singularities.

Theorem 2.2. *Let $F : \text{SO}(2)^6 \rightarrow \text{SE}(3)$ be the kinematic map of a robot with 6 joints. Let $P \in \text{SO}(2)^6$. Then the following are equivalent.*

1. P is a kinematic singularity.
2. The Jacobian of F at P is singular.
3. If $P_1, \dots, P_6 \in \mathbb{P}^5(\mathbb{R})$ are the Plücker representation of the axes (lines in \mathbb{P}^3) of the joints of the robot at the configuration point P then the matrix consisting of the Plücker coordinates $(p_{i,j})_{i,j \leq 6}$ ($P_i = (p_{i,1} : p_{i,2} : \dots : p_{i,6})$ for $i = 1, \dots, 6$) is singular.

Proof. The equivalence of the first two items is clear by definition. The equivalence of the first and the third item is found in [30, §4.5.1], [23, §4.1] or [1, §4.5.1] \square

Assume that we have two non-singular points in the configuration set. As explained earlier, we want to decide if these two configurations can be connected by a curve of configurations which avoids the singular hypersurface (see [33] §1.2 for some history on this question). If yes, then an explicit construction of such a curve is also of interest. In order to tackle these problems, we choose parameters for $\text{SO}(2)$ so that the equation of the hypersurface becomes a polynomial. This is not the case when we use the angles $\theta_1, \dots, \theta_n$, because the Jacobian contains trigonometric functions in these angles. One well-known strategy is to parametrize by points on a unit circle, i.e. by two parameters satisfying the equation of the unit circle. This has a clear disadvantage: the number of variables increases, and the singular set has co-dimension greater than one. Another well-known strategy is to replace θ_i by $v_i = \tan \frac{\theta_i}{2}$. The variable v_i ranges over the projective line, and the angle π corresponds to the point at infinity. If we set $v_i = \tan \frac{\theta_i}{2}$ for $i = 1, \dots, n$, then we obtain, in general, a polynomial in v_2, \dots, v_5 . More precisely, the degree is 2 in v_2 and v_5 and degree 4 in v_3 and in v_4 . The Jacobian does not depend on the joint angles θ_1 and θ_6 . This is clear from the third characterization of singularities in Theorem 2.2: only the position of the axes are relevant, and a rotation along the first or the last axis does not change the position of any axis.

We define the *UR Family* to be robots having a similar DH-parameter as the known UR robots (UR5, UR10, etc.). Such *UR robots* are parameterised by the following DH parameters

distances (m.)	$(a_1, \dots, a_6) := (0, a_2, a_3, 0, 0, 0)$
offsets (m.)	$(d_1, \dots, d_6) := (0, 0, 0, d_4, d_5, 0)$
twist angles (rad.)	$(\alpha_1, \dots, \alpha_6) := (\frac{\pi}{2}, 0, 0, \frac{\pi}{2}, -\frac{\pi}{2}, 0)$

For these robots, the determinant of the Jacobian (see [34]), expressed as a polynomial in v_2, \dots, v_5 , is $A = -Bv_3v_5$ with

$$\begin{aligned}
B = & a_2v_2^2v_3^2v_4^2 - a_3v_2^2v_3^2v_4^2 - 2d_5v_2^2v_3^2v_4 - 2d_5v_2^2v_3v_4^2 - 2d_5v_2v_3^2v_4^2 + a_2v_2^2v_3^2 + a_2v_2^2v_4^2 - a_2v_3^2v_4^2 - a_3v_2^2v_3^2 \\
& + a_3v_2^2v_4^2 + 4a_3v_2v_3v_4^2 + a_3v_3^2v_4^2 + 2d_5v_2^2v_3 + 2d_5v_2^2v_4 + 2d_5v_2v_3^2 + 8d_5v_2v_3v_4 + 2d_5v_2v_4^2 + 2d_5v_3^2v_4 \\
& + 2d_5v_3v_4^2 + a_2v_2^2 - a_2v_3^2 - a_2v_4^2 + a_3v_2^2 + 4a_3v_2v_3 + a_3v_3^2 - a_3v_4^2 - 2d_5v_2 - 2d_5v_3 - 2d_5v_4 - a_2 - a_3
\end{aligned}$$

Note that there is a degree drop in three of the four cases: the degree in v_3 is only 3, and not 4, and the degree in v_4 is only 2, and not 4, and the degree in v_5 is only 1, and not

2. The drop in the degree means that the homogeneous form of the Jacobian has a linear factor that vanishes if and only if the value of the variable whose degree drops is infinity, or equivalently, that the corresponding angle is π . Since we are interested in the complement of the singular space, we may assume that none of these three angles is equal to π , and we can use the parameters v_3, v_4, v_5 without worrying about paths crossing infinity.

For the angle θ_2 , the situation is different. There is no degree drop, hence there are configurations with $\theta_2 = \pi$ in the non singular configuration set. If we use the parametrization by half angle, then we have to take paths in the projective line into account that cross infinity, or, in other words, consider this variable in the projective space $\mathbb{P}^1(\mathbb{R})$. But, to take advantage on algorithms acting on semi-algebraic sets, one needs variables that range over \mathbb{R} .

Hence, we instead use parameters $s_2 = \sin(\theta_2)$ and $c_2 = \cos(\theta_2)$ and add the additional equation $s_2^2 + c_2^2 = 1$, obtaining a polynomial in the variables s_2, c_2, v_3, v_4, v_5 with coefficients depending on the parameters a_2, a_3, d_4, d_5 . Of course, the reason for this more costly treatment for θ_2 is just necessary if we use the ROADMAP algorithm subsequently. For an alternative analysis not using it, it is still better to use the half tangent ranging over the projective line.

3 Analysis of the UR5 Robot

We gave in Example 2.1 the Denavit-Hartenberg parameters of the UR5 robot. These values are used to instantiate a_2, a_3 and d_5 in the above polynomial B ; the specialized polynomial is then denoted by \tilde{B} and we let $\tilde{A} = \tilde{B}v_3v_5$. Recall that v_2 ranges over \mathbb{P}^1 , while v_3, v_4, v_5 range only over the affine line.

We investigate the discriminant of \tilde{B} with respect to the variable v_2 (thus the projection of the critical set to the (v_3, v_4) -plane). The discriminant of \tilde{B} with respect to the variable v_2 we denote as $b \in \mathbb{R}[v_3, v_4]$. This discriminant is still factorisable in $\mathbb{C}[v_3, v_4]$. In fact, one checks, that it is the factor of two complex conjugates of some polynomial in $\mathbb{R}[v_3, v_4]$. This implies that $b = c^2 + d^2$ is the sum of two squares of real polynomials $c, d \in \mathbb{R}[v_3, v_4]$. These two polynomials are given by

$$c = \frac{1577212v_3 - 3561263v_4 - 14850585}{\sqrt{2006237}}$$

$$d = \frac{(\sqrt{2006237}v_4 + 1239915 - 7144712)v_3 + 16090500v_4}{\sqrt{2006237}}$$

Thus, b can have only two real roots (i.e. two pairs (v_3, v_4)), i.e the vanishing set of b in \mathbb{R}^2 is finite, namely they points that are the zeros of both c and d . We solve this as floating numbers to have an idea of their vicinity in an affine chart of the ambient space of the kinematic singularity. The roots are

$$q_1 = (v_3 \simeq -9.140975564 \quad , \quad v_4 \simeq -8.218388067)$$

$$q_2 = (v_3 \simeq 9.140975563 \quad , \quad v_4 \simeq -1216783622)$$

For the two special values q_1 and q_2 in the (v_3, v_4) -plane, all three coefficients of \tilde{B} with respect to v_2 are zero.

Now, since the discriminant b is positive except at these two points and since \tilde{B} itself is quadratic with respect to v_2 we conclude that the preimage of the projection (to (v_3, v_4) -plane) are two real points in the variety defined by \tilde{B} . Thus, the variety defined by \tilde{B} is composed of two sheets (above any two points (v_3, v_4) except q_1 and q_2). Let X be the complement of the vanishing points of \tilde{B} in $\mathbb{P}^1 \times \mathbb{A}^2$. Set

$$Y := \mathbb{A}^2 \setminus (\{q_1, q_2\} \cup \{(0, v_4) \mid v_4 \in \mathbb{R}\})$$

So we have a canonical projection (to the (v_3, v_4) -plane) from $X \cap (\mathbb{P}^1 \times Y)$ to Y . The fiber of this projection is a projective line without two distinct points. Hence, every fiber has two components. The sign of \tilde{B} is different for the two components of each fiber. Then, we have two components of X for each component of Y . Obviously, Y has two components, hence we have a total number of 4 components.

For the non singular set, which is the complement of the zero set of \tilde{A} , we get 8 components: for each component of X , we have one component where v_5 is positive and one where v_5 is negative.

Now assume that we have two non singular configuration points x, y in the same component, and we want to construct a path connecting them. The projections to Y have to lie in the same component of Y , and because Y is the plane without a line and two points, it is easy to connect the images of the projections in Y : in most cases, a straight line segment is fine; if the straight line segment connecting the two image points contains q_1 or q_2 , we have to do a random detour via a third point. The zero set of \tilde{B} is a two-sheeted covering of Y . So, for any value of Y , we have two points in the zero set of \tilde{B} projecting to it. If we look at these points as points in $\text{SO}(2)$, then it is clear that there are two ‘‘midpoints’’ in the zero set of \tilde{B} , which have equal angle distance to these two points. The value of \tilde{B} is positive for one of the two midpoints and negative for the other one. The sign of $\tilde{B}(x)$ and $\tilde{B}(y)$, however, must be the same because the two points are in the same connected component. Suppose, without loss of generality, that $\tilde{B}(x)$ and $\tilde{B}(y)$ are both positive. Then we first connect x to the midpoint over the projection of x to Y with positive sign, by a curve in the fiber. Next, we lift the path in Y , connecting the projections of x and y in the same component, to a path of midpoints with positive sign, arriving at the midpoint with positive sign lying over the projection of y . Finally, we connect this midpoint to y by the other fiber.

Below, we show the sheets in Figure 1 to illustrate that:

- (i) the regions above and below the sheets can be connected
- (ii) the region between the two sheets is the other connected component
- (iii) the two points q_2 and q_3 are points in the projection where the sheets get connected (see asymptotes in Fig. 1). Thus, the variety describing the two sheets is connected.




Figure 1: The two sheets of \tilde{B}

4 UR series

We can make a general statement for robots belonging to the UR family (e.g. UR10, UR3 etc.). We define the *UR Family* to be robots which have a similar DH-parameter as the known UR robots (UR5, UR10 etc.), a robot in this family we shall call a *UR robot*. Namely they are parameterised by the following DH parameters :

$$\begin{aligned} \text{distances (m.)} & & (a_1, \dots, a_6) & := (0, a_2, a_3, 0, 0, 0) \\ \text{offsets (m.)} & & (d_1, \dots, d_6) & := (0, 0, 0, d_4, d_5, 0) \\ \text{twist angles (rad.)} & & (\alpha_1, \dots, \alpha_6) & := (\frac{\pi}{2}, 0, 0, \frac{\pi}{2}, -\frac{\pi}{2}, 0) \end{aligned}$$

i.e. these robots are parameterised by 4 parameters: a_2, a_3, d_4, d_5 .

We can write the largest (in number of terms and in degree) polynomial factor of the polynomial whose vanishing points is the kinematic singularity in configuration space of a UR robot as

$$\begin{aligned} B = & a_2 v_2^2 v_3^2 v_4^2 - a_3 v_2^2 v_3^2 v_4^2 - 2d_5 v_2^2 v_3^2 v_4 - 2d_5 v_2^2 v_3 v_4^2 - 2d_5 v_2 v_3^2 v_4^2 + a_2 v_2^2 v_3^2 + a_2 v_2^2 v_4^2 - a_2 v_3^2 v_4^2 - a_3 v_2^2 v_3^2 \\ & + a_3 v_2^2 v_4^2 + 4a_3 v_2 v_3 v_4^2 + a_3 v_3^2 v_4^2 + 2d_5 v_2^2 v_3 + 2d_5 v_2^2 v_4 + 2d_5 v_2 v_3^2 + 8d_5 v_2 v_3 v_4 + 2d_5 v_2 v_4^2 + 2d_5 v_3^2 v_4 \\ & + 2d_5 v_3 v_4^2 + a_2 v_2^2 - a_2 v_3^2 - a_2 v_4^2 + a_3 v_2^2 + 4a_3 v_2 v_3 + a_3 v_3^2 - a_3 v_4^2 - 2d_5 v_2 - 2d_5 v_3 - 2d_5 v_4 - a_2 - a_3 \end{aligned}$$

Note that d_4 does not affect the singularity of the robot. Taking the discriminant of B with respect to v_2 yields the sum of two squares i.e. the product of two quadratic complex conjugate polynomials $\text{disc}(B, v_2) = g\bar{g}$.

$$g = (-a_2 v_3 v_4 + a_3 v_3 v_4 + d_5 v_3 + d_5 v_4 + a_2 + a_3) + (-d_5 v_3 v_4 + a_2 v_3 + a_2 v_4 - a_3 v_3 + a_3 v_4 + d_5)i$$

For a robot determined by some real quadruple $u \in \mathbb{R}^4$, let A_u, B_u, g_u be the polynomials obtained by instantiating in A, B, g the variables a_2, a_3, d_4, d_5 by the corresponding real values in the quadruple. Let $f : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be the projection $(v_2, v_3, v_5) \mapsto (v_3, v_5)$. Let $Y_u \subset \mathbb{R}^2$ be the complement of (the union of the line $v_3 = 0$ and the common zero set of $\text{Re } g_u$ and $\text{Im } g_u$). Then the real zero set of B_u in \mathbb{R}^3 intersected with $f^{-1}(Y_u)$ projects to surjectively to Y_u , in such a way that there are two sheets, each projecting homeomorphically to Y_u .

For general robot u , the real set of g , which is meaning the set of all points in the real (v_3, v_4) -plane such that both the real part and the imaginary part of g is equal to zero, is a finite subset of \mathbb{R}^2 . All arguments from the previous sections work in this case as well. Hence we get 8 components for these parameters' values. Moreover, we have paths connecting points in the same component, as in the previous section.

It remains to treat the non-general robots where the real zero set of g is one-dimensional. This is the case if and only if $d_5 = a_2^2 - a_3^2 = 0$. The even more special case $d_5 = a_2 = a_3 = 0$ is easy to analyze: here, the determinant of the Jacobian A is identically zero, which means that there are no non singular configurations. Excluding that case, we have two families of robots, and in each family, up to the value of d_4 , the parameters are unique up to scaling. Without loss of generality, we can reduce to exactly two non-general robots $u' = (1, 1, 0, 0)$ and $u'' = (1, -1, 0, 0)$. Then the polynomial $B_{u'}$ has a factor is $C' := v_3v_4 - v_3 - v_4 - 1$, and the polynomial $B_{u''}$ has a factor $C'' := v_3v_4 + v_3 + v_4 - 1$. Apart from that complication, the analysis proceeds similar as in the general case: the set $Y_{u'}$ is the plane minus the line $v_3 = 0$ minus the hyperbola with equation C' , and the set $Y_{u''}$ is the plane minus the hyperbola with equation C'' . In both cases, the number of components of Y is 5, as it can be seen in Figure 2. Consequently, we have 20 components in total. The paths between points in the same component can be constructed similarly as in the general case.

5 Connectivity and roadmaps

We explain the ROADMAP algorithm for the special case where the semi-algebraic set S is given as a subset of some vector space \mathbb{R}^N , $N \in \mathbb{N}$, defined by an equation $f(x_1, \dots, x_N) = 0$ and an inequation $g(x_1, \dots, x_N) \neq 0$. We assume that the algebraic set defined by $f = 0$ is smooth. This is sufficient for our application: the inequation is the determinant of the Jacobian of the kinematic map A , and the equality is $s_2^2 + c_2^2 - 1 = 0$.

One first reduces the problem to one where the semi-algebraic set we consider is bounded. Note that there exists $R > 0$ large enough such that the connected components of S are in one-to-one correspondence with the intersection of S with the hyper-ball defined by $\mathcal{N}_R \leq 0$ where $\mathcal{N}_R = x_1^2 + \dots + x_n^2 - R$. We denote this intersection by S' . Note that a roadmap of S' provides a roadmap of S .

Determining such a large enough real number R is done by choosing it larger than the largest critical value of the restriction of the map $\mathbf{x} \rightarrow \|\mathbf{x}\|^2$ to each regular strata of the the Euclidean closure of S . This leads us to compute critical values of that map restricted to the hypersurface defined by $f = 0$ and next take the limits of the critical values of the sets defined by $g = \pm\varepsilon$ and $f = g \pm \varepsilon = 0$ when $\varepsilon \rightarrow 0$.

Next, we compute the critical values $\eta_1 < \dots < \eta_s$ of the restriction of the map $\mathbf{x} \rightarrow g(\mathbf{x})$ to the semi-algebraic set defined by $f = 0$ and $\mathcal{N}_R \leq 0$. Following Thom's isotopy lemma [13],




Figure 2: Left (resp. right) shows the components of $v_3(v_3v_4 + v_3 + v_4 - 1) \neq 0$ (resp. $v_3(v_3v_4 - v_3 - v_4 - 1) \neq 0$) in Y

when e is chosen between 0 and $\min(|\eta_i|, 1 \leq i \leq s)$, the connected components of the semi-algebraic set S_e^+ (resp. S_e^-) defined by $\mathcal{N}_R \leq 0, f = g - e = 0$ (resp. $\mathcal{N}_R \leq 0, f = g + e = 0$) are in one-to-one correspondence with the connected components of the semi-algebraic set defined by $\mathcal{N}_R \leq 0, f = 0, g > 0$ (resp. $\mathcal{N}_R \leq 0, f = 0, g > 0$). Besides, $S_e^+ \subset S'$ (resp. $S_e^- \subset S'$). Then a roadmap of S' is obtained by taking the union of a roadmap of S_e^+ with the roadmap of S_e^- . Hence, we have performed a reduction to computing roadmaps in the compact semi-algebraic sets S_e^+ and S_e^- .

In our application, the algebraic sets defined by the vanishing of all subsets of the defining polynomials of S_e^+ and S_e^- are smooth. Hence, we can rely on a slight modification of the roadmap algorithm given in [12] where we replace computations with multivariate resultants for solving polynomial systems by computations of Gröbner bases.

The algorithm in [12] then takes as input a polynomial system defining a closed and bounded semi-algebraic set S and proceeds as follows. The core idea is to start by computing a curve \mathcal{C} which has a non-empty intersection with each connected component of S . That curve will be typically the critical locus on the (x_1, x_2) -plane when one is in generic coordinates (else, one just needs to change linearly generically the coordinate system). A few remarks are in order here. When S is defined by $f_1 = \dots = f_p = 0$ and $g_1 \geq 0, \dots, g_s \geq 0$, to define the critical locus of the projection on the (x_1, x_2) -plane restricted to S one takes the union of the critical loci of that projection restricted to the real algebraic sets defined for all $\{i_1, \dots, i_\ell\} \subset \{1, \dots, s\}$, by $f_1 = \dots = f_p = g_{i_1} = \dots = g_{i_\ell} = 0$ and intersect this union of critical loci with S (see [12]).

That way, one obtains curves that intersect all connected components of S but these intersections may not be connected. To repair these connectivity failures, Canny's algorithm finds appropriate slices of S . Let π_1 be the canonical projection $(x_1, \dots, x_n) \rightarrow x_1$. This basically consists in finding $\alpha_1 < \dots < \alpha_k$ in \mathbb{R} such that the union of $\cup_{i=1}^k S \cap \pi_1^{-1}(\alpha_i)$ with the critical curve \mathcal{C} has a non-empty and connected intersection with each connected component of S .

The way Canny proposes to find those α_i 's is to compute the critical values of the restriction of π_1 to \mathcal{C} . By the algebraic Sard's theorem (see e.g. [28, Appendix B]), these values are in finite number and Canny proposes to take $\alpha_1, \dots, \alpha_k$ as those critical values. This leads to compute with real algebraic numbers which can be encoded with their minimal polynomials and isolating intervals. Since these minimal polynomials may have large degrees (singly exponential in n), that step can be prohibitive for practical computations. We use then the technique introduced in [22] which consists in replacing $\alpha_1 < \dots < \alpha_k$ with rational numbers $\rho_1 < \dots < \rho_{k-1}$ with $\alpha_i < \rho_i < \alpha_{i+1}$. We refer to [22] for the rationale justifying this trick. All in all, one obtains a recursive algorithm with a decreasing number of variables at each recursive call. Combined with efficient Gröbner bases engines, we illustrate in Section 7 that the ROADMAP algorithm (with the modifications introduced above) can be used in practice to answer connectivity queries in semi-algebraic sets in concrete applications.

The concept of roadmap and the algorithm computing it, described above, may seem cumbersome and unnecessarily sophisticated, especially when compared with the much more direct CAD approach [29]. The CAD algorithm is also a recursive algorithm, producing its recursive instance by projecting the hypersurface to \mathbb{R}^{n+1} and analyzing the discriminant.

This leads to an iteration of discriminants, and it is easy to see that the degree of the iterated discriminants grows double exponentially in n : roughly, the degree of the discriminant is squared in every iteration. There lies the motivation for all the sophistication of the ROADMAP algorithm: for each instance in the all recursive calls, the degree of the input polynomial is exactly the same as the degree of the initially given polynomial f . This leads to an asymptotic complexity which is only single exponential in n^2 . We refer to [27, 9, 28, 8] for more recent algorithms improving the complexity of roadmap computations.

6 Parametric polynomial systems

Let $F = (f_1, \dots, f_p)$ and $G = (g_1, \dots, g_q)$ in $\mathbb{Q}[\mathbf{x}, \mathbf{y}]$ with $\mathbf{x} = (x_1, \dots, x_n)$ and $\mathbf{y} = (y_1, \dots, y_t)$. We consider further \mathbf{y} as a sequence of parameters and the polynomial system

$$f_1 = \dots = f_p = 0, \quad g_1 \sigma_1 0, \dots, g_q \sigma_q 0$$

with $\sigma_i \in \{>, \geq\}$. We let $S \subset \mathbb{R}^n \times \mathbb{R}^t$ be the semi-algebraic set defined by this system. For $y \in \mathbb{R}^t$, we denote by F_y and G_y the sequences of polynomials obtained after instantiating \mathbf{y} to y in F and G respectively. Also, we denote by $S_y \subset \mathbb{R}^n$ the semi-algebraic set defined by the above system when \mathbf{y} is specialized to y . The algebraic set defined by the simultaneous vanishing of the entries of F (resp. F_y) is denoted by $V(F) \subset \mathbb{C}^{n+t}$ (resp. $V(F_y) \subset \mathbb{C}^n$). We describe an algorithm for solving such a parametric polynomial system *without* assuming that for a *generic* point y in \mathbb{C}^t , $V(F_y)$ is finite. In that situation, solving such a parametric polynomial system may consist in partitioning the parameters' space \mathbb{R}^t into semi-algebraic sets T_1, \dots, T_r such that, for $1 \leq i \leq r$, the number of connected components of S_y is invariant for any choice of y in T_i . We prove below that such an algorithmic problem makes sense.

Proposition 6.1. *Let $S \subset \mathbb{R}^n \times \mathbb{R}^t$ be a semi-algebraic set and π be the canonical projection*

$$(x_1, \dots, x_n, y_1, \dots, y_t) \rightarrow (y_1, \dots, y_t).$$

There exist semi-algebraic sets T_1, \dots, T_r in \mathbb{R}^t such that

- $\mathbb{R}^t = T_1 \cup \dots \cup T_r$,
- *there exists $b_i \in \mathbb{N}$ such that for any $y \in T_i$, the number of connected components of S_y is b_i .*

Proof. Observe that the restriction of π to S is semi-algebraically continuous. From Hardt's semi-algebraic triviality theorem [10, Theorem 9.3.2], there exists a finite partition of \mathbb{R}^t into semi-algebraic sets T_1, \dots, T_r and for each $1 \leq i \leq r$, a trivialization $\vartheta_i : T_i \times E_i \rightarrow \pi^{-1}(T_i) \cap S$ (where E_i is a fiber $\pi^{-1}(y) \cap S$ for some $y \in T_i$). Fix i and choose an arbitrary point $y' \in T_i$. Observe that we are done once we have proved that $\pi^{-1}(y') \cap S$ and E_i have the same number of connected components. Recall that, by definition of a trivialization (see [10, Definition 9.3.1]), $\theta_i : T_i \times E_i \rightarrow \pi^{-1}(T_i) \cap S$ is a semi-algebraic homeomorphism and for any $(y', x) \in T_i \times E_i$, $\pi \circ \theta_i(y', x) = y'$. Hence, we deduce that E_i is homeomorphic to $\pi^{-1}(y') \cap S$. As a consequence, they both have the same number of connected components. \square

Instead of computing a partition of the parameters' space into semi-algebraic sets T_1, \dots, T_r as above, one will consider non-empty *disjoint* open semi-algebraic sets U_1, \dots, U_ℓ in \mathbb{R}^t such that the complement of $U_1 \cup \dots \cup U_\ell$ in \mathbb{R}^t is a semi-algebraic set of dimension less than t and such that for $1 \leq i \leq t$, there exists $b_i \in \mathbb{N}$ such that b_i is the number of connected components of S_y for any $y \in U_i$. For instance, one can take U_1, \dots, U_ℓ as the non-empty interiors (for the Euclidean topology) of T_1, \dots, T_r .

Our strategy to solve this problem is to first compute a polynomial Δ in $\mathbb{Q}[\mathbf{y}] - \{0\}$ defining a Zariski closed set $\mathcal{D} \subset \mathbb{C}^t$ such that \mathcal{D} contains $\mathbb{R}^t - (U_1 \cup \dots \cup U_\ell)$. The next lemma is immediate.

Lemma 6.2. *Let $\mathcal{E} \subset \mathbb{R}^t$ be a finite set of points which has a non-empty intersection with any of the connected components of the semi-algebraic set defined by $\Delta \neq 0$. For $1 \leq i \leq \ell$, $\mathcal{E} \cap U_i$ is not empty.*

Hence, computing sample points in each connected component of the set defined by $\Delta \neq 0$ (e.g. using the algorithm in [26] applied to the set defined by $z\Delta - 1 = 0$ where z is a new variable) is enough to obtain at least one point per connected component of U_1, \dots, U_ℓ . Finally, for each such a point y , it remains to count the number of connected components of the set S_y by using a roadmap algorithm.

We call *partial semi-algebraic resolution* of (F, G) the data $(b_1, \eta_1), \dots, (b_k, \eta_k)$ where b_i is the number of connected components of S_{η_i} and $\{\eta_1, \dots, \eta_k\}$ has a non-empty intersection with each connected component of $U_1 \cup \dots \cup U_\ell$.

Hence, our algorithm relies on three subroutines. The first one, which we call **Eliminate**, takes as input F and G , as well as \mathbf{x} and \mathbf{y} and outputs $\Delta \in \mathbb{Q}[\mathbf{y}]$ as above ; we let $\mathcal{D} = V(\Delta)$. The second one, which we call **SamplePoints** takes as input Δ and outputs a finite set of sample points $\{\eta_1, \dots, \eta_k\}$ (with $\eta_i \in \mathbb{Q}^t$) which meets each connected component of $\mathbb{R}^t - \mathcal{D}$. The last one, which we call **NumberOfConnectedComponents** takes F_η and G_η and for some $\eta \in \mathbb{Q}^t$ and computes the number of connected components of the semi-algebraic set S_η . The algorithm is described hereafter.

Algorithm 1: ParametricSolve($F, G, \mathbf{x}, \mathbf{y}$)

Data: Finite sequences F and G in $\mathbb{Q}[\mathbf{x}, \mathbf{y}]$ with $\mathbf{x} = (x_1, \dots, x_n)$ and $\mathbf{y} = (y_1, \dots, y_t)$.

Result: a partial semi-algebraic resolution of (F, G)

- 1 $\Delta \leftarrow \text{Eliminate}(F, G, \mathbf{x}, \mathbf{y})$
 - 2 $\{\eta_1, \dots, \eta_k\} \leftarrow \text{SamplePoints}(\Delta \neq 0)$
 - 3 **for** i from 1 to k **do**
 - 4 $b_i = \text{NumberOfConnectedComponents}(F_{\eta_i}, G_{\eta_i})$
 - 5 **end**
 - 6 **return** $\{(b_1, \eta_1), \dots, (b_k, \eta_k)\}$.
-

While the rationale of algorithm `ParametricSolve` is mostly straightforward, detailing each of its subroutines is less. The easiest ones are `SamplePoints` and `NumberOfConnectedComponents`: they rely on known algorithms using the critical point method [5, 7], polar varieties [26, 24, 4, 3] and for computing roadmaps [6, 9, 28, 27].

The most difficult one is subroutine `Eliminate`. We provide a detailed description of it under the following regularity assumption. We say that (F, G) satisfies assumption (A)

- (A) for any $\{i_1, \dots, i_s\}$ in $\{1, \dots, q\}$, the Jacobian matrix associated to $(f_1, \dots, f_p, g_{i_1}, \dots, g_{i_s})$ has maximal rank at any complex solution to

$$f_1 = \dots = f_p = g_{i_1} = \dots = g_{i_s} = 0$$

Note that using the Jacobian criterion [14, Chap. 16], it is easy to decide whether (A) holds. Note also that it holds generically.

For $\mathbf{i} = \{i_1, \dots, i_s\} \subset \{1, \dots, q\}$, under assumption (A), the algebraic set $V_{\mathbf{i}} \subset \mathbb{C}^{n+t}$ defined by

$$f_1 = \dots = f_p = g_{i_1} = \dots = g_{i_s} = 0.$$

are smooth and equidimensional and these systems generate radical ideals (applying the Jacobian criterion [14, Theorem 16.19]). Besides, the tangent space to $z \in V_{\mathbf{i}}$ coincides with the the (left) kernel of the Jacobian matrices associated to $(f_1, \dots, f_p, g_{i_1}, \dots, g_{i_s})$ at z .

Let I be the ideal generated by $(f_1, \dots, f_p, g_{i_1}, \dots, g_{i_s})$ and the maximal minors of the truncated Jacobian matrix associated to $(f_1, \dots, f_p, g_{i_1}, \dots, g_{i_s})$ obtained by removing the columns corresponding to the partial derivatives w.r.t. the \mathbf{y} -variables. Under assumption (A), one can compute the set of critical values of the restriction of the projection π to the algebraic set $V_{\mathbf{i}}$ by eliminating the variables \mathbf{x} from I .

Hence, using elimination algorithms, which include Gröbner bases [15, 16] with elimination monomial orderings, or triangular sets (see e.g. [32, 2]) or geometric resolution algorithms [20, 18, 19], one can compute a polynomial $\Delta_{\mathbf{i}} \in \mathbb{Q}[\mathbf{y}]$ whose vanishing set is the set of critical values of the restriction of π to $V_{\mathbf{i}}$. By the algebraic Sard's theorem (see e.g. [28, App. A]), $\Delta_{\mathbf{i}}$ is not identically zero (the critical values are contained in a Zariski closed subset of \mathbb{C}^t).

Under assumption (A), we define the set of critical points (resp. values) of the restriction of π to the Euclidean closure of S as the union of the set of critical points (resp. values) of the restriction of π to $V_{\mathbf{i}} \cap \mathbb{R}^{n+t}$ when \mathbf{i} ranges over the subsets of $\{1, \dots, q\}$. We denote the Euclidean closure of S by \overline{S} , the set of critical points (resp. values) of the restriction of π to \overline{S} by $\mathscr{W}(\pi, \overline{S})$ (resp. $\mathscr{D}(\pi, \overline{S})$).

We say that S satisfies a *properness* assumption (P) if:

- (P) the restriction of π to \overline{S} is proper ($\forall y \in \mathbb{R}^t$, there exists a ball $B \ni y$ s.t. $\pi^{-1}(B) \cap \overline{S}$ is closed and bounded).

Our interest in critical points and values is motivated by the semi-algebraic version of Thom's isotopy lemma (see [13]) which states the following, under assumption (P). Take an open semi-algebraic subset $U \subset \mathbb{R}^t$ which does not meet the set of critical values of

the restriction of π to \overline{S} , $y \in U$ and $E = \pi^{-1}(y) \cap S$. Then, there exists a semi-algebraic trivialization $\vartheta : U \times E \rightarrow \pi^{-1}(U) \cap S$.

Hence, $\cup_{i \in \{1, \dots, q\}} \mathcal{D}(\pi, V_i)$ contains the boundaries of the open disjoint semi-algebraic set U_1, \dots, U_ℓ . Recall that by Sard's theorem it has co-dimension ≥ 1 . This leads to the following algorithm.

Algorithm 2: EliminateProper($F, G, \mathbf{x}, \mathbf{y}$)

Data: Finite sequences F and G in $\mathbb{Q}[\mathbf{x}, \mathbf{y}]$ with $\mathbf{x} = (x_1, \dots, x_n)$ and $\mathbf{y} = (y_1, \dots, y_t)$, defining a semi-algebraic set $S \subset \mathbb{R}^n \times \mathbb{R}^t$.

Assumes that assumptions (A) and (P) hold.

Result: $\Delta \in \mathbb{Q}[\mathbf{y}]$ such that π realizes a fibration over all connected components of $\mathbb{R}^t - \{\Delta = 0\}$

- 1 **for** all subsets i in $\{1, \dots, q\}$ **do**
 - 2 $\mathcal{M} \leftarrow$ maximal minors of $\text{jac}([F, G_i], \mathbf{x})$
 - 3 $\Delta_i \leftarrow$ AlgebraicElimination($[F, G_i, \mathcal{M}], \mathbf{x}$)
 - 4 **end**
 - 5 $\Delta \leftarrow \prod_i \Delta_i$.
 - 6 **return** Δ .
-

Lemma 6.3. *On input (F, G) in $\mathbb{Q}[\mathbf{x}, \mathbf{y}]$ satisfying (A), algorithm EliminateProper is correct.*

For some applications, deciding if (P) holds is easy (e.g. when the inequalities in G define a box). However, in general, one needs to generalize EliminateProper to situations where (P) does not hold.

To do so, we use a classical technique from effective real algebraic geometry. Let ε be an infinitesimal and $\mathbb{R}\langle\varepsilon\rangle$ be the field of Puiseux series in ε with coefficients in \mathbb{R} . By [7, Chap. 2], $\mathbb{R}\langle\varepsilon\rangle$ is a real closed field and one can define semi-algebraic sets over $\mathbb{R}\langle\varepsilon\rangle^{n+t}$. In particular the set solutions in $\mathbb{R}\langle\varepsilon\rangle^{n+t}$ to the system defining S is a semi-algebraic set which we denote by $\text{ext}(S, \mathbb{R}\langle\varepsilon\rangle)$. We refer to [7] for properties of real Puiseux series fields and semi-algebraic sets defined over such field. We make use of the notions of bounded points of $\mathbb{R}\langle\varepsilon\rangle^n$ over \mathbb{R} (those whose all coordinates have non-negative valuation) and their limits in \mathbb{R} (when $\varepsilon \rightarrow 0$). We denote by \lim_0 the operator taking the limits of such points. For $a = (a_1, \dots, a_n)$, we consider the intersection of $\text{ext}(S, \mathbb{R}\langle\varepsilon\rangle)$ with the semi-algebraic set defined by

$$\Phi^{(a)} = a_1 x_1^2 + \dots + a_n x_n^2 - 1/\varepsilon \leq 0$$

where $a_i > 0$ in \mathbb{R} for $1 \leq i \leq n$. We denote by S'_ε this intersection. Since $a_i > 0$ for all $1 \leq i \leq n$, S'_ε satisfies (P).

Lemma 6.4. *Assume that (F, G) satisfies (A). There exists a non-empty Zariski open set $\mathcal{A} \subset \mathbb{C}^n$ such that for any choice of $a = (a_1, \dots, a_n) \in \mathcal{A}$, $(F, G^{(a)})$ satisfies (A) with $G^{(a)} = G \cup \{\Phi^{(a)}\}$.*

Proof. Let $\mathbf{i} = \{i_1, \dots, i_s\} \subset \{1, \dots, q\}$. We prove below that there exists a non-empty Zariski open set $\mathcal{A}_{\mathbf{i}} \subset \mathbb{C}^n$ such that for $(a_1, \dots, a_n) \in \mathcal{A}_{\mathbf{i}}$, the following property $(\mathbf{A})_{\mathbf{i}}$ holds. Denoting by $G^{(a), \mathbf{i}}$ the sequence $(g_{i_1}, \dots, g_{i_s}, \Phi^{(a)})$, the Jacobian matrix of $(F, G^{(a), \mathbf{i}})$ has maximal rank at any point of $V(F, G^{(a), \mathbf{i}})$. Taking the intersection of the (finitely many) $\mathcal{A}_{\mathbf{i}}$'s is then enough to define \mathcal{A} .

Consider new indeterminates $\alpha_1, \dots, \alpha_n$ and the polynomial $\Phi^{(\alpha)} = \alpha_1 x_1^2 + \dots + \alpha_n x_n^2 - 1/\varepsilon$. Let Ψ be the map

$$\Psi : (x, a) \rightarrow F(x), g_{i_1}(x), \dots, g_{i_s}(x), \Phi^{(a)}(x)$$

Observe that $\mathbf{0}$ is a regular value for Ψ since (F, G) satisfies (\mathbf{A}) . Hence, Thom's weak transversality theorem (see e.g. [28, App. B]) implies that there exists $\mathcal{A}_{\mathbf{i}}$ such that $(\mathbf{A})_{\mathbf{i}}$ for any $a \in \mathcal{A}_{\mathbf{i}}$. \square

Assume for the moment that (F, G') satisfies assumption (\mathbf{A}) . Observe that the coefficients of F and G' lie in $\mathbb{Q}(\varepsilon)$. Hence, applying the subroutine `EliminateProper` to (F, G') and the above inequality will output a polynomial $\Delta_\varepsilon \in \mathbb{Q}(\varepsilon)[\mathbf{y}]$ such that the restriction of π to $\overline{S'_\varepsilon}$ realizes a trivialization over each connected component of $\mathbb{R}\langle\varepsilon\rangle^t - \{\Delta_\varepsilon = 0\}$. Without loss of generality, one can assume that $\Delta_\varepsilon \in \mathbb{Q}[\varepsilon][\mathbf{y}]$ and has content 1. In other words, one can write $\Delta_\varepsilon = \Delta_0 + \varepsilon \tilde{\Delta}$ with $\Delta_0 \in \mathbb{Q}[\mathbf{y}]$ and $\tilde{\Delta} \in \mathbb{Q}[\varepsilon][\mathbf{y}]$.

Lemma 6.5. *Let U be a connected component of $\mathbb{R}^t - \{\Delta_0 = 0\}$. Then, there exists a semi-algebraically connected component U_ε of $\mathbb{R}\langle\varepsilon\rangle^t - \{\Delta_\varepsilon = 0\}$ such that $\text{ext}(U, \mathbb{R}\langle\varepsilon\rangle) \subset U_\varepsilon$.*

Proof. Let y and y' be two distinct points in U . Since U is a semi-algebraically connected component of $\mathbb{R}^t - \{\Delta_0 = 0\}$, there exists a semi-algebraic continuous function $\gamma : [0, 1] \rightarrow U$ with $\gamma(0) = y$ and $\gamma(1) = y'$ such that Δ_0 is sign invariant over $\gamma([0, 1])$ (assume, without loss of generality that it is positive). Note also for all $t \in [0, 1]$, $\Delta_0(\gamma(t)) \in \mathbb{R}$. We deduce that $\Delta_\varepsilon(\gamma(t)) > 0$ for all $t \in [0, 1]$. Now, take $\vartheta \in \text{ext}([0, 1], \mathbb{R}\langle\varepsilon\rangle)$. Observe that ϑ is bounded over \mathbb{R} and then $\lim_0 \vartheta$ exists and lies in $[0, 1]$. We deduce that $\Delta_\varepsilon(\lim_0 \vartheta) > 0$ and its limit when $\varepsilon \rightarrow 0$ is $\Delta_0(\lim_0 \vartheta) > 0$ in \mathbb{R} . We deduce that $\Delta_\varepsilon(\vartheta) > 0$. Hence, Δ_ε is sign invariant over $\text{ext}(\gamma([0, 1]), \mathbb{R}\langle\varepsilon\rangle)$ and then y and y' both lie in the same semi-algebraically connected component of $\mathbb{R}\langle\varepsilon\rangle^t - \{\Delta_\varepsilon = 0\}$. \square

We deduce that there exists $b' \in \mathbb{N}$ such that for all $y \in U$, the number of semi-algebraically connected components of $S'_\varepsilon \cap \pi^{-1}(y)$ is b' . Using the transfer principle as in [9], we deduce that there exists $e' \in \mathbb{R}$ positive and small enough such that, the following holds. There exists $b \in \mathbb{N}$ such that for all $e \in]0, e'[$ the number of connected components of $S \cap \{a_1 x_1^2 + \dots + a_n x_n^2 \leq \frac{1}{e}\} \cap \pi^{-1}(y)$ is b when y ranges over U . This proves the following lemma.

Lemma 6.6. *Let U be as above. Then the number of connected components of S_y is invariant when y ranges over U .*

Finally, we can describe the subroutine `Eliminate` whose correctness follows from the previous lemma.

Algorithm 3: Eliminate($F, G, \mathbf{x}, \mathbf{y}$)

Data: Finite sequences F and G in $\mathbb{Q}[\mathbf{x}, \mathbf{y}]$ with $\mathbf{x} = (x_1, \dots, x_n)$ and $\mathbf{y} = (y_1, \dots, y_t)$, defining a semi-algebraic set $S \subset \mathbb{R}^n \times \mathbb{R}^t$.

Assumes that (F, G) satisfies assumption (A).

Result: $\Delta \in \mathbb{Q}[\mathbf{y}]$ such that the number of connected components of S_y is invariant when y ranges over a connected component of $\mathbb{R}^t - \{\Delta = 0\}$

- 1 Choose $a_1 > 0, \dots, a_n > 0$ in \mathbb{Q} randomly and let $g \leftarrow a_1 x_1^2 + \dots + a_n x_n^2 \leq \frac{1}{\varepsilon}$
 - 2 $\Delta \leftarrow \text{EliminateProper}(F, G \cup g, \mathbf{x}, \mathbf{y})$
 - 3 $\Delta \leftarrow \text{Normalize}(\Delta)$ return Δ_0 .
-

7 Computations

We have implemented several variants of the roadmap algorithms sketched in Section 5 as well as variants of the algorithm `ParametricSolve`. To perform algebraic elimination, we use Gröbner bases implemented in the FGB library by J.-C. Faugère [17]. The roadmap algorithm and the routines for computing sample points in semi-algebraic sets are implemented in the RAGLIB library [25].

We have not directly applied the most general version of `ParametricSolve` to the polynomial B . Indeed, since its variables v_2, v_3, v_4 lie in the Cartesian product $\mathbb{P}^1(\mathbb{R}) \times \mathbb{P}^1(\mathbb{R}) \times \mathbb{P}^1(\mathbb{R})$ (which is compact), the projection on the parameter's space is proper and it suffices to compute critical loci of that projection. There is one technical (but easy) difficulty to overcome: polynomial B actually admits a positive dimensional singular locus. But an easy computation shows that this singular locus has one purely complex component (which satisfies $v_4^2 + 1$) which can then be forgotten. The other component has a projection on the parameters' space which Zariski closed (it is contained in the set satisfied by $a_2 a_3 = 0$). This way, we directly obtain the following polynomial for Δ by computing the critical locus and consider additionally the set defined by $a_2 a_3 = 0$.

$$a_2 a_3 d_5 (a_2 + a_3 + d_5) (a_2 + a_3 - d_5)$$

Computing Δ as above does not take more than 3 sec. on a standard laptop using FGB. Getting sample points in the set defined by $\Delta \neq 0$ is trivial. We obtain the following 10 sample points using RAGLIB

$$\begin{aligned} & \{a_2 = -1, a_3 = -3, d_5 = 3\}, \{a_2 = -1, a_3 = -1, d_5 = 3\}, \{a_2 = -1, a_3 = 2, d_5 = 3\}, \{a_2 = \\ & -1, a_3 = 5, d_5 = 3\}, \{a_2 = -1, a_3 = \frac{1}{2}, d_5 = 3\}, \{a_2 = 1, a_3 = -120, d_5 = 118\}, \{a_2 = 1, a_3 = \\ & -118, d_5 = 118\}, \{a_2 = 1, a_3 = 1, d_5 = 118\}, \{a_2 = 1, a_3 = 118, d_5 = 118\}, \{a_2 = 1, a_3 = -1/2, d_5 = 118\} \end{aligned}$$

Our implementation allows us to compute a roadmap for one sample point within 20 minutes on a standard laptop. Analyzing the connectivity of these roadmaps is longer as it takes 40 min. All in all, approximately 10 hours are required to handle this positive dimensional parametric system. The data we computed are available at <http://ecarp.lip6.fr/papers/materials/issac20/>. These computations allow to retrieve the conclusions of our theoretical analysis of the UR family. They illustrate that prototype implementations of our algorithms are becoming efficient enough to tackle automated kinematic singularity analysis in robotics.

Acknowledgments. The three authors are supported by the joint ANR-FWF ANR-19-CE48-0015, FWF I 4452-N ECARP project. Mohab Safey El Din is supported by the ANR grants ANR-18-CE33-0011 SESAME and ANR-19-CE40-0018 DE RERUM NATURA, the PGMO grant CAMISADO and the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement N. 813211 (POEMA).

References

- [1] ANGELES, J. *Fundamentals of Robotic Mechanical Systems, Theory, Methods, and Algorithms*. Springer, 2007.
- [2] AUBRY, P., LAZARD, D., AND MAZA, M. M. On the theories of triangular sets. *Journal of Symbolic Computation* 28, 1-2 (1999), 105–124.
- [3] BANK, B., GIUSTI, M., HEINTZ, J., AND MBAKOP, G. Polar varieties and efficient real equation solving: the hypersurface case. *J. of Complexity* 13, 1 (1997), 5–27.
- [4] BANK, B., GIUSTI, M., HEINTZ, J., SAFEY EL DIN, M., AND SCHOST, E. On the geometry of polar varieties. *Applicable Algebra in Engineering, Communication and Computing* 21, 1 (2010), 33–83.
- [5] BASU, S., POLLACK, R., AND ROY, M.-F. On computing a set of points meeting every cell defined by a family of polynomials on a variety. *Journal of Complexity* 13, 1 (1997), 28 – 37.
- [6] BASU, S., POLLACK, R., AND ROY, M.-F. Computing roadmaps of semi-algebraic sets on a variety. *Journal of the American Mathematical Society* 13, 1 (2000), 55–82.
- [7] BASU, S., POLLACK, R., AND ROY, M.-F. *Algorithms in real algebraic geometry*. Springer-Verlag, 2003.
- [8] BASU, S., AND ROY, M.-F. Divide and conquer roadmap for algebraic sets. *Discrete & Computational Geometry* 52, 2 (2014), 278–343.
- [9] BASU, S., ROY, M.-F., SAFEY EL DIN, M., AND SCHOST, É. A baby step–giant step roadmap algorithm for general algebraic sets. *Foundations of Computational Mathematics* 14, 6 (2014), 1117–1172.
- [10] BOCHNAK, J., COSTE, M., AND ROY, M.-F. *Real Algebraic Geometry*. Springer-Verlag, 1998.
- [11] CANNY, J. *The complexity of robot motion planning*. MIT press, 1988.
- [12] CANNY, J. Computing roadmaps of general semi-algebraic sets. *The Computer Journal* 36, 5 (1993), 504–514.

- [13] COSTE, M., AND SHIOTA, M. Thom's first isotopy lemma: a semialgebraic version, with uniform bound. In *Real Analytic and Algebraic Geometry: Proc. of the International Conference, Trento (1995)*, Walter de Gruyter, p. 83.
- [14] EISENBUD, D. *Commutative Algebra: with a view toward algebraic geometry*, vol. 150. Springer Science & Business Media, 2013.
- [15] FAUGÈRE, J.-C. A new efficient algorithm for computing gröbner bases (f4). *Journal of pure and applied algebra* 139, 1-3 (1999), 61–88.
- [16] FAUGÈRE, J. C. A new efficient algorithm for computing gröbner bases without reduction to zero (f 5). In *Proc. of the 2002 international symposium on Symbolic and algebraic computation (2002)*, ACM, pp. 75–83.
- [17] FAUGÈRE, J.-C. Fgb: A library for computing gröbner bases. In *Mathematical Software - ICMS 2010 (Berlin, Heidelberg, September 2010)*, K. Fukuda, J. Hoeven, M. Joswig, and N. Takayama, Eds., vol. 6327 of *Lecture Notes in Computer Science*, Springer, pp. 84–87.
- [18] GIUSTI, M., HEINTZ, J., MORAIS, J.-E., MORGENSTERN, J., AND PARDO, L.-M. Straight-line programs in geometric elimination theory. *Journal of Pure and Applied Algebra* 124 (1998), 101–146.
- [19] GIUSTI, M., HEINTZ, J., MORAIS, J.-E., AND PARDO, L.-M. When polynomial equation systems can be solved fast? In *AAECC-11 (1995)*, vol. 948 of *LNCS*, Springer, pp. 205–231.
- [20] GIUSTI, M., LECERF, G., AND SALVY, B. A gröbner free alternative for polynomial system solving. *Journal of Complexity* 17, 1 (2001), 154 – 211.
- [21] GOURNAY, L., AND RISLER, J.-J. Construction of roadmaps in semi-algebraic sets. *Applicable Algebra in Engineering, Communication and Computing* 4, 4 (1993), 239–252.
- [22] MEZZAROBBA, M., AND SAFEY EL DIN, M. Computing roadmaps in smooth real algebraic sets. In *Proc. of Transgressive Computing (2006)*, J.-G. Dumas, Ed., pp. 327–338.
- [23] MURRAY, R., LI, Z., AND SASTRY, S. *A Mathematical Introduction to Robotic Manipulation*. CRC Press Taylor & Francis Group, 1994.
- [24] SAFEY EL DIN, M. Finding sampling points on real hypersurfaces is easier in singular situations. *MEGA (Effective Methods in Algebraic Geometry) Electronic proceedings (2005)*.
- [25] SAFEY EL DIN, M. Real algebraic geometry library. available at <http://www-polsys.lip6.fr/~safey>, 2007.

- [26] SAFEY EL DIN, M., AND SCHOST, E. Polar varieties and computation of one point in each connected component of a smooth real algebraic set. In *Proc. of the 2003 Int. Symp. on Symb. and Alg. Comp.* (NY, USA, 2003), ISSAC'03, ACM, pp. 224–231.
- [27] SAFEY EL DIN, M., AND SCHOST, E. A baby steps/giant steps probabilistic algorithm for computing roadmaps in smooth bounded real hypersurface. *Disc. Comput. Geom.* 45, 1 (2011), 181–220.
- [28] SAFEY EL DIN, M., AND SCHOST, É. A nearly optimal algorithm for deciding connectivity queries in smooth and bounded real algebraic sets. *Journal of the ACM (JACM)* 63, 6 (2017), 48.
- [29] SCHWARTZ, J. T., AND SHARIR, M. Algorithmic motion planning in robotics. In *Algorithms and Complexity*. Elsevier, 1990, pp. 391–430.
- [30] SELIG, J. *Geometric Fundamentals of Robotics*. Monographs in Computer Science. Springer, 2005.
- [31] SPONG, M., HUTCHINSON, S., AND VIDYASAGAR, M. *Robot Dynamics and Control*, 2nd ed. Monographs in Computer Science. John Wiley & Sons, 2005.
- [32] WANG, D. *Elimination methods*. Springer Science & Business Media, 2001.
- [33] WENGER, P. Cuspidal robots. In *Singular Configurations of Mechanisms and Manipulators*, Z. D. Müller A., Ed. Springer International Publishing, 2019, pp. 67–100.
- [34] WEYRER, M., BRANDSTÖTTER, M., AND HUSTY, M. Singularity avoidance control of a non-holonomic mobile manipulator for intuitive hand guidance. *Robotics* 8, 1 (2019).