



**HAL**  
open science

## Low rates of immediate coronary angiography among young adults resuscitated from sudden cardiac arrest

Victor Waldmann, Nicole Karam, Julien Rischard, Wulfran Bougouin, Ardalan Sharifzadehgan, Florence Dumas, Kumar Narayanan, Georgios Sideris, Sebastian Voicu, Estelle Gandjbakhch, et al.

### ► To cite this version:

Victor Waldmann, Nicole Karam, Julien Rischard, Wulfran Bougouin, Ardalan Sharifzadehgan, et al.. Low rates of immediate coronary angiography among young adults resuscitated from sudden cardiac arrest. *Resuscitation*, 2020, 147, pp.34-42. 10.1016/j.resuscitation.2019.12.005 . hal-02948006

HAL Id: hal-02948006

<https://hal.sorbonne-universite.fr/hal-02948006>

Submitted on 21 Jul 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

# 1 **Low Rates of Immediate Coronary Angiography among Young Adults** 2 **Resuscitated From Sudden Cardiac Arrest**

3  
4 Victor Waldmann<sup>1,2,3</sup>, MD, MPH; Nicole Karam<sup>1,2,3</sup>, MD, PhD; Julien Rischard<sup>1,2</sup>, MD;  
5 Wulfran Bougouin<sup>2,3,4</sup>, MD, PhD; Ardalan Sharifzadehgan<sup>1,2,3</sup>, MD, MPH; Florence  
6 Dumas<sup>2,3,5</sup>, MD, PhD; Kumar Narayanan<sup>2,6</sup>, MD; Georgios Sideris<sup>7</sup>, MD, PhD; Sebastian  
7 Voicu<sup>8</sup>, MD; Estelle Gandjbakhch<sup>9,10</sup>, MD, PhD; Daniel Jost<sup>11</sup>, MD; Lionel Lamhaut<sup>12</sup>, MD,  
8 PhD; Bertrand Ludes<sup>13</sup>, MD; Isabelle Plu<sup>14</sup>, MD; Frankie Beganton<sup>2</sup>, MS; Karim Wahbi<sup>3,15</sup>,  
9 MD; Olivier Varenne<sup>3,15</sup>, MD, PhD; Bruno Megarbane<sup>8</sup>, MD, PhD; Vincent Algalarrondo<sup>10,16</sup>,  
10 MD, PhD; Fabrice Extramiana<sup>10,16</sup>, MD, PhD; Nicolas Lellouche<sup>10,17</sup>, MD; David S.  
11 Celermajer<sup>18</sup>, MBBS, FRACP; Christian Spaulding<sup>1,3</sup>, MD, PhD; Antoine Lafont<sup>1,3</sup>, MD,  
12 PhD; Alain Cariou<sup>2,3,19</sup>, MD, PhD; Xavier Jouven<sup>1,2,3\*</sup>, MD, PhD; Eloi Marijon<sup>1,2,3,10\*</sup>, MD,  
13 PhD – On Behalf Paris-SDEC investigators

14  
15 <sup>1</sup> AP-HP, European Georges Pompidou Hospital, Cardiology Department, Paris, France  
16 <sup>2</sup> Sudden Death Expertise Center, INSERM U970, Paris Cardiovascular Research Center (PARCC),  
17 European Georges Pompidou Hospital, Paris, France  
18 <sup>3</sup> Paris University, Paris, France  
19 <sup>4</sup> Ramsay Générale de Santé, Hôpital privé Jacques Cartier, Intensive Care Unit, Massy, France  
20 <sup>5</sup> AP-HP, Cochin-Hotel Hospital, Emergency Department, Paris, France  
21 <sup>6</sup> Maxcure Hospitals, Hyderabad, India  
22 <sup>7</sup> AP-HP, Lariboisière Hospital, Cardiology Department, Paris, France  
23 <sup>8</sup> AP-HP, Lariboisière Hospital, Intensive Care Unit, Paris, France  
24 <sup>9</sup> AP-HP, La Pitié Salpêtrière University Hospital, Cardiology Department, Paris, France  
25 <sup>10</sup> Groupe Parisien Universitaire de Rythmologie (G.P.U.R.)  
26 <sup>11</sup> Paris Firefighters Brigade, Paris, France  
27 <sup>12</sup> AP-HP, SAMU de Paris, Necker Hospital, Paris, France  
28 <sup>13</sup> Forensic Medical Institute, Paris, France  
29 <sup>14</sup> AP-HP, La Pitié Salpêtrière University Hospital, Anatomopathology Department, Paris, France  
30 <sup>15</sup> AP-HP, Cochin Hospital, Cardiology Department, Paris, France  
31 <sup>16</sup> AP-HP, Bichat-Claude-Bernard Hospital, Cardiology Department, Paris, France  
32 <sup>17</sup> AP-HP, Henri Mondor Hospital, Cardiology Department, Créteil, France  
33 <sup>18</sup> Faculty of Medicine and Health, Sydney, Australia  
34 <sup>19</sup> AP-HP, Cochin Hospital, Intensive Care Unit, Paris, France

35  
36 \*Equally contributed to the manuscript

## 37 **Corresponding author**

38 Eloi Marijon MD, PhD  
39 Hôpital européen Georges Pompidou  
40 Département de Cardiologie  
41 20-40 Rue Leblanc 75908 Paris Cedex 15, France  
42 Phone: +33662833848  
43 Fax : +33156093047  
44 Email: [eloi.marijon@aphp.fr](mailto:eloi.marijon@aphp.fr)

## 45 **Original Paper – Resuscitation**

46  
47 **Words Count: 2853**  
48  
49  
50

1 **Abstract (250 w)**

2 **Aim**

3 Coronary artery disease (CAD) has recently been emphasized as a major cause of sudden  
4 cardiac arrest (SCA) in young adults. We aim to assess the rate of immediate coronary  
5 angiography performance in young patients resuscitated from SCA.

6 **Methods**

7 From May 2011 to May 2017, all cases of out-of-hospital SCA aged 18 to 40 years alive at  
8 hospital admission were prospectively included in 48 hospitals of the Great Paris area.  
9 Cardiovascular causes of SCA were centrally adjudicated, and management including  
10 immediate coronary angiography performance was assessed.

11 **Results**

12 Out of 3,579 SCA admitted alive, 409 (11.4%) patients were under 40 years of age (32.3±6.2  
13 years, 69.7% males), with 244 patients having a definite cause identified. Among those, CAD  
14 accounted for 72 (29.5%) cases, of which 64 (88.9%) were acute coronary syndromes. The  
15 rate of immediate coronary angiography was only 41.7% compared to 65.1% among those ≥  
16 40-years (P<0.001). During the study period, while the rate of immediate coronary  
17 angiography increased from 60.5% to 70.3% (P <0.001) in patients aged ≥ 40 years, the rate  
18 in patients aged less than 40 years remained stable (43.5% to 45.3%, P =0.795). Patients  
19 younger than 40 years were significantly less likely to undergo immediate coronary  
20 angiography (OR=0.34, 95% CI: 0.25-0.47), although early angiography was associated with  
21 survival at hospital discharge (OR=2.68, 95% CI: 1.21-6.00).

22 **Conclusion**

23 CAD is the first cause of SCA in young adults aged less than 40 years. The observed low  
24 rates of immediate coronary angiography suggest a missed opportunity for early intervention.

1 **Key words** – Sudden cardiac death; cardiac arrest; coronary artery disease; acute coronary  
2 syndrome; epidemiology; percutaneous coronary intervention

3

4

5

6

7

## 1 **Introduction**

2 Sudden Cardiac Arrest (SCA) is defined as a natural, unexpected and sudden pulseless event,  
3 without an obvious non-cardiac cause [1]. It is a significant public health concern, accounting  
4 for approximately 50% of cardiovascular deaths, with 300,000 cases estimated annually in  
5 Europe [2].

6 Coronary Artery Disease (CAD) accounts for around 80% of SCA cases [3]. Along  
7 with bystander interventions and early defibrillation, immediate percutaneous coronary  
8 intervention (PCI) is associated with improved outcomes after resuscitation from SCA  
9 [4][5][6][7][8]. Previous reports have shown an increase in the rate of coronary angiography  
10 performance in patients hospitalized alive after SCA [9][10]. In young patients, non-ischemic  
11 structural heart disease (especially cardiomyopathies) and electrical cardiac disorders are  
12 alternative diagnoses that were traditionally considered more prevalent [11][12]. However, the  
13 epidemiology of CAD has substantially changed over the last two decades, with significant  
14 changes in cardiovascular risk factors in early adulthood [13][14][15][16]. The proportion of  
15 young adults among STEMI victims has considerably increased over the last years [17].  
16 Recent studies have reported that CAD was a major cause of SCA in young adults, in  
17 postmortem examination [18] and in SCA survivors [19]. This has potentially important  
18 implications for emergency decision-making in young subjects, especially with regard to early  
19 coronary angiography and PCI.

20 We undertook a population-based study of SCA patients admitted alive to hospital and  
21 hypothesized that young adults would be less likely to be referred for immediate coronary  
22 angiography at hospital admission due to perceptions of low CAD risk.

23

## 24 **Methods**

### 25 *Study setting*

1 The Paris-Sudden Death Expertise Center (SDEC) registry is an ongoing study which has  
2 been described previously [20][21][22][23][24]. Briefly, it is a comprehensive, prospective  
3 population-based registry comprising Paris and suburbs, encompassing a residential  
4 population of 6.7 million (approximately 10% of the total French population) and covering  
5 762 km<sup>2</sup>. In Paris, management of out-of-hospital cardiac arrest involves mobile emergency  
6 units (Service d'Aide Médicale Urgente), and firefighters (Brigade de Sapeurs Pompiers de  
7 Paris), with at least one trained emergency physician on board. Patients in whom a return of  
8 spontaneous circulation is achieved are then referred to a center with an intensive care unit  
9 (ICU) and coronary intervention facilities. Owing to a close collaboration with all the pre-  
10 hospital Emergency Medical Services (EMS), 48 hospitals, and forensic units, every case of  
11 out-of-hospital cardiac arrest aged  $\geq 18$  years occurring in the area is systematically enrolled  
12 in the SDEC registry, since May 2011. Exclusion criteria include age less than 18 years and  
13 cardiac arrest occurring outside the geographical area of interest. Regular external audits on  
14 the registry have shown that 99 % of cardiac arrest cases admitted alive to the hospital were  
15 detected [20].

16 The study was conducted in compliance with Good Clinical Practice, French Law and  
17 the French data protection law in accordance with the ethical standards laid down in the 1964  
18 Declaration of Helsinki and its later amendments. Data file of the Paris-SDEC registry was  
19 declared to and authorized by the French data protection committee (Commission Nationale  
20 Informatique et Liberté, CNIL, DR-2012-445 authorization n°912309).

### 21 ***Study population***

22 All SCA admitted alive to hospital were included from 15 May 2011 to 15 May 2017.

23 According to definitions from the last consensus, SCA was defined as an unexpected out-of-  
24 hospital cardiac arrest without obvious non-cardiac cause, occurring with a rapid witnessed  
25 collapse within 1 hour after the onset of symptoms, or if unwitnessed, within the 24 hours

1 after the last contact [1]. Those likely due to non-cardiac circumstances (such as trauma,  
2 drowning, hanging etc.) or prior terminal condition were not included. The present study  
3 specifically focused on young adults aged less than 40 years old.

#### 4 ***Data collection***

5 General data included demographic characteristics and location of SCA (residential or public  
6 place). Utstein templates for resuscitation information reporting were followed [25]. Pre-  
7 hospital data recorded included SCA circumstances, presence of a witness, witness-  
8 cardiopulmonary resuscitation before EMS arrival, presence of shockable rhythm before  
9 advanced life support, epinephrine dose injected by EMS, ST segment elevation on first  
10 electrocardiogram (ECG) recorded, and delays from collapse to basic life support and from  
11 call for EMS to arrival of EMS. In addition to the data from EMS and the medical examiner, a  
12 working group of the SDEC collected and assessed the lifetime past medical history. The  
13 information was gathered by the local medical staff, but two investigators thoroughly  
14 reviewed each medical report for data completion and validity, and provided final central  
15 adjudication (assigned diagnosis). In cases of divergent opinion, a third expert was asked to  
16 arbitrate. In survivors without definite diagnosis after the initial work-up performed in ICU,  
17 etiological medical investigations were carried out in cardiology, including cardiac magnetic  
18 resonance imaging, pharmacological tests, vasospasm provocative test, Holter-ECG  
19 recording, exercise stress test as well as genetic screening, all of which were actively  
20 encouraged by the coordinators of the study. The diagnosis of idiopathic ventricular  
21 fibrillation was made among survivors when eventually no phenotype was identified. Survival  
22 at hospital discharge and neurological status were also recorded. Favorable neurological  
23 prognosis was defined by a cerebral performance category (CPC) score 1 or 2, with 1  
24 representing full recovery or mild disability and 2, moderate disability but independent in  
25 activities of daily living.

## 1 ***Definitions***

2 Immediate coronary angiography was defined as that performed within the first two hours  
3 after ICU admission [26][27]. Significant CAD was defined by the presence of a stenosis  
4 producing > 50% narrowing in at least one coronary artery. Flow in coronary arteries was  
5 assessed using the thrombolysis in myocardial infarction (TIMI) classification. A coronary  
6 occlusion was defined as TIMI grade 0 to 1 flow. The occlusion was considered recent if  
7 collaterals were absent and if the occlusion was easily crossable by the wire during  
8 subsequent angioplasty. A culprit lesion was defined by the presence of a thrombus in the  
9 artery, a flow reduction < TIMI 3, and/or an acute coronary occlusion, requiring a  
10 revascularization by PCI (aspiration thrombectomy and/or angioplasty). Chronic ischemic  
11 heart disease-related SCA was defined as significant CAD diagnosed during coronary  
12 angiography in the absence of culprit lesion criteria or other non-cardiac SCA cause.

## 13 ***Statistical analysis***

14 Continuous data were reported as mean  $\pm$  standard deviation (SD) or median and interquartile  
15 range (IQR) for normally and non-normally distributed data respectively. Categorical data  
16 were reported as numbers and percentages. Comparisons used the  $\chi^2$  or Fisher's exact test for  
17 categorical variables and Student's t test or Mann–Whitney–Wilcoxon test, when appropriate,  
18 for continuous variables. Multiple logistic regressions were used (i) to compare the rate of  
19 immediate coronary angiography at hospital admission according to patient's age, and (ii) to  
20 assess the association between immediate coronary angiography performance and survival  
21 rate at hospital discharge. The main known pre-hospital outcome predictors were included as  
22 explicative variables in the former model (age, gender, location of SCA, witness presence and  
23 witness-cardiopulmonary resuscitation, delays from collapse to basic life support and from  
24 call for EMS to arrival of EMS, presence of a shockable rhythm, epinephrine use), as well as  
25 ST segment elevation on first recorded ECG. Main comorbidities were also included in the



1 second model. Variables significantly associated ( $P < 0.20$ ) in univariate analysis were  
2 assessed in multivariate logistic regression. Linear time trends analysis on the rate of  
3 immediate coronary angiography were tested with the use of logistic regression dividing this  
4 6-year study in 3 periods of 24 months. Missing data on study variables were no more than  
5 10%, except for ST segment elevation on first ECG (12.0%) and interval from collapse to  
6 basic life support (12.3%), and were handled using case-complete analysis. Results were  
7 considered statistically significant at  $P < 0.05$ . All analyses were two-tailed. Statistical  
8 analysis was performed using R software, version 3.3.2 (R Project for Statistical Computing).

9

## 10 **Results**

### 11 *Population*

12 From May 15<sup>th</sup>, 2011 to May 15<sup>th</sup>, 2017, among 3,579 comatose patients admitted alive to 48  
13 different hospitals after SCA, 409 (11.4%) were aged less than 40 years. Clinical and  
14 demographic characteristics of the young SCA group are summarized in Table 1. Mean age  
15 was  $32.3 \pm 6.2$  years with 285 (69.7%) males. Compared to patients older than 40, younger  
16 patients had fewer cardiovascular risk factors, comorbidities, or previously diagnosed heart  
17 disease (15.8% vs. 35.3%,  $P < 0.001$ ), especially CAD (1.1% vs. 20.1%,  $P < 0.001$ ). ST-segment  
18 elevation was less frequently recorded in the initial ECG (23.5% vs. 34.2%,  $P < 0.001$ ), family  
19 history of SCA was more frequent (8.0% vs. 1.8%,  $P < 0.001$ ), and SCAs were more often  
20 sports-related (12.2% vs 4.1%,  $P < 0.001$ ). The survival rate at hospital discharge was  
21 comparable between the two groups (31.6% vs 27.4%,  $P = 0.103$ ).

### 22 *Causes of SCA among the Young*

23 Among young SCA patients, the cause was uncertain in 165 (40.3%) cases due to early death  
24 and negative or incomplete initial work-up in ICU. Among 244 young adults with a definite  
25 etiology identified, a non-cardiac cause was diagnosed in 72 (29.5%) patients, and a cardiac

1 cause was identified in 172 (70.5%) patients (Figure 1). CAD was the most frequent etiology,  
2 diagnosed in 72 (29.5%) subjects, of which 64 (88.9%) presented acute coronary syndromes.  
3 Non-ischemic structural heart disease and non-structural heart disease were identified in 62  
4 (25.4%) and 38 (15.6%) patients, respectively. In the 30 to 40 year age group, CAD  
5 represented 40.9% of etiologies identified, and non-ischemic structural heart diseases were  
6 the main cause of SCA under 30 years (38.8%) (Figure 2).

### 7 ***Characteristics of CAD-Related SCA***

8 Among CAD-related SCA cases, acute coronary syndromes were more frequent in younger  
9 patients (100.0% before 30 years, 87.7% between 30 and 40 years, and 76.3% after the age of  
10 40-years, overall  $P=0.033$ ), compared to chronic ischemic heart disease (without an acute  
11 culprit lesion) (Figure 2). Young CAD patients had more cardiovascular risk factors ( $\geq 1$ ) than  
12 young non-CAD patients (84.3% vs. 50.5%,  $P<0.001$ ), in particular current smoking (65.7%  
13 vs. 28.3%,  $P<0.001$ ), dyslipidemia (14.5% vs. 1.0%,  $P=0.001$ ) and family history of CAD  
14 (20.3% vs. 7.1%,  $P=0.03$ ). These patients also had a different distribution of risk factors  
15 compared to older CAD cases, with a higher prevalence of active smoking (65.7% vs. 43.5%,  
16  $P<0.001$ ) and family history of CAD (20.3% vs. 9.3%,  $P=0.006$ ), and a lower prevalence of  
17 other factors. Lastly, subjects with CAD-related SCA under 40 years less often had previously  
18 known heart disease (11.6%) compared to young non-CAD patients (37.0%) or older CAD-  
19 related cases (36.4%) (both  $P<0.001$ ).

### 20 ***Early Coronary Intervention***

21 Among all 409 patients younger than 40, 170 (41.7%) had immediate coronary angiography  
22 (vs. 65.1% in patients aged  $\geq 40$  years,  $P<0.001$ ), and 50 (29.4%) of these patients underwent  
23 immediate PCI at admission (vs. 48.0% in patients aged  $\geq 40$  years,  $P<0.001$ ). Additionally,  
24 among patients with a final diagnosis of acute coronary syndrome related to a culprit lesion,  
25 7.8% did not undergo immediate coronary angiography (vs. 1.6% in older patients,  $P=0.007$ ),

1 and none of these patients presented with typical ST segment elevation on ECG. Compared to  
2 young patients not referred, those who underwent immediate coronary angiography were  
3 older ( $33.3\pm 5.8$  vs.  $31.5\pm 6.3$  years,  $P=0.003$ ), more frequently males ( $75.3\%$  vs.  $66.0\%$ ,  
4  $P=0.049$ ), a greater proportion had ST segment elevation ( $44.0\%$  vs.  $4.1\%$ ,  $P < 0.001$ ), and  
5 presented with better prognostic indicators (Table 2). During the study period, while the rate  
6 of immediate coronary angiography increased from  $60.5\%$  to  $70.3\%$  ( $P$  for trend  $< 0.001$ ) in  
7 patients aged  $\geq 40$  years, the rate in patients aged less than 40 years remained stable ( $43.5\%$  to  
8  $45.3\%$ ,  $P$  for trend  $= 0.795$ ) (Figure 3). On multivariable analysis, patients younger than 40  
9 years were significantly less likely to undergo immediate coronary angiography at hospital  
10 admission (OR=0.34, 95% CI: 0.25-0.47,  $P < 0.001$ ) (Table 3), although immediate coronary  
11 angiography was found to be significantly associated with survival at hospital discharge in  
12 these young patients (OR=2.68, 95% CI: 1.21-6.00,  $P=0.015$ ).

13

## 14 **Discussion**

15 In this contemporary large population-based study, young SCA patients were significantly  
16 less likely to undergo immediate coronary angiography at hospital admission. However, CAD  
17 –especially acute coronary syndrome– was the main identified cause of SCA. Overall, our  
18 findings underline the lack of timely and systematic investigation of SCA in young adults  
19 admitted alive at hospital, suggesting an important missed opportunity for early intervention.

20 In the field of SCA, a threshold of 35 or 40 years is traditionally used to distinguish  
21 younger patients, in whom CAD is considered as less prevalent and in whom greater  
22 emphasis is usually laid on alternative diagnoses, in particular, inherited cardiomyopathies  
23 and electrical disorders (channelopathies)[11][12]. However, CAD epidemiology has changed  
24 over the last two decades, with an increase in CAD prevalence in the young population  
25 [13][14][15][16]. It has been shown recently that the rate of acute myocardial infarction is

1 increasing in young patients [17], which might suggest a potential increase in the rate of  
2 CAD-related SCA among the youth.

3         Few systematic evaluations of SCA in the young have been conducted in the general  
4 population; most studies have only focused on particular subgroups, such as young  
5 competitive athletes [28]. However, underlying mechanisms and causes associated with SCA  
6 occurring during sports may be unique and not generalizable to all young SCA. While  
7 hypertrophic cardiomyopathy, myocarditis, arrhythmogenic right ventricular cardiomyopathy,  
8 or channelopathies have been considered for a long time to be the main etiologies underlying  
9 SCA during competitive sports, CAD has been shown to be an important cause when  
10 considering sports-related SCA in the general population (including recreational sports in the  
11 young) [29]. Outside the particular sports setting, data regarding CAD in young SCA victims  
12 is scarce. Our findings, however, reveal that CAD remains the single most common cause of  
13 SCA identified before the age of 40 years in the general population. These data from a  
14 population of SCA patients admitted alive to the hospital are consistent with the findings of a  
15 recent autopsy-based study of SCA [18].

16         Younger patients present more frequently with ST-segment elevation acute coronary  
17 syndrome (STEMI) [30], where ventricular fibrillation is more frequent than in non-ST  
18 elevation acute coronary events [31]. Although coronary plaque rupture determinants remain  
19 poorly elucidated, it may be hypothesized that shear forces induced by a greater degree of  
20 physical activity may provoke plaque fissuring and that the higher proportion of active  
21 smoking among younger patients could also predispose to plaque erosion and thrombosis,  
22 triggering ventricular arrhythmias [32]. Moreover, a higher risk of SCA during STEMI has  
23 been reported at a younger age, probably due to abrupt coronary artery occlusion in the  
24 absence of a developed collateral circulation as compared to older patients with chronic CAD  
25 [33].

1           Immediate coronary revascularization is associated with survival after resuscitated  
2 SCA [4][5][6][7][8]. In our study, patients younger than 40 years were less frequently  
3 subjected to an early invasive strategy. Moreover, a higher proportion of SCA actually due to  
4 an acute coronary syndrome (with an identified culprit lesion later) did not undergo  
5 immediate coronary angiography, when compared to older patients. While a recent study (in  
6 older populations) did not support immediate angiography in patients without ST-segment  
7 elevation [27], none of these patients had significant ECG changes, suggesting as already  
8 reported that the predictive value of the first recorded ECG in survivors of SCA is poor.  
9 Coronary lesions requiring PCI have been reported in nearly one-third of cases without initial  
10 ST segment elevation [34]. Although overall CAD-related SCA were more frequent in the  
11 elderly, the proportion of acute coronary syndromes was much higher among younger  
12 patients, compared to chronic CAD-related SCA. Most importantly, immediate coronary  
13 angiography was independently associated with improved survival at hospital discharge.  
14 These data suggest that CAD may be underappreciated as a cause of SCA in younger patients  
15 and support the consideration of prompt coronary angiography in young SCA patients as well,  
16 in the absence of another obvious cause.

17           The present study has several limitations, which need to be acknowledged. First, the  
18 number of patients < 40 years with a certain etiology is relatively modest. In most parts of the  
19 world, low autopsy rates in this setting constitute a considerable bottleneck in ascertaining  
20 definitive cause of SCA, and explain the relatively high proportion of cases without  
21 established diagnosis. Given this scenario, a systematic description of survivors represents the  
22 next best approach. SCA patients with a cause identified may not have the same distribution  
23 of causes compared with patients with undetermined cause or patients who do not survive the  
24 initial resuscitation attempt; however our data are consistent with a recent autopsy study of  
25 non-survivors [18], and the proportion of CAD among older patients in our population is

1 congruent with the existing literature. Second, although our data are based on a large  
2 population with numerous centers involved, regional disparities may potentially exist and  
3 caution has to be exercised in generalizing these results. In particular, cardiovascular risk  
4 factor distribution may vary significantly between countries, for instance, smoking which is  
5 particularly prevalent in the young population in France [35]. Third, ascertaining causal  
6 relationship between SCA and CAD is sometimes difficult and debatable, particularly with  
7 chronic CAD-related SCA, in the absence of acute coronary occlusion. However, the majority  
8 of CAD-related SCA in young adults were due to acute coronary syndromes with  
9 identification of a culprit lesion. Lastly, the association identified between coronary  
10 angiography performance and survival at hospital discharge in young patients in our  
11 observational study cannot be interpreted as a causal relationship despite the multivariable  
12 analysis, due to possible confounding non-measured variables.

13

## 14 **Conclusions**

15 This population-based study demonstrates a low rate of immediate coronary  
16 angiography in young adults aged less than 40 years resuscitated from SCA, suggesting that  
17 CAD as a cause of SCA in this population may be underappreciated by the medical  
18 community. The high proportion of CAD in these patients, in particular acute coronary  
19 syndromes, underlines important missed opportunity for early intervention.

20

21

22

23

24

25

26

1

2 **Conflict of interest**

3 None.

4

5 **Acknowledgments**

6 SDEC Executive Committee is part of the ESCAPE-NET project, a €10 million funding from  
7 the European Commission (Horizon2020 program) to develop research on sudden cardiac  
8 death in Europe. Paris-SDEC 2018 Investigators Listing (Data Supplement).

9

10 **Funding Source**

11 The Paris-SDEC activities are supported by the Institut National de la Santé et de la  
12 Recherche Médicale (INSERM), Paris University, Fondation Coeur et Artères, Global Heart  
13 Watch, Fédération Française de Cardiologie, Société Française de Cardiologie, Fondation  
14 Recherche Médicale, as well as unrestricted grants from industrial partners (Medtronic, St  
15 Jude Medical, Boston Scientific, Microport and Biotronik).

16

## 1 References

- 2 [1] Fishman GI, Chugh SS, Dimarco JP, Albert CM, Anderson ME, Bonow RO, et al.  
3 Sudden cardiac death prediction and prevention: report from a National Heart, Lung, and  
4 Blood Institute and Heart Rhythm Society Workshop. *Circulation* 2010;122:2335–48.  
5 <https://doi.org/10.1161/CIRCULATIONAHA.110.976092>.
- 6 [2] Atwood C, Eisenberg MS, Herlitz J, Rea TD. Incidence of EMS-treated out-of-hospital  
7 cardiac arrest in Europe. *Resuscitation* 2005;67:75–80.  
8 <https://doi.org/10.1016/j.resuscitation.2005.03.021>.
- 9 [3] Zipes DP, Wellens HJ. Sudden cardiac death. *Circulation* 1998;98:2334–51.
- 10 [4] Spaulding CM, Joly LM, Rosenberg A, Monchi M, Weber SN, Dhainaut JF, et al.  
11 Immediate coronary angiography in survivors of out-of-hospital cardiac arrest. *N Engl J*  
12 *Med* 1997;336:1629–33. <https://doi.org/10.1056/NEJM199706053362302>.
- 13 [5] Dumas F, Cariou A, Manzo-Silberman S, Grimaldi D, Vivien B, Rosencher J, et al.  
14 Immediate percutaneous coronary intervention is associated with better survival after  
15 out-of-hospital cardiac arrest: insights from the PROCAT (Parisian Region Out of  
16 hospital Cardiac Arrest) registry. *Circ Cardiovasc Interv* 2010;3:200–7.  
17 <https://doi.org/10.1161/CIRCINTERVENTIONS.109.913665>.
- 18 [6] Wissenberg M, Lippert FK, Folke F, Weeke P, Hansen CM, Christensen EF, et al.  
19 Association of national initiatives to improve cardiac arrest management with rates of  
20 bystander intervention and patient survival after out-of-hospital cardiac arrest. *JAMA*  
21 2013;310:1377–84. <https://doi.org/10.1001/jama.2013.278483>.
- 22 [7] Jaeger D, Dumas F, Escutnaire J, Sadoune S, Lauvray A, Elkhoury C, et al. Benefit of  
23 immediate coronary angiography after out-of-hospital cardiac arrest in France: A  
24 nationwide propensity score analysis from the RéAC Registry. *Resuscitation*  
25 2018;126:90–7. <https://doi.org/10.1016/j.resuscitation.2018.03.003>.
- 26 [8] Aissaoui N, Bougouin W, Dumas F, Beganton F, Chocron R, Varenne O, et al. Age and  
27 benefit of early coronary angiography after out-of-hospital cardiac arrest in patients  
28 presenting with shockable rhythm: Insights from the Sudden Death Expertise Center  
29 registry. *Resuscitation* 2018;128:126–31.  
30 <https://doi.org/10.1016/j.resuscitation.2018.05.006>.
- 31 [9] Patterson T, Perkins GD, Hassan Y, Moschonas K, Gray H, Curzen N, et al. Temporal  
32 Trends in Identification, Management, and Clinical Outcomes After Out-of-Hospital  
33 Cardiac Arrest: Insights From the Myocardial Ischaemia National Audit Project  
34 Database. *Circ Cardiovasc Interv* 2018;11:e005346.  
35 <https://doi.org/10.1161/CIRCINTERVENTIONS.117.005346>.
- 36 [10] Patel N, Patel NJ, Macon CJ, Thakkar B, Desai M, Rengifo-Moreno P, et al. Trends and  
37 Outcomes of Coronary Angiography and Percutaneous Coronary Intervention After Out-  
38 of-Hospital Cardiac Arrest Associated With Ventricular Fibrillation or Pulseless  
39 Ventricular Tachycardia. *JAMA Cardiol* 2016;1:890–9.  
40 <https://doi.org/10.1001/jamacardio.2016.2860>.
- 41 [11] Liberthson RR. Sudden Death from Cardiac Causes in Children and Young Adults. *N*  
42 *Engl J Med* 1996;334:1039–44. <https://doi.org/10.1056/NEJM199604183341607>.
- 43 [12] Eckart RE, Shry EA, Burke AP, McNear JA, Appel DA, Castillo-Rojas LM, et al.  
44 Sudden death in young adults: an autopsy-based series of a population undergoing active  
45 surveillance. *J Am Coll Cardiol* 2011;58:1254–61.  
46 <https://doi.org/10.1016/j.jacc.2011.01.049>.
- 47 [13] Briffa T, Nedkoff L, Peeters A, Tonkin A, Hung J, Ridout SC, et al. Discordant age and  
48 sex-specific trends in the incidence of a first coronary heart disease event in Western



- 1 Australia from 1996 to 2007. *Heart* 2011;97:400–4.  
2 <https://doi.org/10.1136/hrt.2010.210138>.
- 3 [14] Wilmot KA, O’Flaherty M, Capewell S, Ford ES, Vaccarino V. Coronary Heart Disease  
4 Mortality Declines in the United States From 1979 Through 2011: Evidence for  
5 Stagnation in Young Adults, Especially Women. *Circulation* 2015;132:997–1002.  
6 <https://doi.org/10.1161/CIRCULATIONAHA.115.015293>.
- 7 [15] Puymirat E, Simon T, Cayla G, Cottin Y, Elbaz M, Coste P, et al. Acute Myocardial  
8 Infarction: Changes in Patient Characteristics, Management, and 6-Month Outcomes  
9 Over a Period of 20 Years in the FAST-MI Program (French Registry of Acute ST-  
10 Elevation or Non-ST-elevation Myocardial Infarction) 1995 to 2015. *Circulation* 2017.  
11 <https://doi.org/10.1161/CIRCULATIONAHA.117.030798>.
- 12 [16] Mentias A, Hill E, Barakat AF, Raza MQ, Youssef D, Banerjee K, et al. An alarming  
13 trend: Change in the risk profile of patients with ST elevation myocardial infarction over  
14 the last two decades. *Int J Cardiol* 2017;248:69–72.  
15 <https://doi.org/10.1016/j.ijcard.2017.05.011>.
- 16 [17] Arora Sameer, Stouffer George A. (Rick), Kucharska-Newton Anna, Qamar Arman,  
17 Vaduganathan Muthiah, Pandey Ambarish, et al. Twenty Year Trends and Sex  
18 Differences in Young Adults Hospitalized with Acute Myocardial Infarction: The ARIC  
19 Community Surveillance Study. *Circulation* n.d.;0.  
20 <https://doi.org/10.1161/CIRCULATIONAHA.118.037137>.
- 21 [18] Bagnall RD, Weintraub RG, Ingles J, Duflou J, Yeates L, Lam L, et al. A Prospective  
22 Study of Sudden Cardiac Death among Children and Young Adults. *N Engl J Med*  
23 2016;374:2441–52. <https://doi.org/10.1056/NEJMoa1510687>.
- 24 [19] Waldmann V, Karam N, Bougouin W, Sharifzadehgan A, Dumas F, Narayanan K, et al.  
25 Burden of Coronary Artery Disease as a Cause of Sudden Cardiac Arrest in the Young. *J*  
26 *Am Coll Cardiol* 2019;73:2118–20. <https://doi.org/10.1016/j.jacc.2019.01.064>.
- 27 [20] Bougouin W, Lamhaut L, Marijon E, Jost D, Dumas F, Deye N, et al. Characteristics  
28 and prognosis of sudden cardiac death in Greater Paris: population-based approach from  
29 the Paris Sudden Death Expertise Center (Paris-SDEC). *Intensive Care Med*  
30 2014;40:846–54. <https://doi.org/10.1007/s00134-014-3252-5>.
- 31 [21] Jabre P, Bougouin W, Dumas F, Carli P, Antoine C, Jacob L, et al. Early Identification  
32 of Patients With Out-of-Hospital Cardiac Arrest With No Chance of Survival and  
33 Consideration for Organ Donation. *Ann Intern Med* 2016;165:770–8.  
34 <https://doi.org/10.7326/M16-0402>.
- 35 [22] Karam N, Marijon E, Dumas F, Offredo L, Beganton F, Bougouin W, et al.  
36 Characteristics and outcomes of out-of-hospital sudden cardiac arrest according to the  
37 time of occurrence. *Resuscitation* 2017;116:16–21.  
38 <https://doi.org/10.1016/j.resuscitation.2017.04.024>.
- 39 [23] Waldmann V, Bougouin W, Karam N, Dumas F, Sharifzadehgan A, Gandjbakhch E, et  
40 al. Characteristics and clinical assessment of unexplained sudden cardiac arrest in the  
41 real-world setting: focus on idiopathic ventricular fibrillation. *Eur Heart J*  
42 2018;39:1981–7. <https://doi.org/10.1093/eurheartj/ehy098>.
- 43 [24] Pechmajou L, Sharifzadehgan A, Bougouin W, Dumas F, Beganton F, Jost D, et al. Does  
44 occurrence during sports affect sudden cardiac arrest survival? *Resuscitation*  
45 2019;141:121–7. <https://doi.org/10.1016/j.resuscitation.2019.06.277>.
- 46 [25] Perkins GD, Jacobs IG, Nadkarni VM, Berg RA, Bhanji F, Biarent D, et al. Cardiac  
47 Arrest and Cardiopulmonary Resuscitation Outcome Reports: Update of the Utstein  
48 Resuscitation Registry Templates for Out-of-Hospital Cardiac Arrest: A Statement for  
49 Healthcare Professionals From a Task Force of the International Liaison Committee on  
50 Resuscitation (American Heart Association, European Resuscitation Council, Australian

- 1 and New Zealand Council on Resuscitation, Heart and Stroke Foundation of Canada,  
2 InterAmerican Heart Foundation, Resuscitation Council of Southern Africa,  
3 Resuscitation Council of Asia); and the American Heart Association Emergency  
4 Cardiovascular Care Committee and the Council on Cardiopulmonary, Critical Care,  
5 Perioperative and Resuscitation. *Resuscitation* 2015;96:328–40.  
6 <https://doi.org/10.1016/j.resuscitation.2014.11.002>.
- 7 [26] Amsterdam EA, Wenger NK, Brindis RG, Casey DE, Ganiats TG, Holmes DR, et al.  
8 2014 AHA/ACC guideline for the management of patients with non-ST-elevation acute  
9 coronary syndromes: a report of the American College of Cardiology/American Heart  
10 Association Task Force on Practice Guidelines. *Circulation* 2014;130:e344-426.  
11 <https://doi.org/10.1161/CIR.000000000000134>.
- 12 [27] Lemkes JS, Janssens GN, van der Hoeven NW, Jewbali LSD, Dubois EA, Meuwissen  
13 M, et al. Coronary Angiography after Cardiac Arrest without ST-Segment Elevation. *N*  
14 *Engl J Med* 2019;380:1397–407. <https://doi.org/10.1056/NEJMoa1816897>.
- 15 [28] Corrado D, Basso C, Pavei A, Michieli P, Schiavon M, Thiene G. Trends in sudden  
16 cardiovascular death in young competitive athletes after implementation of a  
17 preparticipation screening program. *JAMA* 2006;296:1593–601.  
18 <https://doi.org/10.1001/jama.296.13.1593>.
- 19 [29] Marijon E, Tafflet M, Celermajer DS, Dumas F, Perier M-C, Mustafic H, et al. Sports-  
20 related sudden death in the general population. *Circulation* 2011;124:672–81.  
21 <https://doi.org/10.1161/CIRCULATIONAHA.110.008979>.
- 22 [30] Rosengren A, Wallentin L, Simoons M, Gitt AK, Behar S, Battler A, et al. Age, clinical  
23 presentation, and outcome of acute coronary syndromes in the Euroheart acute coronary  
24 syndrome survey. *Eur Heart J* 2006;27:789–95. <https://doi.org/10.1093/eurheartj/ehi774>.
- 25 [31] Bougouin W, Marijon E, Puymirat E, Defaye P, Celermajer DS, Le Heuzey J-Y, et al.  
26 Incidence of sudden cardiac death after ventricular fibrillation complicating acute  
27 myocardial infarction: a 5-year cause-of-death analysis of the FAST-MI 2005 registry.  
28 *Eur Heart J* 2014;35:116–22. <https://doi.org/10.1093/eurheartj/ehi453>.
- 29 [32] Burke AP, Farb A, Malcom GT, Liang Y, Smialek JE, Virmani R. Plaque rupture and  
30 sudden death related to exertion in men with coronary artery disease. *JAMA*  
31 1999;281:921–6.
- 32 [33] Karam N, Bataille S, Marijon E, Giovannetti O, Tafflet M, Savary D, et al. Identifying  
33 Patients at Risk for Prehospital Sudden Cardiac Arrest at the Early Phase of Myocardial  
34 Infarction: The e-MUST Study (Evaluation en Médecine d’Urgence des Stratégies  
35 Thérapeutiques des infarctus du myocarde). *Circulation* 2016;134:2074–83.  
36 <https://doi.org/10.1161/CIRCULATIONAHA.116.022954>.
- 37 [34] Dumas F, Bougouin W, Geri G, Lamhaut L, Rosencher J, Pène F, et al. Emergency  
38 Percutaneous Coronary Intervention in Post-Cardiac Arrest Patients Without ST-  
39 Segment Elevation Pattern: Insights From the PROCAT II Registry. *JACC Cardiovasc*  
40 *Interv* 2016;9:1011–8. <https://doi.org/10.1016/j.jcin.2016.02.001>.
- 41 [35] GBD 2015 Tobacco Collaborators. Smoking prevalence and attributable disease burden  
42 in 195 countries and territories, 1990–2015: a systematic analysis from the Global  
43 Burden of Disease Study 2015. *Lancet* 2017;389:1885–906.  
44 [https://doi.org/10.1016/S0140-6736\(17\)30819-X](https://doi.org/10.1016/S0140-6736(17)30819-X).
- 45  
46

1 **Figures legends**

2

3 **Figure 1.** Causes of SCA in young adults

4 *Details of SCA causes identified in 244 patients < 40 years admitted alive to hospital.*

5 *ARVC, Arrhythmogenic right ventricular cardiomyopathy; LVNC, left ventricle non-compaction; SADS,*  
6 *sudden arrhythmic death syndrome; SCA, sudden cardiac arrest; SUDS, sudden unexplained death*  
7 *syndrome; VF, ventricular fibrillation; WPW, Wolff-Parkinson-White*

8

9 **Figure 2.** SCA etiologies identified by age groups (dark blue representing acute coronary syndrome  
10 and light blue chronic coronary artery disease).

11 *Etiologies of SCA by age group (<30 years, 30-40 years, ≥40 years).*

12

13 **Figure 3.** Evolution of the rate of immediate coronary angiography at hospital admission in patients  
14 ≥ 40 years (n = 3,170) compared with patients < 40 years (n = 409).

15

1 **Tables**

2

3 **Table 1.** General characteristics and resuscitation data of SCA admitted alive to hospital.

Sudden Cardiac Arrests	All n/3,579 (%)	18-40 y n/409 (%)	≥ 40 y n/3,170 (%)	p
Age (years±SD)	59.3±15.6	32.3±6.2	62.8±12.8	<0.001
Male gender	2,616 (73.1)	285 (69.7)	2,331 (73.5)	0.111
Prior known cardiac disease	1,099 (33.1)	59 (15.8)	1,040 (35.3)	<0.001
Coronary artery disease	592 (17.9)	4 (1.1)	588 (20.1)	<0.001
Non-ischemic heart disease	300 (8.5)	41 (10.1)	259 (8.3)	0.251
AF or flutter	338 (10.2)	6 (1.6)	332 (11.3)	<0.001
Pacemaker	89 (2.7)	4 (1.1)	85 (2.9)	0.040
Implantable cardioverter defibrillator	28 (0.8)	2 (0.5)	26 (0.9)	0.763
≥ 1 cardiovascular risk factor	2,715 (81.6)	202 (53.6)	2,513 (85.2)	<0.001
Hypertension	1,271 (38.3)	24 (6.4)	1,247 (42.4)	<0.001
Overweight (BMI > 25 kg/m <sup>2</sup> )	1,189 (35.6)	88 (23.2)	1,101 (37.2)	<0.001
Current smoking	1,128 (34.1)	134 (35.4)	994 (34.0)	0.641
Dyslipidemia	745 (22.5)	15 (4.0)	730 (24.9)	<0.001
Diabetes mellitus	583 (17.6)	8 (2.1)	575 (19.6)	<0.001
Family history of CAD	184 (5.6)	24 (6.4)	160 (5.5)	0.471
Chronic respiratory failure	215 (6.5)	4 (1.1)	211 (7.2)	<0.001
Chronic renal failure	178 (5.4)	7 (1.9)	171 (5.8)	0.001
Stroke	176 (5.3)	4 (1.1)	172 (5.9)	<0.001
Family history of SCA	84 (2.5)	30 (8.0)	54 (1.8)	<0.001
Sports-related	176 (5.0)	49 (12.2)	127 (4.1)	<0.001
Public place (vs home)	1,625 (45.5)	189 (46.2)	1,436 (45.4)	0.785
Witnessed SCA	3,262 (91.2)	367 (89.7)	2,895 (91.4)	0.319
Witnessed-CPR	2,347 (71.8)	271 (73.8)	2,076 (71.5)	0.383
Time from collapse to basic life support, median (IQR)	3.0 (0.0-8.3)	4.0 (0.0-10.0)	3.0 (0.0-8.0)	0.253
Time from EMS call to EMS arrival, median (IQR)	9.0 (7.0-12.0)	9.0 (7.0-12.0)	9.0 (7.0-12.0)	1
Initial shockable rhythm	1,905 (55.7)	231 (58.2)	1,674 (55.4)	0.314
Epinephrine use	2,149 (62.1)	264 (67.2)	1,885 (61.4)	0.030
ST segment elevation on first ECG	1,043 (33.1)	77 (23.5)	966 (34.2)	<0.001
Discharged alive	996 (27.9)	129 (31.6)	868 (27.4)	0.103
CPC score 1-2 at discharge	931 (26.0)	119 (29.1)	812 (25.6)	0.224

4

5 *AF, atrial fibrillation; BMI, body mass index; CAD, coronary artery disease; CPR,*

6 *cardiopulmonary resuscitation; ECG, electrocardiogram; EMS, emergency medical service;*

7 *SCA, sudden cardiac arrest; SD, standard deviation*

8

9

1  
2  
3  
4  
5  
6  
7  
8

**Table 2.** Characteristics of young patients referred or not for immediate coronary angiography

<b>Sudden Cardiac Arrests</b>	<b>&lt; 40 y Coronary angiography - n / 238 (%)</b>	<b>&lt; 40 y Coronary angiography + n / 170 (%)</b>	<b>p</b>
<b>Age (years±SD)</b>	31.5±6.3	33.3±5.8	0.003
<b>Male gender</b>	157 (66.0)	128 (75.3)	0.049
<b>Sports-related</b>	18 (7.7)	31 (18.5)	0.002
<b>Public location</b>	98 (41.2)	91 (53.5)	0.018
<b>Witnessed SCA</b>	208 (87.4)	158 (92.9)	0.098
<b>Witnessed-CPR</b>	141 (67.8)	130 (82.3)	0.003
<b>Time from collapse to basic life support, median (IQR)</b>	4.5 (0.0-10.0)	3.0 (0.0-7.0)	0.015
<b>Time from EMS call to EMS arrival, median (IQR)</b>	9.0 (7.0-12.0)	9.0 (7.0-11.0)	0.002
<b>Initial shockable rhythm</b>	95 (41.1)	135 (81.8)	<0.001
<b>Epinephrine use</b>	179 (77.8)	84 (51.9)	<0.001
<b>ST segment elevation on first ECG</b>	7 (4.1)	70 (44.0)	<0.001

*CPR, cardiopulmonary resuscitation; ECG, electrocardiogram; EMS, emergency medical service; IQR: interquartile range; SCA, sudden cardiac arrest; SD: standard deviation*

1 **Table 3.** Factors associated with immediate coronary angiography performance (among 3,579  
 2 SCA patients admitted alive at hospital).

	Univariate analysis		Multivariate analysis	
	OR (95% CI)	p	OR (95% CI)	p
<b>Age, &lt; 40 years</b>	0.38 (0.31-0.47)	<0.001	0.34 (0.25-0.47)	<0.001
<b>Male gender</b>	1.91 (1.65-2.22)	<0.001	1.49 (1.18-1.87)	<0.001
<b>Public location</b>	1.74 (1.52-2.00)	<0.001	1.29 (1.04-1.60)	0.023
<b>Witnessed SCA</b>	1.89 (1.50-2.38)	<0.001	0.91 (0.19-4.79)	0.902
<b>Witnessed-CPR</b>	1.37 (1.17-1.61)	<0.001	1.28 (0.98-1.69)	0.075
<b>Time from collapse to basic life support, &gt; 3 min</b>	0.87 (0.75-1.01)	0.055	1.31 (1.02-1.69)	0.038
<b>Time from EMS call to EMS arrival, &gt; 9 min</b>	0.87 (0.75-0.99)	0.043	0.97 (0.78-1.19)	0.762
<b>Initial shockable rhythm</b>	5.19 (4.47-6.05)	<0.001	3.35 (2.70-4.17)	<0.001
<b>Epinephrine use</b>	0.34 (0.29-0.40)	<0.001	0.43 (0.34-0.54)	<0.001
<b>ST segment elevation on first ECG</b>	9.71 (7.72-12.37)	<0.001	7.14 (5.36-9.67)	<0.001

3

4

5 *CI, confidence interval; CPR, cardiopulmonary resuscitation; ECG, electrocardiogram; EMS,*  
 6 *emergency medical service; SCA, sudden cardiac arrest*

7

8

9

10


11

12

13

14

15




2  
3 **Figure 1.** Causes of SCA in young adults

4 *Details of SCA causes identified in 244 patients < 40 years admitted alive to hospital.*

5 *ARVC, Arrhythmogenic right ventricular cardiomyopathy; LVNC, left ventricle non-compaction; SADS,*  
 6 *sudden arrhythmic death syndrome; SCA, sudden cardiac arrest; SUDS, sudden unexplained death*  
 7 *syndrome; VF, ventricular fibrillation; WPW, Wolff-Parkinson-White*

8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20

1  
2  
3




4  
5

6 **Figure 2.** SCA etiologies identified by age groups (dark blue representing acute coronary syndrome  
7 and light blue chronic coronary artery disease).

8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21



1  
2



4

5 **Figure 3.** Evolution of the rate of immediate coronary angiography at hospital admission in patients

6  $\geq 40$  years (n = 3,170) compared with patients < 40 years (n = 409).

7

8

9

## **Data Supplement– List of Paris-Sudden Death Expertise Center Investigators**

- 1  
2
- 3 **Adnet F**, AP-HP, Paris 13 University, Avicenne Hospital, Emergency Medical Services  
4 (SAMU) 93, Bobigny, France
- 5 **Agostinucci JM**, AP-HP, Paris 13 University, Avicenne Hospital, Emergency Medical  
6 Services (SAMU) 93, Bobigny, France
- 7 **Aissaoui-Balanant N**, AP-HP, Georges Pompidou European Hospital, Medical Intensive  
8 Care Unit, Paris, France
- 9 **Algalarrondo V**, AP-HP, Antoine Beclere University Hospital, Department of Cardiology,  
10 Clamart, France
- 11 **Alla F**, French National Health Insurance (CNAMTS), Paris, France
- 12 **Alonso C**, Clinique Ambroise Paré, Department of Cardiology, Neuilly, France
- 13 **Amara W**, Hospital Group of Montfermeil – Le Raincy, Department of Cardiology,  
14 Montfermeil, France
- 15 **Annane D**, AP-HP, INSERM U1173 (Laboratory of Inflammation and Infection ), University  
16 of Versailles - Saint Quentin en Yvelines, Raymond Poincare Hospital, General  
17 Intensive Care Unit, Garches, France
- 18 **Antoine C**, Biomedecine agency, Saint-Denis-la-Plaine, France
- 19 **Aubry P**, AP-HP, Bichat Hospital, Department of Cardiology, Paris, France
- 20 **Azoulay E**, AP-HP, Paris Diderot Sorbonne University, Saint Louis Hospital, Medical  
21 Intensive Care Unit, Paris, France
- 22 **Beganton F**, INSERM U970 - PARCC, Paris, France
- 23 **Benhamou D**, AP-HP, Bicetre Hospital, Department of Anaesthesia and Intensive Care  
24 Medicine, Le Kremlin-Bicetre, France
- 25 **Billon C**, AP-HP, Georges Pompidou European Hospital, Department of Genetic, Paris,  
26 France
- 27 **Bougouin W**, INSERM U970 - PARCC, Paris, France
- 28 **Boutet J**, AP-HP, Raymond Poincare Hospital, Emergency Medical Services (SAMU) 92,  
29 Garches, France
- 30 **Bruel C**, Saint Joseph Hospital, Medical-Surgical Intensive Care Unit, Paris, France
- 31 **Bruneval P**, AP-HP, Georges Pompidou European Hospital, Pathology Department, Paris,  
32 France
- 33 **Cariou A**, AP-HP, Cochin Hospital, Medical Intensive Care Unit, Paris, France
- 34 **Carli P**, AP-HP, Paris Descartes University, Necker - Enfants Malades Hospital, Emergency  
35 Medical Services (SAMU) 75, Intensive Care Unit, Paris, France
- 36 **Casalino E**, AP-HP, Emergency Medical Services (SAMU) 92, Clichy, France
- 37 **Cerf C**, Foch Hospital, Intensive Care Unit, Suresnes, France
- 38 **Chaib A**, Andre Gregoire Hospital, Department of Cardiology, Montreuil, France
- 39 **Cholley B**, AP-HP, Georges Pompidou European Hospital, Department of Anaesthesia and  
40 Surgical Intensive Care Medicine, Paris, France
- 41 **Cohen Y**, AP-HP, Avicenne Hospital, Medical-Surgical Intensive Care Unit, Bobigny,  
42 France
- 43 **Combes A**, AP-HP, Pitié-Salpetriere Hospital, Cardiology Institute (ICAN), Intensive Care  
44 Unit, Paris, France
- 45 **Crahes M**, AP-HP, Georges Pompidou European Hospital, Pathology Department, Paris,  
46 France
- 47 **Da Silva D**, Delafontaine Hospital, Intensive Care Unit, Saint-Denis, France
- 48 **Das V**, Andre Gregoire Hospital, Medical-Surgical Intensive Care Unit, Montreuil, France

1 **Demoule A**, AP-HP, Pitie Salpetriere Hospital, Medical Intensive Care Unit and Respiratory  
2 Division, Paris, France  
3 **Denjoy I**, AP-HP, Bichat Hospital, Department of Cardiology, Paris, France  
4 **Deye N**, AP-HP, Lariboisiere Hospital, Intensive Care Unit, Paris, France  
5 **Dhonneur G**, AP-HP, Henri Mondor Hospital, Department of Anaesthesia and Surgical  
6 Intensive Care Medicine, Creteil, France  
7 **Diehl JL**, AP-HP, Georges Pompidou European Hospital, Medical Intensive Care Unit, Paris,  
8 France  
9 **Dinanian S**, AP-HP, Antoine Beclere Hospital, Department of Cardiology, Clamart, France  
10 **Domanski L**, Brigade de Sapeurs Pompiers de Paris (BSPP), Paris, France  
11 **Dreyfuss D**, AP-HP, Louis Mourier Hospital, Intensive Care Unit, Colombes, France  
12 **Duboc D**, AP-HP, Cochin Hospital, Department of Cardiology, Paris, France  
13 **Dubois-Rande JL**, AP-HP, Henri Mondor Hospital, Department of Cardiology, Creteil,  
14 France  
15 **Dumas F**, AP-HP, Cochin Hospital, Department of Emergency, Paris, France  
16 **Empana JP**, INSERM U970 - PARCC, Paris, France  
17 **Extramiana F**, AP-HP, Bichat Hospital, Department of Cardiology, Paris, France  
18 **Fartoukh M**, AP-HP, Sorbonne University (Paris 6), Tenon Hospital, Intensive Care Unit,  
19 Paris, France  
20 **Fioux F**, CHI Montfermeil, Intensive Care Unit, France  
21 **Gabbas M**, French National Health Insurance (CNAMTS), Paris, France  
22 **Gandjbakhch E**, AP-HP, Pitie-Salpetriere Hospital, Cardiology Institute (ICAN),  
23 Department of Cardiology, Paris, France  
24 **Geri G**, AP-HP, Ambroise Pare Hospital, Intensive Care Unit, Boulogne-Billancourt, France  
25 **Guidet B**, AP-HP, Sorbonne University (Pierre et Marie Curie University – Paris 6), Saint  
26 Antoine Hospital, Intensive Care Unit, Paris, France  
27 **Halimi F**, Private Hospital of Parly 2, Department of Cardiology, Le Chesnay, France  
28 **Henry P**, AP-HP, Sorbonne Paris Cite University (Paris Diderot University), Lariboisiere  
29 Hospital, Department of Cardiology, Paris, France  
30 **Hidden Lucet F**, AP-HP, Pitie-Salpetriere Hospital, Cardiology Institute (ICAN),  
31 Department of Cardiology, Paris, France  
32 **Jabre P**, INSERM U970 – PARCC, Emergency Medical Services (SAMU) 75, Paris, France  
33 **Jacob L**, AP-HP, Saint Louis Hospital, Department of Anaesthesia and Surgical Intensive  
34 Care Medicine, Paris, France  
35 **Joseph L**, Bicetre Hospital, Transplant Coordination, Le Kremlin-Bicetre, France  
36 **Jost D**, Brigade de Sapeurs Pompiers de Paris (BSPP), Paris, France  
37 **Jouven X**, AP-HP, Georges Pompidou European Hospital, Department of Cardiology, Paris,  
38 France  
39 **Karam N**, AP-HP, Georges Pompidou European Hospital, Department of Cardiology, Paris,  
40 France  
41 **Kassim H**, INSERM U970 - PARCC, Paris, France  
42 **Lacotte J**, Private Hospital Jacques Cartier, Department of Cardiology, Massy, France  
43 **Lahlou-Laforet K**, AP-HP, Georges Pompidou European Hospital, Department of  
44 Psychiatry, Paris, France  
45 **Lamhaut L**, AP-HP, Paris Descartes University, Necker - Enfants Malades Hospital,  
46 Emergency Medical Services (SAMU) 75, Intensive Care Unit, Paris, France  
47 **Lanceleur A**, Foch Hospital, Intensive Care Unit, Suresnes, France  
48 **Langeron O**, AP-HP, Pitie-Salpetriere Hospital, Department of Anaesthesia and Surgical  
49 Intensive Care Medicine, Paris, France

1 **Lavergne T**, AP-HP, Georges Pompidou European Hospital, Department of Cardiology,  
2 Paris, France  
3 **Lecarpentier E**, AP-HP, Henri Mondor Hospital, Emergency Medical Services (SAMU) 94,  
4 Creteil, France  
5 **Leenhardt A**, AP-HP, Bichat Hospital, Department of Cardiology, Paris, France  
6 **Lellouche N**, AP-HP, Henri Mondor Hospital, Department of Cardiology, Creteil, France  
7 **Lemiale V**, AP-HP, Paris Diderot University, Saint Louis Hospital, Medical Intensive Care  
8 Unit, Paris, France  
9 **Lemoine F**, Brigade de Sapeurs Pompiers de Paris (BSPP), Paris, France  
10 **Linval F**, AP-HP, Paris 13 University, Avicenne Hospital, Emergency Medical Services  
11 (SAMU) 93, Bobigny, France  
12 **Loeb T**, AP-HP, Raymond Poincare Hospital, Emergency Medical Services (SAMU) 92,  
13 Garches, France  
14 **Ludes B**, Institute of Legal Medicine, Paris Descartes University, Paris, France  
15 **Luyt CE**, AP-HP, Pitie-Salpetriere Hospital, Cardiology Institute (ICAN), Intensive Care  
16 Unit, Paris, France  
17 **Maltret A**, AP-HP, Necker - Enfants Malades Hospital, Department of Pediatric Cardiology,  
18 Paris, France  
19 **Mansencal N**, Ambroise Pare Hospital, Department of Cardiology, Boulogne-Billancourt,  
20 France  
21 **Mansouri N**, AP-HP, Henri Mondor Hospital, Emergency Medical Services (SAMU) 94,  
22 Creteil, France  
23 **Marijon E**, AP-HP, Georges Pompidou European Hospital, Department of Cardiology, Paris,  
24 France  
25 **Marty J**, AP-HP, Henri Mondor Hospital, Emergency Medical Services (SAMU) 94, Creteil,  
26 France  
27 **Maury E**, AP-HP, Sorbonne University (Pierre et Marie Curie University – Paris 6), Saint  
28 Antoine Hospital, Intensive Care Unit, Paris, France  
29 **Maxime V**, AP-HP, Raymond Poincare Hospital, General Intensive Care Unit, Garches,  
30 France  
31 **Megarbane B**, AP-HP, Lariboisiere Hospital, Intensive Care Unit, Paris, France  
32 **Mekontso-Dessap A**, AP-HP, DHU A-TVVB, CARMAS Research Group, Henri Mondor  
33 Hospital, Intensive Care Unit, Creteil, France  
34 **Mira JP**, AP-HP, Cochin Hospital, Medical Intensive Care Unit, Paris, France  
35 **Monnet X**, AP-HP, Bicetre Hospital, Medical Intensive Care Unit, Le Kremlin-Bicetre,  
36 France  
37 **Narayanan K**, INSERM U970 - PARCC, Paris, France  
38 **Ngoyi N**, AP-HP, Paris 7 University, Beaujon Hospital, Emergency Medical Services  
39 (SAMU) 92, Clichy, France  
40 **Perier MC**, INSERM U970 - PARCC, Paris, France  
41 **Piot O**, Centre Cardiologique du Nord, Department of Cardiology, Saint-Denis, France  
42 **Pirracchio R**, AP-HP, Georges Pompidou European Hospital, Intensive Care Unit, Paris,  
43 France  
44 **Plaisance P**, Emergency Medical Services (SAMU) 75, Lariboisière SMUR, Paris, France  
45 **Plu I**, AP-HP, Pitie-Salpetriere Hospital, Neuropathology Escourolle, Paris, France  
46 **Raux M**, AP-HP, Pitié Salpêtrière Hospital, Department of Anesthesiology and Critical Care,  
47 Paris, France  
48 **Revaux F**, AP-HP, Henri Mondor Hospital, Emergency Medical Services (SAMU) 94,  
49 Creteil, France  
50 **Ricard JD**, AP-HP, Louis Mourier Hospital, Intensive Care Unit, Colombes, France

1 **Richard C**, AP-HP, Bicetre Hospital, Medical Intensive Care Unit, Le Kremlin-Bicetre,  
2 France  
3 **Riou B**, AP-HP, Pitie-Salpetriere Hospital, Sorbonne University (Pierre et Marie Curie  
4 University – Paris 6), Department of Emergency Medicine and Surgery, Paris,  
5 France  
6 **Roussin F**, AP-HP, Saint Louis Hospital, Department of Anaesthesia and Surgical Intensive  
7 Care Medicine, Paris, France  
8 **Santoli F**, Robert Ballanger Hospital, Intensive Care Unit, Aulnay-sous-Bois, France  
9 **Schortgen F**, Centre Hospitalier Intercommunal Créteil, Intensive Care Unit, Creteil, France  
10 **Sharifzadehgan A**, AP-HP, Georges Pompidou European Hospital, Department of  
11 Cardiology, Paris, France  
12 **Sideris G**, AP-HP, Sorbonne Paris Cite University (Paris Diderot University), Lariboisiere  
13 Hospital, Department of Cardiology, Paris, France  
14 **Spaulding C**, AP-HP, Georges Pompidou European Hospital, Department of Cardiology,  
15 Paris, France  
16 **Teboul JL**, AP-HP, Bicetre Hospital, Medical Intensive Care Unit, Le Kremlin-Bicetre,  
17 France  
18 **Timsit JF**, AP-HP, Bichat Hospital, Medical Intensive Care Unit, Paris, France  
19 **Tourtier JP**, Brigade de Sapeurs Pompiers de Paris (BSPP), Paris, France  
20 **Tuppin P**, French National Health Insurance (CNAMTS), Paris, France  
21 **Ursat C**, AP-HP, Raymond Poincare Hospital, Emergency Medical Services (SAMU) 92,  
22 Garches, France  
23 **Varenne O**, AP-HP, Cochin Hospital, Department of Cardiology, Paris, France  
24 **Veillard-Baron A**, AP-HP, Ambroise Pare Hospital, Intensive Care Unit, Boulogne-  
25 Billancourt, France  
26 **Voicu S**, AP-HP, Lariboisiere Hospital, Intensive Care Unit, Paris, France  
27 **Wahbi K**, AP-HP, Cochin Hospital, Department of Cardiology, Paris, France  
28 **Waldmann V**, AP-HP, Georges Pompidou European Hospital, Department of Cardiology,  
29 Paris, France  
30  
31  
32