



**HAL**  
open science

## Toward reliable habitat suitability and accessibility models in an era of multiple environmental stressors

Hanne de Kort, Michel Baguette, Jonathan Roger Michel Henri Lenoir,  
Virginie Marie Stevens

► **To cite this version:**

Hanne de Kort, Michel Baguette, Jonathan Roger Michel Henri Lenoir, Virginie Marie Stevens. Toward reliable habitat suitability and accessibility models in an era of multiple environmental stressors. *Ecology and Evolution*, 2020, 10 (20), pp.10937-10952. 10.1002/ece3.6753 . hal-02958923

**HAL Id: hal-02958923**

<https://hal.sorbonne-universite.fr/hal-02958923v1>

Submitted on 6 Oct 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## REVIEW ARTICLE

# Toward reliable habitat suitability and accessibility models in an era of multiple environmental stressors

Hanne De Kort<sup>1</sup> | Michel Baguette<sup>2,3</sup> | Jonathan Lenoir<sup>4</sup> | Virginie M. Stevens<sup>2</sup>

<sup>1</sup>Plant Conservation and Population Biology, Biology Department, University of Leuven, Leuven, Belgium

<sup>2</sup>Station d'Ecologie Théorique et Expérimentale (UMR 5321 SETE), National Center for Scientific Research (CNRS), Université Toulouse III – Paul Sabatier, Moulis, France

<sup>3</sup>Institut de Systématique, Evolution, Biodiversité (UMR 7205), Muséum National d'Histoire Naturelle, Paris, France

<sup>4</sup>UR "Ecologie et Dynamique des Systèmes Anthropisés" (EDYSAN, UMR 7058 CNRS-UPJV), Université de Picardie Jules Verne, Amiens Cedex 1, France

**Correspondence**

Hanne De Kort, Plant Conservation and Population Biology, Biology Department, University of Leuven, Kasteelpark Arenberg 31, 3001 Leuven, Belgium.  
Email: hanne.dekort@kuleuven.be

**Funding information**

HDK holds a postdoctoral fellowship funded by the FWO (Research Foundation Flanders, 12P6517N).

**Abstract**

Global biodiversity declines, largely driven by climate and land-use changes, urge the development of transparent guidelines for effective conservation strategies. Species distribution modeling (SDM) is a widely used approach for predicting potential shifts in species distributions, which can in turn support ecological conservation where environmental change is expected to impact population and community dynamics. Improvements in SDM accuracy through incorporating intra- and interspecific processes have boosted the SDM field forward, but simultaneously urge harmonizing the vast array of SDM approaches into an overarching, widely adoptable, and scientifically justified SDM framework. In this review, we first discuss how climate warming and land-use change interact to govern population dynamics and species' distributions, depending on species' dispersal and evolutionary abilities. We particularly emphasize that both land-use and climate change can reduce the accessibility to suitable habitat for many species, rendering the ability of species to colonize new habitat and to exchange genetic variation a crucial yet poorly implemented component of SDM. We then unite existing methodological SDM practices that aim to increase model accuracy through accounting for multiple global change stressors, dispersal, or evolution, while shifting our focus to model feasibility. We finally propose a roadmap harmonizing model accuracy and feasibility, applicable to both common and rare species, particularly those with poor dispersal abilities. This roadmap (a) paves the way for an overarching SDM framework allowing comparison and synthesis of different SDM studies and (b) could advance SDM to a level that allows systematic integration of SDM outcomes into effective conservation plans.

**KEYWORDS**

Anthropocene, biological conservation, ecological niche modeling, global change, habitat suitability modeling, management, species range shifts, stacked species distribution model

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Ecology and Evolution* published by John Wiley & Sons Ltd.


## 1 | INTRODUCTION

Biodiversity is under threat across the globe, and its preservation requires transparent and effective guidelines for policy, conservation practitioners, and educators based on realistic assessments of biodiversity–environment relations (Elith & Leathwick, 2009; Kok et al., 2017; Pereira et al., 2010; Titeux, Henle, Mihoub, & Brotons, 2016). Species distribution modeling (SDM) has been a popular toolbox for studying relationships between environmental change stressors and spatial shifts in species' suitable habitat and for predicting the potential distribution of single species, communities, and ecosystems through user-defined environmental change scenarios (Alvarado-Serrano & Knowles, 2014; Peterson, 2003; Weber, Stevens, Diniz-Filho, & Grelle, 2017). The overall SDM framework is not just an interesting tool for identifying areas of local conservation concern or areas not yet occupied but potentially suitable; it has the potential to contribute substantially to the global protection of biodiversity and ecosystem services threatened by multiple environmental stressors, including land-use change and habitat fragmentation, climate change, invasive alien species, pollution, and overexploitation (Franklin, 2013; Kok et al., 2017; Wiens, Stralberg, Jongsomjit, Howell, & Snyder, 2009).

Ignoring the joint effects of multiple environmental stressors can be highly misleading; they have been found to give rise to ecological outcomes unpredicted by single environmental stressors (Bellard, Leclerc, & Courchamp, 2015; Fournier, Barbet-Massin, Rome, & Courchamp, 2017; Guo, Lenoir, & Bonebrake, 2018; Marshall et al., 2017; Peterson & Nakazawa, 2007; Segan, Murray, & Watson, 2016; Visconti et al., 2016,) and can trigger evolutionary responses that

differ from expectations assumed by single stressor evolution (Kelly, DeBiaise, Villela, Roberts, & Cecola, 2016; McClanahan, Graham, & Darling, 2014). Accounting for multiple environmental changes and potential evolutionary responses can greatly improve model accuracy (Bellard et al., 2015; Titeux et al., 2016), providing SDM outcomes that better represent the potential distribution and ultimately the occupied distributional area of the species or community under study (Figure 1). Indeed, part of the suitable habitat of the potential distribution is often not reachable due to dispersal limitation and spatially variable habitat connectivity (e.g., physical barriers), sizing the potential distribution down to the accessible habitat (Barve et al., 2011; Peterson, Papeş, & Soberón, 2015; Pulliam, 2000). Where dispersal of the focal species strongly depends on the distribution and dispersal of interacting species, the biotic context is an additional determinant of the occupied distributional area of the focal species (Figure 1). Finally, decreased connectivity among suitable habitat patches may promote evolution toward reduced dispersal (Cote et al., 2017; Graae et al., 2018), further reducing the size of the occupied distributional area (Figure 1).

In this review, we pinpoint how joint environmental changes drive population, species, and community dynamics in comparison with single stressors, focusing on climate and land-use change as two of the most prominent threats to biodiversity. Predominantly driven by the ongoing biodiversity crisis, a universal urge for large-scale land-use restoration, and the existence of user-friendly implementation tools, scientists increasingly study the impacts of both climate- and land-use change on biodiversity redistribution using SDM (Araújo et al., 2019; Harrison & Gassner 2020; Milanesi, Rocca, & Robinson, 2020; Titeux et al., 2016). This ongoing SDM



**FIGURE 1** Schematic summary of the elements contributing to species distribution modeling (SDM) reliability and utility (a), the eco-evolutionary processes underlying distribution dynamics and SDM performance (b), and a hypothetical representation of the predicted occupied distributional area of a species (*Phengaris arion*) that relies on the presence of host plants (c). Colors represent the accessible and suitable habitat (purple), habitat unsuitable due to the predicted absence of host plant (gray), habitat with host plant but inaccessible (dark purple), inaccessible habitat without host plant (dark red), and environmentally unsuitable habitat (white background). Patches occupied by larger butterflies (representing better dispersers) are predicted to be accessible due to dispersal evolution (after DeKort, Prunier, et al., 2018)

### Box 1 The main sources of uncertainty that can compromise species distribution modeling (SDM)

The main outcomes (predicted range maps) of the whole SDM framework are frequently misleading and sometimes misused (see Carlson, Chipperfield, Benito, Telford, & O'Hara, 2020) due to several main sources of uncertainty that the user, especially the beginner user, needs to have in mind when performing SDM. First, the correlative nature of SDM inherently prevents this toolbox from building on causal relationships between environmental input variables and the occurrence of a species. This *causality limitation* is unavoidable because SDM inevitably requires predictors able to reflect or capture presumed causal mechanisms from natural history knowledge. The lack of information about the causal link between the predictor variables and the response variable, usually a binary variable representing species presence-absence or presence-background data, is clearly the main limitation that the user needs to constantly keep in mind when running SDM. For instance, every single SDM method, and especially the most advanced ones such as machine learning methods, will almost always be able to find a statistical link between spatially structured predictors and the response variable used to measure species distribution. Even when predictors are known to be completely unrelated to the species distribution, such as the caricatural but very eloquent use of paintings to predict species distributions (Fourcade, Besnard, & Secondi, 2018), SDM will allow the user to draw nice-looking maps with high prediction accuracy according to the metrics used by state-of-the-art SDM methods. This suggests another important limitation hampering the use of SDM techniques which is the *misuse or overconfidence in metrics* used to measure the performance of SDM outcomes (Lobo, Jiménez-Valverde, & Real, 2008). Indeed, typical SDM model performance metrics such as the area under the receiving operating characteristic curve (AUC), the Kappa statistic, and the true skill statistic (TCC) have been heavily criticized, as they tend to overpredict model performance under the influence of sample prevalence, consequently compromising model validation accuracy and model comparability (e.g., Leroy et al., 2018; Morán-Ordóñez, Lahoz-Monfort, Elith, & Wintle, 2017). Hence, the reason why these metrics are misleading measures of the performance of predictive distribution models partly relates to one last limitation to have in mind when running SDM which is the *data quality of the response variable*. Unless reliable and ground-truth distribution data are available, such model validation parameters are extremely questionable. In situ validation of the presence or abundance data at locations for a range of predicted probability estimates therefore is highly recommended for an unbiased perspective of model performance. The final but equally important issue regarding data quality revolves around *absence data*. All the metrics used to measure the performance of SDM outcomes somehow rely on absence information, whether obtained from field observations or from a random selection strategy of background data, also known as pseudo-absences. Clearly, collection of field-validated true absence data across the species range is highly encouraged over pseudo-absences selection strategies to improve SDM reliability (Leroy et al., 2018; Lobo, 2016). The sampling strategy used to select pseudo-absences, such as considerations on the spatial extent to be used, is known to highly influence SDM outcomes (Barbet-Massin, Jiguet, Albert, & Thuiller, 2012; Lobo, Jiménez-Valverde, & Hortal, 2010; Van Der Wal, Shoo, Graham, & Williams, 2009). Even absence data recorded during field surveys are tricky to use and should be carefully handled depending on the aim of the undertaken SDM exercise (Hattab et al., 2017): mapping the occupied distributional area or modeling the potential distribution. If the user aims at mapping the occupied distributional area, which is more or less a spatial interpolation of the suitable and accessible locations for the focal species, then all absence data recorded during field surveys are useful for model calibration and validation: whether it represents true environmental absences or dispersal-limited absences reflecting sites out of dispersal reach but potentially suitable regarding the abiotic and biotic conditions. In such cases, absence data bearing dispersal limitation information are very important and should be incorporated in the SDM framework together with predictor variables reflecting dispersal constraints, such as species-specific dispersal kernel, so as to improve the mapping of the occupied distributional area (Lobo et al., 2010; Meentemeyer, Anacker, Mark, & Rizzo, 2008; Václavík & Meentemeyer, 2009). However, if the user aims at modeling the potential distribution, which implies spatial extrapolations beyond the actual occupied distributional area, then it is recommended to exclude dispersal-limited absence from both model calibration and validation (Hattab et al., 2017). Indeed, using dispersal-limited absences to validate the potential distribution predicted from SDM will inevitably and mathematically lead to low AUC or TSS values, which results in models misclassified as having poor predictive performance. Hence, one needs to carefully think about the meaning of the available absence data and how to use it for SDM calibration and validation steps. Modeling the potential distribution or mapping the occupied distributional area are two very different SDM exercises requiring a different thinking on the use of absence data (Hattab et al., 2017).

boost opens the door for comparing and synthesizing published SDM studies for answering taxon-wide and large-scale research questions, including the role of species' traits and evolutionary potential in driving general species distribution shifts in response to land-use change. The lack of an overarching, streamlined, and

widely adoptable SDM framework containing all tools required for model preparation, parameterization, and selection, however, likely strongly refrains SDM from its full potential (Araújo et al., 2019). Moreover, a standardized and science-based SDM toolbox would pave the way for practitioners to more systematically use

SDM for conservation decision making while being aware of the potential pitfalls (Box 1). Strikingly, however, while land use has begun its debut into SDM, scientists do not generally quantify or map the amount of habitat that will likely be colonized by the species under study. To meet this concern, we specifically elaborate on the impacts of reduced habitat connectivity, a major side effect of various environmental changes, on SDM outcomes. We also investigate how dispersal is limited both by the effects of the landscape structure and configuration, and by changes in biotic interactions within communities. We then discuss why evolutionary processes should be incorporated to increase the reliability of SDM (Bush et al., 2016), and finally propose an SDM roadmap that incorporates multiple environmental stressors, dispersal, and evolution into a feasible, more reliable, and streamlined modeling framework. We stress that even for data-insufficient systems, it is possible to incorporate the processes described above. For rare and poor-disperser species in particular, realistic scenarios that take into account the various drivers of population and range dynamics can have crucial conservation implications. An SDM roadmap could also motivate nonexperts in the modeling field and young scientists to apply SDM to their target species, thereby facilitating the implementation of SDM in applied sciences and conservation practices. We stress, however, that expert evaluation of any SDM study is vital to a correct interpretation of SDM outcomes (see Box 1 for potential pitfalls).

## 2 | TOWARD ACCURATE AND MORE RELIABLE SDM

### 2.1 | The climatic niche: ecological implications

The majority of SDM studies still relies on a single environmental stressor (or group of related stressors), with bioclimatic stressors (e.g., warmer and dryer conditions) representing, by far, the most popular environmental predictor used to forecast species' distribution changes (Titeux et al., 2016). These studies observed drastic reductions in the modeled climatic suitability of currently occupied habitat for macroinvertebrates (up to 65%, Domisch et al., 2013; Parmesan et al., 1999), vertebrates (up to 80%, Warren, Wright, Seifert, & Shaffer, 2014), and plants (up to 90%, Aguirre-Gutiérrez, van Treuren, Hoekstra, & van Hintum, 2017; Kane et al., 2017). In addition, poleward and upward shifts of species distributions are widely observed (Perry, Reid, Ibanez, Lindley, & Edwards, 2005; Kelly & Goulden, 2008; Chen, Hill, Ohlemuller, Roy, & Thomas, 2011; Lenoir & Svenning 2013) and predicted (Aguirre-Gutiérrez et al., 2017; Barton, Irwin, Finkel, & Stock, 2016; Inoue & Berg, 2017), yet the velocity of species range shifts is generally thought to be inferior to the velocity of climate change (Bertrand et al., 2011; Chivers, Walne, & Hays, 2017; Corlett & Westcott, 2013; Devictor et al., 2012; Liang, Duveneck, Gustafson, Serra-Diaz, & Thompson, 2017; Schloss, Nunez, & Lawler, 2012; Zhu, Woodall, & Clark, 2012).

The effective impact of climate change on biodiversity goes beyond direct climate–occurrence relations, also involving disruption of habitat connectivity and of species interactions within communities (Bertrand et al., 2016; Garcia, Cabeza, Rahbek, & Araujo, 2014; Walther et al., 2002). First, through reducing the amount of suitable habitat, climate change increases isolation between remaining patches, consequently inhibiting gene flow across the landscape and impairing population dynamics (Graae et al., 2018; Inoue & Berg, 2017; Razgour et al., 2018). Reduced gene flow increases local inbreeding risk and extinction rates, and reduces the exchange of adaptive variation (Razgour et al., 2018; Slatkin, 1987). Second, climate change renders habitat more prone to alien species' invasions (Bellard et al., 2013; Hulme, 2017) and can alter community composition and ecosystem processes (Carroll et al., 2015; García Molinos et al., 2015; Pearson et al., 2013; Perring et al., 2016; Sheldon, Yang, & Tewksbury, 2011; Sunday, Bates, & Dulvy, 2012). Global change may therefore not only alter species' distributions through direct abiotic environment–occurrence interactions but also indirectly through shaping the biotic context (e.g., prey, competitors, and pollinators) (Carroll et al., 2015; González-Varo et al., 2013; Warren & Bradford, 2014; Wisz et al., 2013).

### 2.2 | The potential distribution is shaped by a joint environmental niche

Over half of the terrestrial, ice-free surface is transformed through ongoing land-use change and habitat loss and fragmentation, considerably adding to the impacts of climate change on terrestrial biodiversity (Aguar et al., 2016; Hansen et al., 2013; Newbold et al., 2015; Reino, Beja, Araújo, Dray, & Segurado, 2013; Titeux et al., 2016; Vitousek, 1994, but see Warren et al., 1999 for antagonistic effects). Assuming unchanged (static) land use in SDM thus renders future projections questionable (Ay, Guillemot, Martin-StPaul, Doyen, & Leadley, 2017; Fournier et al., 2017; Perring et al., 2016; Titeux et al., 2016), yet despite the continuous recognition that land-use change constitutes a major threat to global biodiversity (Perring et al., 2016; Titeux et al., 2016), the translation of this awareness into SDM keeps lagging behind. Given the extent and magnitude of climate and land-use change, their combined impact on natural ecosystems is expected to be complex (Guo et al., 2018), urging for a transparent framework allowing the identification of areas susceptible to combined global change threats.

Co-occurrence of climate and land-use change has been shown to have interactive and often synergistic effects on biodiversity and species redistribution (Jetz, Wilcove, & Dobson, 2007; Marshall et al., 2017; Pereira et al., 2010; Visconti et al., 2016; Zwiener et al., 2017). First, land-use change reinforces climate warming when it is associated with livestock breeding and deforestation, which considerably boost greenhouse gas emissions (Naudts et al., 2016; Reisinger & Clark, 2017). Second, land-use change increases the amount of suitable habitat edges that are sensitive to climate change due to the absence of a protective microclimate (Brook, Sodhi, & Bradshaw,

2008; Lembrechts, Nijs, & Lenoir, 2018; Suggitt et al., 2020, Vanneste et al., 2020). Typical examples are the increased risk of forest fires due to increased wind exposure of forest remnant edges and inward desiccation of natural grasslands due to conversion of surrounding habitat into intensively managed agricultural land (e.g., Alencar, Brando, Asner, & Putz, 2015; Tuff, Tuff, & Davies, 2016). Third, deforestation can accelerate the rate of upslope range shifts in tropical regions whereas the opposite has been demonstrated for higher latitudes (Guo et al., 2018), indicating the confounding effects of land-use change on climate-driven range shifts. Fourth, reduced gene flow across the remaining suitable landscape jeopardizes the exchange of adaptive genetic variation that could otherwise allow evolutionary adaptation to a changing climate. Inhibited gene flow also compromises genetic diversity and fitness within remaining populations, increasing their vulnerability to environmental stressors (Frankham, Ballou, Briscoe, & Ballou, 2002; Markert et al., 2010; Schrieber & Lachmuth, 2017). A global study assessing the interactive impacts of climate and habitat loss on vertebrate diversity predicted that nearly half of the ecoregions worldwide, mainly including tropical forest, savannahs, and wetlands, will be impacted by a synergy between habitat loss and climate change during the 21st century (Segan et al., 2016).

Species distribution modeling studies increasingly include variables representing both climate and land-use change in their models and have found varying support for land-use change impacts on predicted species distributions, depending on the thematic resolution of the land-use variables and the species under study (Brodie, 2016; Marshall et al., 2017; Martin, Van Dyck, Dendoncker, & Titeux, 2013; Scheller & Mladenoff, 2005; Zamora-Gutierrez, Pearson, Green, & Jones, 2018). With the availability of a 250-m resolution regional land-use map and IUCN distribution maps for 286 mammal species, Brodie (2016) was able to study the interactive effects of climate change and land-use change (through oil palm plantations) on mammal diversity in South-East Asia. Mammal species were found to be robust to changes in climate alone, but responded dramatically to the combined effects of climate and land-use change, with median habitat suitability losses of 47.5% (low carbon emission scenario) up to 67.7% (high emission). In a similar vein, SDM and conservation prioritization testing based on 2,255 woody plant species of the Brazilian Atlantic Forest showed that contrasting climate change scenarios did not shape conservation prioritizations, while management strategies aiming to reduce habitat fragmentation were found to be indispensable for the long-term conservation of Atlantic Forest diversity (Zwiener et al., 2017). We stress that while the majority of SDM studies demonstrates adverse effects of climate and land-use change on the extent of species' distributions, a carefully implemented SDM framework could reveal the true extent of positive global change impacts on species' distributions. For instance, forest fragmentation is likely to be detrimental to the redistribution of forest plant species—known to be poor dispersers—under climate change while it may favor the spread of more generalist plant species that are less constrained by their dispersal abilities and benefit from more open conditions inside forest edges.

## 2.3 | The occupied distributional area is shaped by the accessibility of suitable habitat

The potential distribution as predicted by SDM unconstrained by dispersal limitations is fundamentally different from the occupied distributional area, that is, the distribution actually occupied by the species, requiring the integration of species' dispersal abilities and dispersal barriers into SDM. In other words, the occupied distributional area of a given species (also referred to as the realized distribution) can be seen as a spatial interpolation of the suitable and accessible locations occupied by the focal species at a given moment in time: an instantaneous map of the real spatial occupation of the focal species. By contrast, the potential distribution can be interpreted as a spatial extrapolation, albeit not an environmental extrapolation beyond the species environmental niche, of where the species could find suitable environmental conditions to occur if it would be able to reach that location, that is, unlimited by its own dispersal abilities or by dispersal barriers.

Many studies reporting poleward or upward range shifts in the occupied distributional area also show that actual expansion rates of the studied organisms lag behind the displacement of their climatic envelopes (i.e., the potential distribution) (e.g., Bertrand et al., 2011; Bertrand et al., 2016), most likely due to dispersal and establishment lags at the leading edge (Alexander et al., 2018). Two nonexclusive factors can explain dispersal and establishment lags: (a) The displacements of individuals are slowed down by low habitat connectivity, and (b) individuals are struggling to settle viable populations in new habitat due to their dependence on biotic interactions. A basic understanding of these processes is required to assist with the implementation of dispersal-informed SDM (Alexander et al., 2018).

### 2.3.1 | Low habitat connectivity impedes the accessibility to suitable habitat

Habitat connectivity describes how dispersal of individuals across the landscape is facilitated or impeded by landscape structure and configuration (Taylor, Fahrig, Henein, & Merriam, 1993). Dispersal, that is, movements potentially leading to gene flow among populations (Ronce, 2007), is thus key for species to track suitable habitat shifts (Berg, Julliard, & Baguette, 2010). The study of dispersal in ecology and evolution is a swiftly evolving field of investigation since almost two decades (Bowler & Benton, 2005), generating findings that are crucial for understanding the role of dispersal in SDM. Changes in landscape structure and configuration entail high dispersal costs and hence strongly affect the fitness of dispersing individuals (Bonte et al., 2012). Accordingly, theoretical models predict that dispersal will be most generally counterselected if its costs increase along habitat fragmentation gradients and exceed its expected benefits (Cote et al., 2017; Duputie & Massol, 2013; Heino & Hanski, 2001; Mathias, Kisdí, & Olivieri, 2001; Travis & Dytham, 1999). Empirical studies confirm that habitat fragmentation can decrease dispersal propensity (the probability that an

individual leaves a habitat patch) and increase dispersal efficiency by reducing dispersal costs either through a reduced search time and/or through the selection of safer dispersal routes (Baguette, Blanchet, Legrand, Stevens, & Turlure, 2013; Baguette & Van Dyck, 2007). However, under particular conditions of landscape configuration and habitat suitability, theory also predicts the emergence of dispersal polymorphism within populations, in which high dispersal phenotypes with a generalist strategy coexist with low dispersal specialist phenotypes (Mathias et al., 2001). Taken together, habitat connectivity can have various ecological and evolutionary consequences, and fully ignoring this important component of the accessibility to suitable habitat may therefore strongly interfere with SDM outcomes.

To increase our understanding of the mismatch between the potential distribution (as modeled through SDM unconstrained by dispersal limitations or habitat connectivity) and the effectively occupied distributional area, the integration of dispersal and habitat connectivity metrics into SDM studies was put forward as a priority a decade ago (Araújo & Guisan, 2006; Hirzel & Le Lay, 2008). Although a growing number of studies discussed or implemented the effects of dispersal in SDM, the proportion of SDM studies implementing dispersal has remained steady for years (Bateman, Murphy, Reside, Mokany, & VanDerWal, 2013; Holloway & Miller, 2017). One of the most striking examples of the spatial lapse between potential distribution and the occupied distributional area was provided by an SDM study on Bornean Orangutans (Struebig et al., 2015). The authors predicted a loss of ca. 74% (within current range) and 84% (outside current range) of the potential Orangutan distribution by 2080s due to both climate change and direct habitat loss. However, given the sedentary lifestyle of the females, it is unlikely that the species would shift its distribution toward all suitable habitat (not all suitable habitat is accessible, even within the current range) with the predicted pace of environmental change. Hence, conservation corridors or assisted translocation is required to merge suitable and accessible distributions and to ensure long-term persistence of this endangered species (Struebig et al., 2015). Such extreme examples of jeopardized dispersal clearly show the urgency of a paradigm shift in conservation biology distinguishing suitable from accessible. In the same vein, Albert, Rayfield, Dumitru, and Gonzalez (2017) evidenced that accounting for connectivity in spatial prioritization of protected areas for 14 focal vertebrate species strongly modified conservation priorities.

How climate change interacts with landscape features to affect dispersal is another key question, and the fine-scale consequences of the interplay of climate and land-use change on the spatial distribution of suitable habitat may explain dispersal lagging behind climate change (Lembrechts et al., 2018). Mestre, Risk, Mira, Beja, and Pita (2017) accordingly showed that range shift predictions consecutive to climate alterations were overly optimistic when using SDM disregarding habitat connectivity. In a very interesting study, Fordham et al. (2017) showed that models incorporating both habitat connectivity and climate suitability provided better predictions of the range shifts observed between 1970 and 2017 for 20 British bird species, than do models based on climate suitability changes alone. Such integrated

modeling scenarios could be greatly simplified by translating climate changes directly into habitat connectivity changes after assessing how changes in climatic conditions modify both habitat suitability and resistance to individual movements (see also Inoue & Berg, 2017; Razgour et al., 2017). The end point of this procedure is a single model that incorporates the effects of climate change as one of the drivers of changes in habitat suitability and connectivity. Accordingly, this parsimonious approach is biologically more relevant than the production of competing models that consider the effects of either climate change or land-use change on the accessibility of suitable habitat, whereas these two factors are clearly not independent.

### 2.3.2 | Biotic interactions modulate the probability that accessible and suitable habitat is occupied

Organisms are affected by complex networks of interactions in communities that are often neglected in SDM (Wisz et al., 2013). The higher conservation value of communities than single species is another reason for SDM to stack predictions of multiple species distributions (Guisan & Rahbek 2011). The simplest study cases are couples of species living in obligatory positive (e.g., symbiotic) or negative (e.g., parasitic) relationships. SDM targeting one partner has been shown to contribute considerably to the predictive power of the presence of its obligatory associate (e.g., DeKort, Prunier, et al., 2018; Gutiérrez, Fernández, Seymour, & Jordano, 2005; Hanspach et al., 2014). Accounting for the presence of specialist species' partners in SDM can therefore have strong impacts on the probability that a suitable and accessible habitat is actually occupied by the focal specialist species, and usually results in more restricted species distributions as compared to SDM based on a single species (Dormann et al., 2018; Hanspach et al., 2014; Hof, Jansson, & Nilsson, 2012). In a context of shifting abiotic conditions, such close interactions can be disrupted by differences in dispersal abilities among interacting species further contributing to the observed lag in biotic responses to environmental stressors (Alexander et al., 2018). In general, effective dispersal of specialists depends on dispersal abilities of their least dispersive interactor (see also Supporting Information S1). For instance, the expansion rate of specialist butterflies to accessible and suitable habitat patches under contrasted scenarios of habitat connectivity has been shown to be severely curbed by the dispersal rates of their only host plants (DeKort, Prunier, et al., 2018; Schweiger, Settele, Kudrna, Klotz, & Kühn, 2008). Conversely, the dispersal lag between a weakly mobile parasitoid and its highly dispersive butterfly host provides the latter with enemy-free accessible and suitable habitat patches in highly fragmented habitats (Bergerot et al., 2010).

### 2.4 | Evolvability of the accessible and suitable habitat

The vast majority of SDM studies implicitly considers the relationship between an organism's ability to survive and the environmental

conditions as fixed in time and space. However, the probability for a species to overcome environmental changes increases, by definition, with its adaptive potential. The ecological niche of a given species can thus evolve (Visser, 2008, Wasof et al., 2013), and local adaptation can be seen as an alternative or complementary solution for dispersal under changing environmental conditions (Graae et al., 2018). Correspondingly, phenological shifts were reported for a variety of organisms, ranging from plants (e.g., Franks, Sim, & Weis, 2007) over birds (e.g., Charmantier & Gienapp, 2014) to arthropods (e.g., Van Asch, Salis, Holleman, van Lith, & Visser, 2013). Other traits, including color morphs (Karell, Ahola, Karstinen, Valkama, & Brommer, 2011), dispersal ability (Travis et al., 2013) and the thermal niche (Rolland et al., 2018) have been shown to evolve under climate change.

The key elements to phenotypic evolution and thus the evolution of the ecological niche under climate change are local genetic additive variance (underlying interindividual variation), life history (affecting both the pace of natural evolution and the feedbacks among traits), and the interplay between dispersal and the landscape structure (balancing genetic mixing and drift). Evolutionary potential may thus differ considerably within and between species. Using a dynamic eco-evolutionary model coupled to correlative niche projections, Cotto et al. (2017) showed that evolutionary adaptation is unlikely to prevent the predicted range contraction of long-lived perennial alpine plant species under predicted climate change. On the contrary, evolutionary rescue was reported for insects with short generation time and fast growth (Kearney, Porter, Williams, Ritchie, & Hoffmann, 2009). Similarly, using a hybrid SDM approach that incorporates dispersal and evolutionary dynamics into correlative SDM (see Supporting information S3), Bush et al. (2016) showed that evolutionary adaptation reduces projected range losses up to 33% for 17 species of *Drosophila*. Although evolution will probably not be adequate to ensure population persistence for most species under the current pace of climate change, slowing down the pace of climate change is expected to promote evolutionary rescue (Cotto et al., 2017).

Evolution of phenotypic traits, including dispersal evolution (see Supporting information S2), thus plays a major role in shaping and conserving the potential distribution of many species.

### 3 | TOWARD FEASIBLE SDM INTEGRATING LAND USE, DISPERSAL, AND EVOLUTION

#### 3.1 | Data collection


Researchers are increasingly motivated to share *sampling locations* on online archives, resulting in continuously growing databases. For example, the European Vegetation Archive (EVA) counted 1,529,550 vegetation plots of species co-occurrence data most of them with geographical coordinates (Chytry et al., 2016, see also sPlot: Bruelheide, Dengler, Jiménez-Alfaro, & Purschke, 2018a, Bruelheide et al., 2018b). In addition, the Global Biodiversity Information

Facility (GBIF) currently contains 1,074,675,056 georeferenced occurrence records of microorganisms, plants, and animals. Exploring such databases for the species of interest in combination with information obtained through regional nature conservation organizations and publications can improve the detail of the spatial extent of the respective species (e.g., Cardador, Carrete, Gallardo, & Tella, 2016; Diniz-Filho et al., 2016) (Figure 2.1). In parallel, researchers should continue to share their species survey data on publicly available repositories, while increasing computational power and data accessibility might increase the number of species for which mechanistic SDM can be parameterized. For instance, in birds, a high number of telemetry datasets were made available in public repositories such as Movebank (<https://www.movebank.org>) that may help parameterizing dispersal in trait-based SDM. The integration of ecology and physiology required for trait-based approaches will also certainly be favored by the routine integration of metabolic data into life-history database such as Pantheria (Jones et al., 2009).

A struggling point complicating SDM for many species is the lack of sufficient absence and occurrence data, which is problematic because species detectability often varies in space and time, resulting in poor model performance and biased SDM outcomes (Guillera-Aroita, 2017; Merow, Wilson, & Jetz, 2017; Morán-Ordóñez et al., 2017; van Proosdij, Sosef, Wieringa, & Raes, 2016) (see also Box 1). To overcome *imperfect detection*, informative field surveys should be combined with a modeling approach accounting for the detection process. This could be realized through cropping the environmental input maps to the regions that were surveyed prior to extrapolation of the environment–occurrence probability relations, or through including model parameters describing the detection conditions (Acevedo, Jiménez-Valverde, Lobo, & Real, 2012; Guillera-Aroita, 2017; Pennino et al., 2019). The choice of absence data could furthermore have strong implications on model outcomes (Hattab et al., 2017) (see also Box 1) and are ideally categorized into: (a) environmental absence data; reflecting unsuitable habitat; (b) dispersal-limited absences, reflecting inaccessible but suitable habitat; and where tight species' interactions occur, (c) community-limited absences, reflecting accessible and suitable habitat but lacking the species upon which the focal species obligatory depends. Although studies implementing field surveys that cover all types of absence data are more likely to discriminate between the potential distribution and the occupied distributional area, this distinction may be arbitrary and the selection of field-confirmed absences in the broad sense already considerably improves model reliability (Guillera-Aroita et al., 2015 Leroy et al., 2018) (Figure 2.2).

Environmental predictor maps (Figure 2.1) are often freely downloadable and can be merged with occurrence points using a basic GIS application. Global climate data are widely available for researchers to model past, current, and future climate projections (e.g., WorldClim and CHELSA), rendering climate niche modeling an attractive approach for global change research relative to land-use modeling. Yet, also regional and global land-use maps have become accessible (e.g., GlobCover, MODIS2005, CORINE) and can be converted from a vector to an SDM-friendly raster format using any





**FIGURE 2** Circular roadmap presenting the necessary steps toward science-based species distribution modeling (SDM) that is feasible for most species and implementers, with the emphasis on poor dispersers as a particularly vulnerable and valuable group of species. Note that step 4 has been given priority over step 5 because of its direct conservation implications. Step 7 represents the completion of the suitable and accessible species distribution as predicted by the SDM framework. See Figure 1 for color scheme of hypothetical distribution. We refer to the roadmap steps in the main text using bold references (e.g., Figure 2.2 refers to step 2 of the roadmap), where specific directions are provided for the respective steps. Please carefully read Box 1 to understand the importance of the “further considerations” presented in the blue rectangle of Figure 2

geographical information system (GIS rasterizing) to model land-use and habitat connectivity. Moreover, while the short-term nature of land-use maps and land-use change may complicate land-use change SDM, simple land-use change scenarios could be tested in a sensitivity analysis for a given near-future climate change scenario. Where land cover maps generally lack resolutions beyond the dominant land cover types (e.g., “grasslands” and “urban area”), there are several ways that allow more ecologically relevant land-use mapping. Information on the protection level of areas across the globe (e.g., through the protected area network) (Kremen et al., 2008) can be merged with a land cover map to indicate the level of management of specific land cover types (Rodrigues et al., 2004). Grasslands and forests are, for example, more likely to be intensively managed or exploited when unprotected. Alternatively, human population density maps and road maps can be used to fine-tune the intensity of land disturbance (e.g., Newbold et al., 2015), while a forest cover map can be integrated to model high-resolution forest change for species strongly depending on the absence or presence of forests (e.g., Hansen et al., 2013). Finally, inclusion of microclimatic variables such as topography not only allows increasing the spatial resolution and accuracy of SDM predictions, it also has been demonstrated to play an underappreciated role in shaping the trajectories of species evolution and redistribution (De Kort et al., 2020; Suggitt et al., 2020).

The choice of environmental predictors is crucial for SDM, as noncausal and redundant predictors unnecessarily increase model complexity and frequently give rise to misleading model performance estimates and flawed projections (Brodie et al., 2020; Fourcade et al., 2018; Warren, Matzke, & Iglesia, 2020). Therefore, SDM

should only be used when the presumed causality and ecological relevance of tested predictors are carefully considered (Figure 2.1) (see also Box 1). When soil water availability, for example, has been suggested to be deterministic for the presence of the species under study, a wetness index map or equivalent, should evidently be incorporated during modeling.

### 3.2 | Model parameterization and in silico validation

Although the technical aspects underlying model evaluation lie beyond the scope of this review (see, e.g., Aiello-Lammens, Boria, Radosavljevic, Vilela, & Anderson, 2015; Muscarella, Galante, Soley-Guardia, & Boria, 2014; Peterson et al., 2007; Phillips & Dudík, 2008; Radosavljevic & Anderson, 2014, for technical SDM considerations), we highlight the importance of model tuning and parameterization as a key criterion for realistic modeling (Box 1). The predictive performance of SDMs is usually evaluated through cross-validation, using training data to fit models and a set of testing data that is spatially independent of the training data to evaluate these models (Hijmans, 2012). This model evaluation approach is, however, highly sensitive to spatial autocorrelation of environmental variables between training and testing data, and only tests the set of known occurrence data. These drawbacks artificially inflate cross-validated model statistics (Hijmans, 2012; Morán-Ordóñez et al., 2017) and stress the importance of in situ validation (see below).

We recommend SDM users to consistently model at least three scenarios: climate only, land use only, and climate and land use

together, in addition to a full model that incorporates potential additional predictors such as soil moisture and biotic interactions (Figure 2.3). Such a standardized SDM framework allows (a) addressing model complexity issues (see Brun et al., 2020; Gregor, Palacios, Thompson, & Chan, 2019) and (b) comparing the contribution of major environmental change stressors to predicted shifts in species distributions between independent SDM studies. Although more efforts are needed for true cross-species and cross-study comparison of SDM outcomes (e.g., the development of an overarching, decision tree-based SDM environment that systematically informs users about parameter and predictor choice and more detailed land-use scenarios), we emphasize the promise that standardizing SDM holds (a for uncovering overall effects of dispersal, evolution, life history, and anthropogenic stressors on the projected distribution of many species and communities, and (b) for framing each study into a much larger and significant context.

### 3.3 | Integrate evolutionary potential

Integrating *evolution* into SDM without assumptions on mutation and demographic rates (Supporting information S3), which can be particularly challenging for rare species, can be achieved if the evolutionary potential of populations across the study area is quantified (Gotelli & Stanton-Geddes, 2015; Ikeda et al., 2017). Quantitative genetic screening of phenotypic traits has long been thought to provide the most accurate information on the evolutionary potential of traits and life-history syndromes. Such quantitative genetic research is now, however, questioned due to drawbacks related to limited statistical power, high time consumption, unrealistic assumptions on the genetic architecture underlying adaptive traits and poor representation of natural conditions (Hoffmann, Sgrò, & Kristensen, 2017; Wood, Yates, & Fraser, 2016). More recently, the use of genomic markers representing neutral and/or adaptive genetic variation has been proposed and tested for modeling local adaptation and adaptive potential in SDM (Fitzpatrick & Keller, 2015; Ikeda et al., 2017; Marcer, Méndez-Vigo, Alonso-Blanco, & Picó, 2016). As a consequence of local adaptation, many populations develop a local genetic signature shaped by historical and current environmental conditions (Watanabe & Monaghan, 2017). Variation in population responses and environmental change is thus expected, and shown, to affect the ability of SDMs that do not incorporate genetic structure to predict future species' distributions (Ikeda et al., 2017; Marcer et al., 2016).

In addition to the implementation of neutral genetic structure, explicitly incorporating genetic variation underlying adaptive traits can be achieved through ecology-informed genome screening (e.g., De Kort & Honnay, 2017). As phenotypic traits respond to environmental change through shifts in the underlying genes, associations between genetic and environmental variation are assumed to result from local adaptation (Manel, Schwartz, Luikart, & Taberlet, 2003; Storfer et al., 2007). Even without a sequenced reference genome, landscape genomic analysis of genetic markers, allowing the identification of genetic patterns associated with environmental variation

(e.g., climate or land use), is a feasible strategy for any species (Frichot, Schoville, Bouchard, & François, 2013; Manel & Holderegger, 2013; Rellstab, Gugerli, Eckert, Hancock, & Holderegger, 2015; Sork et al., 2016). High variation at genetic variants associated with temperature or habitat fragmentation may therefore indicate a high potential to adapt to future climate and land-use change. While the integration of this *adaptive potential* into SDM is under development (e.g., Peterson, Doak, & Morris, 2019), simple correlations between neutral and adaptive genetic diversity on the one hand, and SDM habitat suitability estimates on the other hand, indicates the extent to which adaptive evolution may affect SDM projections. High adaptive potential at the rear edge, for example, may prevent local extinctions despite a projected northward shift with climate change (Erichsen et al., 2018; Exposito-Alonso et al., 2018). Low genetic diversity at suitable locations, on the other hand, may be indicative of imminent local extinction and poor connectivity, and/or of relatively recent colonization after a period of reduced habitat suitability (e.g., Gutiérrez-Rodríguez, Barbosa, & Martínez-Solano, 2017; DeKort, Baguette, et al., 2018; DeKort, Prunier, et al., 2018). Conservation actions focusing on expanding or connecting SDM-based suitable patches holding populations of low genetic diversity may consequently increase options for dispersal and evolution. We therefore recommend conservation prioritization based on a combination of SDM and genetic marker assessment (Figures 2.2 and 2.4). Although the collection of genetic material for genetic marker analysis may be particularly challenging for rare species, noninvasive sampling such as fecal, hair, and eggshell sampling may overcome this issue (e.g., Beja-Pereira, Oliveira, Alves, Schwartz, & Luikart, 2009) (Figure 2.2).

### 3.4 | Run global change scenarios

Depending on the scale of the study, *land-use change scenarios* (Figure 2.5) could rely on local managers that are aware of ongoing land-use developments, regional storylines related to demands of arable production, livestock number, urbanization, and/or international socioeconomic parameters (e.g., DeKort, Prunier, et al., 2018; Dullinger et al., 2020; La Sorte et al., 2017; Marshall et al., 2018), or global land-use scenarios as outlined by the Intergovernmental Panel on Climate Change (IPCC). Land-use variables can subsequently be manipulated (e.g., partial conversion of extensive grasslands into forest, or forest into built-up area) using basic GIS applications to generate SDM scenarios integrating both climate change and habitat fragmentation (e.g., Lehsten et al., 2015; Marshall et al., 2018; Martin et al., 2013). *Climate change scenarios* have been outlined by the IPCC and are freely available at [chelsa-climate.org](http://chelsa-climate.org) and [WorldClim.org](http://WorldClim.org).

At this point, all data are available for modeling, using *dismo* or another SDM framework (Figure 2.5). Models are first trained and tested using the occurrence points and (cropped) environmental maps, finally providing a habitat suitability map that reflects the present distribution. The model results are then extrapolated to predict future distributions under the provided scenarios. These projections do not account for dispersal and evolution, and require

fine-tuning and contextualization based on species' life-history traits (through partial dispersal modeling, Figure 2.6) and genetic markers (through discussion of the adaptive potential, Figure 2.4).

### 3.5 | Integrate dispersal into SDM

Dispersal-informed SDM evidently requires a basic understanding of *dispersal behavior* in the species under study. Obtaining this information could be particularly challenging for species that are hard to monitor or for plant species that depend on vectors for seed dispersal. For plants, dispersal syndrome (wind, animal, ant, and ballistic or no syndrome) and growth form (tree, shrub, and herb) provide reasonable predictions of maximum dispersal distance (MacLean & Beissinger, 2017; Tamme et al., 2014; Thomson, Letten, Tamme, Edwards, & Moles, 2018), allowing integration of dispersal into SDM through simple field observations (e.g., Midgley, Hughes, Thuiller, & Rebelo, 2006; Peyre et al., 2020). For animals, movement ability, longevity, and habitat breadth are important predictors of dispersal distance and climate change-induced range shifts (MacLean & Beissinger, 2017; Stevens et al., 2013). These *life-history traits* can therefore be considered to define a partial dispersal SDM approximating the occupied distributional area to a more accurate extent than a SDM assuming no dispersal (Bateman et al., 2013). Alternatively, it is possible to create a species-specific dispersal kernel and use it as a predictor variable capturing the impact of dispersal limitations on the occupied distributional area (Hattab et al., 2017; Meentemeyer et al., 2008; Václavík & Meentemeyer, 2009) (see Box 1). Even relatively simple partial dispersal models, where the potential distribution has been clipped down to accessible distribution based on estimates of maximum dispersal distances, have been shown to improve distribution projections under environmental change (DeKort, Prunier, et al., 2018; Fitzpatrick, Gove, Sanders, & Dunn, 2008; Meier et al., 2012; Midgley et al., 2006). The implementation comfort and the limited number of assumptions related to demographic rates make this type of partial-dispersal SDM the preferred option for many species (Bateman et al., 2013). The most reliable and informative partial dispersal SDMs are expected for species with poor dispersal capacities, because *poor dispersers* (a) are often of high conservation concern, (b) facilitate integration of dispersal into SDM through assuming limited dispersal between suitable patches (Figure 2.6), and (c) provide conservative estimates of patch accessibility for associated species. Although we do not specifically recommend to focus on poor dispersers, we do believe that this important target group should receive particular attention in future SDM studies aiming to develop dispersal- and evolution-informed conservation strategies. Although model accuracy improves considerably in partial dispersal SDM, they still not fully reflect real conditions. A more mechanistic approach, for example, through hybrid models integrating both correlative and mechanistic principles, could further increase model reliability (see Supporting information S3).

### 3.6 | In situ model validation

In silico model parameterization and validation should be complemented with in situ model evaluation in unsampled regions, through extracting a set of suitable and unsuitable habitat coordinates from model output and empirically evaluate occurrence in the field (Araujo, Pearson, Thuiller, & Erhard, 2005; DeKort, Prunier, et al., 2018) (Figure 2.8). Among the rare examples of studies using in situ SDM validation, Williams et al. (2009) were able to find 24 new localities (out of 36 checkpoint sites) shared among four rare plant species across the Rattlesnake Creek Terrane in California. Area under the curve (AUC) of the receiver operating characteristic (Hanley & McNeil, 1982), a commonly used model validation statistic retrieved from the cross-validation approach, ranged between 0.94 and 0.98, commonly interpreted as nearly perfect predictive performance (but see Box 1 for pitfalls related to this model validation metrics like the AUC statistic). Two important conclusions can be drawn from this study. First, there can be considerable inconsistency between in silico (simulated) and in situ (real) model validation, which may reflect (a) the drawbacks of in silico model validation methods and (b) the (in)accessibility of suitable habitat patches. Second, SDM studies can be highly suitable for conservation purposes, given that (a) validation was performed both in silico using independent calibration and testing data and in situ, (b) relevant environmental maps and scenarios are generated, and (c) dispersal and evolution are implemented. We finally recommend projected shifts in distributions to be followed up in situ at regular time intervals (Figure 2.7) (see, e.g., Areias Guerreiro, Mira, & Barbosa, 2016; Barbet-Massin, Rome, Villemant, & Courchamp, 2018; West, Kumar, Brown, Stohlgren, & Bromberg, 2016). This has two major advantages, including the ability to test the power of SDM approaches to predict distribution shifts and to assess the impact of conservation actions on projected shifts in SDM.

#### ACKNOWLEDGMENTS

We thank the referees and editor for handling this manuscript.

#### CONFLICT OF INTEREST

No competing interests.

#### AUTHOR CONTRIBUTION

**Hanne De Kort:** Conceptualization (equal); Writing-original draft (lead); Writing-review & editing (equal). **Michel Baguette:** Conceptualization (equal); Funding acquisition (lead); Writing-original draft (equal); Writing-review & editing (equal). **Jonathan Lenoir:** Conceptualization (supporting); Supervision (equal); Writing-review & editing (equal). **Virginie Stevens:** Supervision (lead); Writing-review & editing (equal).

#### DATA AVAILABILITY STATEMENT

All data used were accessed through referenced literature.

#### ORCID

Hanne De Kort  <https://orcid.org/0000-0003-2516-0134>

## REFERENCES

- Acevedo, P., Jiménez-Valverde, A., Lobo, J. M., & Real, R. (2012). Delimiting the geographical background in species distribution modelling. *Journal of Biogeography*, *39*, 1383–1390. <https://doi.org/10.1111/j.1365-2699.2012.02713.x>
- Aguiar, A. P. D., Vieira, I. C. G., Assis, T. O., Dalla-Nora, E. L., Toledo, P. M., Oliveira Santos-Junior, R. A., ... Ometto, J. P. H. (2016). Land use change emission scenarios: Anticipating a forest transition process in the Brazilian Amazon. *Global Change Biology*, *22*, 1821–1840. <https://doi.org/10.1111/gcb.13134>
- Aguirre-Gutiérrez, J., van Treuren, R., Hoekstra, R., & van Hintum, T. J. L. (2017). Crop wild relatives range shifts and conservation in Europe under climate change. *Diversity and Distributions*, *23*, 739–750. <https://doi.org/10.1111/ddi.12573>
- Aiello-Lammens, M. E., Boria, R. A., Radosavljevic, A., Vilela, B., & Anderson, R. P. (2015). spThin: An R package for spatial thinning of species occurrence records for use in ecological niche models. *Ecography*, *38*, 541–545.
- Albert, C. H., Rayfield, B., Dumitru, M., & Gonzalez, A. (2017). Applying network theory to prioritize multispecies habitat networks that are robust to climate and land-use change. *Conservation Biology*, *31*, 1383–1396. <https://doi.org/10.1111/cobi.12943>
- Alencar, A. A., Brando, P. M., Asner, G. P., & Putz, F. E. (2015). Landscape fragmentation, severe drought, and the new Amazon forest fire regime. *Ecological Applications*, *25*, 1493–1505.
- Alexander, J. M., Chalmandrier, L., Lenoir, J., Burgess, T. I., Essl, F., Haider, S., ... Pellissier, L. (2018). Lags in the response of mountain plant communities to climate change. *Global Change Biology*, *24*, 563–579. <https://doi.org/10.1111/gcb.13976>
- Alvarado-Serrano, D. F., & Knowles, L. L. (2014). Ecological niche models in phylogeographic studies: Applications, advances and precautions. *Molecular Ecology Resources*, *14*, 233–248.
- Araújo, M. B., & Guisan, A. (2006). Five (or so) challenges for species distribution modelling. *Journal of Biogeography*, *33*, 1677–1688. <https://doi.org/10.1111/j.1365-2699.2006.01584.x>
- Araújo, M. B., Anderson, R. P., Márcia Barbosa, A., Beale, C. M., Dormann, C. F., Early, R., ... Rahbek, C. (2019). Standards for distribution models in biodiversity assessments. *Science Advances*, *5*, eaat4858.
- Araujo, M. B., Pearson, R. G., Thuiller, W., & Erhard, M. (2005). Validation of species-climate impact models under climate change. *Global Change Biology*, *11*, 1504–1513. <https://doi.org/10.1111/j.1365-2486.2005.01000.x>
- Areias Guerreiro, J., Mira, A., & Barbosa, M. (2016). How well can models predict changes in species distributions? A 13-year-old otter model revisited. *Hystrix*, [//hdl.handle.net/10174/19995](http://hdl.handle.net/10174/19995)
- Ay, J.-S., Guillemot, J., Martin-StPaul, N., Doyen, L., & Leadley, P. (2017). The economics of land use reveals a selection bias in tree species distribution models. *Global Ecology and Biogeography*, *26*, 65–77. <https://doi.org/10.1111/geb.12514>
- Baguette, M., Blanchet, S., Legrand, D., Stevens, V. M., & Turlure, C. (2013). Individual dispersal, landscape connectivity and ecological networks. *Biological Reviews*, *88*, 310–326.
- Baguette, M., & Van Dyck, H. (2007). Landscape connectivity and animal behavior: Functional grain as a key determinant for dispersal. *Landscape Ecology*, *22*, 1117–1129. <https://doi.org/10.1007/s10980-007-9108-4>
- Barbet-Massin, M., Jiguet, F., Albert, C. H., & Thuiller, W. (2012). Selecting pseudo-absences for species distribution models: How, where and how many? *Methods in Ecology and Evolution*, *3*, 327–338. <https://doi.org/10.1111/j.2041-210X.2011.00172.x>
- Barbet-Massin, M., Rome, Q., Villemant, C., & Courchamp, F. (2018). Can species distribution models really predict the expansion of invasive species? *PLoS One*, *13*(3), e0193085. <https://doi.org/10.1371/journal.pone.0193085>
- Barton, A. D., Irwin, A. J., Finkel, Z. V., & Stock, C. A. (2016). Anthropogenic climate change drives shift and shuffle in North Atlantic phytoplankton communities. *Proceedings of the National Academy of Sciences USA*, *113*, 2964–2969. <https://doi.org/10.1073/pnas.1519080113>
- Barve, N., Barve, V., Jiménez-Valverde, A., Lira-Noriega, A., Maher, S. P., Peterson, A. T., ... Villalobos, F. (2011). The crucial role of the accessible area in ecological niche modeling and species distribution modeling. *Ecol. Modell.*, *222*, 1810–1819. <https://doi.org/10.1016/j.ecolmodel.2011.02.011>
- Bateman, B. L., Murphy, H. T., Reside, A. E., Mokany, K., & VanDerWal, J. (2013). Appropriateness of full-, partial- and no-dispersal scenarios in climate change impact modelling. *Diversity and Distributions*, *19*, 1224–1234.
- Beaugrand, G., Reid, P. C., Ibanez, F., Lindley, J. A., & Edwards, M. (2005). Reorganization of North Atlantic marine copepod biodiversity and climate. *Science*, *308*, 1692–1694.
- Bellard, C., Leclerc, C., & Courchamp, F. (2015). Combined impacts of global changes on biodiversity across the USA. *Scientific Reports*, *5*, 11828. <https://doi.org/10.1038/srep11828>
- Bellard, C., Thuiller, W., Leroy, B., Genovesi, P., Bakkenes, M., & Courchamp, F. (2013). Will climate change promote future invasions? *Global Change Biology*, *19*, 3740–3748. <https://doi.org/10.1111/gcb.12344>
- Bertrand, R., Lenoir, J., Piedallu, C., Riofrío-Dillon, G., de Ruffray, P., Vidal, C., ... Gégout, J.-C. (2011). Changes in plant community composition lag behind climate warming in lowland forests. *Nature*, *479*, 517–520.
- Bertrand, R., Riofrío-Dillon, G., Lenoir, J., Drapier, J., de Ruffray, P., Gégout, J.-C., & Loreau, M. (2016). Ecological constraints increase the climatic debt in forests. *Nature Communications*, *7*, 12643.
- Bergerot, B., Julliard, R., & Baguette, M. (2010). Metacommunity dynamics: decline of functional relationship along a habitat fragmentation gradient. *PLoS One*, *5*, e11294. <https://doi.org/10.1371/journal.pone.0011294>
- Beja-Pereira, A., Oliveira, R., Alves, P. C., Schwartz, M. K., & Luikart, G. (2009). Advancing ecological understandings through technological transformations in noninvasive genetics. *Molecular Ecology Resources*, *9*, 1297–1301.
- Bonte, D., Van Dyck, H., Bullock, J. M., Coulon, A., Delgado, M., Gibbs, M., ... Travis, J. M. J. (2012). Costs of dispersal. *Biological Reviews*, *87*, 290–312. <https://doi.org/10.1111/j.1469-185X.2011.00201.x>
- Bowler, D. E., & Benton, T. G. (2005). Causes and consequences of animal dispersal strategies: Relating individual behaviour to spatial dynamics. *Biological Reviews*, *80*, 205–225. <https://doi.org/10.1017/S1464793104006645>
- Brodie, J. F. (2016). Synergistic effects of climate change and agricultural land use on mammals. *Frontiers in Ecology and the Environment*, *14*, 20–26. <https://doi.org/10.1002/16-0110.1>
- Brodie, S. J., Thorson, J. T., Carroll, G., Hazen, E. L., Bograd, S., Haltuch, M. A., ... Selden, R. L. (2020). Trade-offs in covariate selection for species distribution models: A methodological comparison. *Ecography*, *43*(1), 11–24. <https://doi.org/10.1111/ecog.04707>
- Brook, B., Sodhi, N., & Bradshaw, C. (2008). Synergies among extinction drivers under global change. *Trends in Ecology & Evolution*, *23*, 453–460. <https://doi.org/10.1016/j.tree.2008.03.011>
- Brun, P., Thuiller, W., Chauvier, Y., Pellissier, L., Wüest, R. O., Wang, Z., & Zimmermann, N. E. (2020). Model complexity affects species distribution projections under climate change. *Journal of Biogeography*, *47*, 130–142. <https://doi.org/10.1111/jbi.13734>
- Bruelheide, H., Dengler, J., Jiménez-Alfaro, B., Purschke, O. (2019). splot - a new tool for global vegetation analyses. *Journal of Vegetation Science*, *30*, 161–186.
- Bruelheide, H., Dengler, J., Purschke, O., Lenoir, J., Jiménez-Alfaro, B., Hennekens, S. M., ... Jandt, U. (2018). Global trait-environment relationships of plant communities. *Nature Ecology and Evolution*, *2*, 1906–1917.

- Bush, A., Mokany, K., Catullo, R., Hoffmann, A., Kellermann, V., Sgrò, C., ... Ferrier, S. (2016). Incorporating evolutionary adaptation in species distribution modelling reduces projected vulnerability to climate change. *Ecology Letters*, 19, 1468–1478. <https://doi.org/10.1111/ele.12696>
- Cardador, L., Carrete, M., Gallardo, B., & Tella, J. L. (2016). Combining trade data and niche modelling improves predictions of the origin and distribution of non-native European populations of a globally invasive species. *Journal of Biogeography*, 43, 967–978. <https://doi.org/10.1111/jbi.12694>
- Carlson, C. J., Chipperfield, J. D., Benito, B. M., Telford, R. J., & O'Hara, R. B. (2020). Species distribution models are inappropriate for COVID-19. *Nature Ecology and Evolution*, 4(6), 770–771. <https://doi.org/10.1038/s41559-020-1212-8>
- Carroll, M. J., Heinemeyer, A., Pearce-Higgins, J. W., Dennis, P., West, C., Holden, J., ... Thomas, C. D. (2015). Hydrologically driven ecosystem processes determine the distribution and persistence of ecosystem-specialist predators under climate change. *Nature Communications*, 6, 7851. <https://doi.org/10.1038/ncomms8851>
- Charmantier, A., & Gienapp, P. (2014). Climate change and timing of avian breeding and migration: Evolutionary versus plastic changes. *Evolutionary Applications*, 7, 15–28. <https://doi.org/10.1111/eva.12126>
- Chen, I.-C., Hill, J. K., Ohlemüller, R., Roy, D. B., & Thomas, C. D. (2011). Rapid range shifts of species associated with high levels of climate warming. *Science*, 333, 1024–1026. <https://doi.org/10.1126/science.1206432>
- Chivers, W. J., Walne, A. W., & Hays, G. C. (2017). Mismatch between marine plankton range movements and the velocity of climate change. *Nature Communications*, 8, 14434. <https://doi.org/10.1038/ncomms14434>
- Corlett, R. T., & Westcott, D. A. (2013). Will plant movements keep up with climate change? *Trends in Ecology & Evolution*, 28, 482–488. <https://doi.org/10.1016/j.tree.2013.04.003>
- Cote, J., Bestion, E., Jacob, S., Travis, J., LeGrand, D., & Bagueette, M. (2017). Evolution of dispersal strategies and dispersal syndromes in fragmented landscapes. *Ecography*, 40, 56–73. <https://doi.org/10.1111/ecog.02538>
- Cotto, O., Wessely, J., Georges, D., Klöner, G., Schmid, M., Dullinger, S., ... Guillaume, F. (2017). A dynamic eco-evolutionary model predicts slow response of alpine plants to climate warming. *Nature Communications*, 8(8), 15399. <https://doi.org/10.1038/ncomms15399>
- De Kort, H., & Honnay, O. (2017). Assessing evolutionary potential in tree species through ecology-informed genome screening. In P. Pontarotti (Ed.), *Evolutionary biology: Self/nonsel evolution, species and complex traits evolution, methods and concepts*, (313–327). Cham, Switzerland: Springer.
- De Kort, H., Panis, B., Helsen, K., Douzet, R., Janssens, S. B., & Honnay, O. (2020). Pre-adaptation to climate change through topography-driven phenotypic plasticity. *Journal of Ecology*, 108(4), 1465–1474. <https://doi.org/10.1111/1365-2745.13365>
- De Kort, H., Bagueette, M., Prunier, J. G., Tessier, M., Monsimet, J., Turlure, C., & Stevens, V. (2018). Genetic costructure in a meta-community under threat of habitat fragmentation. *Molecular Ecology*, 27(9), 2193–2203. <https://doi.org/10.1111/mec.14569>
- De Kort, H., Prunier, J. G., Tessier, M., Turlure, C., Bagueette, M., & Stevens, V. (2018). Interacting grassland species under threat of multiple global change drivers. *Journal of Biogeography*, 45(9), 2133–2145. <https://doi.org/10.1111/jbi.13397>
- Devictor, V., van Swaay, C., Brereton, T., Brotons, L., Chamberlain, D., Heliölä, J., ... Jiguet, F. (2012). Differences in the climatic debts of birds and butterflies at a continental scale. *Nature Climate Change*, 2, 121–124. <https://doi.org/10.1038/nclimate1347>
- Diniz-Filho, J. A., Barbosa, A. C., Collevatti, R. G., Chaves, L. J., Terribile, L. C., Lima-Ribeiro, M. S., & Telles, M. P. (2016). Spatial autocorrelation analysis and ecological niche modeling allows inference of range dynamics driving the population genetic structure of a Neotropical savanna tree. *Journal of Biogeography*, 43, 167–177.
- Domisch, S., Araújo, M. B., Bonada, N., Pauls, S. U., Jähnig, S. C., & Haase, P. (2013). Modelling distribution in European stream macroinvertebrates under future climates. *Global Change Biology*, 19, 752–762. <https://doi.org/10.1111/gcb.12107>
- Dullinger, I., Gatttringer, A., Wessely, J., Moser, D., Plutzer, C., Willner, W., ... Dullinger, S. (2020). A socio-ecological model for predicting impacts of land-use and climate change on regional plant diversity in the Austrian Alps. *Global Change Biology*, 26(4), 2336–2352. <https://doi.org/10.1111/gcb.14977>
- Duputie, A., & Massol, F. (2013). An empiricist's guide to theoretical predictions on the evolution of dispersal. *Interface Focus*, 3, 20130028. <https://doi.org/10.1098/rsfs.2013.0028>
- Elith, J., & Leathwick, J. R. (2009). Species distribution models: Ecological explanation and prediction across space and time. *Annual Review of Ecology Evolution and Systematics*, 40, 677–697. <https://doi.org/10.1146/annurev.ecolsys.110308.120159>
- Erichsen, E. O., Budde, K. B., Sagheb-Talebi, K. B., Bagnoli, F., Vendramin, G. G., & Hansen, O. K. (2018). Hyrcanian forests-Stable rear-edge populations harbouring high genetic diversity of *Fraxinus excelsior*, a common European tree species. *Diversity and Distributions*, 24, 1521–1533.
- Exposito-Alonso, M., Vasseur, F., Ding, W., Wang, G., Burbano, H. A., & Weigel, D. (2018). Genomic basis and evolutionary potential for extreme drought adaptation in *Arabidopsis thaliana*. *Nature Ecology and Evolution*, 2, 352–358.
- Fitzpatrick, M. C., Gove, A. D., Sanders, N. J., & Dunn, R. R. (2008). Climate change, plant migration, and range collapse in a global biodiversity hotspot: The Banksia (Proteaceae) of Western Australia. *Global Change Biology*, 14, 1337–1352.
- Fitzpatrick, M. C., & Keller, S. R. (2015). Ecological genomics meets community-level modelling of biodiversity: Mapping the genomic landscape of current and future environmental adaptation. *Ecology Letters*, 18, 1–16. <https://doi.org/10.1111/ele.12376>
- Fordham, D. A., Bertelsmeier, C., Brook, B. W., Early, R., Neto, D., Brown, S. C., ... Araújo, M. B. (2018). How complex should models be? Comparing correlative and mechanistic range dynamics models. *Global Change Biology*, 24, 1357–1370. <https://doi.org/10.1111/gcb.13935>
- Fourcade, Y., Besnard, A. G., & Secondi, J. (2018). Paintings predict the distribution of species, or the challenge of selecting environmental predictors and evaluation statistics. *Global Ecology and Biogeography*, 27, 245–256. <https://doi.org/10.1111/geb.12684>
- Fournier, A., Barbet-Massin, M., Rome, Q., & Courchamp, F. (2017). Predicting species distribution combining multi-scale drivers. *Global Ecology and Conservation*, 12, 215–226. <https://doi.org/10.1016/j.gecco.2017.11.002>
- Frankham, R., Ballou, S. E., Briscoe, D. A., & Ballou, J. D. (2002). *Introduction to conservation genetics*. Cambridge, UK: Cambridge University Press.
- Franklin, J. (2013). Species distribution models in conservation biogeography: Developments and challenges. *Diversity and Distributions*, 19, 1217–1223. <https://doi.org/10.1111/ddi.12125>
- Franks, S. J., Sim, S., & Weis, A. E. (2007). Rapid evolution of flowering time by an annual plant in response to a climate fluctuation. *Proceedings of the National Academy of Sciences USA*, 104, 1278–1282. <https://doi.org/10.1073/pnas.0608379104>
- Frichot, E., Schoville, S. D., Bouchard, G., & François, O. (2013). Testing for associations between loci and environmental gradients using latent factor mixed models. *Molecular Biology and Evolution*, 30, 1687–1699. <https://doi.org/10.1093/molbev/mst063>
- García, R. A., Cabeza, M., Rahbek, C., & Araújo, M. B. (2014). Multiple dimensions of climate change and their implications for biodiversity. *Science*, 344, 1247579. <https://doi.org/10.1126/science.1247579>

- García Molinos, J., Halpern, B. S., Schoeman, D. S., Brown, C. J., Kiessling, W., Moore, P. J., ... Burrows, M. T. (2015). Climate velocity and the future global redistribution of marine biodiversity. *Nature Climate Change*, 6, 83–88. <https://doi.org/10.1038/nclimate2769>
- González-Varo, J. P., Biesmeijer, J. C., Bommarco, R., Potts, S. G., Schweiger, O., Smith, H. G., ... Vilà, M. (2013). Combined effects of global change pressures on animal-mediated pollination. *Trends in Ecology & Evolution*, 28, 524–530. <https://doi.org/10.1016/j.tree.2013.05.008>
- Gotelli, N. J., & Stanton-Geddes, J. (2015). Climate change, genetic markers and species distribution modelling. *Journal of Biogeography*, 42, 1577–1585.
- Graae, B. J., Vandvik, V., Armbruster, W. S., Eiserhardt, W. L., Svenning, J. C., Hylander, K., ... Milbau, A. (2017). Stay or go – how topographic complexity influences alpine plant population and community responses to climate change. *Perspectives in Plant Ecology, Evolution and Systematics*, 30, 41–50.
- Gregg, E. J., Palacios, D. M., Thompson, A., & Chan, K. M. A. (2019). Why less complexity produces better forecasts: An independent data evaluation of kelp habitat models. *Ecography*, 42, 428–443. <https://doi.org/10.1111/ecog.03470>
- Guillera-Aroita, G. (2017). Modelling of species distributions, range dynamics and communities under imperfect detection: Advances, challenges and opportunities. *Ecography*, 40, 281–295.
- Guisan, A., & Rahbek, C. (2011). SESAM - a new framework integrating macroecological and species distribution models for predicting spatio-temporal patterns of species assemblages. *Journal of Biogeography*, 38, 1433–1444.
- Guillera-Aroita, G., Lahoz-Monfort, J. J., Elith, J., Gordon, A., Kujala, H., Lentini, P. E., ... Wintle, B. A. (2015). Is my species distribution model fit for purpose? Matching data and models to applications. *Global Ecology and Biogeography*, 24, 276–292.
- Guo, F., Lenoir, J., & Bonebrake, T. C. (2018). Land-use change interacts with climate to determine elevational species redistribution. *Nature Communications*, 9, 1315.
- Gutiérrez, D., Fernández, P., Seymour, A. S., & Jordano, D. (2005). Habitat distribution models: Are mutualist distributions good predictors of their associates? *Ecological Applications*, 15, 3–18. <https://doi.org/10.1890/03-5344>
- Gutiérrez-Rodríguez, J., Barbosa, A. M., & Martínez-Solano, I. (2017). Integrative inference of population history in the Ibero-Maghrebian endemic *Pleurodeles waltl* (Salamandridae). *Molecular Phylogenetics and Evolution*, 112, 122–137.
- Hanley, J. A., & McNeil, B. J. (1982). The meaning and use of the area under a receiver operating characteristic (ROC) curve. *Radiology*, 143, 29–36. <https://doi.org/10.1148/radiology.143.1.7063747>
- Hansen, M. C., Potapov, P. V., Moore, R., Hancher, M., Turubanova, S. A., Tyukavina, A., ... Townshend, J. R. G. (2013). High-resolution global maps of 21st-century forest cover change. *Science*, 342, 850–853. <https://doi.org/10.1126/science.1244693>
- Hanspach, J., Schweiger, O., Kühn, I., Plattner, M., Pearman, P. B., Zimmermann, N. E., & Settele, J. (2014). Host plant availability potentially limits butterfly distributions under cold environmental conditions. *Ecography*, 37, 301–308. <https://doi.org/10.1111/j.1600-0587.2013.00195.x>
- Harrison, R. D., & Gassner, A. (2020). Agricultural lands key to mitigation and adaptation. *Science*, 367, 518.
- Hattab, T., Garzón-López, C. X., Ewald, M., Skowronek, S., Aerts, R., Horen, H., ... Lenoir, J. (2017). A unified framework to model the potential and realized distributions of invasive species within the invaded range. *Diversity and Distributions*, 23, 806–819. <https://doi.org/10.1111/ddi.12566>
- Heino, M., & Hanski, I. (2001). Evolution of migration rate in a spatially realistic metapopulation model. *American Naturalist*, 157, 495–511. <https://doi.org/10.1086/319927>
- Hijmans, R. J. (2012). Cross-validation of species distribution models: Removing spatial sorting bias and calibration with a null model. *Ecology*, 93, 679–688. <https://doi.org/10.1890/11-0826.1>
- Hirzel, A. H., & Le Lay, G. (2008). Habitat suitability modelling and niche theory. *Journal of Applied Ecology*, 45, 1372–1381. <https://doi.org/10.1111/j.1365-2664.2008.01524.x>
- Hof, A. R., Jansson, R., & Nilsson, C. (2012). How biotic interactions may alter future predictions of species distributions: Future threats to the persistence of the arctic fox in Fennoscandia. *Diversity and Distributions*, 18, 554–562. <https://doi.org/10.1111/j.1472-4642.2011.00876.x>
- Hoffmann, A. A., Sgrò, C. M., & Kristensen, T. N. (2017). Revisiting adaptive potential, population size, and conservation. *Trends in Ecology & Evolution*, 32, 506–517.
- Holloway, P., & Miller, J. A. (2017). A quantitative synthesis of the movement concepts used within species distribution modelling. *Ecological Modelling*, 356, 91–103. <https://doi.org/10.1016/j.ecolmodel.2017.04.005>
- Hulme, P. E. (2017). Climate change and biological invasions: Evidence, expectations, and response options. *Biological Reviews*, 92, 1297–1313.
- Ikeda, D. H., Max, T. L., Allan, G. J., Lau, M. K., Shuster, S. M., & Whitham, T. G. (2017). Genetically informed ecological niche models improve climate change predictions. *Global Change Biology*, 23, 164–176. <https://doi.org/10.1111/gcb.13470>
- Inoue, K., & Berg, D. J. (2017). Predicting the effects of climate change on population connectivity and genetic diversity of an imperiled freshwater mussel, *Cumberlandia monodonta* (Bivalvia: Margaritiferidae), in riverine systems. *Global Change Biology*, 23, 94–107.
- Jetz, W., Wilcove, D. S., & Dobson, A. P. (2007). Projected impacts of climate and land-use change on the global diversity of birds (GM Mace, Ed.). *PLoS Biology*, 5, e157.
- Jones, K. E., Bielby, J., Cardillo, M., Fritz, S. A., O'Dell, J., Orme, C. D., ... Connolly, C. (2009). PanTHERIA: A species-level database of life history, ecology, and geography of extant and recently extinct mammals. *Ecology*, 90, 2648.
- Kane, K., Debinski, D. M., Anderson, C., Scasta, J. D., Engle, D. M., & Miller, J. R. (2017). Using regional climate projections to guide grassland community restoration in the face of climate change. *Frontiers in Plant Science*, 8, 730. <https://doi.org/10.3389/fpls.2017.00730>
- Karell, P., Ahola, K., Karstinen, T., Valkama, J., & Brommer, J. E. (2011). Climate change drives microevolution in a wild bird. *Nature Communications*, 2, 207–208. <https://doi.org/10.1038/ncomms1213>
- Kearney, M., Porter, W. P., Williams, C., Ritchie, S., & Hoffmann, A. A. (2009). Integrating biophysical models and evolutionary theory to predict climatic impacts on species' ranges: The dengue mosquito *Aedes aegypti* in Australia. *Functional Ecology*, 23, 528–538.
- Kelly, A. E., & Goulden, M. L. (2008). Rapid shifts in plant distribution with recent climate change. *Proceedings of the National Academy of Sciences USA*, 105, 11823–11826. <https://doi.org/10.1073/pnas.0802891105>
- Kelly, M. W., DeBiasse, M. B., Villela, V. A., Roberts, H. L., & Cecola, C. F. (2016). Adaptation to climate change: Trade-offs among responses to multiple stressors in an intertidal crustacean. *Evolutionary Applications*, 9, 1147–1155. <https://doi.org/10.1111/eva.12394>
- Kok, M. T. J., Kok, K., Peterson, G. D., Hill, R., Agard, J., & Carpenter, S. R. (2017). Biodiversity and ecosystem services require IPBES to take novel approach to scenarios. *Sustainability Science*, 12, 177–181. <https://doi.org/10.1007/s11625-016-0354-8>
- Kremen, C., Cameron, A., Moilanen, A., Phillips, S. J., Thomas, C. D., Beentje, H., ... Zjhra, M. L. (2008). Aligning conservation priorities across taxa in Madagascar with high-resolution planning tools. *Science*, 320, 222–226. <https://doi.org/10.1126/science.1155193>
- La Sorte, F. A., Fink, D., Blancher, P. J., Rodewald, A. D., Ruiz-Gutierrez, V., Rosenberg, K. V., ... Kelling, S. (2017). Global change and the distributional dynamics of migratory bird populations wintering in Central America. *Global Change Biology*, 23, 5284–5296. <https://doi.org/10.1111/gcb.13794>

- Lehsten, V., Sykes, M. T., Scott, A. V., Tzanopoulos, J., Kallimanis, A., Mazaris, A., ... Vogiatzakis, I. (2015). Disentangling the effects of land-use change, climate and CO<sub>2</sub> on projected future European habitat types. *Global Ecology and Biogeography*, *24*, 653–663.
- Lembrechts, J. J., Nijs, I., & Lenoir, J. (2018). Incorporating microclimate into species distribution models. *Ecography*, *42*, 1267–1279.
- Lenoir, J., & Svenning, J. C. (2013). Latitudinal and elevational range shifts under contemporary climate change. *Encyclopedia of Biodiversity*, (Vol. 4, pp. 599–611). San Diego, CA: Academic Press.
- Leroy, B., Delsol, R., Hugué, B., Meynard, C. N., Barhoumi, C., Barbet-Massin, M., & Bellard, C. (2018). Without quality presence-absence data, discrimination metrics such as TSS can be misleading measures of model performance. *Journal of Biogeography*, *45*, 1994–2002.
- Liang, Y. U., Duveneck, M. J., Gustafson, E. J., Serra-Diaz, J. M., & Thompson, J. R. (2017). How disturbance, competition, and dispersal interact to prevent tree range boundaries from keeping pace with climate change. *Global Change Biology*, *24*(1), e335–e351. <https://doi.org/10.1111/gcb.13847>
- Lobo, J. (2016). The use of occurrence data to predict the effects of climate change on insects. *Current Opinion in Insect Science*, *17*, 62–68. <https://doi.org/10.1016/j.cois.2016.07.003>
- Lobo, J. M., Jiménez-Valverde, A., & Hortal, J. (2010). The uncertain nature of absences and their importance in species distribution modelling. *Ecography*, *33*, 103–114. <https://doi.org/10.1111/j.1600-0587.2009.06039.x>
- Lobo, J. M., Jiménez-Valverde, A., & Real, R. (2008). AUC: A misleading measure of the performance of predictive distribution models. *Global Ecology and Biogeography*, *17*, 145–151. <https://doi.org/10.1111/j.1466-8238.2007.00358.x>
- MacLean, S. A., & Beissinger, S. R. (2017). Species' traits as predictors of range shifts under contemporary climate change: A review and meta-analysis. *Global Change Biology*, *23*, 4094–4105. <https://doi.org/10.1111/gcb.13736>
- Manel, S., & Holderegger, R. (2013). Ten years of landscape genetics. *Trends in Ecology & Evolution*, *28*, 614–621. <https://doi.org/10.1016/j.tree.2013.05.012>
- Manel, S., Schwartz, M. K., Luikart, G., & Taberlet, P. (2003). Landscape genetics: Combining landscape ecology and population genetics. *Trends in Ecology & Evolution*, *18*, 189–197. [https://doi.org/10.1016/S0169-5347\(03\)00008-9](https://doi.org/10.1016/S0169-5347(03)00008-9)
- Marcer, A., Méndez-Vigo, B., Alonso-Blanco, C., & Picó, F. X. (2016). Tackling intraspecific genetic structure in distribution models better reflects species geographical range. *Ecology and Evolution*, *6*, 2084–2097. <https://doi.org/10.1002/ece3.2010>
- Markert, J. A., Champlin, D. M., Gutjahr-Gobell, R., Grear, J. S., Kuhn, A., McGreevy, T. J., ... Nacci, D. E. (2010). Population genetic diversity and fitness in multiple environments. *BMC Evolutionary Biology*, *10*, 205. <https://doi.org/10.1186/1471-2148-10-205>
- Marshall, L., Biesmeijer, J. C., Rasmont, P., Vereecken, N. J., Dvorak, L., Fitzpatrick, U., ... Dendoncker, N. (2018). The interplay of climate and land use change affects the distribution of EU bumblebees. *Global Change Biology*, *24*, 101–116. <https://doi.org/10.1111/gcb.13867>
- Martin, Y., Van Dyck, H., Dendoncker, N., & Titeux, N. (2013). Testing instead of assuming the importance of land use change scenarios to model species distributions under climate change. *Global Ecology and Biogeography*, *22*, 1204–1216. <https://doi.org/10.1111/geb.12087>
- Mathias, A., Kisi, E. A., & Olivieri, I. (2001). Divergent evolution of dispersal in a heterogeneous landscape. *BioOne*, *55*, 246–259. <https://doi.org/10.1111/j.0014-3820.2001.tb01290.x>
- McClanahan, T. R., Graham, N. A. J., & Darling, E. S. (2014). Coral reefs in a crystal ball: Predicting the future from the vulnerability of corals and reef fishes to multiple stressors. *Current Opinion in Environmental Sustainability*, *7*, 59–64. <https://doi.org/10.1016/j.cosust.2013.11.028>
- Meentemeyer, R. K., Anacker, B. L., Mark, W., & Rizzo, D. M. (2008). Early detection of emerging forest disease using dispersal estimation and ecological niche modeling. *Ecological Applications*, *18*, 377–390. <https://doi.org/10.1890/07-1150.1>
- Meier, E. S., Lischke, H., Schmatz, D. R., & Zimmermann, N. E. (2012). Climate, competition and connectivity affect future migration and ranges of European trees. *Global Ecology and Biogeography*, *21*, 164–178.
- Merow, C., Wilson, A. M., & Jetz, W. (2017). Integrating occurrence data and expert maps for improved species range predictions. *Global Ecology and Biogeography*, *26*, 243–258. <https://doi.org/10.1111/geb.12539>
- Mestre, F., Risk, B. B., Mira, A., Beja, P., & Pita, R. (2017). A metapopulation approach to predict species range shifts under different climate change and landscape connectivity scenarios. *Ecological Modelling*, *359*, 406–414. <https://doi.org/10.1016/j.ecolmodel.2017.06.013>
- Midgley, G. F., Hughes, G. O., Thuiller, W., & Rebelo, A. G. (2006). Migration rate limitations on climate change-induced range shifts in Cape Proteaceae. *Diversity and Distributions*, *12*, 555–562. <https://doi.org/10.1111/j.1366-9516.2006.00273.x>
- Milanesi, P., Rocca, F. D., & Robinson, R. A. (2020). Integrating dynamic environmental predictors and species occurrences: Toward true dynamic species distribution models. *Ecology and Evolution*, *10*, 1087–1092.
- Morán-Ordóñez, A., Lahoz-Monfort, J. J., Elith, J., & Wintle, B. A. (2017). Evaluating 318 continental-scale species distribution models over a 60-year prediction horizon: What factors influence the reliability of predictions? *Global Ecology and Biogeography*, *26*, 371–384. <https://doi.org/10.1111/geb.12545>
- Muscarella, R., Galante, P. J., Soley-Guardia, M., & Boria M. A. (2014). ENMeval: An R package for conducting spatially independent evaluations and estimating optimal model complexity for Maxent ecological niche models. *Methods in Ecology and Evolution*, *5*, 1198–1205.
- Naudts, K., Chen, Y., McGrath, M. J., Ryder, J., Valade, A., Otto, J., & Luysaert, S. (2016). Europe's forest management did not mitigate climate warming. *Science*, *351*, 597–600. <https://doi.org/10.1126/science.aad7270>
- Newbold, T., Hudson, L. N., Hill, S. L., Contu, S., Lysenko, I., Senior, R. A., ... Day, J. (2015). Global effects of land use on local terrestrial biodiversity. *Nature*, *520*, 45–50.
- Parmesan, C., Ryrholm, N., Stefanescu, C., Hill, J. K., Thomas, C. D., Descimon, H., ... Warren, M. (1999). Poleward shifts in geographical ranges of butterfly species associated with regional warming. *Nature*, *399*, 579–583. <https://doi.org/10.1038/21181>
- Pearson, R. G., Phillips, S. J., Lorant, M. M., Beck, P. S. A., Damoulas, T., Knight, S. J., & Goetz, S. J. (2013). Shifts in Arctic vegetation and associated feedbacks under climate change. *Nature Climate Change*, *3*, 673–677. <https://doi.org/10.1038/nclimate1858>
- Pennino, M. G., Paradinas, I., Illian, J. B., Muñoz, F., Bellido, J. M., López-Quílez, A., & Conesa, D. (2019). Accounting for preferential sampling in species distribution models. *Ecology and Evolution*, *9*, 653–663. <https://doi.org/10.1002/ece3.4789>
- Pereira, H. M., Leadley, P. W., Proença, V., Alkemade, R., Scharlemann, J. P. W., Fernandez-Manjarrés, J. F., ... Walpole, M. (2010). Scenarios for global biodiversity in the 21st century. *Science*, *330*, 1496–1501. <https://doi.org/10.1126/science.1196624>
- Perring, M. P., De Frenne, P., Baeten, L., Maes, S. L., Depauw, L., Blondeel, H., ... Verheyen, K. (2016). Global environmental change effects on ecosystems: The importance of land-use legacies. *Global Change Biology*, *22*, 1361–1371. <https://doi.org/10.1111/gcb.13146>
- Peterson, T. A., Papeş, M., & Eaton, M. (2007). Transferability and model evaluation in ecological niche modeling: A comparison of GARP and Maxent. *Ecography*, *30*, 550–560. <https://doi.org/10.1111/j.0906-7590.2007.05102.x>
- Peterson, A. T. (2003). Predicting the geography of species' invasions via ecological niche modeling. *The Quarterly Review of Biology*, *78*, 419–433. <https://doi.org/10.1086/378926>

- Peterson, A. T., & Nakazawa, Y. (2007). Environmental data sets matter in ecological niche modelling: An example with *Solenopsis invicta* and *Solenopsis richteri*. *Global Ecology and Biogeography*, 17(1), 135–144. <https://doi.org/10.1111/j.1466-8238.2007.00347.x>
- Peterson, A. T., Papeš, M., & Soberón, J. (2015). Mechanistic and correlative models of ecological niches. *European Journal of Ecology*, 1, 28–38. <https://doi.org/10.1515/eje-2015-0014>
- Peterson, M. L., Doak, D. F., & Morris, W. F. (2019). Incorporating local adaptation into forecasts of species' distribution and abundance under climate change. *Global Change Biology*, 25(3), 775–793. <https://doi.org/10.1111/gcb.14562>
- Peyre, G., Lenoir, J., Karger, D. N., Gomez, M., Gonzalez, A., Broennimann, O., & Guisan, A. (2020). The fate of páramo plant assemblages in the sky islands of the northern Andes. *Journal of Vegetation Science*. Accepted. <https://doi.org/10.1111/jvs.12898>
- Phillips, S. J., & Dudík, M. (2008). Modeling of species distributions with Maxent: New extensions and a comprehensive evaluation. *Ecography*, 31, 161–175. <https://doi.org/10.1111/j.0906-7590.2008.5203.x>
- Pulliam, H. R. (2000). On the relationship between niche and distribution. *Ecology Letters*, 3, 349–361. <https://doi.org/10.1046/j.1461-0248.2000.00143.x>
- Radosavljevic, A., & Anderson, R. P. (2014). Making better Maxent models of species distributions: Complexity, overfitting and evaluation. *Journal of Biogeography*, 41, 629–643.
- Razgour, O., Taggart, J. B., Manel, S., Juste, J., Ibáñez, C., Rebelo, H., ... Park, K. (2018). An integrated framework to identify wildlife populations under threat from climate change. *Molecular Ecology Resources*, 18, 18–31. <https://doi.org/10.1111/1755-0998.12694>
- Reino, L., Beja, P., Araújo, M. B., Dray, S., & Segurado, P. (2013). Does local habitat fragmentation affect large-scale distributions? The case of a specialist grassland bird. *Diversity and Distributions*, 19, 423–432. <https://doi.org/10.1111/ddi.12019>
- Reisinger, A., & Clark, H. (2017). How much do direct livestock emissions actually contribute to global warming? *Global Change Biology*, 24(4), 1749–1761. <https://doi.org/10.1111/gcb.13975>
- Rellstab, C., Gugerli, F., Eckert, A. J., Hancock, A. M., & Holderegger, R. (2015). A practical guide to environmental association analysis in landscape genomics. *Molecular Ecology*, 24, 4348–4370. <https://doi.org/10.1111/mec.13322>
- Rodrigues, A. S. L., Andelman, S. J., Bakarr, M. I., Boitani, L., Brooks, T. M., Cowling, R. M., ... Yan, X. (2004). Effectiveness of the global protected area network in representing species diversity. *Nature*, 428, 640–643. <https://doi.org/10.1038/nature02422>
- Rolland, J., Silvestro, D., Schluter, D., Guisan, A., Broennimann, O., & Salamin, N. (2018). The impact of endothermy on the climatic niche evolution and the distribution of vertebrate diversity. *Nature Ecology & Evolution*, 2(3), 459–464.
- Ronce, O. (2007). How does it feel to be like a rolling stone? ten questions about dispersal evolution. *Annual Review of Ecology and Systematics*, 38, 231–253. <https://doi.org/10.1146/annurev.ecolsys.38.091206.095611>
- Scheller, R. M., & Mladenoff, D. J. (2005). A spatially interactive simulation of climate change, harvesting, wind, and tree species migration and projected changes to forest composition and biomass in northern Wisconsin, USA. *Global Change Biology*, 11, 307–321. <https://doi.org/10.1111/j.1365-2486.2005.00906.x>
- Schloss, C. A., Nunez, T. A., & Lawler, J. J. (2012). Dispersal will limit ability of mammals to track climate change in the Western Hemisphere. *Proceedings of the National Academy of Sciences USA*, 109, 8606–8611. <https://doi.org/10.1073/pnas.1116791109>
- Schrieber, K., & Lachmuth, S. (2017). The Genetic Paradox of Invasions revisited: The potential role of inbreeding × environment interactions in invasion success. *Biological Reviews*, 92, 939–952. <https://doi.org/10.1111/brv.12263>
- Schweiger, O., Settele, J., Kudrna, O., Klotz, S., & Kühn, I. (2008). Climate change can cause spatial mismatch of trophically interacting species. *Ecology*, 89, 3472–3479. <https://doi.org/10.1890/07-1748.1>
- Segan, D. B., Murray, K. A., & Watson, J. E. M. (2016). A global assessment of current and future biodiversity vulnerability to habitat loss–climate change interactions. *Global Ecology and Conservation*, 5, 12–21. <https://doi.org/10.1016/j.gecco.2015.11.002>
- Sheldon, K. S., Yang, S., & Tewksbury, J. J. (2011). Climate change and community disassembly: Impacts of warming on tropical and temperate montane community structure. *Ecology Letters*, 14, 1191–1200. <https://doi.org/10.1111/j.1461-0248.2011.01689.x>
- Slatkin, M. (1987). Gene flow and the geographic structure of natural populations. *Science*, 236, 787–792. <https://doi.org/10.1126/science.3576198>
- Sork, V. L., Squire, K., Gugger, P. F., Steele, S. E., Levy, E. D., & Eckert, A. J. (2016). Landscape genomic analysis of candidate genes for climate adaptation in a California endemic oak, *Quercus lobata*. *American Journal of Botany*, 103, 33–46.
- Storfer, A., Murphy, M. A., Evans, J. S., Goldberg, C. S., Robinson, S., Spear, S. F., ... Waits, L. P. (2007). Putting the 'landscape' in landscape genetics. *Heredity*, 98, 128–142. <https://doi.org/10.1038/sj.hdy.6800917>
- Struebig, M. J., Fischer, M., Gaveau, D. L. A., Meijaard, E., Wich, S. A., Gonner, C., ... Kramer-Schadt, S. (2015). Anticipated climate and land-cover changes reveal refuge areas for Borneo's orangutans. *Glob. Chang. Biol.*, 21, 2891–2904. <https://doi.org/10.1111/gcb.12814>
- Suggitt, A. J., Wilson, R. J., Isaac, N. J. B., Beale, C. M., Auffret, A. G., August, T., ... Maclean, I. M. D. (2020). Extinction risk from climate change is reduced by microclimatic buffering. *Nature Climate Change*, 8, 713–717. <https://doi.org/10.1038/s41558-018-0231-9>
- Sunday, J. M., Bates, A. E., & Dulvy, N. K. (2012). Thermal tolerance and the global redistribution of animals. *Nature Climate Change*, 2, 686–690.
- Stevens, V. M., Trochet, A., Blanchet, S., Moulherat, S., Clobert, J., & Baguette, M. (2013). Dispersal syndromes and the use of life-histories to predict dispersal. *Evolutionary Applications*, 6, 630–642.
- Tamme, R., Götzenberger, L., Zobel, M., Bullock, J. M., Hooftman, D. A. P., Kaasik, A., & Pärtel, M. (2014). Predicting species' maximum dispersal distances from simple plant traits. *Ecology*, 95, 505–513. <https://doi.org/10.1890/13-1000.1>
- Taylor, P. D., Fahrig, L., Henein, K., & Merriam, G. (1993). Connectivity is a vital element of landscape structure. *Oikos*, 68, 571. <https://doi.org/10.2307/3544927>
- Thomson, F. J., Letten, A. D., Tamme, R., Edwards, W., & Moles, A. T. (2018). Can dispersal investment explain why tall plant species achieve longer dispersal distances than short plant species? *New Phytologist*, 217, 407–415. <https://doi.org/10.1111/nph.14735>
- Titeux, N., Henle, K., Mihoub, J.-B., & Brotons, L. (2016). Climate change distracts us from other threats to biodiversity. *Frontiers in Ecology and the Environment*, 14, 291. <https://doi.org/10.1002/fee.1303>
- Travis, J. M. J., Delgado, M., Bocedi, G., Baguette, M., Bartoň, K., Bonte, D., ... Bullock, J. M. (2013). Dispersal and species' responses to climate change. *Oikos*, 122, 1532–1540. <https://doi.org/10.1111/j.1600-0706.2013.00399.x>
- Travis, J. M. J., & Dytham, C. (1999). Habitat persistence, habitat availability and the evolution of dispersal. *Proceedings of the Royal Society B-Biological Sciences*, 266, 723–728.
- Tuff, K. T., Tuff, T., & Davies, K. F. (2016). A framework for integrating thermal biology into fragmentation research. *Ecology Letters*, 19, 361–374. <https://doi.org/10.1111/ele.12579>
- Václavík, T., & Meentemeyer, R. K. (2009). Invasive species distribution modeling (iSDM): Are absence data and dispersal constraints needed to predict actual distributions? *Ecological Modelling*, 220, 3248–3258. <https://doi.org/10.1016/j.ecolmodel.2009.08.013>
- van Asch, M., Salis, L., Holleman, L. J. M., van Lith, B., & Visser, M. E. (2013). Evolutionary response of the egg hatching date of a



- herbivorous insect under climate change. *Nature Climate Change*, 3, 244–248. <https://doi.org/10.1038/nclimate1717>
- van Proosdij, A. S. J., Sosef, M. S. M., Wieringa, J. J., & Raes, N. (2016). Minimum required number of specimen records to develop accurate species distribution models. *Ecography*, 39, 542–552. <https://doi.org/10.1111/ecog.01509>
- VanDerWal, J., Shoo, L. P., Graham, C., & Williams, S. E. (2009). Selecting pseudo-absence data for presence-only distribution modeling: How far should you stray from what you know? *Ecological Modelling*, 220, 589–594. <https://doi.org/10.1016/j.ecolmodel.2008.11.010>
- Vanneste, T., Govaert, S., Spicher, F., Brunet, J., Cousins, S. A. O., Decocq, G., ... De Frenne, P. (2020). Contrasting microclimates among hedgerows and woodlands across temperate Europe. *Agricultural and Forest Meteorology*, 281, 107818. <https://doi.org/10.1016/j.agrfor.2019.107818>
- Visconti, P., Bakkenes, M., Baisero, D., Brooks, T., Butchart, S. H. M., Joppa, L., ... Rondinini, C. (2016). Projecting global biodiversity indicators under future development scenarios. *Conservation Letters*, 9, 5–13. <https://doi.org/10.1111/conl.12159>
- Visser, M. E. (2008). Keeping up with a warming world; assessing the rate of adaptation to climate change. *Proceedings of the Royal Society B-Biological Sciences*, 275, 649–659. <https://doi.org/10.1098/rspb.2007.0997>
- Vitousek, P. M. (1994). Beyond global warming: Ecology and global change. *Ecology*, 75, 1861–1876. <https://doi.org/10.2307/1941591>
- Walther, G.-R., Post, E., Convey, P., Menzel, A., Parmesan, C., Beebee, T. J. C., ... Bairlein, F. (2002). Ecological responses to recent climate change. *Nature*, 416, 389–395. <https://doi.org/10.1038/416389a>
- Warren, D. L., Matzke, N. J., & Iglésias, T. L. (2020). Evaluating presence-only species distribution models with discrimination accuracy is uninformative for many applications. *Journal of Biogeography*, 47, 167–180. <https://doi.org/10.1111/jbi.13705>
- Warren, D. L., Wright, A. N., Seifert, S. N., & Shaffer, H. B. (2014). Incorporating model complexity and spatial sampling bias into ecological niche models of climate change risks faced by 90 California vertebrate species of concern. *Diversity and Distributions*, 20, 334–343. <https://doi.org/10.1111/ddi.12160>
- Warren, M. S., Hill, J. K., Thomas, J. A., Asher, J., Fox, R., Huntley, B., ... Thomas, C. D. (1999). Rapid responses of British butterflies to opposing forces of climate and habitat change. *Nature*, 414, 65–69. <https://doi.org/10.1038/35102054>
- Warren, R. J., & Bradford, M. A. (2014). Mutualism fails when climate response differs between interacting species. *Global Change Biology*, 20, 466–474. <https://doi.org/10.1111/gcb.12407>
- Watanabe, K., & Monaghan, M. T. (2017). Comparative tests of the species-genetic diversity correlation at neutral and nonneutral loci in four species of stream insect. *Evolution*, 71, 1755–1764. <https://doi.org/10.1111/evo.13261>
- Weber, M. M., Stevens, R. D., Diniz-Filho, J. A. F., & Grelle, C. E. V. (2017). Is there a correlation between abundance and environmental suitability derived from ecological niche modelling? A meta-analysis. *Ecography*, 40, 817–828. <https://doi.org/10.1111/ecog.02125>
- West, A. M., Kumar, S., Brown, C. S., Stohlgren, T. J., & Bromberg, J. (2016). Field validation of an invasive species Maxent model. *Ecological Informatics*, 36, 126–134. <https://doi.org/10.1016/j.ecoinf.2016.11.001>
- Wiens, J. A., Stralberg, D., Jongsomjit, D., Howell, C. A., & Snyder, M. A. (2009). Niches, models, and climate change: Assessing the assumptions and uncertainties. *Proceedings of the National Academy of Sciences USA*, 106(Suppl 2), 19729–19736.
- Williams, J. N., Seo, C., Thorne, J., Nelson, J. K., Erwin, S., O'Brien, J. M., & Schwartz, M. W. (2009). Using species distribution models to predict new occurrences for rare plants. *Diversity and Distributions*, 15, 565–576. <https://doi.org/10.1111/j.1472-4642.2009.00567.x>
- Wisz, M. S., Pottier, J., Kissling, W. D., Pellissier, L., Lenoir, J., Damgaard, C. F., ... Svenning, J.-C. (2013). The role of biotic interactions in shaping distributions and realised assemblages of species: implications for species distribution modelling. *Biological Reviews*, 88, 15–30.
- Wood, J. L. A., Yates, M. C., & Fraser, D. J. (2016). Are heritability and selection related to population size in nature? Meta-analysis and conservation implications. *Evolutionary Applications*, 9, 640–657. <https://doi.org/10.1111/eva.12375>
- Zamora-Gutierrez, V., Pearson, R. G., Green, R. E., & Jones, K. E. (2018). Forecasting the combined effects of climate and land use change on Mexican bats. *Diversity and Distributions*, 24, 363–374. <https://doi.org/10.1111/ddi.12686>
- Zhu, K., Woodall, C. W., & Clark, J. S. (2012). Failure to migrate: Lack of tree range expansion in response to climate change. *Global Change Biology*, 18, 1042–1052. <https://doi.org/10.1111/j.1365-2486.2011.02571.x>
- Zwiener, V. P., Padial, A. A., Marques, M. C. M., Faleiro, F. V., Loyola, R., & Peterson, A. T. (2017). Planning for conservation and restoration under climate and land use change in the Brazilian Atlantic Forest. *Diversity and Distributions*, 23, 955–966. <https://doi.org/10.1111/ddi.12588>

## SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

**How to cite this article:** De Kort H, Bague M, Lenoir J, Stevens VM. Toward reliable habitat suitability and accessibility models in an era of multiple environmental stressors. *Ecol Evol*. 2020;00:1–16. <https://doi.org/10.1002/ece3.6753>