



HAL
open science

First estimation of the spontaneous mutation rate in Diatoms

Marc Krasovec, Sophie Sanchez-Brosseau, Gwenael Piganeau

► **To cite this version:**

Marc Krasovec, Sophie Sanchez-Brosseau, Gwenael Piganeau. First estimation of the spontaneous mutation rate in Diatoms. *Genome Biology and Evolution*, In press, 10.1093/gbe/evz130 . hal-02978190v1

HAL Id: hal-02978190

<https://hal.sorbonne-universite.fr/hal-02978190v1>

Submitted on 26 Jun 2019 (v1), last revised 26 Oct 2020 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

First estimation of the spontaneous mutation rate in Diatoms

Marc Krasovec^{1,2*}, Sophie Sanchez-Brosseau¹, Gwenael Piganeau^{1*}

¹ Sorbonne Universités, UPMC Univ Paris 06, CNRS, Biologie Intégrative des Organismes Marins (BIOM), Observatoire Océanologique, F-66650 Banyuls/Mer, France

² Current address: Department of Plant Sciences, University of Oxford, Oxford OX1 3RB, UK.

* Corresponding authors: marc.krasovec@plants.ox.ac.uk ; gwenael.piganeau@obs-banyuls.fr

Keywords: Spontaneous mutation rate, Mutation accumulation, Mutation spectrum, *Phaeodactylum tricornutum*, diatoms.

Running title: Mutation rate in *Phaeodactylum tricornutum*

ABSTRACT

Mutations are the origin of genetic diversity, and the mutation rate is a fundamental parameter to understand all aspects of molecular evolution. The combination of mutation-accumulation experiments and high throughput sequencing enabled the estimation of mutation rates in most model organisms, but several major eukaryotic lineages remain unexplored. Here, we report the first estimation of the spontaneous mutation rate in a model unicellular eukaryote from the Stramenopile kingdom, the diatom *Phaeodactylum tricornutum* (strain RCC2967). We sequenced 36 mutation accumulation lines for an average of 181 generations per line and identified 156 *de novo* mutations. The base substitution mutation rate per site per generation is $\mu_{bs} = 4.77 \times 10^{-10}$ and the insertion-deletion mutation rate is $\mu_{id} = 1.58 \times 10^{-11}$. The mutation rate varies as a function of the nucleotide context and is biased towards an excess of mutations from GC to AT, consistent with previous observations in other species. Interestingly, the mutation rates between the genomes of organelles and the nucleus differ, with a significantly higher mutation rate in the mitochondria. This confirms previous claims based on indirect estimations of the mutation rate in mitochondria of photosynthetic eukaryotes that acquired their plastid through a secondary endosymbiosis. This novel estimate enables us to infer the effective population size of *P. tricornutum* to be $N_e \sim 8.72 \times 10^6$.

INTRODUCTION

The direct estimation of the spontaneous mutation rate (μ) is one of the most exciting possibilities in evolutionary biology since the development of high-throughput “evolve and resequence” experiments (Katju and Bergthorsson 2019). The spontaneous mutation rate determines the frequency of *de novo* mutations introduced into a population, allowing adaptation by selection and the renewal of standing genetic variation. Therefore, estimation of μ in a large number of species is necessary to understand the origin of its variation at different scales. Our current knowledge points to a high variation in the mutation rate between species (Lynch et al. 2016; Katju and Bergthorsson 2019). The drift barrier hypothesis is the most accepted explanation of this variation (Sung et al. 2012; Lynch et al. 2016). Under this hypothesis, the mutation load due to deleterious mutations (Charlesworth and Charlesworth 1998) leads to the selection of the lowest possible mutation rate, and species with large N_e are thus expected to have a lower mutation rate than species with low N_e . Another parameter affecting the mutation rate is the GC nucleotide content. A bias from GC to AT nucleotide mutations has been reported across most branches of the tree of life (Hershberg and Petrov 2010; Ossowski et al. 2010; Denver et al. 2012; Schrider et al. 2013; Krasovec et al. 2017), but the strength of this bias is not equal between species. The difference between the current and the expected GC content reflecting the observed mutation bias between GC and AT nucleotides may explain some of the variation in the mutation rate observed between species (Krasovec et al. 2017) and within a genome (Kiktev et al. 2018). Despite significant advances in understanding of the origin of variations in mutation rates, our knowledge is limited to a few biological models. Within eukaryotes, mutation rate estimates are almost exclusively available for organisms in two kingdoms (Figure 1), the Archaeplastida (plants and green algae) and the Unikont (including fungi and metazoans). Outside these two kingdoms, eukaryotic mutation rates are available in Alveolata: in three species from *Paramecium* (*P. tetraurelia*, *P. sexaurelia*, *P. biaurelia* (Long et al. 2018)), in *Tetrahymena thermophila* (Long et al. 2016) and in *Plasmodium falciparum*

(Hamilton et al. 2017). In *Paramecium*, the mutation rate is extremely low; one to two orders of magnitude lower than in other unicellular species (Figure 1).

To broaden our knowledge on mutation rate over a wider phylogenetic range, we performed a mutation accumulation experiment (MA) with the model diatom *Phaeodactylum tricornutum* (strain RCC2967 from the Roscoff Culture Collection (Vaulot et al. 2004)), belonging to the Stramenopile (or Heterokonta) eukaryotic kingdom. Estimates of the mutation rate of ecological key species such as phytoplankton are scarce and are only available for coastal green algae species (Krasovec et al. 2017; Krasovec et al. 2018), notably in the model species *Ostreococcus tauri*. Diatoms (Bacillariophyta) are one of the most successful phytoplankton group with a worldwide distribution both in freshwater and marine ecosystems (de Vargas et al. 2015) and comprise about 200,000 species (Mann and Droop 1996). At a global scale, diatoms produce one-fifth of the primary production on Earth and play a fundamental role in carbon and silica bio-geochemical cycles (Falkowski et al. 1998; Kemp et al. 2006; Armbrust 2009; Bowler et al. 2010; Tréguer and De La Rocha 2013) through the long term carbon sequestration in the sea floor and the production of frustules, silica structures forming the external cell wall of diatoms. After cell death, the frustules sink into the water column and contribute significantly to the formation of siliceous sediments. *Phaeodactylum tricornutum*, and particularly the strain RCC2967 (also known as CCAP1055), originally isolated in 1956 in the North Atlantic Ocean (U.K. coast, Blackpool), is one of the few well studied diatoms species. For decades, it has been used as a model species for studying the evolution of diatoms (Bowler et al. 2008), physiology and diverse metabolic pathways (Allen et al. 2008; Kroth et al. 2008). Furthermore, several genetic tools have been developed for this species (Zhang and Hu 2014), including CRISPR/Cas9 (Nymark et al. 2016; Serif et al. 2018), and several biotechnological applications are underway (Molina Grima et al. 2003). Its complete genome is diploid with a 27.45 Mb haploid genome size, divided into 33 chromosomes with an average GC content of 49% (Bowler et al. 2008). Estimating the spontaneous mutation rate of a species belonging to the

Stramenopile kingdom is essential to expand knowledge about the mutation rate of eukaryotes and brings important insights into phytoplankton diversity and evolution.

RESULTS

Mutation rates

We applied a previously described mutation accumulation (MA) experiment protocol in liquid medium (Krasovec et al. 2016) to 36 MA lines derived from one ancestral cell. MA lines were maintained over ~181 generations on average (Table S1) at a low effective population size ($N_e \sim 7$) through single cell bottlenecks to ensure minimal selection against deleterious mutations. After sequencing the ancestral and the MA lines, about ~98.5% of the genome could be screened for mutations. We identified 156 *de novo* mutations (all leading to heterozygosity, Tables S1 and S2): 151 in the nuclear and 5 in the mitochondrial genome, respectively. Within the 86 mutations occurring in protein coding sequences from the nuclear genome, 26 were synonymous, 57 non-synonymous, and 3 were indels (1 frameshift, 2 codons insertions). The 77 Kb mitochondrial genome (35% GC content) contained 3 mutations (intergenic region) and 2 indels (codon deletions). No mutations were detected on the 117 Kb chloroplast genome (33% GC content). The twenty-two mutations randomly selected for independent confirmation by Sanger sequencing have all been validated.

The per nucleotide mutation rates (*bs* for base substitutions, *id* for insertion-deletions) are $\mu_{bs} = 4.77 \times 10^{-10}$ (Poisson CI 95%: $4.04 \times 10^{-10} - 5.61 \times 10^{-10}$) and $\mu_{id} = 1.58 \times 10^{-11}$ (Poisson CI 95%: $5.13 \times 10^{-12} - 3.69 \times 10^{-11}$) corresponding to a genome wide mutation rate of $U_{bs} = 0.0125$ and $U_{id} = 0.00041$ mutations per haploid genome. The mutation rate in the mitochondria equals $\mu_{mt_bs} = 1.1 \times 10^{-9}$, $\mu_{mt_id} = 7.3 \times 10^{-10}$ and is significantly higher than the nuclear mutation rate (Chi-squared test, P-value < 0.001). The upper limit of the mutation rate of the chloroplast can be estimated to be below 1.0×10^{-10} for both μ_{chl_bs} and μ_{chl_id} . Because mutations are rare events, the mutation distribution between lines is expected to follow a Poisson distribution (Cutler 2000). Otherwise the distribution is considered as over-dispersed, which means that the mutation rates between the lines differ significantly. In our

dataset, the number of mutations per line ranged between 1 to 11 with an average of 4.3 (Figure S1). This is consistent with a Poisson distribution (Chi-Square test, X-squared = 59.53, df = 63, p-value = 0.60) with a distribution index of 0.62. Figure 1 shows the nuclear mutation rate of *P. tricornutum* in relation to the genome wide mutation rates available for other species. The data are largely biased towards the two Eukaryote kingdoms, which include the traditional animal and plant model species, highlighting the lack of mutation rate estimates in many lineages.

As *P. tricornutum* is diploid, the rate of heterozygosity along its genome provides an estimation of the level of polymorphism in the population. Single Nucleotide Polymorphism (SNP) analysis revealed 239,478 SNPs for 25,613,219 positions, which corresponds to a level of heterozygosity of 0.9%. In protein coding sequences, there are 57,890 synonymous polymorphisms out of 3,479,077 synonymous sites, and 50,925 non-synonymous polymorphisms out of 11,024,020 non-synonymous sites, corresponding to $\pi_s=0.0166$ and $\pi_n/\pi_s\sim 0.278$.

Mutation spectrum

The synonymous and non-synonymous mutation rates (Table 1) did not deviate from the neutral expectation (Chi-squared test, NS), supporting a low selection against non-synonymous deleterious mutations during the MA experiment. One multiple nucleotide mutation event was detected on chromosome one (3 mutations within 4 nucleotides), which corresponds to ~0.6% of total mutation events. No bias in the distribution of mutations was observed between protein coding and intergenic sequences (Chi-squared test, NS, Table 1), nor between chromosomes (Chi-squared test, NS). Mutated sites had similar RNAseq coverage to constant sites, suggesting that the transcription rate has no effect on the mutation rate in this species (transcription data from Levering (Levering et al. 2017)). However, there were two biases in the distribution of mutations. First, the dinucleotide context had a significant effect on mutation rates (Chi-squared test, P-value=0.0002); CG/CC/TG dinucleotides were mutagenic, while AT/CA/GA/AA displayed a lower mutation

rate on the nucleotide at the second position (Figure S2). Second, there was a mutation bias between G or C and A or T nucleotides (Figure 2): the mutation rate from G or C to A or T was 2.21 times that from A or T to G or C (Table 1). Consequently, the expected equilibrium GC content, e.g. the GC content resulting from the mutational process alone, equals 31.2%, while the observed GC content is 48.8%. Using a previously described calculation of the expected mutation rate at the equilibrium GC content (Krasovec et al. 2017), the expected mutation rate equals 4.13×10^{-10} mutations per site. The difference between the observed GC content and the equilibrium GC content therefore leads to a 15.4% increase in the mutation rate in this species.




Figure 1. Spontaneous mutation rates from mutation accumulation and pedigree studies, left to right: *Homo sapiens* (Besenbacher et al. 2016); *Clupea harengus* (Feng et al. 2017); *Mus musculus* (Uchimura et al. 2015); *Ficedula albicollis* (Smeds et al. 2016); *Caenorhabditis elegans*, *C. briggsae* (Denver et al. 2012); *Pristionchus pacificus* (Weller et al. 2014); *Daphnia pulex* (Flynn et al. 2017); *Drosophila melanogaster* (Schridder et al. 2013); *Heliconius melpomene* (Keightley et al. 2015); *Bombus terrestris*, *Apis mellifera* (Liu et al. 2017); *Chironomus riparius* (Oppold and Pfenninger 2017); *Saccharomyces cerevisiae* (Zhu et al. 2014); *Schizosaccharomyces pombe* (Farlow et al. 2015); *Dictyostelium discoideum* (Saxer et al. 2012); *Sphaeroforma arctica* (Long et al. 2018); *Arabidopsis thaliana* (Ossowski et al. 2010); *Silene latifolia* (Krasovec et al. 2018); *Prunus persica* (Xie et al. 2016); *Ostreococcus tauri*, *O. mediterraneus*, *Bathycoccus prasinos*, *Micromonas pusilla* (Krasovec et al. 2017); *Chlamydomonas reinhardtii* (Ness et al. 2015); *Picochlorum costaverdella* (Krasovec et al. 2018); *Pt: Phaeodactylum tricorutum*; *Paramecium tetraurelia* (Sung et al. 2012); *P. biaurelia* (Long et al. 2018); *P. sexaurelia* (Long et al. 2018); *Tetrahymena thermophila* (Long et al. 2016); *Plasmodium falciparum* (Hamilton et al. 2017); *Bacillus subtilis* (Sung et al. 2015); *Escherichia coli* (Lee et al. 2012); *Mesoplasma florum* (Sung et al. 2012); *Burkholderia cenocepacia* (Dillon et al. 2015); *Pseudomonas aeruginosa* (Dettman et al. 2016); *Salmonella typhimurium* (Lind and Andersson 2008); *Mycobacterium tuberculosis* (Ford et al. 2011); *M. smegmatis* (Kucukyildirim et al. 2016); *Deinococcus radiodurans* (Long et al. 2015); *Vibrio cholerae*, *V. fischeri* (Dillon et al. 2017); *Ruegeria pomeroyi* (Sun et al. 2017); *Arthrobacter* sp., *Flavobacterium* sp., *Janthinobacterium lividum*, *Micrococcus* sp., *Caulobacter crescentus*, *Rhodobacter sphaeroides*, *Staphylococcus aureus*,

Kineococcus radiotolerans, *Colwellia psychrerythraea*, *Lactococcus lactis*, *Gemmata obscuriglobus* (Long et al. 2018).




Figure 2. Distribution of the 146 *de novo* base substitution mutations in the nuclear genome across the 36 MA lines.

Table 1. Nuclear mutation rates. *N* is the number of *de novo* mutations, μ is the mutation rate per site per generation.

Substitution	N	μ
GC -> AT	80	4.82E-10
AT -> GC	38	2.18E-10
GC -> CG	24	1.44E-10
AT -> TA	4	2.30E-11
Intergenic	57	3.88E-10
Intron	5	
Non-syn	57	
Genic	26	5.00E-10
Syn	26	
UTR	3	
Frame shift	1	
Codon insertion	2	

DISCUSSION

This first estimation of a spontaneous mutation rate in a species belonging to the Stramenopile kingdom is consistent with previous estimates in four phytoplanktonic species from the Mamiellophyceae class (green algae). It is similar to the spontaneous mutation rates of 4.79×10^{-10} mutations per site per generation reported in the model species *Ostreococcus tauri* (Krasovec et al. 2017). This novel estimation places the average mutation rate of unicellular eukaryotes in the range of 10^{-10} mutations per site per generation, with the notable exception of ciliates (Figure 1).

P. tricornutum, as well as *O. tauri*, belong to phytoplankton communities isolated in coastal areas in the North Atlantic Ocean and the North West Mediterranean Sea, respectively. Coastal phytoplankton species contribute to primary production supporting the base of the food web in ocean ecosystems. For decades, estuarine and coastal ecosystems have been subjected to high anthropogenic pressures (Jackson et al. 2001), reducing their ecological services like carbon sequestration, raw material source, food or water purification (Barbier et al. 2011). Knowledge about the mutation rate and the effective population sizes of phytoplankton species is crucial to evaluate and understand their adaptive potential to any selection pressure or environmental change. From the equation $\pi_s = 4 \cdot N_e \cdot \mu$ and the estimated π_s value $\pi_s = 0.0166$, we can deduce the effective population size of *P. tricornutum* RCC2967 is $\sim 8.7 \times 10^6$ (Poisson CI 95%: $7.4 \times 10^6 - 1.0 \times 10^7$). This is similar to the effective population size in *O. tauri* $N_e \sim 1.2 \times 10^7$ (Blanc-Mathieu et al. 2017) one of the rare available estimates in phytoplankton. In diatoms, a second estimate of $N_e \sim 16.5 \times 10^7$ has been inferred for the Southern Ocean species *Fragilariopsis cylindrus*, using a mutation rate value of $\mu = 1 \times 10^{-10}$ as a proxy (Mock et al. 2017). Applying the *P. tricornutum* mutation rate, the *F. cylindrus* N_e estimation decreases to $\sim 3.46 \times 10^7$. This suggests that the effective population size of *F. cylindrus* is about 4 times larger than that of *P. tricornutum*. If the effective population size is correlated to the census population in a similar way in these two species,

this is consistent with a previous report based on environmental sequencing. Indeed, *Fragilariopsis* has been estimated to be the second most abundant diatom genus after *Chaetoceros* (Malviya et al. 2016) from 46 sampling sites of the TARA Ocean expedition, while *Phaedactylum* could not be detected in this dataset. On the contrary, *P. tricornutum* is a littoral species and may thus be expected to have a smaller habitat range. However, the application of macro-ecological concepts like species ranges on microorganisms, which can reach concentrations of 10^6 individuals per ml of seawater, has to be interpreted with caution. The effective population size can vary significantly depending on the life cycle and the frequency of sexual reproduction. This has been demonstrated in wild yeast populations (Tsai et al. 2008), where a population with lower sexual reproduction frequency had a smaller effective population size. In diatoms, the primary mode of reproduction is asexual, so that the frequency of the sexual reproduction may also induce a variation in the effective population size, which corresponds to the number of clones in the population. The effective population size is ideally calculated from population genomics data, but such studies are scarce in phytoplankton species. Here, we have estimated the effective population size of *P. tricornutum* from the level of heterozygosity of the strain RCC2967. This strain was isolated in 1956, and a monoclonal culture with fusiform morphotype was generated in 2004 (Martino et al. 2007), and maintained as culture CCAP1055/1 (=RCC2967 or CCMP2561) for 14 years, from which the T_0 line of our MA experiment was derived. This history could have affected the selection of the initial strain and the level of heterozygosity estimated here may not be an accurate estimate of the heterozygosity within the natural population. Lab conditions have their own sources of stress, which may influence the mutation rate. Consistent with this, the microsatellite mutation rate was reported to be higher in recently established lines of cultures of the diatoms species *Pseudo-nitzschia multistriata* (Tesson et al. 2013). Last but not least, the mutation rate can be affected by the environment (Baer 2008) and stress (Jiang et al. 2014). As a consequence, any estimation of experimental mutation rates may differ from the long-term average mutation rate in the natural environment.

Mutation rates in primary and secondary organelles

The mitochondrial and chloroplast genomes display a ~10-fold variation in their mutation rates in *P. tricornutum*. Large variations in organelle mutation rates have been observed in eukaryotes and have been linked to their evolutionary history, in particular in relation to the acquisition of plastids following a primary or secondary endosymbiosis (Smith 2015; Smith and Keeling 2015), phylogenetic constraint, genome structure, and non-adaptive processes (Lynch et al. 2006). Most current data on organellar mutation rates are derived from synonymous substitutions rates (Wolfe et al. 1987; Smith and Keeling 2015). Available evidence from photosynthetic species support that species which evolved after a secondary endosymbiosis event have a higher mitochondrion substitution rate compared to the chloroplast and nuclear mutation rates (Smith and Keeling 2012). Our data on *P. tricornutum* provides direct evidence for a higher mitochondrial mutation rate compared to the chloroplast and nucleus in a microalga evolved from a secondary endosymbiosis event. In addition to this study, direct estimates from mutation accumulation experiment are available in the green algae *Chlamydomonas reinhardtii* (Ness et al. 2016) and in a few metazoans such as *Caenorhabditis elegans* and *Daphnia pulex* (Denver et al. 2000; Xu et al. 2012; Konrad et al. 2017). Mitochondrial mutation-rate estimates in metazoans are 100 times higher than those for the nuclear genomes in the same lineages (Denver et al. 2000; Xu et al. 2012; Konrad et al. 2017). Conversely, in land plants, the mitochondrial mutation rate is generally lower than the nuclear mutation rate (Drouin et al. 2008). This might be the result of greater accuracy of a specific DNA polymerase involved in organelle replication and repair (Parent et al. 2011; Gualberto et al. 2014). However, this DNA polymerase family is not specific to land plants and the genome of *P. tricornutum* encodes a gene belonging to the same orthologous gene family (DNA polymerase A domain, ORTHO000433 orthologous gene family indexed in the picoPLAZA database (Vandepoele et al. 2013)). The three genome compartments seem to have similar mutation rates in green algae, as observed in

Chlamydomonas reinhardtii (Hua et al. 2012; Ness et al. 2016) and *Volvox carteri* (Smith and Lee 2010). Previous mutation accumulation experiments on five species of marine green algae did not report any *de novo* organellar mutations (Krasovec et al. 2017; Krasovec et al. 2018). Given the mitochondrial and chloroplast genome sizes in *O. tauri* (Mamiellophyceae), it follows μ are less than 8.28×10^{-10} and 1.45×10^{-9} in the chloroplast and the mitochondria, respectively. The application of the same reasoning to *Picochlorum costavermella* (Trebouxiophyceae) leads to the conclusion that the mutation rates are less than 8.43×10^{-9} and 1.8×10^{-8} in the chloroplast and the mitochondria for a nuclear mutation rate of 1.01×10^{-9} mutations per site. These maximum organellar mutation rate estimates would be consistent with a higher mutation rates in organelles than in the nuclear genomes in these lineages. Precise estimations, implying mutation accumulation experiments over a higher number of generations are obviously needed to test this hypothesis.

Mutation rate variation

The difference between the observed GC and the equilibrium GC content has a moderate effect on the mutation rate (+15.4%) in *P. triconutum*. This effect is variable between species; in *A. thaliana* and *C. reinhardtii*, it leads to a mutation rate increase of 64% and 42%, respectively (see Table S12 of (Krasovec et al. 2017)). In the bacteria *Mesoplasma florum*, which appears to have the highest mutation rate measured in bacteria (Figure 1), the GC bias effect is very high with an observed μ ~263% its expected value at equilibrium GC content. Within the Mamiellophyceae, the mutation rate is increased by ~2 to ~12% depending on the species. The difference between the observed and equilibrium GC content may be explained by three processes, either a recent change in the mutation bias between AT and GC nucleotides, or GC biased gene conversion (Duret and Galtier 2009), or selection on GC (Eyre-Walker 1999; Hildebrand et al. 2010). A change in the mutation bias may be due to a change in the rate of oxidation of the G nucleotide to 8-oxoguanine (Cooke et al. 2003) or a change in the proportion of C nucleotide methylation, which has an increased probability of mutation towards T (Holliday and Grigg 1993). Cytosine methylation

has been documented in *P. tricornutum*, where about 6% of the genome is methylated (Huff and Zilberman 2014; Veluchamy et al. 2014). This may explain some of the GC bias, even moderate, observed in this species.

The mutation rate of *P. tricornutum* is not homogeneous within the genome. The effect of the nucleotide context has already been observed in others species such as the green algae *Chlamydomonas reinhardtii* (Ness et al. 2015) and the bacteria *Bacillus subtilis* (Sung et al. 2015). The link between mutation probability and nucleotide context is of primary importance for molecular evolution on synonymous versus non-synonymous sites. In the bacteria *Bacillus subtilis* (Sung et al. 2015), the differences in the probability of mutation between sites may induce a difference in the rate of evolution between codons. In *P. tricornutum*, the most mutable dinucleotide is CG->CN. A codon starting with CG codes for the amino acid arginine: this amino-acid therefore has a higher probability of mutating into another amino-acid. Since G undergoes more frequent mutations than A, it is expected that arginine codons will be more frequently replaced by histidine or glutamine. Similarly, codons ending in CG (serine, proline, alanine and threonine) will evolve faster than synonymous codons ending with GA. In conclusion, knowledge of the mutation spectrum is not only essential to predict the rate of adaptation to environmental change, but also to infer the evolutionary history through accurate calibration of the molecular clock.

MATERIALS AND METHODS

MA experiment

We performed a mutation accumulation experiment in liquid medium using the method described by Krasovec et al. 2016). Briefly, one single cell from the strain RCC2967 (monoclonal culture obtained in 2004 from a harvested strain in 1956 (Martino et al. 2007)) from the Roscoff Culture Collection (Vaulot et al. 2004) was selected from 10 cells sampled from the culture. The 10 cells were obtained by pipetting the corresponding volume estimated from the cell concentration by flow cytometry. Resampling 6 cells out of 10 into 6 wells to choose one well greatly decreases the probability of sampling zero cells (Krasovec et al. 2016). This sampling procedure was repeated to inoculate one single cell for each MA line from this T_0 culture in 24-wells plate in L1 liquid medium with a life cycle of 8h light - 16h dark at 20°C. 36 MA lines were kept during 98 to 196 days corresponding to 154 to 210 generations. Cell concentrations were estimated by flow cytometry at each bottleneck time (every 14 days) to estimate the population size N_t and to isolate one single cell for re-inoculation by dilution. Single cell re-inoculation reduces effective population size and thus selection on deleterious spontaneous mutations. N_t was used to estimate the effective population size (N_e) of MA lines as the harmonic mean of cell number, and to estimate the number of generations (d) from the following equation:

$$d = e^{\left[\ln\left(\frac{N_t}{N_0}\right)/t\right]} \quad (1)$$

with $t=14$ days, $N_0=1$ cell and N_t the cell concentration at each bottleneck time.

Mutation calling and spectrum

The genomes of the 36 MA lines and the ancestral T_0 genome were sequenced by Illumina MiSeq technology (125 bp paired-end reads). We checked the raw reads quality with *Trimmomatic* v0.36 (Bolger et al. 2014) and then aligned them to the reference genome (NCBI reference: GCA_000150955.2) using *BWA mem* v0.7.15 with standard parameters (Li

and Durbin 2010). Resulting BAM files were treated with *SAMtools* v0.1.19 (Li et al. 2009) and mutation calling was done with and *HaplotypeCaller* from GATK v3.5 following the best practice recommendations (*BaseRecalibrator*, *RealignerTargetCreator*, *IndelRealigner*) (McKenna et al. 2010) and a homemade C code to check the called mutations in the mpileup files. We used similar criteria to identify true *de novo* mutations as used in previous MA studies in diploid species (Keightley et al. 2015; Krasovec et al. 2018). Several criteria enable to discriminate *de novo* mutations from heterozygotic SNPs: (i) Callable sites were considered above a threshold of 40 mapping quality (MQ value provided by the vcf GATK file) and (ii) a minimum read coverage of 30 and (iii) a maximum coverage of 2.5 times of the average coverage (to exclude false positives due to repetitive sequences) both in the MA lines and the initial T_0 genome; (iv) A *de novo* mutation was called only if the polymorphism appeared in a unique MA line, was covered by a minimum of 10 reads or 30% of the total coverage; (v) each mutation candidate was inspected visually in the *mpileup* files and in the bam file with IGV (Robinson et al. 2011) both in the MA lines and the T_0 genome; Last, (vi) we randomly selected 22 of the final *de novo* mutation candidates for independent Sanger sequencing in MA lines and in the initial line to validate the candidates. All of the 22 mutations were validated by PCR. Mutation effects (synonymous, non-synonymous, intergenic) were predicted with *snpEff* v3.6 (Cingolani et al. 2012).

The distribution of the mutations along the genome was compared to the random and independent distributions of mutations with a Chi-squared test. First, we tested the distribution between protein coding, non-protein coding, intergenic, intronic, exonic and UTR regions and between chromosomes. Second, we investigated the nucleotide context effect to detect the impact of the previous (5 prime) nucleotide on the probability of mutation (NX). Last, we aligned RNA sequence data from Levering (Levering et al. 2017) in standard condition (SRA numbers SRS3629289, SRS3629290 and SRS3629291) to test the correlation between transcription and mutation rates. RNA data were aligned against the reference using STAR with standard parameters (Dobin et al. 2013) and expression level was

analysed with HTSeq (Anders et al. 2015). Statistical analyses were performed with R v3.1.1. The mutation spectrum was obtained using the equations below (Sueoka 1962):

$$R_1 = \frac{(GC \rightarrow AT)}{GC_n}, R_2 = \frac{(AT \rightarrow GC)}{AT_n}, R_3 = \frac{(AT \rightarrow AT)}{AT_n}, R_4 = \frac{(GC \rightarrow GC)}{GC_n} \quad (2)$$

$$GC_{eq} = \frac{R_2}{R_1 + R_2} \quad (3)$$

with GC_{eq} the GC content of the genome under mutation processes alone. The level of polymorphism of *P. tricornutum* was estimated from the level of heterozygosity of the T_0 genome using the following SNP calling thresholds; a minimum coverage of 20, a minimum mapping quality of 40 and the position defined as diploid (0/1) in the GATK vcf file.

Mutation rate estimation

The mutation rate was calculated from the whole experiment. First, the total number of callable sites was calculated for each MA line using the callable sites quality threshold (see above). The callable site represented ~98.5% of the genome on average and the average number of generations per line is ~181, summing up to ~6,500. Then, the mutation rate was calculated by dividing the total number of mutations by twice of the total number of callable sites, as *P. tricornutum* is diploid. The effective population size of the strain was estimated from the equation $\pi_s = 4.N_e.\mu$ (Nei and Tajima 1981) with π_s estimated from the T_0 genome. The π_n/π_s ratio was estimated from the T_0 genome considering the number of synonymous and non-synonymous SNPs relative to the synonymous and non-synonymous callable sites. The number of mitochondria and chloroplast genomes was estimated by dividing the average coverage of each organelle by half the average coverage of the nuclear genome. This calculation inferred ~11 mitochondria and ~26 chloroplasts per cell. To deal with heteroplasmy, *de novo* mutations in organelles were considered true if supported by at least the coverage corresponding to one copy, e.g. 1/11 and 1/26 of the total coverage of the mitochondria and the chloroplast, respectively.

DATA AVAILABILITY

All raw reads are available under the bio-project PRJNA478011. The bio-samples are from SAMN09489389 to SAMN09489425.

ACKNOWLEDGEMENT

We are grateful to all members of the GENOPHY Lab for discussion and technical support. Special acknowledgements to Elodie Desgranges, Christophe Salmeron and David Pecqueur from the BIOPIC platform for experimental support on flow cytometry, and BIO2MAR platform for Sanger sequencing. We thank the GenoToul Bioinformatics platform from Toulouse, France, for cluster availability and support on bioinformatics analysis on cluster. This work was funded by ANRJJC-SVSE6-2013-0005 to GP and SSB.

COMPETING INTERESTS

The authors declare no conflicts of interest.

AUTHOR CONTRIBUTIONS

GP, SSB and MK designed the experiment. MK performed the experiment and analysed the data. MK, GP and SSB wrote the paper.

BIBLIOGRAPHY

Allen AE, LaRoche J, Maheswari U, Lommer M, Schauer N, Lopez PJ, Finazzi G, Fernie AR, Bowler C. 2008. Whole-cell response of the pennate diatom *Phaeodactylum tricornutum* to iron starvation. *Proc. Natl. Acad. Sci. U. S. A.* 105:10438–10443.

Anders S, Pyl PT, Huber W. 2015. HTSeq--a Python framework to work with high-throughput sequencing data. *Bioinforma. Oxf. Engl.* 31:166–169.

Armbrust EV. 2009. The life of diatoms in the world's oceans. *Nature.* 459:185–192

Baer CF. 2008. Does Mutation Rate Depend on Itself. *PLOS Biol.* 6:e52.

Barbier EB, Hacker SD, Kennedy C, Koch EW, Stier AC, Silliman BR. 2011. The value of estuarine and coastal ecosystem services. *Ecol. Monogr.* 81:169–193.

- Besenbacher S, Sulem P, Helgason A, Helgason H, Kristjansson H, Jonasdottir Aslaug, Jonasdottir Adalbjorg, Magnusson OT, Thorsteinsdottir U, Masson G, et al. 2016. Multi-nucleotide de novo Mutations in Humans. *PLoS Genet.* 12:e1006315.
- Blanc-Mathieu R, Krasovec M, Hebrard M, Yau S, Desgranges E, Martin J, Schackwitz W, Kuo A, Salin G, Donnadiou C, et al. 2017. Population genomics of picophytoplankton unveils novel chromosome hypervariability. *Sci. Adv.* 3:e1700239.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120.
- Bowler C, Allen AE, Badger JH, Grimwood J, Jabbari K, Kuo A, Maheswari U, Martens C, Maumus F, Otillar RP, et al. 2008. The *Phaeodactylum* genome reveals the evolutionary history of diatom genomes. *Nature* 456:239–244.
- Bowler C, Vardi A, Allen AE. 2010. Oceanographic and biogeochemical insights from diatom genomes. *Annu. Rev. Mar. Sci.* 2:333–365.
- Charlesworth B, Charlesworth D. 1998. Some evolutionary consequences of deleterious mutations. *Genetica* 102–103:3–19.
- Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, Land SJ, Lu X, Ruden DM. 2012. A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster* strain w1118; iso-2; iso-3. *Fly (Austin)* 6:80–92.
- Cooke MS, Evans MD, Dizdaroglu M, Lunec J. 2003. Oxidative DNA damage: mechanisms, mutation, and disease. *FASEB J. Off. Publ. Fed. Am. Soc. Exp. Biol.* 17:1195–1214.
- Cutler DJ. 2000. Understanding the overdispersed molecular clock. *Genetics* 154:1403–1417.
- Denver DR, Morris K, Lynch M, Vassilieva LL, Thomas WK. 2000. High direct estimate of the mutation rate in the mitochondrial genome of *Caenorhabditis elegans*. *Science* 289:2342–2344.
- Denver DR, Wilhelm LJ, Howe DK, Gafner K, Dolan PC, Baer CF. 2012. Variation in base-substitution mutation in experimental and natural lineages of *Caenorhabditis nematodes*. *Genome Biol. Evol.* 4:513–522.
- Dettman JR, Sztepanacz JL, Kassen R. 2016. The properties of spontaneous mutations in the opportunistic pathogen *Pseudomonas aeruginosa*. *BMC Genomics.* 5;17:27
- Dillon MM, Sung W, Lynch M, Cooper VS. 2015. The Rate and Molecular Spectrum of Spontaneous Mutations in the GC-Rich Multichromosome Genome of *Burkholderia cenocepacia*. *Genetics* 200:935–946.
- Dillon MM, Sung W, Sebra R, Lynch M, Cooper VS. 2017. Genome-Wide Biases in the Rate and Molecular Spectrum of Spontaneous Mutations in *Vibrio cholerae* and *Vibrio fischeri*. *Mol. Biol. Evol.* 34:93–109.

- Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinforma. Oxf. Engl.* 29:15–21.
- Drouin G, Daoud H, Xia J. 2008. Relative rates of synonymous substitutions in the mitochondrial, chloroplast and nuclear genomes of seed plants. *Mol. Phylogenet. Evol.* 49:827–831.
- Duret L, Galtier N. 2009. Biased gene conversion and the evolution of mammalian genomic landscapes. *Annu. Rev. Genomics Hum. Genet.* 10:285–311.
- Eyre-Walker A. 1999. Evidence of selection on silent site base composition in mammals: potential implications for the evolution of isochores and junk DNA. *Genetics* 152:675–683.
- Falkowski null, Barber null, Smetacek null. 1998. Biogeochemical Controls and Feedbacks on Ocean Primary Production. *Science* 281:200–207.
- Farlow A, Long H, Arnoux S, Sung W, Doak TG, Nordborg M, Lynch M. 2015. The Spontaneous Mutation Rate in the Fission Yeast *Schizosaccharomyces pombe*. *Genetics* 201:737–744.
- Feng C, Pettersson M, Lamichhaney S, Rubin C-J, Rafati N, Casini M, Folkvord A, Andersson L. Moderate nucleotide diversity in the Atlantic herring is associated with a low mutation rate. *eLife* 30:6: e23907.
- Flynn JM, Chain FJJ, Schoen DJ, Cristescu ME. 2017. Spontaneous Mutation Accumulation in *Daphnia pulex* in Selection-Free vs. Competitive Environments. *Mol. Biol. Evol.* 34:160–173.
- Ford CB, Lin PL, Chase MR, Shah RR, Iartchouk O, Galagan J, Mohaideen N, Ioerger TR, Sacchettini JC, Lipsitch M, et al. 2011. Use of whole genome sequencing to estimate the mutation rate of *Mycobacterium tuberculosis* during latent infection. *Nat. Genet.* 43:482–486.
- Gualberto JM, Mileshina D, Wallet C, Niazi AK, Weber-Lotfi F, Dietrich A. 2014. The plant mitochondrial genome: Dynamics and maintenance. *Biochimie* 100:107–120.
- Hamilton WL, Claessens A, Otto TD, Kekre M, Fairhurst RM, Rayner JC, Kwiatkowski D. 2017. Extreme mutation bias and high AT content in *Plasmodium falciparum*. *Nucleic Acids Res.* 45:1889–1901.
- Hershberg R, Petrov DA. 2010. Evidence That Mutation Is Universally Biased towards AT in Bacteria. *PLoS Genet* 6:e1001115.
- Hildebrand F, Meyer A, Eyre-Walker A. 2010. Evidence of Selection upon Genomic GC-Content in Bacteria. *PLoS Genet.* 9:6:e1001107
- Holliday R, Grigg GW. 1993. DNA methylation and mutation. *Mutat. Res.* 285:61–67.
- Hua J, Smith DR, Borza T, Lee RW. 2012. Similar relative mutation rates in the three genetic compartments of *Mesostigma* and *Chlamydomonas*. *Protist* 163:105–115.

- Huff JT, Zilberman D. 2014. Dnmt1-Independent CG Methylation Contributes to Nucleosome Positioning in Diverse Eukaryotes. *Cell* 156:1286–1297.
- Jackson JBC, Kirby MX, Berger WH, Bjorndal KA, Botsford LW, Bourque BJ, Bradbury RH, Cooke R, Erlandson J, Estes JA, et al. 2001. Historical overfishing and the recent collapse of coastal ecosystems. *Science* 293:629–638.
- Jiang C, Mithani A, Belfield EJ, Mott R, Hurst LD, Harberd NP. 2014. Environmentally responsive genome-wide accumulation of de novo *Arabidopsis thaliana* mutations and epimutations. *Genome Res.* 24:1821–1829.
- Katju V, Bergthorsson U. 2019. Old Trade, New Tricks: Insights into the Spontaneous Mutation Process from the Partnering of Classical Mutation Accumulation Experiments with High-Throughput Genomic Approaches. *Genome Biol. Evol.* 11:136–165.
- Keightley PD, Pinharanda A, Ness RW, Simpson F, Dasmahapatra KK, Mallet J, Davey JW, Jiggins CD. 2015. Estimation of the spontaneous mutation rate in *Heliconius melpomene*. *Mol. Biol. Evol.* 32:239–243.
- Kemp AES, Pearce RB, Grigorov I, Rance J, Lange CB, Quilty P, Salter I. 2006. Production of giant marine diatoms and their export at oceanic frontal zones: Implications for Si and C flux from stratified oceans. *Glob. Biogeochem. Cycles.* 20.
- Kiktev DA, Sheng Z, Lobachev KS, Petes TD. 2018. GC content elevates mutation and recombination rates in the yeast *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci.* 115:E7109–E7118.
- Konrad A, Thompson O, Waterston RH, Moerman DG, Keightley PD, Bergthorsson U, Katju V. 2017. Mitochondrial Mutation Rate, Spectrum and Heteroplasmy in *Caenorhabditis elegans* Spontaneous Mutation Accumulation Lines of Differing Population Size. *Mol. Biol. Evol.* 34:1319–1334.
- Krasovec M, Chester M, Ridout K, Filatov DA. 2018. The Mutation Rate and the Age of the Sex Chromosomes in *Silene latifolia*. *Curr. Biol.* 28:1832-1838.e4.
- Krasovec M, Eyre-Walker A, Grimsley N, Salmeron C, Pecqueur D, Piganeau G, Sanchez-Ferandin S. 2016. Fitness Effects of Spontaneous Mutations in Picoeukaryotic Marine Green Algae. G3 Bethesda Md.
- Krasovec M, Eyre-Walker A, Sanchez-Ferandin S, Piganeau G. 2017. Spontaneous Mutation Rate in the Smallest Photosynthetic Eukaryotes. *Mol. Biol. Evol.* 34:1770–1779.
- Krasovec M, Sanchez-Brosseau S, Grimsley N, Piganeau G. 2018. Spontaneous mutation rate as a source of diversity for improving desirable traits in cultured microalgae. *Algal Res.* 35:85–90.
- Kroth PG, Chiovitti A, Gruber A, Martin-Jezequel V, Mock T, Parker MS, Stanley MS, Kaplan A, Caron L, Weber T, et al. 2008. A Model for Carbohydrate Metabolism in the Diatom *Phaeodactylum tricornutum* Deduced from Comparative Whole Genome Analysis. *PLOS ONE* 3:e1426.

Kucukyildirim S, Long H, Sung W, Miller SF, Doak TG, Lynch M. 2016. The Rate and Spectrum of Spontaneous Mutations in *Mycobacterium smegmatis*, a Bacterium Naturally Devoid of the Post-replicative Mismatch Repair Pathway. *G3 GenesGenomesGenetics*:g3.116.030130.

Lee H, Popodi E, Tang H, Foster PL. 2012. Rate and molecular spectrum of spontaneous mutations in the bacterium *Escherichia coli* as determined by whole-genome sequencing. *Proc. Natl. Acad. Sci. U. S. A.* 109:E2774-2783.

Levering J, Dupont CL, Allen AE, Palsson BO, Zengler K. 2017. Integrated Regulatory and Metabolic Networks of the Marine Diatom *Phaeodactylum tricornutum* Predict the Response to Rising CO₂ Levels. *mSystems*. 14:2(1). e00142-16

Li H, Durbin R. 2010. Fast and accurate long-read alignment with Burrows–Wheeler transform. *Bioinformatics* 26:589–595.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000 Genome Project Data Processing Subgroup. 2009. The Sequence Alignment/Map format and SAMtools. *Bioinforma. Oxf. Engl.* 25:2078–2079.

Lind PA, Andersson DI. 2008. Whole-genome mutational biases in bacteria. *Proc. Natl. Acad. Sci. U. S. A.* 105:17878–17883.

Liu H, Jia Y, Sun X, Tian D, Hurst LD, Yang S. 2017. Direct Determination of the Mutation Rate in the Bumblebee Reveals Evidence for Weak Recombination-Associated Mutation and an Approximate Rate Constancy in Insects. *Mol. Biol. Evol.* 34:119–130.

Long H, Doak TG, Lynch M. 2018. Limited Mutation-Rate Variation Within the *Paramecium aurelia* Species Complex. *G3 GenesGenomesGenetics* 8:2523–2526.

Long H, Kucukyildirim S, Sung W, Williams E, Lee H, Ackerman M, Doak TG, Tang H, Lynch M. 2015. Background Mutational Features of the Radiation-Resistant Bacterium *Deinococcus radiodurans*. *Mol. Biol. Evol.* 32:2383–2392.

Long H, Sung W, Kucukyildirim S, Williams E, Miller SF, Guo W, Patterson C, Gregory C, Strauss C, Stone C, et al. 2018. Evolutionary determinants of genome-wide nucleotide composition. *Nat. Ecol. Evol.* 2:237–240.

Long H, Winter DJ, Chang AY-C, Sung W, Wu SH, Balboa M, Azevedo RBR, Cartwright RA, Lynch M, Zufall RA. 2016. Low Base-Substitution Mutation Rate in the Germline Genome of the Ciliate *Tetrahymena thermophil*. *Genome Biol. Evol.* 8:3629–3639.

Lynch M, Ackerman MS, Gout J-F, Long H, Sung W, Thomas WK, Foster PL. 2016. Genetic drift, selection and the evolution of the mutation rate. *Nat. Rev. Genet.* 17:704–714.

Lynch M, Koskella B, Schaack S. 2006. Mutation pressure and the evolution of organelle genomic architecture. *Science* 311:1727–1730.

Malviya S, Scalco E, Audic S, Vincent F, Veluchamy A, Poulain J, Wincker P, Iudicone D, de Vargas C, Bittner L, et al. 2016. Insights into global diatom distribution and diversity in the world's ocean. *Proc. Natl. Acad. Sci. U. S. A.* 113:E1516-1525.

- Mann DG, Droop SJM. 1996. 3. Biodiversity, biogeography and conservation of diatoms. *Hydrobiologia* 336:19–32.
- Martino AD, Meichenin A, Shi J, Pan K, Bowler C. 2007. Genetic and phenotypic characterization of *Phaeodactylum tricornutum* (Bacillariophyceae) accessions. *J. Phycol.* 43:992–1009.
- McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K, Altshuler D, Gabriel S, Daly M, et al. 2010. The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res.* 20:1297–1303.
- Mock T, Otiillar RP, Strauss J, McMullan M, Paajanen P, Schmutz J, Salamov A, Sanges R, Toseland A, Ward BJ, et al. 2017. Evolutionary genomics of the cold-adapted diatom *Fragilariopsis cylindrus*. *Nature* 541:536–540.
- Molina Grima E, Belarbi E-H, Ación Fernández FG, Robles Medina A, Chisti Y. 2003. Recovery of microalgal biomass and metabolites: process options and economics. *Biotechnol. Adv.* 20:491–515.
- Nei M, Tajima F. 1981. Genetic drift and estimation of effective population size. *Genetics* 98:625–640.
- Ness RW, Kraemer SA, Colegrave N, Keightley PD. 2016. Direct Estimate of the Spontaneous Mutation Rate Uncovers the Effects of Drift and Recombination in the *Chlamydomonas reinhardtii* Plastid Genome. *Mol. Biol. Evol.* 33:800–808.
- Ness RW, Morgan AD, Vasanthakrishnan RB, Colegrave N, Keightley PD. 2015. Extensive de novo mutation rate variation between individuals and across the genome of *Chlamydomonas reinhardtii*. *Genome Res.* 25:1739–1749.
- Nymark M, Sharma AK, Sparstad T, Bones AM, Winge P. 2016. A CRISPR/Cas9 system adapted for gene editing in marine algae. *Sci. Rep.* 6:24951.
- Oppold A-M, Pfenninger M. 2017. Direct estimation of the spontaneous mutation rate by short-term mutation accumulation lines in *Chironomus riparius*. *Evol. Lett.* 1:86–92.
- Ossowski S, Schneeberger K, Lucas-Lledó JI, Warthmann N, Clark RM, Shaw RG, Weigel D, Lynch M. 2010. The rate and molecular spectrum of spontaneous mutations in *Arabidopsis thaliana*. *Science* 327:92–94.
- Parent J-S, Lepage E, Brisson N. 2011. Divergent Roles for the Two PolII-Like Organelle DNA Polymerases of *Arabidopsis*. *Plant Physiol.* 156:254–262.
- Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, Mesirov JP. 2011. Integrative Genomics Viewer. *Nat. Biotechnol.* 29:24–26.
- Saxer G, Havlak P, Fox SA, Quance MA, Gupta S, Fofanov Y, Strassmann JE, Queller DC. 2012. Whole genome sequencing of mutation accumulation lines reveals a low mutation rate in the social amoeba *Dictyostelium discoideum*. *PLoS One* 7:e46759.

- Schrider DR, Houle D, Lynch M, Hahn MW. 2013. Rates and genomic consequences of spontaneous mutational events in *Drosophila melanogaster*. *Genetics* 194:937–954.
- Serif M, Dubois G, Finoux A-L, Teste M-A, Jallet D, Daboussi F. 2018. One-step generation of multiple gene knock-outs in the diatom *Phaeodactylum tricornutum* by DNA-free genome editing. *Nat. Commun.* 9: 3924.
- Smeds L, Qvarnstrom A, Ellegren H. 2016. Direct estimate of the rate of germline mutation in a bird. *Genome Res.*
- Smith DR. 2015. Mutation rates in plastid genomes: they are lower than you might think. *Genome Biol. Evol.* 7:1227–1234.
- Smith DR, Keeling PJ. 2012. Twenty-Fold Difference in Evolutionary Rates between the Mitochondrial and Plastid Genomes of Species with Secondary Red Plastids. *J. Eukaryot. Microbiol.* 59:181–184.
- Smith DR, Keeling PJ. 2015. Mitochondrial and plastid genome architecture: Reoccurring themes, but significant differences at the extremes. *Proc. Natl. Acad. Sci.* 112:10177–10184.
- Smith DR, Lee RW. 2010. Low nucleotide diversity for the expanded organelle and nuclear genomes of *Volvox carteri* supports the mutational-hazard hypothesis. *Mol. Biol. Evol.* 27:2244–2256.
- Sueoka N. 1962. On the genetic basis of variation and heterogeneity of DNA base composition. *Proc. Natl. Acad. Sci. U. S. A.* 48:582–592.
- Sun Y, Powell KE, Sung W, Lynch M, Moran MA, Luo H. 2017. Spontaneous mutations of a model heterotrophic marine bacterium. *ISME J.* 11(7):1713-1718.
- Sung W, Ackerman MS, Gout J-F, Miller SF, Williams E, Foster PL, Lynch M. 2015. Asymmetric Context-Dependent Mutation Patterns Revealed through Mutation–Accumulation Experiments. *Mol. Biol. Evol.* 32:1672–1683.
- Sung W, Ackerman MS, Miller SF, Doak TG, Lynch M. 2012. Drift-barrier hypothesis and mutation-rate evolution. *Proc. Natl. Acad. Sci. U. S. A.* 109:18488–18492.
- Sung W, Tucker AE, Doak TG, Choi E, Thomas WK, Lynch M. 2012. Extraordinary genome stability in the ciliate *Paramecium tetraurelia*. *Proc. Natl. Acad. Sci. U. S. A.* 109:19339–19344.
- Tesson SVM, Legrand C, van Oosterhout C, Montresor M, Kooistra WHCF, Procaccini G. 2013. Mendelian inheritance pattern and high mutation rates of microsatellite alleles in the diatom *Pseudo-nitzschia multistriata*. *Protist* 164:89–100.
- Tréguer PJ, De La Rocha CL. 2013. The world ocean silica cycle. *Annu. Rev. Mar. Sci.* 5:477–501.
- Tsai IJ, Bensasson D, Burt A, Koufopanou V. 2008. Population genomics of the wild yeast *Saccharomyces paradoxus*: Quantifying the life cycle. *Proc. Natl. Acad. Sci.* 105:4957–4962.

Uchimura A, Higuchi M, Minakuchi Y, Ohno M, Toyoda A, Fujiyama A, Miura I, Wakana S, Nishino J, Yagi T. 2015. Germline mutation rates and the long-term phenotypic effects of mutation accumulation in wild-type laboratory mice and mutator mice. *Genome Res.* 25:1125–1134.

Vandepoele K, Van Bel M, Richard G, Van Landeghem S, Verhelst B, Moreau H, Van de Peer Y, Grimsley N, Piganeau G. 2013. pico-PLAZA, a genome database of microbial photosynthetic eukaryotes. *Environ. Microbiol.* 15:2147–2153.

de Vargas C, Audic S, Henry N, Decelle J, Mahé F, Logares R, Lara E, Berney C, Le Bescot N, Probert I, et al. 2015. Ocean plankton. Eukaryotic plankton diversity in the sunlit ocean. *Science* 348:1261605.

Vaulot D, Gall FL, Marie D, Guillou L, Partensky F. 2004. The Roscoff Culture Collection (RCC): a collection dedicated to marine picoplankton. *Nova Hedwigia* 79(1-2):49-70.

Veluchamy A, Lin X, Maumus F, Rivarola M, Bhavsar J, Creasy T, O'Brien K, Sengamalay NA, Tallon LJ, Smith AD, et al. 2014. Corrigendum: Insights into the role of DNA methylation in diatoms by genome-wide profiling in *Phaeodactylum tricornutum*. *Nat. Commun.* 5:3028.

Weller AM, Rödelberger C, Eberhardt G, Molnar RI, Sommer RJ. 2014. Opposing forces of A/T-biased mutations and G/C-biased gene conversions shape the genome of the nematode *Pristionchus pacificus*. *Genetics* 196:1145–1152.

Wolfe KH, Li WH, Sharp PM. 1987. Rates of nucleotide substitution vary greatly among plant mitochondrial, chloroplast, and nuclear DNAs. *Proc. Natl. Acad. Sci. U. S. A.* 84:9054–9058.

Xie Z, Wang Long, Wang Lirong, Wang Z, Lu Z, Tian D, Yang S, Hurst LD. 2016. Mutation rate analysis via parent-progeny sequencing of the perennial peach. I. A low rate in woody perennials and a higher mutagenicity in hybrids. *Proc. Biol. Sci.* 283.

Xu S, Schaack S, Seyfert A, Choi E, Lynch M, Cristescu ME. 2012. High mutation rates in the mitochondrial genomes of *Daphnia pulex*. *Mol. Biol. Evol.* 29:763–769.

Zhang C, Hu H. 2014. High-efficiency nuclear transformation of the diatom *Phaeodactylum tricornutum* by electroporation. *Mar. Genomics* 16:63–66.

Zhu YO, Siegal ML, Hall DW, Petrov DA. 2014. Precise estimates of mutation rate and spectrum in yeast. *Proc. Natl. Acad. Sci. U. S. A.* 111:E2310-2318.