



HAL
open science

Final size and convergence rate for an epidemic in heterogeneous population

Luís Almeida, Pierre-Alexandre Bliman, Grégoire Nadin, Benoît Perthame,
Nicolas Vauchelet

► **To cite this version:**

Luís Almeida, Pierre-Alexandre Bliman, Grégoire Nadin, Benoît Perthame, Nicolas Vauchelet. Final size and convergence rate for an epidemic in heterogeneous population. *Mathematical Models and Methods in Applied Sciences*, 2021, 31 (05), pp.1021-1051. hal-02981952v1

HAL Id: hal-02981952

<https://hal.sorbonne-universite.fr/hal-02981952v1>

Submitted on 28 Oct 2020 (v1), last revised 1 Mar 2023 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Final size and convergence rate for an epidemic in heterogeneous population

Luis Almeida* Pierre-Alexandre Bliman† Grégoire Nadin* Benoît Perthame*‡
Nicolas Vauchelet§

October 29, 2020

Abstract

We formulate a general SEIR epidemic model in a heterogeneous population characterized by some trait in a discrete or continuous subset of a space \mathbb{R}^d . The incubation and recovery rates governing the evolution of each homogeneous subpopulation depend upon this trait, and no restriction is assumed on the contact matrix that defines the probability for an individual of a given trait to be infected by an individual with another trait. Our goal is to derive and study the final size equation fulfilled by the limit distribution of the population. We show that this limit exists and satisfies the final size equation. The main contribution is to prove the uniqueness of this solution among the distributions smaller than the initial condition. We also establish that the dominant eigenvalue of the next-generation operator (whose initial value is equal to the basic reproduction number) decreases along every trajectory until a limit smaller than 1. The results are shown to remain valid in presence of diffusion term. They generalize previous works corresponding to finite number of traits (including metapopulation models) or to rank 1 contact matrix (modeling e.g. susceptibility or infectivity presenting heterogeneity independently of one another).

2010 *Mathematics Subject Classification.*: 35Q92; 35R09; 45C05; 45G15; 92D30

Keywords: SIR model; Epidemic spread; Structured population equations; Next generation operator; Long term asymptotics;

1 Introduction

Observing the complex behaviour of the COVID-19 [6, 7, 9, 10, 13, 1] reveals the importance of the population heterogeneity in the dynamics of an outbreak. Inspired by this question, but in contrast with recent papers aiming at reproducing the dynamics of the pandemic observed through various data, we adopt here an abstract point of view and investigate issues related to the concepts of epidemic final size and herd immunity in an ample setting.

*Sorbonne Université, CNRS, Université de Paris, Inria, Laboratoire Jacques-Louis Lions UMR7598, F-75005 Paris, France. Email : {almeida,gregoire.nadin,benoit.perthame}@ljl.math.upmc.fr

†Sorbonne Université, Inria, CNRS, Université de Paris, Laboratoire Jacques-Louis Lions UMR7598, Equipe MAMBA, F-75005 Paris, France. Email : pierre-alexandre.bliman@inria.fr

‡B.P. has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 740623).

§LAGA, UMR 7539, CNRS, Université Sorbonne Paris Nord, 99 avenue Jean-Baptiste Clément, 93430 Villetaneuse, France. Email : vauchelet@math.univ-paris13.fr

Before explaining in more details the context and contribution, we present the system under study. We consider here the general SEIR epidemic model (1), which describes the spread of a disease in a heterogeneous population characterized by a “trait” $x \in \Omega$, for a given open subset $\Omega \subset \mathbb{R}^d$. The underlying structure described by this trait can be very varied, typical examples being one or several of the following characteristics: propensity to have social contacts and hazardousness of the latter (in the sense of the transmission disease), susceptibility, infectivity, characteristics of the immunological response, age, spatial location, etc.

$$\partial_t S(t, x) = -S(t, x) \int_{\Omega} \beta(x, y) I(t, y) dy, \quad S(0, x) = S_0(x) \quad (1a)$$

$$\partial_t E(t, x) = S(t, x) \int_{\Omega} \beta(x, y) I(t, y) dy - \alpha(x) E(t, x), \quad E(0, x) = E_0(x), \quad (1b)$$

$$\partial_t I(t, x) = \alpha(x) E(t, x) - \gamma(x) I(t, x), \quad I(0, x) = I_0(x), \quad (1c)$$

$$\partial_t R(t, x) = \gamma(x) I(t, x), \quad R(0, x) = R_0(x). \quad (1d)$$

Classical notations are used : S denotes the proportion of susceptible individuals, E the proportion of those who have been exposed to the disease, I the proportion of infected individuals, R the proportion of removed individuals. The component $\beta(x, y)$ of the so-called (infinite-dimensional) *contact matrix* β is the product of the average contact rate between individuals of traits x and y and average transmission probability from an individual with trait y to individual with trait x . The force of infection $\int_{\Omega} \beta(x, y) I(t, y) dy$ that applies to the individuals of trait x thus obeys the law of mass-action. The average incubation rate is denoted $\alpha(x)$ and the average recovery rate is denoted $\gamma(x)$: both quantities may depend upon the trait x .

As can be seen, the dynamics of the outbreak is assumed to be sufficiently fast to neglect the effects of the demography in the population. A particular case of system (1) is when the parameters α, β, γ are independent of the trait: the classical (“homogeneous”) SEIR model, which reads

$$\dot{S} = -\beta SI, \quad \dot{E} = \beta SI - \alpha E, \quad \dot{I} = \alpha E - \gamma I \quad (2)$$

is then recovered after integration over Ω . A variant of (1) with diffusion operator in the infected compartment capable to account for possible mutations or spatial displacement of the infected population is also considered, see equation (7) below.

For an epidemic model where the effects of demography are neglected, the number of exposed and infected individuals is expected to go to zero after reaching an epidemic peak (if the initial proportion of susceptible is sufficient). An important question is then to determine what will be the so-called *epidemic final size*, that is the cumulative number (or proportion) of individuals infected, and thus ultimately removed, during the outbreak. For the homogeneous SEIR model (2), the answer to this question is well-known. Indeed, due to the invariance of the function $S(t) + E(t) + I(t) - \frac{\gamma}{\beta} \ln S(t)$ along the trajectory, the asymptotic value S_{∞} of $S(t)$ is necessarily a root of the equation $S_{\infty} - \frac{\gamma}{\beta} \ln S_{\infty} = S_0 + E_0 + I_0 - \frac{\gamma}{\beta} \ln S_0$. Indeed, one may show that it is the unique root of this equation smaller than the so-called *herd immunity threshold* $\frac{\gamma}{\beta}$ —the value of S below which the number $E + I$ of exposed and infected necessarily decreases.

The study of the final size of an epidemic has been addressed in several works since its appearance in early contributions [17, 4]. Reference [19] was among the first papers to unveil the generality of this concept, extending it to models with several distinct infectious stages, or with arbitrarily distributed mean contact rate, or again having spatially heterogeneous contact structures. Multiple susceptible classes were considered in [3]. Reference [5] established final size equation for age-of-infection model.

Distributed susceptibility or distributed infectivity has been studied in [22], and [2] studied an epidemic model within a population of n different groups. Analysis in a general framework was presented in [21], establishing sufficient conditions under which epidemic final size equation exists. Sharp estimates and bounds are obtained in [15] for general model with heterogeneous susceptibility. Also, the case of a two-group SIR model has been studied in [20].

An important quantity to describe the dynamics of an epidemic is the *basic reproduction number* \mathcal{R}_0 . From a biological point of view, the latter represents the expected number of secondary cases directly generated by an individual in a completely susceptible population during its entire period of infectiousness. A famous threshold property states that the disease can invade (in the sense that it will spread and reach an epidemic peak) if $\mathcal{R}_0 > 1$ and the initial number of susceptible is sufficient, whereas it cannot if $\mathcal{R}_0 < 1$. Mathematically, this number has been defined in full generality in the seminal paper [8] as the dominant eigenvalue of the so-called *next-generation operator*.

Our aim in the present paper is to extend some of these previous results and completely characterize the final size of an epidemic in a heterogeneous population whose dynamics is governed by the general systems (1) or (7). On this occasion, we revisit and present in a unified setting the notions of herd immunity and basic reproductive number, and demonstrate the centrality and powerfulness of the concept of next-generation operator.

The outline of this work is the following. We present in Section 2 the assumptions used in the sequel, and provide the key results, Theorems 2.1 and 2.2. They correspond respectively to epidemic spreading without and with diffusion. Their proof is given in Sections 3 and 4. Section 5 shows how the present framework includes several examples previously considered in the literature.

2 Assumptions and main results

Heterogeneous populations without diffusion. To begin with, we investigate the final size of an epidemic in a heterogeneous population whose dynamics is modelled by system (1). Denote $L_+^\infty(\Omega)$ the set of measurable functions which are bounded and nonnegative a.e. on Ω , and assume that

$$\begin{cases} \alpha \in L_+^\infty(\Omega) & \text{and} & \underline{\alpha} := \inf_{x \in \Omega} \alpha(x) > 0 & \beta \in L_+^\infty \cap L^2(\Omega \times \Omega), & \text{and} & \beta > 0, \\ \gamma \in L_+^\infty(\Omega) & \text{and} & \underline{\gamma} := \inf_{x \in \Omega} \gamma(x) > 0. \end{cases} \quad (3)$$

In order for the disease to start, we assume also that the initial proportions of susceptible individuals, and of infected and exposed individuals, are nonzero :

$$S_0, E_0, I_0, R_0 \in L_+^\infty(\Omega), \quad S_0 > 0, \quad E_0 + I_0 \neq 0, \quad \int_{\Omega} (S_0 + E_0 + I_0 + R_0) dx = 1. \quad (4)$$

The normalization assumption contained in the last identity materializes the fact that the variables constitute proportions of a given population whose overall value is conserved along time.

The first result concerns system (1).

Theorem 2.1 *Assume that assumptions (3) and (4) hold, then for the solution of problem (1), $t \mapsto S(t, \cdot)$ is a decreasing function and there exists $S_\infty \in L_+^\infty(\Omega) \cap L^1(\Omega)$ such that, for a.e. $x \in \Omega$,*

$$\lim_{t \rightarrow +\infty} S(t, x) = S_\infty(x), \quad \text{and} \quad \lim_{t \rightarrow +\infty} I(t, x) = \lim_{t \rightarrow +\infty} E(t, x) = 0, \quad (5)$$

with exponential rates. Moreover, S_∞ is the unique solution such that $S_\infty \leq S_0$ of the final size equation

$$\ln S_\infty(x) - \int_{\Omega} \frac{\beta(x, y)}{\gamma(y)} S_\infty(y) dy = \ln S_0(x) - \int_{\Omega} \frac{\beta(x, y)}{\gamma(y)} (S_0(y) + E_0(y) + I_0(y)) dy. \quad (6)$$

An alternative form of (6), useful in the sequel, may be expressed as follows:

$$S_\infty(x) = S_0(x) \exp \left(\int_\Omega \frac{\beta(x, y)}{\gamma(y)} (S_\infty(y) - (S_0(y) + E_0(y) + I_0(y))) dy \right).$$

The proof of Theorem 2.1 is the subject of Section 3. The main difficulty in this demonstration is to prove uniqueness of the solution of the final size equation (6), in the set of those S at most equal to the initial condition S_0 . It uses as central tool an irreducibility property related to the assumption $\beta > 0$. It is possible to relax this hypothesis by only assuming that the next-generation operator (defined in (13) below) is strongly positive —this is indeed the case if $\beta > 0$. On the contrary, $\beta \geq 0$ is not sufficient to deduce uniqueness, see a counterexample at the end of the present Section.

Heterogeneous populations with diffusion. Diffusion is introduced in the infected compartment for the second result, allowing to consider possible additional natural phenomenon as mutations or spatial diffusion of infected individuals. There is a long history on this problem, in particular to study the effect of diffusion on the basic reproduction number, the spread or extinction of the epidemic and the existence of propagating waves, see [16, 23, 11, 12] and the references therein. Here, we are interested in characterizing the final size of the epidemic in a bounded domain.

The corresponding system reads, for $t \geq 0$, $x \in \Omega$,

$$\partial_t S(t, x) = -S(t, x) \int_\Omega \beta(x, y) I(t, y) dy, \quad S(0, x) = S_0(x), \quad (7a)$$

$$\partial_t E(t, x) = S(t, x) \int_\Omega \beta(x, y) I(t, y) dy - \alpha(x) E(t, x), \quad E(0, x) = E_0(x), \quad (7b)$$

$$\begin{cases} \partial_t I(t, x) = \alpha(x) E(t, x) - \gamma(x) I(t, x) + \Delta I(t, x), & I(0, x) = I_0(x), \\ \partial_n I(t, x) = 0 & \text{on } (0, \infty) \times \partial\Omega, \end{cases} \quad (7c)$$

$$\partial_t R(t, x) = \gamma(x) I(t, x), \quad R(0, x) = R_0(x). \quad (7d)$$

With the initial data, one can build the solution of

$$\begin{cases} -\Delta \Phi_0 + \gamma \Phi_0 = S_0 + E_0 + I_0, \\ \partial_n \Phi_0 = 0 & \text{on } (0, \infty) \times \partial\Omega. \end{cases} \quad (8)$$

Essentially the same assumptions than Theorem 2.1 allow to obtain the similar result.

Theorem 2.2 *Assume that (3) and (4) hold. Assume moreover that Ω is a smooth, bounded open set. Then, for the solution of (7), $t \mapsto S(t, \cdot)$ is a nondecreasing function and there exists $S_\infty \in L_+^\infty(\Omega) \cap L^1(\Omega)$ such that (5) holds for a.e. $x \in \Omega$, with exponential rates.*

Moreover, S_∞ is characterized by

$$S_\infty = S_0 \exp \left(\int_\Omega \beta(x, y) (\Phi_\infty(y) - \Phi_0(y)) dy \right), \quad (9a)$$

where Φ_∞ is the unique solution such that $\Phi_\infty \leq \Phi_0$ (with Φ_0 defined in (8)) of

$$\begin{cases} -\Delta \Phi_\infty + \gamma(x) \Phi_\infty = S_0 \exp \left(\int_\Omega \beta(x, y) (\Phi_\infty(y) - \Phi_0(y)) dy \right), \\ \partial_n \Phi_\infty = 0 & \text{on } (0, \infty) \times \partial\Omega. \end{cases} \quad (9b)$$

Equation (9) constitutes, together with (8), the final size equation for equation (7). It appears analogous to (6) when ignoring the diffusion term. It may also be written as

$$-\Delta\Phi_\infty + \gamma\Phi_\infty = S_\infty.$$

Again, we can relax the hypothesis $\beta > 0$ by only assuming that the modified next-generation operator (defined in (25) below) is strongly positive, which is of course true when $\beta > 0$. Another possible set of assumption guaranteeing this strong positivity, and thus the availability of Theorem 2.2, is $\beta \geq 0$ and Ω connected. In that case, even if $\beta = 0$ in some parts of Ω , the diffusion ensures that the infection reaches such parts. If β is not positive and Ω is not connected, it is easy to construct counter-examples where a connected part of Ω is invaded whereas another is not.

Finally, we mention that these results may be adapted straightforwardly when the exposed compartment is neglected, that is for the heterogeneous SIR model. Also they may be extended to other operators than the diffusion, as the fractional Laplacian or integral operators, as long as a standard spectral theory is available. Last, diffusion could be included on the Exposed compartment with the same results and methods.

Nonuniqueness and positivity of β In order to illustrate the necessity of assumption $\beta > 0$ in (3), we provide a simple example for the discrete case with piecewise constant coefficient. More precisely, assume that for some disjoint sets Ω_1, Ω_2 for which $\Omega = \Omega_1 \cup \Omega_2$, and some positive constants $\alpha_k, \beta_k, \gamma_k$, $k = 1, 2$ and β_{12} , the coefficients verify:

$$\begin{aligned} \alpha(x) &= \alpha_1 \mathbf{1}_{\Omega_1}(x) + \alpha_2 \mathbf{1}_{\Omega_2}(x), & \gamma(x) &= \gamma_1 \mathbf{1}_{\Omega_1}(x) + \gamma_2 \mathbf{1}_{\Omega_2}(x), \\ \beta(x, y) &= \begin{cases} \beta_k, & \text{if } (x, y) \in \Omega_k \times \Omega_k, \quad k = 1, 2 \\ \beta_{12}, & \text{if } (x, y) \in \Omega_1 \times \Omega_2 \\ 0, & \text{if } (x, y) \in \Omega_2 \times \Omega_1. \end{cases} \end{aligned}$$

Then, denoting

$$S_k(t) := \int_{\Omega_k} S(t, x) dx, \quad E_k(t) := \int_{\Omega_k} E(t, x) dx, \quad I_k(t) := \int_{\Omega_k} I(t, x) dx$$

for $k = 1, 2$, system (1) implies (after integration in space) that

$$\begin{aligned} \dot{S}_1 &= -S_1(\beta_1 I_1 + \beta_{12} I_2), & \dot{S}_2 &= -\beta_2 S_2 I_2, \\ \dot{E}_1 &= S_1(\beta_1 I_1 + \beta_{12} I_2) - \alpha_1 E_1, & \dot{E}_2 &= \beta_2 S_2 I_2 - \alpha_2 E_2, \\ \dot{I}_1 &= \alpha_1 E_1 - \gamma_1 I_1, & \dot{I}_2 &= \alpha_2 E_2 - \gamma_2 I_2, \end{aligned}$$

complemented with initial data S_k^0, E_k^0, I_k^0 , for $k = 1, 2$ (with the notations $S_k^0 = \int_{\Omega_k} S_0(x) dx$, and so on). Denoting $S_{k\infty}$ the limit of $S_k(t)$ when $t \rightarrow +\infty$, $k = 1, 2$, the final state is characterized, in this particular setting, by the system of two scalar identities

$$\ln S_{1\infty} - \frac{\beta_1}{\gamma_1} S_{1\infty} - \frac{\beta_{12}}{\gamma_2} S_{2\infty} = \ln S_1^0 - \frac{\beta_1}{\gamma_1} (S_1^0 + E_1^0 + I_1^0) - \frac{\beta_{12}}{\gamma_2} (S_2^0 + E_2^0 + I_2^0), \quad (10a)$$

$$\ln S_{2\infty} - \frac{\beta_2}{\gamma_2} S_{2\infty} = \ln S_2^0 - \frac{\beta_2}{\gamma_2} (S_2^0 + E_2^0 + I_2^0). \quad (10b)$$

If $E_2^0 + I_2^0 = 0$, then (10b) admits two solutions, namely S_2^0 (corresponding to absence of disease in Ω_2), and another solution, denoted \underline{S}_2 , which is strictly smaller than S_2^0 (corresponding to a spread of the epidemic in Ω_2). If now $E_1^0 + I_1^0 > 0$, for each of these solutions, solving (10a) for $S_{1\infty} \leq S_1^0$

yields a unique solution. We thus obtain two distinct solutions of equation (10), smaller or equal to (S_0^1, S_0^2) . Notice that, for any initial condition of (6) constant on each of the two sets Ω_1, Ω_2 , the solution itself retains this property. The previous considerations thus allows to exhibit for such a choice of initial condition, two distinct solutions of the final size equation (6). As a conclusion, the uniqueness property stated in Theorem 2.1 may not hold if $\beta \not\equiv 0$.

3 General contact matrices in absence of diffusion

We investigate here system (1). After introducing some adequate notions and results, Theorem 2.1 is proved. Some conserved quantities are first studied in Section 3.1. The notions of *basic reproduction number* and *herd immunity domain* are then introduced in this general setting in Section 3.2, based on the next-generation operator. The limit behaviour is established in Section 3.3, together with the properties of the solution of the final size equation, formally achieving the proof of Theorem 2.1. Section 3.4 then exploits the tools introduced during this section to disentangle the link between herd immunity and equilibrium stability.

3.1 Conserved quantities and long time limit

A central role in the dynamics is played by the integral quantity

$$\Phi(t, x) = \int_{\Omega} \frac{\beta(x, y)}{\gamma(y)} [S(t, y) + E(t, y) + I(t, y)] dy, \quad (11)$$

extending a similar and standard quantity for the SIR model.

Lemma 3.1 *Under assumption (3), the solution of (1) verifies that, for a.e. $x \in \Omega$, the quantities $S(t, x) + E(t, x) + I(t, x) + R(t, x)$ and $\ln S(t, x) - \Phi(t, x)$ are constant with respect to time.*

Proof. Firstly, adding the equations in (1) provides directly the conservation of $S + E + I + R$ with respect to time. Secondly, we compute

$$\begin{aligned} \frac{\partial}{\partial t} [\ln S(t, x) - \Phi(t, x)] &= \frac{\dot{S}(t, x)}{S(t, x)} - \int_{\Omega} \frac{\beta(x, y)}{\gamma(y)} (\dot{S}(t, y) + \dot{E}(t, y) + \dot{I}(t, y)) dy \\ &= - \int_{\Omega} \beta(x, y) I(t, y) dy + \int_{\Omega} \beta(x, y) I(t, y) dy = 0. \end{aligned}$$

We thus deduce the conservation relation

$$\frac{\partial}{\partial t} [\ln S(t, x) - \Phi(t, x)] = 0, \quad (12)$$

and Lemma 3.1 is proved. \square

Lemma 3.2 *Assume that (3) and (4) hold. Then the solution of (1) verifies : there exists $S_{\infty} \in L_{+}^{\infty}(\Omega)$, $S_{\infty} \leq S_0$ such that, for a.e. $x \in \Omega$, (5) holds.*

Proof. We deduce easily from (1) that for a.e. $x \in \Omega$, $t \mapsto S(t, x)$ is decreasing. Being bounded from below by 0, it admits a limit as t goes to $+\infty$. By the same token, $S(\cdot, x) + E(\cdot, x)$ and $S(\cdot, x) + E(\cdot, x) + I(\cdot, x)$ admit similarly a limit. As $S(\cdot, x)$ admits a limit, we deduce that $E(\cdot, x)$ and $I(\cdot, x)$ equally converge when $t \rightarrow +\infty$. Passing to the limit into system (1), one concludes that these two limits are null, which achieves the demonstration of (5), and thus of Lemma 3.2. \square

3.2 Herd immunity and basic reproduction number

Recall that (3) and (4) have been assumed. Following the idea in [8], we define now a key notion.

Definition 1 For any $S(\cdot) \in L^1 \cap L^\infty_+(\Omega)$, we call next-generation operator $K_{S(\cdot)}$ associated to (1), the operator defined over $L^2(\Omega)$ by:

$$K_{S(\cdot)}[\phi](x) = S(x) \int_{\Omega} \frac{\beta(x, y)}{\gamma(y)} \phi(y) dy. \quad (13)$$

Under assumptions (3) and (4), we know that, for any $t \geq 0$, the solution $S(t, \cdot)$ of (1) belongs to $L^\infty_+(\Omega) \cap L^1(\Omega)$. The operator $K_{S(t, \cdot)}$ is thus well-defined and it is a positive and compact operator over $L^2(\Omega)$. This permits to define its norm and its spectral radius in the usual way recalled now.

Definition 2 The spectral radius $r(K_{S(t, \cdot)})$ of $K_{S(t, \cdot)}$ is defined by

$$r(K_{S(t, \cdot)}) = \lim_{n \rightarrow +\infty} \|K_{S(t, \cdot)}^n\|^{1/n}.$$

As β is irreducible thanks to (3) and using Krein-Rutman theory [18, 26], we can identify the spectral radius $r(K_{S(t, \cdot)})$ of $K_{S(t, \cdot)}$ with its principal eigenvalue. The latter is simple, and defined as the unique eigenvalue associated to a positive eigenfunction. We denote $\varphi_{S(t, \cdot)}$ this eigenfunction, so that by definition,

$$K_{S(t, \cdot)}[\varphi_{S(t, \cdot)}](x) = S(t, x) \int_{\Omega} \frac{\beta(x, y)}{\gamma(y)} \varphi_{S(t, \cdot)}(y) dy = r(K_{S(t, \cdot)}) \varphi_{S(t, \cdot)}(x).$$

We also know that $r(K_{S(t, \cdot)}) = r(K_{S(t, \cdot)}^*)$, where $K_{S(t, \cdot)}^*$ is the adjoint operator, defined by

$$\langle K_{S(t, \cdot)}^*[\psi], \phi \rangle = \langle \psi, K_{S(t, \cdot)}[\phi] \rangle = \int_{\Omega} \psi(x) S(t, x) \int_{\Omega} \frac{\beta(x, y)}{\gamma(y)} \phi(y) dy dx$$

that is

$$K_{S(t, \cdot)}^*[\psi](y) := \int_{\Omega} S(t, x) \frac{\beta(x, y)}{\gamma(y)} \psi(x) dx. \quad (14)$$

We let $\psi_{S(t, \cdot)}$ be the associated normalized adjoint eigenfunction, defined by

$$K_{S(t, \cdot)}^*[\psi_{S(t, \cdot)}] = r(K_{S(t, \cdot)}) \psi_{S(t, \cdot)}, \quad \text{and} \quad \int_{\Omega} \psi_{S(t, \cdot)}(x) S_0(x) dx = 1.$$

We are now in position to define for (1) the key notions of basic reproduction number and herd immunity, based on the definition of next-generation operator in (13).

Definition 3 Under assumptions (3) and (4), we define:

- basic reproduction number \mathcal{R}_0 the spectral radius of the operator $K_{S_0(\cdot)}$, i.e., $\mathcal{R}_0 = r(K_{S_0(\cdot)})$;
- herd immunity domain the set of those elements \bar{S} of $L^\infty_+(\Omega) \cap L^1(\Omega)$ such that $r(K_{\bar{S}(\cdot)}) \leq 1$.

In contrast to the homogeneous SEIR system where herd immunity is reached for a uniquely defined proportion of the variable S , here the boundary of the herd immunity domain is a set of functions corresponding to various susceptible population distributions. It is hard to determine in which point of the boundary the trajectory will enter this domain, but this necessarily takes place after a certain time, as stated now.

Lemma 3.3 *Assume that (3) and (4) hold, and that $r(K_{S_0(\cdot)}) > 1$. Then, $t \mapsto r(K_{S(t,\cdot)})$ is decreasing and continuous with*

$$\lim_{t \rightarrow +\infty} r(K_{S(t,\cdot)}) < 1, \quad (15)$$

and there exists a unique $T_0 > 0$ such that $r(K_{S(T_0,\cdot)}) = 1$. As a consequence $S(t,\cdot)$ belongs to the herd immunity domain iff $t \geq T_0$.

Proof. We first establish that $t \mapsto r(K_{S(t,\cdot)})$ is non-increasing and continuous. This is a consequence of the fact that for a.e. $x \in \Omega$, $t \mapsto S(t,x)$ is decreasing and continuous. Indeed, by definition (13), for any $\phi \in L^2(\Omega)$, and any $n \in \mathbb{N}^*$, $t \mapsto \|K_{S(t,\cdot)}^n[\phi]\|$ is decreasing and continuous. Hence, $t \mapsto \|K_{S(t,\cdot)}^n\|^{1/n}$ is decreasing and continuous for any $n \in \mathbb{N}^*$. The result follows by passing to the limit $n \rightarrow +\infty$.

Secondly, since $r(K_{S_0(\cdot)}) > 1$, we are left to prove that (15) holds. From Lemma 3.2, we may use the fact that $S_\infty(x)$ is the limit of $S(t,x)$ when $t \rightarrow +\infty$, for a.e. $x \in \Omega$. Denote simply $K_\infty := K_{S_\infty(\cdot)}$ the associated next-generation operator, and by ψ_∞ the principal eigenfunction associated to the adjoint operator K_∞^* .

Adding the second and third equations in (1), multiplying the result by ψ_∞ and integrating, one obtains from (14) taken at the limit,

$$\begin{aligned} & \frac{d}{dt} \int_{\Omega} \psi_\infty(x) (E(t,x) + I(t,x)) dx \\ &= \int_{\Omega \times \Omega} \psi_\infty(x) S(t,x) \beta(x,y) I(t,y) dy dx - \int_{\Omega} \gamma(x) \psi_\infty(x) I(t,x) dx \\ &\geq \int_{\Omega \times \Omega} \psi_\infty(x) S_\infty(x) \beta(x,y) I(t,y) dy dx - \int_{\Omega} \gamma(x) \psi_\infty(x) I(t,x) dx \\ &= \langle K_\infty^*[\psi_\infty], \gamma(\cdot) I(t,\cdot) \rangle - \langle \psi_\infty, \gamma(\cdot) I(t,\cdot) \rangle \\ &= (r(K_\infty^*) - 1) \int_{\Omega} \psi_\infty(y) \gamma(y) I(t,y) dy. \end{aligned}$$

Hence, arguing by contradiction, if $r(K_\infty^*) \geq 1$, then, on the one hand $t \mapsto \int_{\Omega} \psi_\infty(x) (E(t,x) + I(t,x)) dx$ is non-decreasing. On the other hand, we have seen in Lemma 3.2 that $\lim_{t \rightarrow +\infty} (E(t,x) + I(t,x)) = 0$. It is a contradiction. Thus, $r(K_\infty^*) < 1$ and by continuity there exists T_0 such that $r(K_{S(T_0,\cdot)}) = 1$. This achieves the proof of Lemma 3.3. \square

As is the case for the homogenous SEIR model, there exists a tight relation between herd immunity and stability of the equilibrium points of (1). This link is elucidated in Section 3.4 below.

3.3 Long time behaviour and final size equation – Proof of Theorem 2.1

We are now in position to prove Theorem 2.1.

Step 1. Existence of limits. In Lemma 3.2 we already proved the decay of $t \mapsto S(t,x)$ and existence of limits for all variables S, E, I, R . We are left to identify the limit S_∞ .

Integrating (12) between t and $+\infty$, we deduce that the limit $S_\infty(x)$ satisfies necessarily

$$\begin{cases} \ln S_\infty(x) - \Phi_\infty(x) = \ln S(t,x) - \int_{\Omega} \frac{\beta(x,y)}{\gamma(y)} [S(t,y) + E(t,y) + I(t,y)] dy, \\ \Phi_\infty(x) = \int_{\Omega} \frac{\beta(x,y)}{\gamma(y)} S_\infty(y) dy, \end{cases} \quad (16)$$

for any $t \geq 0$. Taking $t = 0$, we obtain that S_∞ is solution of (6).

Step 2. Exponential rate of convergence. From the above formula, and using the dominated convergence theorem, we have

$$\ln \frac{S(t, x)}{S_\infty(x)} = \int_\Omega \frac{\beta(x, y)}{\gamma(y)} [S_\infty(x) - S(t, y) - E(t, y) - I(t, y)] dy \rightarrow 0 \quad \text{as } t \rightarrow \infty.$$

Therefore, for all $\varepsilon > 0$ there is a T_ε such that $S(t) \leq (1 + \varepsilon)S_\infty$ for $t \geq T_\varepsilon$. We also introduce $\theta \in (0, 1)$, to be chosen later, and compute

$$\frac{d}{dt} [E(t, x) + \theta I(t, x)] \leq (1 + \varepsilon)S_\infty(x) \int_\Omega \beta(x, y) I(t, y) dy - (1 - \theta)\underline{\alpha}E(t, x) - \theta\gamma(x)I(t, x), \quad (17)$$

where $\underline{\alpha} > 0$ has been defined in (3). Multiplying again by ψ_∞ , the principal eigenfunction associated to the adjoint operator $K_\infty^* = K_{S_\infty(\cdot)}^*$ introduced in the proof of Lemma 3.3, and using (14), we find for $t \geq T_\varepsilon$

$$\begin{aligned} & \frac{d}{dt} \int_\Omega \psi_\infty(x) [E(t, x) + \theta I(t, x)] dx \\ & \leq (1 + \varepsilon) \int_{\Omega \times \Omega} \psi_\infty(x) S_\infty(x) \beta(x, y) I(t, y) dy dx - \int_\Omega \psi_\infty(x) [(1 - \theta)\underline{\alpha}E(t, x) + \theta\gamma(x)I(t, x)] dx \\ & \leq (1 + \varepsilon)r(K_\infty) \int_\Omega \psi_\infty(y) \gamma(y) I(t, y) dy - \int_\Omega \psi_\infty(x) [(1 - \theta)\underline{\alpha}E(t, x) + \theta\gamma(x)I(t, x)] dx. \end{aligned}$$

Since we know from Lemma 3.3 that $r(K_\infty) < 1$, it remains to choose θ sufficiently close to 1 and ε small enough such that $1 > \theta > (1 + \varepsilon)r(K_\infty)$, in such a way that

$$\lambda := \min \left((1 - \theta)\underline{\alpha}, \left(1 - \frac{(1 + \varepsilon)r(K_\infty)}{\theta} \right) \underline{\gamma} \right) > 0.$$

We obtain

$$\frac{d}{dt} \int_\Omega \psi_\infty(x) [E(t, x) + \theta I(t, x)] dx \leq -\lambda \int_\Omega \psi_\infty(x) [E(t, x) + \theta I(t, x)] dx,$$

and, applying Gronwall lemma, we find exponential decay to 0 with rate λ for $I(t, x)$ and $E(t, x)$.

Finally, using the fact that $S(t, x)$ decreases along time, equation (7a) tells us that

$$\begin{aligned} 0 \leq S(t, x) - S_\infty(x) &= \int_t^\infty S(s, x) \int_\Omega \beta(x, y) I(s, y) dy ds \\ &\leq \frac{S^0(x) \|\beta\|_{L^\infty}}{\inf \psi_\infty} \int_\Omega \psi_\infty(y) I(s, y) dy ds = O(e^{-\lambda t}). \end{aligned}$$

Step 3. Contraction property for t large enough. From Lemma 3.3, there exists $T > 0$ such that $r(K_{S(T, \cdot)}) < 1$. From the first line of equation (16), we deduce the expression for S_∞

$$S_\infty(x) = S(T, x) \exp(\Phi_\infty(x) - \Phi(T, x)),$$

where we use the notation (11). Injecting this expression into the second line of (16), yields the following fixed point problem for the function Φ_∞ :

$$\Phi_\infty(x) = \int_\Omega \frac{\beta(x, y)}{\gamma(y)} S(T, y) \exp(\Phi_\infty(y) - \Phi(T, y)) dy. \quad (18)$$

Notice that since $t \mapsto S(t, x)$ is decreasing from (1a), we deduce that $S_\infty \leq S(T, \cdot)$ and $\Phi_\infty \leq \Phi(T, \cdot)$.

Introduce

$$\mathcal{F}(u)(x) := \int_{\Omega} \frac{\beta(x, y)}{\gamma(y)} S(T, y) \exp(u(y) - \Phi(T, y)) dy,$$

in such a way that the fixed point problem (18) reads $\Phi_\infty = \mathcal{F}(\Phi_\infty)$. Let us denote

$$\mathcal{K} := \{u \in L^1(\Omega), \text{ such that } 0 \leq u \leq \Phi(T, \cdot) \text{ a.e.}\}.$$

Clearly, \mathcal{K} is stable by \mathcal{F} (using the definition of Φ in (11)), and we may prove that \mathcal{F} is a contraction on the set \mathcal{K} . Indeed, we compute, for any u, v in \mathcal{K} ,

$$\begin{aligned} |\mathcal{F}(u) - \mathcal{F}(v)|(x) &\leq \int_{\Omega} \frac{\beta(x, y)}{\gamma(y)} S(T, y) \exp(-\Phi(T, y)) |\exp(u(y)) - \exp(v(y))| dy \\ &\leq \int_{\Omega} \frac{\beta(x, y)}{\gamma(y)} S(T, y) |u(y) - v(y)| dy, \end{aligned} \quad (19)$$

where the last inequality is a consequence of the definition of \mathcal{K} and of the fact that the exponential is 1-Lipschitz in the set of nonpositive real numbers.

At this stage, we introduce on the set \mathcal{K} the norm

$$\|u\|_T := \int_{\Omega} \psi_{S(T, \cdot)}(x) S(T, x) |u(x)| dx,$$

where, as defined in (14), $\psi_{S(T, \cdot)}$ is the principal eigenfunction of the adjoint $K_{S(T, \cdot)}^*$. Multiplying by $\psi_{S(T, \cdot)}$ and $S(T, \cdot)$, we get with the help of (19), that

$$\begin{aligned} \|\mathcal{F}(u) - \mathcal{F}(v)\|_T &= \int_{\Omega} \psi_{S(T, \cdot)}(x) S(T, x) |\mathcal{F}(u)(x) - \mathcal{F}(v)(x)| dx \\ &\leq \int_{\Omega \times \Omega} \psi_{S(T, \cdot)}(x) \frac{\beta(x, y)}{\gamma(y)} S(T, x) S(T, y) |u(y) - v(y)| dy dx \\ &= r(K_{S(T, \cdot)}) \int_{\Omega} \psi_{S(T, \cdot)}(y) S(T, y) |u - v|(y) dy = r(K_{S(t, \cdot)}) \|u - v\|_T, \end{aligned}$$

by definition of the norm $\|\cdot\|_T$. As $r(K_{S(T, \cdot)}) < 1$, we deduce that \mathcal{F} is a contraction on \mathcal{K} . Hence, using Banach contraction mapping principle, there exists a unique solution S_∞ to (18) in \mathcal{K} .

Step 4. Uniqueness below S_0 . It remains to extend the uniqueness result to all function smaller or equal to S_0 . We make use of the following result.

Lemma 3.4 *Among the functions $S \leq S_0$, there is a solution \bar{S} of (6) which is larger than all sub-solutions of (6), and $S_\infty = \bar{S}$.*

Indeed, we deduce from Lemma 3.4 that any other solution \underline{S} of (6) satisfies $\underline{S} \leq S_\infty$. Therefore it also satisfies $\underline{S} \leq S(T)$ for T large enough. Thus by the contraction property of Step 3 it is equal to S_∞ . This achieves the demonstration of Theorem 2.1, with the exception of the proof of Lemma 3.4, which we now complete.

Proof. [Proof of Lemma 3.4] We first build a sequence $S_k(x)$ departing from the initial condition S_0 , by iterating for $k \geq 0$:

$$\ln S_{k+1}(x) = \int_{\Omega} \frac{\beta(x, y)}{\gamma(y)} S_k(y) dy + A(x), \quad A(x) = \ln S_0(x) - \int_{\Omega} \frac{\beta(x, y)}{\gamma(y)} [S_0 + E_0 + I_0](y) dy.$$

Obviously, we have

$$\ln S_1(x) = \int_{\Omega} \frac{\beta(x, y)}{\gamma(y)} S_0(y) dy + A(x) = \ln S_0(x) - \int_{\Omega} \frac{\beta(x, y)}{\gamma(y)} [E_0 + I_0](y) dy < \ln S_0(x).$$

And iterating, if we know that $S_k < S_{k-1}$, we find

$$\ln S_{k+1}(x) = \int_{\Omega} \frac{\beta(x, y)}{\gamma(y)} S_k(y) dy + A(x) < \int_{\Omega} \frac{\beta(x, y)}{\gamma(y)} S_{k-1}(y) dy + A(x) = \ln S_k(x),$$

which gives $S_{k+1} < S_k$. As a consequence S_k is a decreasing sequence which converges to a solution \bar{S} of (6).

Secondly, let $\underline{S} \leq S_0$ be a sub-solution of (6). We have

$$\ln \underline{S}(x) \leq \int_{\Omega} \frac{\beta(x, y)}{\gamma(y)} \underline{S}(y) dy + A(x) \leq \int_{\Omega} \frac{\beta(x, y)}{\gamma(y)} S_0(y) dy + A(x) = \ln S_1(x),$$

therefore $\underline{S} \leq S_1$, and iterating the argument $\underline{S} \leq S_k$ for all $k \in \mathbb{N}$ and thus $\underline{S} \leq \bar{S}$. In particular we have $S_{\infty} \leq \bar{S}$.

Thirdly, we prove that $\bar{S} \leq S_{\infty}$. We know that $\bar{S} < S_0$ and we are going to prove that $\bar{S} < S(t)$ for all $t \geq 0$ from which we conclude, by definition of S_{∞} , that $\bar{S} \leq S_{\infty}$ and thus that $S_{\infty} = \bar{S}$. To prove this, we consider, by contradiction, the first time t_0 when $S(t_0)$ touches \bar{S} , i.e., $\bar{S} \leq S(t_0)$ and for some $x_0 \in \Omega$, $\bar{S}(x_0) = S(t_0, x_0)$. We conclude that

$$0 = \ln \bar{S}(x_0) - \ln S(t_0, x_0) = \int_{\Omega} \frac{\beta(x, y)}{\gamma(y)} [\bar{S}(y) - S(t_0, y) - E(t_0, y) - I(t_0, y)] dy < 0,$$

a contradiction. This achieves the demonstration of Lemma 3.4. \square

3.4 Herd immunity and stability of the equilibrium points

Before closing Section 3 and the study of system (1), we analyze in the following result the link between herd immunity and the stability of the equilibria of this system.

Lemma 3.5 *Assume (3) and (4) hold. The equilibrium points of system (1) are exactly the configurations $(S^*, 0, 0, R^*)$ that fulfil (4). Moreover each of them is stable if $r(K_{S^*(\cdot)}) < 1$, unstable if $r(K_{S^*(\cdot)}) > 1$,*

Proof. The equilibria for (1) are any configuration (S^*, E^*, I^*, R^*) such that, for a.e. $x \in \Omega$,

$$\begin{aligned} -S^*(x) \int_{\Omega} \beta(x, y) I^*(y) dy &= 0, & S^*(x) \int_{\Omega} \beta(x, y) I^*(y) dy - \alpha(x) E^*(x) &= 0, \\ \alpha(x) E^*(x) - \gamma(x) I^*(x) &= 0, & \gamma(x) I^*(x) &= 0. \end{aligned}$$

We deduce from the last two equations that $(E^*, I^*) = (0, 0)$. This condition is also sufficient to have an equilibrium, provided that it fulfils (4).

The stability of such an equilibrium is related to the stability of the linearized system

$$\frac{\partial s}{\partial t} = -S^*(x) \int_{\Omega} \beta(x, y) i(t, y) dy, \quad \frac{\partial e}{\partial t} = S^*(x) \int_{\Omega} \beta(x, y) i(t, y) dy - \alpha(x) e(t, x), \quad (20a)$$

$$\frac{\partial i}{\partial t} = \alpha(x) e(t, x) - \gamma(x) i(t, x), \quad \frac{\partial r}{\partial t} = \gamma(x) i(t, x). \quad (20b)$$

Notice that (20) admits a block-triangular structure: the evolution of (e, i) is independent of the other two components s and r ; while the evolution of s and r only depends upon i .

Assume first that $r(K_{S^*(\cdot)}) < 1$. Let $\psi_{S^*(\cdot)}$ be the principal adjoint eigenvector of the next-generation operator $K_{S^*(\cdot)}$ associated to S^* . Mimicking the computation achieved in Step 2 of Section 3.3, one finds that, for any $\theta \in (0, 1)$ and a.e. $x \in \Omega$,

$$\begin{aligned} & \frac{d}{dt} \int_{\Omega} \psi_{S^*(\cdot)}(x)[e(t, x) + \theta i(t, x)] dx \\ &= \int_{\Omega \times \Omega} \psi_{S^*(\cdot)}(x) S^*(x) \beta(x, y) i(t, y) dy - \int_{\Omega} \psi_{S^*(\cdot)}(x) [(1 - \theta) \alpha(x) e(t, x) + \theta \gamma(x) i(t, x)] dx \\ &\leq r(K_{S^*(\cdot)}) \int_{\Omega} \psi_{S^*(\cdot)}(y) \gamma(y) i(t, y) dy - \int_{\Omega} \psi_{S^*(\cdot)}(x) [(1 - \theta) \underline{\alpha} e(t, x) + \theta \gamma(x) i(t, x)] dx. \end{aligned}$$

Taking $\theta \in (r(K_{S^*(\cdot)}), 1)$ then yields (through application of Gronwall's lemma) exponential decrease of $\int_{\Omega} \psi_{S^*(\cdot)}(x)[e(t, x) + \theta i(t, x)] dx$ when $t \rightarrow +\infty$. One sees easily that the subsystem in (e, i) of (20) is a monotone system. The eigenfunction $\psi_{S^*(\cdot)}$ being positive, the functional $(e(\cdot), i(\cdot)) \mapsto \int_{\Omega} \psi_{S^*(\cdot)}(x)[e(\cdot) + \theta i(\cdot)] dx$ is thus a Lyapunov functional for this subsystem, and the origin of the latter is asymptotically stable. Due to the block-triangular structure mentioned earlier, the complete system (20) is (simply) stable.

Assume now that $r(K_{S^*(\cdot)}) > 1$. Denoting $\varphi_{S^*(\cdot)}$ the principal eigenvector of $K_{S^*(\cdot)}$, define

$$e^*(x) := \frac{r(K_{S^*(\cdot)}) + 1}{2} \frac{\varphi_{S^*(\cdot)}(x)}{\alpha(x)}, \quad i^*(x) := \frac{\varphi_{S^*(\cdot)}(x)}{\gamma(x)}.$$

Let us evaluate the value of the right-hand sides corresponding to $\frac{\partial e}{\partial t}$ and $\frac{\partial i}{\partial t}$ at the point (e^*, i^*) . Then, for any $x \in \Omega$,

$$\begin{aligned} & S^*(x) \int_{\Omega} \beta(x, y) i^*(y) dy - \alpha(x) e^*(x) \\ &= S^*(x) \int_{\Omega} \beta(x, y) \frac{\varphi_{S^*(\cdot)}(y)}{\gamma(y)} dy - \frac{r(K_{S^*(\cdot)}) + 1}{2} \varphi_{S^*(\cdot)}(x) = K_{S^*(\cdot)}[\varphi_{S^*(\cdot)}](x) - \frac{r(K_{S^*(\cdot)}) + 1}{2} \varphi_{S^*(\cdot)}(x) \\ &= \frac{r(K_{S^*(\cdot)}) - 1}{2} \varphi_{S^*(\cdot)}(x) > \frac{r(K_{S^*(\cdot)}) - 1}{r(K_{S^*(\cdot)}) + 1} \underline{\alpha} e^*(x), \end{aligned}$$

while similarly

$$\begin{aligned} \alpha(x) e^*(x) - \gamma(x) i^*(x) &= \frac{r(K_{S^*(\cdot)}) + 1}{2} \varphi_{S^*(\cdot)}(x) - \varphi_{S^*(\cdot)}(x) \\ &= \frac{r(K_{S^*(\cdot)}) - 1}{2} \varphi_{S^*(\cdot)}(x) > \frac{r(K_{S^*(\cdot)}) - 1}{2} \underline{\gamma} i^*(x). \end{aligned}$$

The subsystem describing the evolution of the components (e, i) being a monotone system, this condition is sufficient to ensure that any trajectory of this subsystem departing from a point $(e_0, i_0) \geq \lambda(e^*, i^*)$, $\lambda > 0$, is increasing. This shows the instability of the origin of (20), and achieves the proof of Lemma 3.5. \square

4 General contact matrices with diffusion

We now come to system (7) where diffusion is included, and prove Theorem 2.2. We adapt for this the methodology developed in Section 3.

4.1 Conserved quantities and long time limit

Let us introduce the solution of the nonlinear elliptic equation

$$\begin{cases} -\Delta\Phi(t, x) + \gamma\Phi(t, x) = S(t, x) + E(t, x) + I(t, x), & \text{on } \Omega, \\ \partial_n\Phi(t, x) = 0 & \text{on } (0, \infty) \times \partial\Omega, \end{cases} \quad (21)$$

where n is the outward normal unit at the boundary of Ω .

Lemma 4.1 *Under assumption (3), the solution of (7) verifies that, for a.e. $x \in \Omega$, the quantities $\int_{\Omega}(S(t, x) + E(t, x) + I(t, x) + R(t, x)) dx$ and $\ln S(t, x) - \int_{\Omega} \beta(x, y)\Phi(t, y) dy$ are constant with respect to time.*

Proof. The first conservation relation is obtained straightforwardly by adding and integrating on Ω the equations in (7). For the second, we compute, as usual,

$$\partial_t \ln S(t, x) = - \int_{\Omega} \beta(x, y)I(t, y) dy. \quad (22)$$

Secondly

$$-\Delta\partial_t\Phi(t, x) + \gamma(x)\partial_t\Phi = \frac{\partial}{\partial t}(S + E + I)(t, x) = -\gamma(x)I(t, x) + \Delta I(t, x).$$

Therefore, the uniqueness being ensured by the fact that $\underline{\gamma} > 0$, see (3), we obtain

$$\partial_t\Phi(t, x) = -I(t, x).$$

Inserting this expression of I in equation (22) then yields

$$\frac{\partial}{\partial t} \left[-\ln S(t, x) + \int_{\Omega} \beta(x, y)\Phi(t, y)dy \right] = 0. \quad (23)$$

This achieves the proof of Lemma 4.1. □

As before, we may deduce existence of limits from the previous result, as stated now.

Lemma 4.2 *Assume that (3) and (4) hold. Then the solution of (7) verifies : there exists $S_{\infty} \in L_+^{\infty}(\Omega)$, $S_{\infty} \leq S_0$, such that, for a.e. $x \in \Omega$, (5) holds.*

Moreover, for a.e. $x \in \Omega$, $\Phi(t, x)$ converges to $\Phi_{\infty}(x)$ as t goes to $+\infty$ where

$$\begin{cases} -\Delta\Phi_{\infty}(x) + \gamma\Phi_{\infty}(x) = S_{\infty}(x), \\ \partial_n\Phi_{\infty}(x) = 0 & \text{on } (0, \infty) \times \partial\Omega. \end{cases} \quad (24)$$

Proof. As above, for a.e. $x \in \Omega$, since $t \mapsto S(t, x)$ is decreasing, it has a limit $S_{\infty}(x)$, and because $t \mapsto S(t, x) + E(t, x)$ is decreasing, $t \mapsto E(t, x)$ has a limit which vanishes as can be seen adding from (7a)–(7b). Inserting this information in equation (7c), we conclude that, for a.e. $x \in \Omega$, $I(t, x) \rightarrow 0$ as $t \rightarrow +\infty$ too. Finally, we pass to the limit in equation (21) to get the limit Φ_{∞} of Φ . □

4.2 The next-generation operator

Following the approach of Section 3, we define the next-generation operator.

Definition 4 For any $t \geq 0$ and $S(t, \cdot) \in L^1 \cap L^\infty_+(\Omega)$ solution of (7), we call next-generation operator $K_{S(t, \cdot)}$ associated to (7), the operator defined over $L^2(\Omega)$ by

$$K_{S(t, \cdot)}^\Delta[\phi](x) := S(t, x) \int_\Omega \frac{\beta(x, y)}{\gamma(y)} \phi(y) dy + \Delta \left(\frac{\phi(x)}{\gamma(x)} \right), \quad (25)$$

with Neumann boundary conditions. Its adjoint is defined as

$$K_{S(t, \cdot)}^{\Delta*}[\psi](y) := \int_\Omega S(t, x) \frac{\beta(x, y)}{\gamma(y)} \psi(x) dx + \frac{1}{\gamma(y)} \Delta \psi(y), \quad (26)$$

also with Neumann boundary conditions.

The existence of a principal eigenvalue for the next-generation operator is established in the next result, whose proof is postponed to Appendix A.1.

Lemma 4.3 Let $S \in L^1 \cap L^\infty_+(\Omega)$. The operator $K_{S(\cdot)}^\Delta$ admits a principal eigenvalue, that is, there exists $r(K_{S(\cdot)}^\Delta) > 0$ and functions $\varphi_{S(\cdot)} \in L^2_+(\Omega)$ and $\psi_{S(\cdot)} \in L^2_+(\Omega)$ such that

$$K_{S(\cdot)}^\Delta[\varphi_{S(\cdot)}] = r(K_{S(\cdot)}^\Delta)\varphi_{S(\cdot)}, \quad \text{in } \Omega, \quad \partial_n \varphi_{S(\cdot)} = 0, \quad \text{on } \partial\Omega,$$

and

$$K_{S(\cdot)}^{\Delta*}[\psi_{S(\cdot)}] = r(K_{S(\cdot)}^\Delta)\psi_{S(\cdot)}, \quad \text{in } \Omega, \quad \partial_n \psi_{S(\cdot)} = 0, \quad \text{on } \partial\Omega.$$

The basic reproduction number and the herd immunity domain may then be defined as in Definition 3. Notice that an analogue of Lemma 3.5, which links herd immunity and stability, may also be written when diffusion is present.

From Lemma 4.2, there exists $S_\infty \in L^1 \cap L^\infty(\Omega)$, which is the limit of $S(t, \cdot)$. We use the shorthand notation $K_\infty^\Delta := K_{S_\infty(\cdot)}^\Delta$ and $K_\infty^{\Delta*} := K_{S_\infty(\cdot)}^{\Delta*}$; its principal eigenvalue is denoted $r_\infty^\Delta := r(K_{S_\infty(\cdot)}^\Delta)$ and the associated eigenfunctions are respectively $\varphi_\infty := \varphi_{S_\infty(\cdot)}$ and $\psi_\infty := \psi_{S_\infty(\cdot)}$.

The properties described in Lemma 3.3 still hold in this framework:

Lemma 4.4 Assume that (3), (4) hold, and that $r(K_{S_0(\cdot)}^\Delta) > 1$. Then, $t \mapsto r(K_{S(t, \cdot)}^\Delta)$ is decreasing, continuous; $\lim_{t \rightarrow +\infty} r(K_{S(t, \cdot)}^\Delta) < 1$ and there exists a unique $T_0 > 0$ such that $r(K_{S(T_0, \cdot)}^\Delta) = 1$.

Proof. The first point follows exactly the proof of Lemma 3.3. For the existence of T_0 , as above, we multiply (7b) and (7c) by ψ_∞ , and integrating, one finds

$$\begin{aligned} & \frac{d}{dt} \int_\Omega \psi_\infty(x) (E(t, x) + I(t, x)) dx \\ &= \int_{\Omega \times \Omega} \psi_\infty(x) S(t, x) \beta(x, y) I(t, y) dy dx - \int_\Omega \gamma(x) \psi_\infty(x) I(t, x) dx + \int_\Omega \psi_\infty(x) \Delta I(t, x) dx \\ &= \int_{\Omega \times \Omega} K_{S(t, \cdot)}^{\Delta*}[\psi_\infty](y) \gamma(y) I(t, y) dy - \int_\Omega \psi_\infty(x) \gamma(x) I(t, x) dx \\ &= (r_\infty^\Delta - 1) \int_\Omega \gamma(x) \psi_\infty(x) I(t, x) dx. \end{aligned}$$

As in the proof of Lemma 3.3, this proves that there is $T_0 > 0$ such that herd immunity is reached, that is $r(K_{S(T_0, \cdot)}^\Delta) = 1$. \square

4.3 Long time behavior and final size equation – Proof of Theorem 2.2

We have now the material to prove Theorem 2.2. We follow and adapt to the case at hand the proof in Section 3.

Step 1. Existence of limits. The long time convergence has been studied in Lemma 3.3. Then, integrating (23) between t and $+\infty$, we get

$$\ln S(t, x) - \int_{\Omega} \beta(x, y) \Phi(t, y) dy = \ln S_{\infty}(x) - \int_{\Omega} \beta(x, y) \Phi_{\infty}(y) dy, \quad \text{for a.e. } x \in \Omega, t \geq 0. \quad (27)$$

It may be rewritten straightforwardly

$$S_{\infty}(x) = S(t, x) \exp \left(\int_{\Omega} \beta(x, y) (\Phi_{\infty}(y) - \Phi(t, y)) dy \right), \quad \text{for a.e. } x \in \Omega, t \geq 0. \quad (28)$$

Injecting this identity into the equation for Φ_{∞} , and taking $t = 0$, we deduce that the long time limit is characterized by (9b).

Step 2. Exponential rate of decay. This also follows as in Section 3, we just indicate the main steps. Firstly, from (27), and because $\Phi(t, x)$ converges to $\Phi_{\infty}(x)$ almost everywhere, we conclude that for all ε and for $t \geq T_{\varepsilon}$, we have $S(t, x) \leq (1 + \varepsilon) S_{\infty}(x)$.

Secondly, adapting formula (17) to the present setting and using the definition of the adjoint operator $K_{S(t, \cdot)}^{\Delta*}$ given in (26), we compute, for $\theta < 1$ but close to 1

$$\begin{aligned} & \frac{d}{dt} \int_{\Omega} \psi_{\infty}(x) (E(t, x) + \theta I(t, x)) dx \\ & \leq (1 + \varepsilon) \int_{\Omega \times \Omega} \psi_{\infty}(x) S_{\infty}(x) \beta(x, y) I(t, y) dy dx - (1 - \theta) \int_{\Omega} \psi_{\infty}(x) \alpha(x) E(t, x) dx \\ & \quad - \theta \int_{\Omega} \psi_{\infty}(x) \gamma(x) I(t, x) dx + \theta \int_{\Omega} \psi_{\infty}(x) \Delta I(t, x) dx \\ & \leq \theta \int_{\Omega \times \Omega} K_{S(t, \cdot)}^{\Delta*}[\psi_{\infty}](y) \gamma(y) I(t, y) dy - (1 - \theta) \int_{\Omega} \psi_{\infty}(x) \alpha(x) E(t, x) dx \\ & \quad - \theta \int_{\Omega} \psi_{\infty}(x) \gamma(x) I(t, x) dx + (1 + \varepsilon - \theta) \int_{\Omega \times \Omega} \psi_{\infty}(x) S_{\infty}(x) \beta(x, y) I(t, y) dy dx \\ & \leq -\theta(1 - r(K_{S(t, \cdot)}^{\Delta})) \int_{\Omega} \psi_{\infty}(x) \gamma(x) I(t, x) dx - (1 - \theta) \int_{\Omega} \psi_{\infty}(x) \alpha(x) E(t, x) dx \\ & \quad + (1 + \varepsilon - \theta) \int_{\Omega \times \Omega} \psi_{\infty}(x) S_{\infty}(x) \beta(x, y) I(t, y) dy dx. \end{aligned}$$

Taking $t > T_0$ in such a way that $r(K_{S(t, \cdot)}^{\Delta}) < 1$, choosing $\varepsilon > 0$ small enough and $0 < \theta < 1$ close enough to 1 to have $1 + \varepsilon - \theta < 0$, and using the fact (assumed in (3)) that $\inf_{x \in \Omega} \alpha(x)$ and $\inf_{x \in \Omega} \gamma(x)$ are positive together with the positivity of the principal eigenvector ψ_{∞} , allows to use again Gronwall lemma, and to conclude to the exponential rate of convergence of E and I . Exponential convergence of S is then obtained as in Section 3.

Step 3. Uniqueness for t large enough. From Lemma 4.4 we may choose $T > 0$ large enough such that $r(K_{S(T, \cdot)}^{\Delta}) < 1$. From (28) and (24), we deduce

$$-\Delta \Phi_{\infty}(x) + \gamma(x) \Phi_{\infty}(x) = S(T, x) \exp \left(\int_{\Omega} \beta(x, y) (\Phi_{\infty}(y) - \Phi(T, y)) dy \right),$$

complemented with Neumann boundary conditions. Recall that $\Phi(t, \cdot)$ is defined for any t by (21). We notice that since $t \mapsto S(t, x)$ is decreasing, we have $S_\infty \leq S(T)$ and by the maximum principle applied to (21), we deduce also that $\Phi_\infty \leq \Phi(T)$.

Let us consider the problem

$$\begin{cases} -\Delta\left(\frac{u(x)}{\gamma(x)}\right) + u(x) = S(T, x) \exp\left(\int_\Omega \beta(x, y) \left(\frac{u(y)}{\gamma(y)} - \Phi(T, y)\right) dy\right), \\ \partial_n\left(\frac{u(x)}{\gamma(x)}\right) = 0 \quad \text{on} \quad (0, \infty) \times \partial\Omega. \end{cases} \quad (29)$$

We show the uniqueness of a solution of this problem in the set $\{0 \leq u \leq \gamma\Phi(T, \cdot) \text{ a.e.}\}$. Let us assume that there are two solutions u and v . We have as in Section 3.3, using the fact that the exponential is 1-Lipschitz in the set of nonpositive real numbers,

$$-\Delta\left(\frac{(u-v)(x)}{\gamma(x)}\right) + (u-v)(x) \leq S(T, x) \int_\Omega \frac{\beta(x, y)}{\gamma(y)} |u(y) - v(y)| dy.$$

From Kato's inequality [14], we deduce

$$-\Delta\left(\frac{|u-v|(x)}{\gamma(x)}\right) + |u-v|(x) \leq S(T, x) \int_\Omega \frac{\beta(x, y)}{\gamma(y)} |u(y) - v(y)| dy. \quad (30)$$

We recall that, from Lemma 4.3, there exists $\psi_{S(T, \cdot)}$ such that

$$K_{S(T, \cdot)}^{\Delta*}[\psi_{S(T, \cdot)}](y) = \int_\Omega S(T, x) \frac{\beta(x, y)}{\gamma(y)} \psi_{S(T, \cdot)}(x) dx + \frac{1}{\gamma(y)} \Delta \psi_{S(T, \cdot)}(y) = r(K_{S(T, \cdot)}^\Delta) \psi_{S(T, \cdot)}(y).$$

We use the norm

$$\|u\|_T := \int_\Omega \psi_{S(T, \cdot)}(x) |u(x)| dx.$$

Multiplying (30) by $\psi_{S(T, \cdot)}$ and integrating, we get

$$\begin{aligned} \int_\Omega \psi_{S(T, \cdot)}(x) |u-v|(x) dx &\leq \int_{\Omega \times \Omega} S(T, x) \psi_{S(T, \cdot)}(x) \frac{\beta(x, y)}{\gamma(y)} |u(y) - v(y)| dy dx \\ &\quad + \int_\Omega \Delta \psi_{S(T, \cdot)}(x) \left(\frac{|u-v|(x)}{\gamma(x)}\right) dx \\ &\leq r(K_{S(T, \cdot)}^\Delta) \|u-v\|_T. \end{aligned}$$

Since $r(K_{S(T, \cdot)}^\Delta) < 1$, we conclude to the uniqueness of the solution of (29) in the set $\{0 \leq u \leq \gamma\Phi(T, \cdot)\}$, and thus to the uniqueness of Φ_∞ , obtained by dividing by γ , in the set $\{0 \leq \Phi \leq \Phi(T, \cdot)\}$.

Step 4. Uniqueness below Φ_0 . It remains to extend the uniqueness result to all functions smaller than Φ_0 . Analogously to the argument conducted in Section 3, we make use of the following result.

Lemma 4.5 *Among the functions $\Phi \leq \Phi_0$, there is a solution $\bar{\Phi}$ of (9b) which is larger than all subsolutions of (9b), and $\Phi_\infty = \bar{\Phi}$.*

Lemma 4.5 implies that any other solution $\underline{\Phi}$ of (9b) satisfies $\underline{\Phi} \leq \Phi_\infty$, then obviously $\underline{\Phi} \leq \Phi(T, \cdot)$, but by the uniqueness result below $\Phi(T, \cdot)$ obtained in Step 3, we deduce that $\underline{\Phi} = \Phi_\infty$. To achieve the proof of Theorem 2.2, it now only remains to prove the previous lemma.

Proof. [Proof of Lemma 4.5] The proof follows closely the proof of Lemma 3.4 in Section 3.3. For the sake of completeness, we repeat the argument in this framework. Firstly, we build a sequence $\Phi_k(x)$ departing from Φ_0 and defined by induction by, for $k \geq 0$, $\Phi_{k+1}(x)$ is a solution to

$$\begin{cases} -\Delta\Phi_{k+1} + \gamma\Phi_{k+1} = S_0 \exp\left(\int_{\Omega} \beta(x, y)(\Phi_k(y) - \Phi_0(y)) dy\right), \\ \partial_n\Phi_{k+1} = 0 \quad \text{on} \quad \partial\Omega. \end{cases}$$

Clearly, from (8) and the maximum principle, we deduce that $\Phi_1 \leq \Phi_0$ on Ω . Then iterating we deduce that $\Phi_{k+1} \leq \Phi_k$ for any $k \geq 0$. Hence $(\Phi_k)_k$ is a non-increasing sequence which converges to a solution of (9b) denoted by $\bar{\Phi}$.

Secondly, if $\underline{\Phi}$ is a subsolution of (9b) which verifies $\underline{\Phi} \leq \Phi_0$. We have

$$\begin{aligned} -\Delta\underline{\Phi} + \gamma\underline{\Phi} &\leq S_0 \exp\left(\int_{\Omega} \beta(x, y)(\underline{\Phi}(y) - \Phi_0(y)) dy\right) \\ &\leq S_0 = -\Delta\Phi_1 + \gamma\Phi_1. \end{aligned}$$

By the maximum principle, we deduce that $\underline{\Phi} \leq \Phi_1$. Iterating, we deduce that $\underline{\Phi} \leq \Phi_k$ for any $k \geq 0$. Thus $\underline{\Phi} \leq \bar{\Phi}$. In particular, we have $\Phi_{\infty} \leq \bar{\Phi}$.

Thirdly, we prove that $\bar{\Phi} \leq \Phi_{\infty}$. We have $\bar{\Phi} \leq \Phi_0$. Let us show that $\bar{\Phi} \leq \Phi(t)$ for all $t \geq 0$. By contradiction, if it is not true, there exists a first time t_0 for which $\Phi(t_0)$ touches $\bar{\Phi}$, i.e. $\bar{\Phi} \leq \Phi(t_0)$ and for some $x_0 \in \Omega$, $\bar{\Phi}(x_0) = \Phi(t_0, x_0)$. Then, $\Delta\bar{\Phi}(x_0) \leq \Delta\Phi(t_0, x_0)$ and we have

$$\begin{aligned} S(t_0, x_0) + E(t_0, x_0) + I(t_0, x_0) &= -\Delta\Phi(t_0, x_0) + \gamma\Phi(t_0, x_0) \\ &\leq -\Delta\bar{\Phi}(x_0) + \gamma\bar{\Phi}(x_0) \\ &= S_0 \exp\left(\int_{\Omega} \beta(x_0, y)(\bar{\Phi}(y) - \Phi_0(y)) dy\right). \end{aligned}$$

Moreover, from (23), we deduce

$$\begin{aligned} S(t_0, x_0) &= S_0 \exp\left(\int_{\Omega} \beta(x_0, y)(\Phi(t_0, y) - \Phi_0(y)) dy\right) \\ &\geq S_0 \exp\left(\int_{\Omega} \beta(x_0, y)(\bar{\Phi}(y) - \Phi_0(y)) dy\right). \end{aligned}$$

It is a contradiction which concludes the proof of Lemma 4.5 and thus of uniqueness of Φ_{∞} . \square

5 Examples: finite rank contact matrices

The results of Section 2 apply to finite rank contact matrices which have been studied widely with several applications as distributed susceptibility, [25, 24]. We show in Section 5.1 that for rank-1 contact matrices, the general formulas previously found may be simplified. An alternative approach is presented in Section 5.2 in the case where γ is constant, and generalized to finite rank matrices in Section 5.3.

5.1 SEIR system with rank-1 matrices

The case of rank-1 contact matrices refers to matrices $\beta(x, y)$ which we may be split into

$$\beta(x, y) = \beta(x)p(y).$$

Then system (1) reads

$$\dot{S}(t, x) = -\beta(x)S(t, x) \int_{\Omega} p(y)I(t, y) dy, \quad S(0, x) = S_0(x), \quad (31a)$$

$$\dot{E}(t, x) = \beta(x)S(t, x) \int_{\Omega} p(y)I(t, y) dy - \alpha(x)E(t, x), \quad E(0, x) = E_0(x), \quad (31b)$$

$$\dot{I}(t, x) = \alpha(x)E(t, x) - \gamma(x)I(t, x), \quad I(0, x) = I_0(x), \quad (31c)$$

$$\dot{R}(t, x) = \gamma(x)I(t, x), \quad R(0, x) = R_0(x), \quad (31d)$$

where α and γ satisfy assumptions (3), β and p are bounded and positive, and the initial data are given non-negative functions satisfying the conditions in (4). The result in Theorem 2.1 applies.

We can identify explicitly the next-generation operator $K_{S(t, \cdot)}$ in (13) as

$$K_{S(t, \cdot)}[\phi](x) = \beta(x)S(t, x) \int_{\Omega} \frac{p(y)}{\gamma(y)} \phi(y) dy, \quad K_{S(t, \cdot)}^*[\psi](y) = \frac{p(y)}{\gamma(y)} \int_{\Omega} \beta(x)S(t, x)\psi(x) dx.$$

It is worth noticing that, for this operator, the principal eigenlements are (up to normalisation)

$$\begin{cases} r(K_{S(t, \cdot)}) = \int_{\Omega} \frac{\beta(y)p(y)}{\gamma(y)} S(t, y) dy, \\ \varphi_{S(t, \cdot)}(x) = \beta(x) S(t, x), \quad \psi_{S(t, \cdot)}(y) = \frac{p(y)}{\gamma(y)}. \end{cases}$$

Indeed, both $\varphi_{S(t, \cdot)}$ and $\psi_{S(t, \cdot)}$ are positive and satisfy the eigenvalue property

$$K_{S(t, \cdot)}[\beta S(t, \cdot)](x) = r(K_{S(t, \cdot)}) \beta(x)S(t, x), \quad K_{S(t, \cdot)}^*\left[\frac{p(\cdot)}{\gamma(\cdot)}\right](y) = r(K_{S(t, \cdot)}) \frac{p(y)}{\gamma(y)}.$$

Hence, in this case, the basic reproduction number introduced in Definition 3 just reads:

$$\mathcal{R}_0 := \int_{\Omega} S_0(x) \frac{\beta(x)p(x)}{\gamma(x)} dx, \quad (32)$$

and the herd immunity domain is reached for distributions \bar{S} such that

$$\int_{\Omega} \bar{S}(x) \frac{\beta(x)p(x)}{\gamma(x)} dx < 1.$$

Lastly, the computation of the final size reduces to

$$S_{\infty}(x) = S_0(x) \exp\left(\beta(x) \int_{\Omega} \frac{p(y)}{\gamma(y)} (S_{\infty}(y) - S_0(y) - E_0(y) - I_0(y)) dy\right), \quad \text{with } S_{\infty} \leq S_0.$$

5.2 An alternative approach for SIR system with rank-1 matrices

In the simpler case of the SIR system with finite rank contact matrices, another method can be used to study the final size. We begin with rank-1 matrices, for which the formulas are simpler, and assume that $\gamma(x)$ is constant. Analogously to (31), the system reads

$$\dot{S}(t, x) = -\beta(x)S(t, x) \int_{\Omega} p(y)I(t, y) dy, \quad S(0, x) = S_0(x), \quad (33a)$$

$$\dot{I}(t, x) = \beta(x)S(t, x) \int_{\Omega} p(y)I(t, y) dy - \gamma I(t, x), \quad I(0, x) = I_0(x), \quad (33b)$$

$$\dot{R}(t, x) = \gamma I(t, x), \quad R(0, x) = R_0(x). \quad (33c)$$

In that case, one can reduce the system to a single ODE by introducing the quantity

$$m(t) := \int_0^t \int_{\Omega} p(y) I(s, y) dy ds. \quad (34)$$

As a matter of fact, the equation for $S(t, x)$ can be solved as

$$\ln S(t, x) = \ln S_0(x) - \beta(x)m(t).$$

Inserting expression (34) in the equation (33b) for I , and integrating against the weight $p(\cdot)$, we find

$$\begin{aligned} \frac{d}{dt} \dot{m}(t) &= \frac{d}{dt} \int_{\Omega} p(x) I(t, x) dx = \dot{m}(t) \int_{\Omega} \beta(x) p(x) S(t, x) dx - \gamma \dot{m}(t) \\ &= \dot{m}(t) \int_{\Omega} \beta(x) p(x) S_0(x) e^{-\beta(x)m(t)} dx - \gamma \dot{m}(t). \end{aligned}$$

Integrating once, one finds, since $\dot{m}(t=0) = \int_{\Omega} p(y) I_0(y) dy$,

$$\dot{m}(t) = \int_{\Omega} p(x) [I_0 + S_0](x) dx - \int_{\Omega} p(x) S_0(x) e^{-\beta(x)m(t)} dx - \gamma m(t), \quad m(0) = 0. \quad (35)$$

Herd immunity is reached when $\dot{m} = 0$, that is when

$$\int_{\Omega} p(x) [I_0 + S_0](x) dx = \int_{\Omega} p(x) S_0(x) e^{-\beta(x)m_H} dx + \gamma m_H =: F(m_H)$$

which, for any $I_0 > 0$ has a single root $m_H > 0$ which is attractive. Indeed, F is a smooth and convex function on \mathbb{R}^+ such that $F(0) < \int_{\Omega} p(x) [I_0 + S_0](x) dx$ and $\lim_{m \rightarrow +\infty} F(m) = +\infty$. For I_0 small, m_H may be close or far from S_0 depending on the stability of $I \equiv 0$ which is determined by $\mathcal{R}_0 = 1 - \frac{F'(0)}{\gamma}$.

5.3 SIR system with rank- N matrices

The reduction to a single equation for rank-1 matrices can be extended to rank- N contact matrices, i.e., which may be written

$$\beta(x, y) = \sum_{i=1}^N \beta_i(x) p_i(y).$$

Following the above construction, we introduce the matrix M defined by

$$M_{i,j} := \int_{\Omega} \frac{\beta_i(x) p_j(x)}{\gamma(x)} S_0(x) dx.$$

This matrix has positive coefficients and thus we can define its principal eigenvalue λ and the associated eigenvector $X = (X_i)_{i \in \{1, \dots, N\}}$, with $X_i > 0$ for all $i = 1, \dots, N$.

Lemma 5.1 *The basic reproduction number, defined in Definition 3 is given by*

$$\mathcal{R}_0 = r_{S_0(\cdot)} = \lambda$$

and (up to normalisation) the corresponding eigenfunctions are

$$\phi_{S_0(\cdot)}(x) := \sum_{i=1}^N X_i \beta_i(x) S_0(x), \quad \psi_{S_0(\cdot)}(y) := \sum_{i=1}^N X_i \frac{p_i(y)}{\gamma(y)}.$$

Proof. For the given expression of $\phi_{S_0(\cdot)}$, we compute

$$\begin{aligned} K_{S_0(\cdot)}[\phi_{S_0(\cdot)}](x) &= \sum_{i,j=1}^N S_0(x)\beta_i(x) \int_y \frac{p_i(y)}{\gamma(y)} \beta_j(y) X_j dy \\ &= \sum_{i,j=1}^N S_0(x)\beta_i(x) M_{i,j} X_j \\ &= \lambda \sum_{i=1}^N S_0(x)\beta_i(x) X_i = \lambda \phi_{S_0(\cdot)} \end{aligned}$$

which proves the result for $\phi_{S_0(\cdot)}$, the same calculation can be performed for $\psi_{S_0(\cdot)}$. \square

Notice that the characterization of \mathcal{R}_0 given in the statement extends (32).

For rank- N matrices, it is also noticeable that one can also reduce the SIR system to N differential equations as we did in the rank-1 case. We assume that γ is independent of x and define for $j = 1, \dots, N$,

$$m_j(t) = \int_0^t p_j(y) I(t, y) dy, \quad Q(t, x) = \sum_{k=1}^N \beta_k(x) m_k(t).$$

We have

$$\dot{S}(t, x) = -S(t, x) \dot{Q}(t, x), \quad S(t, x) = S_0(x) e^{-Q(t, x)}.$$

From the equation for infected, we compute

$$\frac{d}{dt} \dot{m}_i(t) = \int_{\Omega} p_i(x) S(t, x) \dot{Q}(t, x) dx - \gamma \dot{m}_i(t) = \int_{\Omega} p_i(x) S_0(x) e^{-Q(t, x)} \dot{Q}(t, x) dx - \gamma \dot{m}_i(t).$$

Integrating once in t , we obtain the equations, for $i = 1, \dots, N$,

$$\dot{m}_i(t) = - \int_{\Omega} p_i(x) S_0(x) e^{-Q(t, x)} dx - \gamma m_i(t) + \int_{\Omega} p_i(x) [S_0(x) + I_0(x)] dx, \quad m_i(0) = 0. \quad (36)$$

Notice that (36) extends (35). Conceptually, this is a system of N differential nonlinear equations, which can be computed at least numerically.

6 Conclusion

We considered an epidemic in a heterogeneous population modelled by a SEIR system with a continuous structure variable and a general contact matrix. We investigated question of the final size of the epidemic both with and without diffusion. In this general framework, our main contribution is to prove uniqueness results for the equation characterizing the final size of the epidemic. The proof combines a contraction property close to the steady state with a monotonicity argument which localized roughly the final state.

Although we presented our results for a SEIR model, they can be extended to other compartmental models, like the SIR model and to more general equations than the simple diffusion.

The main limitation comes from the specific form of the SEIR model excluding birth and loss of immunity. In those models, up to our knowledge, proving merely convergence to a steady state is an open problem.

Appendix

A.1 Proof of Lemma 4.3

In this technical part, we show that the operator defined on $L^2(\Omega)$ by

$$K_{S(\cdot)}^\Delta[\varphi](x) := S(x) \int_{\Omega} \frac{\beta(x, y)}{\gamma(y)} \varphi(y) dy + \Delta \left(\frac{\varphi(x)}{\gamma(x)} \right),$$

admits a principal eigenvalue, that is, that there exist $\lambda > 0$ and a function $\varphi > 0$ such that $K_{S(\cdot)}^\Delta \varphi = \lambda \varphi$ in Ω , with $\partial_n \varphi = 0$ in $\partial\Omega$.

It is more convenient to work with $\phi = \varphi/\gamma$. Let the operator R be defined as

$$R\phi(x) := \frac{S(t, x)}{\gamma(x)} \int_{\Omega} \beta(x, y) \phi(y) dy.$$

We define for all $M > 0$ the operator:

$$T\phi := \left((-\Delta + M)^{-1} \circ R \right) \phi,$$

where $(-\Delta + M)^{-1}f$ is the unique solution u of $-\Delta u + Mu = f$ in Ω , with $\partial_n u = 0$ on $\partial\Omega$.

This operator is clearly compact and strongly positive, from $L^2(\Omega)$ to $L^2(\Omega)$. The Krein-Rutman theorem thus guarantees the existence of a principal eigenvalue Λ_M , associated with a positive eigenfunction ϕ_M that we normalize by $\int_{\Omega} (\phi_M(x))^2 dx = 1$. The identity $T\phi_M = \Lambda_M \phi_M$ rewrites

$$-\Delta \phi_M + M \phi_M = \frac{1}{\Lambda_M} R \phi_M \quad \text{in } \Omega. \tag{A.1}$$

Moreover, the function $M \mapsto \Lambda_M$ is clearly continuous by Kato's regularity theory.

On the one hand, multiplying by ϕ_M and integrating by parts, one finds

$$\begin{aligned} \frac{1}{\Lambda_M} \langle R \phi_M, \phi_M \rangle_{L^2(\Omega)} &= \int_{\Omega} |\nabla \phi_M|^2 dx + M \int_{\Omega} (\phi_M(x))^2 dx \\ &\geq M \int_{\Omega} (\phi_M(x))^2 dx. \end{aligned}$$

On the other hand, one has

$$\begin{aligned} \langle R \phi_M, \phi_M \rangle_{L^2(\Omega)} &= \int_{\Omega \times \Omega} \frac{1}{\gamma(x)} \beta(x, y) S(t, x) \phi_M(x) \phi_M(y) dy dx \\ &\leq C \left(\int_{\Omega} \phi_M(x) dx \right)^2, \end{aligned}$$

for some constant $C > 0$ due to (3) and $S(t, x) \leq S_0(x)$. Hence, using the Cauchy-Schwarz inequality, one gets (increasing C if necessary) $\frac{C}{\Lambda_M} \geq M$ and thus $\Lambda_M \rightarrow 0$ as $M \rightarrow +\infty$.

Moreover, integrating equation (A.1) and using again our hypotheses (3), one gets

$$M \int_{\Omega} \phi_M(x) dx = \frac{1}{\Lambda_M} \int_{\Omega \times \Omega} \frac{1}{\gamma(x)} \beta(x, y) S(t, x) \phi_M(y) dy dx \geq \frac{1}{C \Lambda_M} \int_{\Omega} \phi_M(x) dx$$

for some positive constant $C > 0$. Hence, $CM \geq 1/\Lambda_M$ and thus $\Lambda_M \rightarrow +\infty$ as $M \rightarrow 0$.

We conclude that there exists $M > 0$ such that $\Lambda_M = 1$ by the intermediate value theorem, and thus (A.1) exactly yields, by letting $\varphi = \gamma \phi_M$:

$$K_{S(\cdot)}^\Delta \varphi = M \varphi,$$

with $\varphi > 0$ in Ω . Letting $\lambda := M$ ends the proof.

References

- [1] Ricardo Aguas, Rodrigo M. Corder, Jessica G. King, Guilherme Goncalves, Marcelo U. Ferreira, M. Gabriela M. Gomes, Herd immunity thresholds for SARS-CoV-2 estimated from unfolding epidemics, medRxiv preprint doi: <https://doi.org/10.1101/2020.07.23.20160762> .
- [2] Viggo Andreasen, The final size of an epidemic and its relation to the basic reproduction number, *Bull. Math. Biol.* (2011) 73:2305–2321.
- [3] Julien Arino, Fred Brauer, Pauline van den Driessche, James Watmough, Jianhong Wu, A final size relation for epidemic models, *Math. Biosc. & Eng.*, Volume 4, Number 2, April 2007, pp 159–175.
- [4] Norman T.J. Bailey, *The mathematical theory of infectious diseases* (1975) Griffin, London.
- [5] Fred Brauer, Age-of-infection and the final size relation, *Math. Biosc. & Eng.* **5** (4) (2008), 681–690.
- [6] Tom Britton, Frank Ball, Pieter Trapman, A mathematical model reveals the influence of population heterogeneity on herd immunity to SARS-CoV-2, *Science* 10.1126/science.abc6810 (2020).
- [7] Tom Britton, Frank Ball, Pieter Trapman, The disease-induced herd immunity level for Covid-19 is substantially lower than the classical herd immunity level, preprint arXiv:2005.03085.
- [8] Odo Diekmann, Hans Heesterbeek, and Johann Anton J. Metz, On the definition and the computation of the basic reproduction ratio R_0 in models for infectious diseases in heterogeneous populations, *J. Math. Biol.* (1990) **28**, No. 4, 365–382.
- [9] Francesco Di Lauro, Luc Berthouze, Matthew D. Dorey, Joel C. Miller, István Z. Kiss, The impact of network properties and mixing on control measures and disease-induced herd immunity in epidemic models: a mean-field model perspective, preprint arXiv:2007.06975.
- [10] Jean Dolbeault, Gabriel Turinici, Heterogeneous social interactions and the COVID-19 lockdown outcome in a multi-group SEIR model, *Math. Model. Nat. Phenom.* 15 (2020) 36.
- [11] Romain Ducasse, Threshold phenomenon and traveling waves for heterogeneous integral equations and epidemic models. Preprint 2019. hal-02004866
- [12] Romain Ducasse, Qualitative properties of spatial epidemiological models. Preprint. 2020. hal-02571610
- [13] M. Gabriela M. Gomes, Rodrigo M. Corder, Jessica G. King, Kate E. Langwig, Caetano Souto-Maior, Jorge Carneiro, Guilherme Goncalves, Carlos Penha-Goncalves, Marcelo U. Ferreira, Ricardo Aguas, Individual variation in susceptibility or exposure to SARS-CoV-2 lowers the herd immunity threshold, medRxiv preprint doi 10.1101/2020.04.27.20081893.
- [14] Tosis Kato, Schrödinger operators with singular potentials. *Israel Journal of Mathematics*, (1972) 13(1-2), 135–148.
- [15] Guy Katriel, The size of epidemics in populations with heterogeneous susceptibility, *J. Math. Biol.* (2012) 65:237–262.

- [16] David G. Kendall, Discussion of "measles periodicity and community size" by M.S. Bartlett. *J. Roy. Stat. Soc. A*, 120:64–76, 1957.
- [17] William O. Kermack, Anderson G. McKendrick, Contributions to the mathematical theory of epidemics – I, *Proceedings of the Royal Society* 115A (1927), 700–721.
- [18] Mark Grigor'evich Kreĭn, Mark Aronovich Rutman. Linear operators which leave a cone in Banach space invariant. *Amer. Math. Soc. Transl.* (1950) 1, 10, 199–325.
- [19] Junling Ma, David J.D. Earn, Generality of the final size formula for an epidemic of a newly invading infectious disease, *Bull. Math. Biol.* (2006) **68**: 679–702.
- [20] Pierre Magal, Ousmane Seydi, Glenn Webb, Final size of an epidemic for a two-group SIR model, *SIAM J. Appl. Math.* (2016) Vol. 76, No. 5, pp. 2042–2059.
- [21] Joel C. Miller, A Note on the derivation of epidemic final sizes, *Bull. Math. Biol.* (2012) **74**: 2125–2141.
- [22] Artem S. Novozhilov, On the spread of epidemics in a closed heterogeneous population. *Mathematical Biosciences* 215 (2008) 177–185.
- [23] Horst R. Thieme, A model for the spatial spread of an epidemic. *J. Math. Biol.*, 4(4):337–351, 1977.
- [24] Horst R. Thieme, *Mathematics in population biology*. Woodstock Princeton university press. Princeton (2003).
- [25] Horst R. Thieme, Distributed susceptibility: a challenge to persistence theory in infectious disease models. *DCDS(B)*, Vol. 12, No. 4 (2009).
- [26] Eberhard Zeidler, *Nonlinear Functional Analysis and its Applications, I: Fixed Point Theorems*. Springer-Verlag (1986).