The first damselfly (Odonata, Lestidae) from the upper Eocene of Monteils (Gard, France)

Romain Jattiot, Benjamin Latutrie, André Nel

► To cite this version:

Romain Jattiot, Benjamin Latutrie, André Nel. The first damselfly (Odonata, Lestidae) from the upper Eocene of Monteils (Gard, France). *Zootaxa*, 2020, 4750 (3), pp.432-436. 10.11646/zootaxa.4750.3.9 . hal-02985932

HAL Id: hal-02985932

<https://hal.sorbonne-universite.fr/hal-02985932>

Submitted on 2 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The first damselfly (Odonata, Lestidae) from the upper Eocene of Monteils (Gard, France)

Romain Jattiot^a, Benjamin Latutrie^b and André Nel^c

^a Fachbereich 5 Geowissenschaften, Universität Bremen, Klagenfurter Strasse 4, 28357, Bremen, Germany. E-mail: jattiot@uni-bremen.de

^b La Grange, 9003 En Cros, route de Garrigues, 81500, Lavaur. E-mail: benjamin.latutrie@gmail.com

^c Institut de Systématique, Évolution, Biodiversité, ISYEB - UMR 7205 – CNRS, MNHN, UPMC, EPHE, Muséum national d'Histoire naturelle, Sorbonne Universités, Université des Antilles, 57 rue Cuvier, CP 50, Entomologie, F-75005, Paris, France. E-mail: anel@mnhn.fr

Abstract

The discovery of the first damselfly *Lestes regina* from Monteils (Gard, France) supports the identity of late Eocene age of this outcrop with the historical outcrop of Célas, type locality for the type series of this species. *Lestes regina* is also documented from the late Eocene Isle of Wight basin, confirming the presence of significant contacts between this southern area and the anglo-Parisian lacustrine basin at that time. Nearly all the Eocene and Oligocene fossil *Lestes* from Western Europe have a particular character, viz. the presence of a supplementary row of cells between the veins MP and CuA. This character is much less frequent in extant *Lestes* and is still unknown among Neogene representatives of the genus.

Key words: Odonatoptera, late Eocene, Monteils Formation, paleobiogeography, wing venation.

Introduction

Even though the Lestoidea are recorded from the Lower Cretaceous with *Cretalestes martinae* Jarzembowski *et al.*, 1998, the oldest accurate fossil record of the extant family Lestidae comes from the late Eocene, as the Paleocene *Lestes zaleskyi* Piton, 1940 is based on a poor and lost fossil of Zygoptera of uncertain position (Piton, 1940; Nel & Paicheler, 1994). The oldest known Lestidae have been found in the lacustrine outcrops of the Isles of Wight (UK) and Célas (France). The discovery of new Eocene fossils of this family can be of great interest. Here, we describe the first fossil Odonata found in the upper Eocene outcrop of Monteils (Gard, France). This outcrop is very rich in fossil plants, fishes and insects, although rather few of these have been described (Laurent, 1899, 1910; Feist-Castel, 1971; Nel, 1988; Séméria & Nel, 1990; Rückert-Ülkümen, 2001; Nel *et al.*, 2008; Legalov, 2013; Perrard *et al.*, 2014; Cai *et al.*, 2014; Skartveit & Nel, 2017).

Material and method

The studied specimen was collected from the yellowish late Eocene (Priabonian) shale of Alès-Monteils in Gard, southern France (44°05'9"N, 04°11'39"E, after the BRGM geological maps 1/50 000, numbers 912 – Alès, XXVIII – 40 and 938 - Anduze, XXVIII-41). It was examined using a Leica M300 binocular and photographed using a Nikon D5100 camera equipped with a Nikon AF 60 mm Micro lense.

We follow the wing venation nomenclature of Riek & Kukalová-Peck (1984), modified by Nel *et al.* (1993), Bechly (1996, 2016) and Jacquelin *et al.* (2018).

Abbreviations. Ax, primary antenodal crossvein; arc, arculus; dc, discoidal cell; sn, subnodus; C, Costa; Pt, pterostigma; RA, Radius anterior; RP, Radius posterior; IR, Interradius; MA, Media anterior; MP, Media posterior; CuA, Cubitus anterior. Add: N, nodus, which has indicated in the Fig.2.

Systematic paleontology

Order Odonata Fabricius, 1793

Family Lestidae Calvert, 1901

Genus *Lestes* Leach, 1815

Lestes regina Théobald, 1937

(Figs 1–2)

Material. Specimen UBGD 32320 (imprint, a nearly complete damselfly with all legs and wings preserved, sex unknown), stored at the Université de Bourgogne, Géologie, Dijon, France; counterimprint in Coll. Benjamin Latutrie.

Description. Body black, wings hyaline with pterostigma dark brown. Head deformed as in all fossil Lestidae, with mouthparts anteriorly projected, 2.6 mm long, and 3.5 mm wide. Thorax 6.0 mm long, and 3.4 mm wide. Fore- and hindwing nearly identical, in particular discoidal cells of same shapes and widths; fore wing 19.9 mm long, 4.5 mm wide, ratio L/W 4.4, width at nodus level 3.2 mm; pterostigma 2.3 mm long, and 0.5 mm wide; distance from base to arculus 3.7 mm, from arculus to nodus 3.7 mm, from nodus to pterostigma 9.8 mm, and from subnodus to base of RP2 2.8 mm; oblique vein ‘O’ two cells distal of base of RP2; a supplementary row of 5-6 cells between MP and CuA. Abdomen 27.9 mm long, 1.3 mm wide; cerci present but poorly preserved.

Discussion

The wing venation of this fossil is characteristic of the Lestidae and more precisely of the genus *Lestes*: long pterostigmata; bases of RP3/4 and IR2 midway between arculus and nodus; an oblique crossvein ‘O’; distal discoidal vein MAb very oblique, with distal angle of discoidal cell very acute; MA strongly zigzagged; area between IR2 and RP3/4 distally strongly widened

with three rows of cells between these two veins; fore- and hindwing discoidal cells of the same width and shape. One species of this genus was already described from the uppermost Eocene of Célas (in a lost outcrop at the old trench at the entrance to an abandoned railway tunnel), an outcrop situated at 4 km to and contemporaneous with Monteils (Cavelier, 1979), viz. *Lestes regina* Théobald, 1937. The holotype of this species is probably lost and Nel & Paicheler (1994: 10-11, fig. 5) designated a lectotype and redescribed it. Mentioned below are the very few differences between *L. regina* and the herein described specimen.

The fore wing of the specimen from Monteils is 19.9 mm long, 4.5 mm wide, ratio L/W 4.4, width at nodus level 3.2 mm; while in the holotype of *Lestes regina*, the fore wing is 21.5 mm long, 4.6 mm wide with a ratio L/W 4.7 (Théobald, 1937). In the lectotype, the fore wing is 22.0 mm long, 4.5 mm wide with a ratio L/W 4.8, width at nodus level 3.0 mm (Nel & Paicheler, 1994). The body dimensions are less reliable because of possible deformations due to taphonomic processes. Nevertheless, the abdomen is shorter than in the holotype of *L. regina* (27.9 mm long vs. 33.0 mm), and broader (1.3 mm vs. 0.75 mm). An additional difference could be the presence of three cells between base of RP2 and oblique vein 'O' in the lectotype of *L. regina*, whereas there are only two in the specimen from Monteils. In our opinion, the aforementioned differences are not sufficient to clearly separate this latter from *Lestes regina*. This discovery confirms that the two outcrops of Célas and Monteils are contemporaneous or nearly so, despite the different sediment aspect between them (darker limestone at Célas, compared to the white to grey blue limestones at Monteils).

Conclusion

The occurrence of *Lestes regina* in the three localities of Célas, Monteils and Isle of Wight suggests that there were significant contacts between the region of Alès and the northern Anglo-Parisian lacustrine basin. Feist-Castel (1971) came to the same conclusion based on

charophytes. The presence of the same genus of Orthoptera Tettigoniidae at Célas and at Isle of Wight further strengthens this hypothesis (Nel *et al.*, 2008).

The presence of a supplementary row of cells between MP and CuA is encountered in several Eocene-Oligocene *Lestes* spp. from Europe, viz. *Lestes regina*, *Lestes* aff. *regina* (late Eocene, Isles of Wight, UK, Nel and Fleck, 2014), *L. ceresti* Nel & Papazian, 1985, *Lestes brisaci* Nel *et al.*, 1997 (both from the late Oligocene, France), *L. statzi* Schmidt, 1958 (late Oligocene, Germany), *L. foersteri* Hess, 1895 (early Oligocene, France), and *L. aquisextanus* Nel, 1985 (latest Oligocene, France) (Hess, 1895; Schmidt, 1958; Nel, 1985; Nel & Papazian, 1985; Nel *et al.*, 1997). The two lestad species of uncertain generic affinities *Lestes vicina* Hagen, 1858 (early Oligocene, Germany) and *Lestes plicata* Piton & Théobald, 1939 (early Oligocene, France) also have this supplementary row of cells (Hagen, 1858; Piton & Théobald, 1939; Nel & Paicheler, 1994). Conversely, the Miocene European species of *Lestes* do not exhibit this particular character. Compère-Sart (2014) indicated that this supplementary row of cells between MP and CuA can be found also in some specimens but not in others, of extant *Lestes* species (*L. dryas*), and therefore considered it of minor importance. Yet, it would be somewhat significant to find it in all specimens of *L. ceresti* and *L. regina*. Naturally, these fossils are still too few (ca. 10 specimens of *L. ceresti*) to represent a valid statistical basis. More specimens are necessary to draw any conclusion on the frequency and stability of this character among the Eocene-Oligocene species of *Lestes*. Nevertheless, it seems that this character was more frequent among the Paleogene *Lestes* than in the Neogene ones.

Acknowledgements

We thank a lot two anonymous referees for their helpful remarks on the first version of the paper.

References

- Bechly, G. (1996) Morphologische Untersuchungen am Flügelgeäder der rezenten Libellen und deren Stammgruppenvertreter (Insecta; Pterygota; Odonata), unter besonderer Berücksichtigung der Phylogenetischen Systematik und des Grundplanes der Odonata. *Petalura*, Special Volume 2, 1–402.
- Bechly, G. (2016) Phylogenetic systematics of Odonata. - homepage on Internet site: <https://bechly.lima-city.de/phylosys.htm>
- Cai, C.-y., Clarke, D.J., Huang, D.-y & Nel, A. (2014) A new genus and species of Steninae from the late Eocene of France (Coleoptera, Staphylinidae). *Alcheringa: An Australasian Journal of Palaeontology*, 38, 557–562.
- Calvert, P.P. (1892-1908). Insecta Neuroptera Odonata. pp. 17–420. In: Godman, F.D. (ed.). *Biologia Centrali-Americana*. London.
- Cavelier, C. (1979) La limite "Eocène - Oligocène" en Europe occidentale. *Mémoires des Sciences Géologiques, CNRS*, 54, 1–280.
- Compte-Sart, A. (2014) Nueva especie del género *Lestes* Leach, 1815 (Insecta, Odonata) del Mioceno de Bellver de Cerdanya (Lérida). *Graellsia*, 70, 1–10.
- Fabricius, J.C. (1793) Entomologia systematica emendata et aucta, secundum classes, ordines, genera, species, adjectis synonymis, locis, observationibus, descriptionibus. C.G. Proft, Hafniae [= Copenhagen], 3, 1–487 and 1–349.
- Feist-Castel, M. (1971) Sur les charophytes fossiles du bassin tertiaire d'Alès (Gard). *Geobios*, 4, 157–172.
- Hagen, H.A. (1858) Zwei Libellen aus der Braunkohle von Sieblos. *Palaeontographica*, 5, 121–126.
- Hess, W. (1895) Beitrag zur Kenntnis der tertiären Agrioniden: eine neue *Lestes*-Art aus dem plattigen Steinmergel von Brunstatt bei Mülhausen i. E. *Wissenschaftliche Beilage zum*

Programm der Oberrealschule, Gewerbeschule für das Schuljahr 1894-95, Programm, Mülhausen, 538, 1–15.

Jacquelin, L., Desutter-Grandcolas, L., Chintauan-Marquier, I., Boistel, R., Zheng, D.-r., Prokop, J. & Nel, A. (2018). New insights on basivenal sclerites using 3D tools and homology of wing veins in Odonatoptera (Insecta). *Scientific Reports*, 8 (238), 1–7.

Jarzembowski, E.A., Martinez-Declos, X., Bechly, G., Nel, A., Coram, R. & Escullié, F. (1998) The Mesozoic non-calopterygoid Zygoptera: description of new genera and species from the Lower Cretaceous of England and Brazil and their phylogenetic significance (Odonata, Zygoptera, Coenagrionoidea, Hemiphleboidea, Lestoidea). *Cretaceous Research*, 19, 403–444.

Laurent, L. (1899) Flore des calcaires de Célas. *Annales du Musée d'Histoire Naturelle de Marseille*, (2), 1, 148 pp.

Laurent, L. (1910) Etude phytologique sur le bassin de Célas. *Bulletin de la Société Géologique de France*, (4), 10, 910–913.

Leach, W.E. (1815) Entomology. In: Brewster, D. (ed.): The Edinburgh Encyclopaedia, 9, 57–172.

Legalov, A.A. (2013) New and little known weevils (Coleoptera: Curculionoidea) from the Paleogene and Neogene. *Historical Biology: An International Journal of Paleobiology*, 25, 59–80.

Nel, A. (1985). Sur la présence d'un *Lestes* Leach (1815) fossile de la lignée de *Lestes regina* Théobald dans les calcaires stampiens d'Aix-en-Provence. (Odon., Lestidae). *Entomologica Gallica*, 1, 317–319.

Nel, A. (1988) Une nêpe fossile de l'Eocène supérieur du bassin d'Alès (Hemiptera, Heteroptera, Nepidae). *Nouvelle Revue d'Entomologie*, (N.S.), 5, 273–276.

- Nel, A. & Fleck, G. (2014) Dragonflies and damselflies (Insecta: Odonata) from the Late Eocene of the Isle of Wight. *Earth and Environmental Science, Transactions of the Royal Society of Edinburgh*, 104, 283–306.
- Nel, A., Martínez-Delclòs, X., Paicheler, J.-C. & Henrotay, M. (1993) Les ‘Anisozygoptera’ fossiles. Phylogénie et classification (Odonata). *Martinia*, Numéro Hors Série 3, 1–311.
- Nel, A., Martínez-Delclòs, X., Papier, F. & Oudard, J. (1997) New Tertiary fossil Odonata from France. (Sieblosiidae, Lestidae, Coenagrioniidae, Megapodagrionidae, Libellulidae). *Deutsche Entomologische Zeitschrift*, 44, 231–258.
- Nel, A. & Paicheler, J.-C. (1994) Les Lestoidea (Odonata, Zygoptera) fossiles: un inventaire critique. *Annales de Paléontologie*, 80, 1–59.
- Nel, A. & Papazian, M. (1985) Description d'une nouvelle espèce fossile de *Lestes* leach, 1815, du Stampien de Céreste (Alpes-de-Haute-Provence) (Odon. Lestidae). *Entomologica Gallica*, 1, 275–279.
- Nel, A., Prokop, J. & Ross, A.J. (2008) New genus of leaf-mimicking katydids (Orthoptera: Tettigoniidae) from the Late Eocene - Early Oligocene of France and England. *C.R. Palevol*, 7, 211–216.
- Perrard, A., Nel, A. & Carpenter, J.M. (2014) A new paper wasp from Late Eocene of France (Hymenoptera: Vespidae: Polistinae). *Zootaxa*, 3753, 542–548.
- Piton, L. & Théobald, N. (1939) Poissons, crustacés et insectes fossiles de l'Oligocène du Puy-du-Mur, Auvergne. *Mémoires de la Société Scientifique de Nancy*, 4, 77–123.
- Riek, E.F. & Kukalová-Peck, J. (1984) A new interpretation of dragonfly wing venation based upon Early Carboniferous fossils from Argentina (Insecta: Odonatoidea) and basic characters states in pterygote wings. *Canadian Journal of Zoology*, 62, 1150–1166.

- Rückert-Ülkümen, N. (2001) *Hemitrichas vardinis* (Sauvage, 1883) (Pisces, Teleostei) aus dem Alt-Tertiär (Obereozän) von Gard, Südfrankreich. *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Historische Geologie*, 41, 65–71.
- Schmidt, E. (1958) Bemerkungen über Lestiden. 3. Über *Oligolestes grandis* Statz, 1935 und eine neue *Lestes*-Art aus dem mittel-Oligozän von Rott im Siebengebirge. *Decheniana*, 111, 1–7.
- Séméria, Y. & Nel, A. (1990) *Paleochrysopa monteilsensis* gen. et sp. nov., a new fossil of Chrysopidae from the Upper Eocene formation of Monteils (France with a review of the known Chrysopid fossils (Insecta, Neuroptera). In: Mansell, M.W. & Aspöck H. (eds). *Advances in Neuropterology, Proceedings of the Third International Symposium on Neuropterology, Bergen Dal, Kruger National Park, R.S.A. 1988*. Pretoria, R.S.A., 27–32.
- Skartveit, J. & Nel, A. (2017) Revision of fossil Bibionidae (Insecta: Diptera) from French Oligocene deposits. *Zootaxa*, 4225, 1–83.
- Théobald, N. (1937) Les insectes fossiles des terrains oligocènes de France. *Bulletin Mensuel (Mémoires) de la Société des Sciences de Nancy*, 1, 1–473.

FIGURE 1. *Lestes regina* Théobald, 1937, UBGD 32320, Monteils. General habitus. Scale bar = 10 mm.

FIGURE 2. *Lestes regina* Théobald, 1937, UBGD 32320, Monteils. (A) Fore wing; (B) Hindwing, arrow supplementary row of cells between MP and CuA. Scale bar = 10 mm.