Modes of propagation of continental break-up and associated oblique rift structures

Anthony Jourdon, Laetitia Le Pourhiet, Frédéric Mouthereau, Dave A. May

► To cite this version:

Anthony Jourdon, Laetitia Le Pourhiet, Frédéric Mouthereau, Dave A. May. Modes of propagation of continental break-up and associated oblique rift structures. *Journal of Geophysical Research: Solid Earth*, 2020, 125 (9), 10.1029/2020JB019906 . hal-02985979

HAL Id: hal-02985979

<https://hal.sorbonne-universite.fr/hal-02985979>

Submitted on 2 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/343788283>

Modes of propagation of continental break-up and associated oblique rift structures

Preprint · August 2020

CITATIONS

0

READS

407

4 authors, including:

Anthony Jourdon

Géosciences Environnement Toulouse - Observatoire Midi-Pyrénées

18 PUBLICATIONS 121 CITATIONS

[SEE PROFILE](#)

Laetitia le pourhiet

Sorbonne Université

70 PUBLICATIONS 2,436 CITATIONS

[SEE PROFILE](#)

Frederic Mouthereau

Paul Sabatier University - Toulouse III

144 PUBLICATIONS 4,460 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:

Crustal deformation of the Tien Shan belt (Central Asia) [View project](#)

Flexural state of the lithosphere and strain localisation [View project](#)

Modes of propagation of continental break-up and associated oblique rift structures

Anthony Jourdon^{1,2}, Laetitia Le Pourhiet², Frédéric Mouthereau¹, Dave May³

1- CNRS, GET, Université Paul Sabatier, Toulouse, France

2- Sorbonne Université, CNRS-INSU, Institut des Sciences de la Terre Paris, ISTE-P UMR, Paris, France

3- Oxford University, Department of Earth Sciences

Abstract

V-shaped oceanic propagators are widespread around the world. Their geometry combined with magnetic anomalies associated to their opening shows at first order that ridge propagation in the third dimension occurs by pulses. In this study we use 3D thermomechanical numerical models to show how oblique kinematic boundary conditions control both the intracontinental rift development and the oceanic ridge propagation. To do so, we apply a shortening velocity boundary condition in the direction perpendicular to the extension for “strong” and “weak” crustal rheologies. Numerical models results highlight that three ridge propagation modes can occur. For low out of plane velocities (12% to 15% of the extension rate), the ridge propagation is fast (>1.5 cm/yr) and straight. Higher shortening velocities (15% to 17%) lead to a ridge propagation by pulses alternating between fast propagation (~ 1.5 cm/yr) and stalling phases. Finally, for higher velocities (17% to 20%) a ridge jump propagation mode occurs, localizing a new spreading centre between 100 and 200 km far from the initial ridge. We also show that ridge propagation phases are associated with dip-slip dominated deformation while stalling phases are dominated by strike-slip deformation. These deformation regimes are marked by structures reorientation while kinematic boundary conditions remain constant. We discuss these results in term of plate tectonics reconstructions and regional geological studies.

1 Introduction

Magnetic anomalies record the past direction and rate of seafloor spreading (Vine & Matthews, 1963). As such they are our best constrain for global plate tectonic reconstructions (Müller et al., 2008; 2016; Seton et al., 2012). In perfectly cylindrical condition the orientation of the first magnetic anomaly is parallel to the rifted margin of continents. However, in some areas the first magnetic anomalies intercept the rifted margins of continents at acute angle instead of being parallel to it. These remnants are precious as they allow us to quantify the rate at which continental break-up propagates along the strike of the mid-oceanic ridges (Courillot, 1982). Indeed, using only basic kinematic considerations, it is clear that if the propagation occurs at the same velocity as the spreading rate, the angle between the margin and the magnetic anomaly should be roughly 45° . Faster propagation leads to the parallelisation of the magnetic anomaly with the margin of the oceanic basin matching with the 2D cylindrical model. Slower propagation produces magnetic anomalies that normalise to the rifted margin, until the limit case of a transform margin (no ridge propagation at all).

In map, these phases of sluggish ridge propagation produce V-shaped margins. V-shaped propagators are found in many places on Earth (Figure 1) and seafloor age map compilation by Müller et al. (2008) is often sufficient to unravel the first order features of the timing of continental break-up at regional or more global scale. Examples displayed in Figure 1 show propagators separated by very long segments

where magnetic anomalies are parallel to the margin. This indicates that pulsation in the rate of continental break-up propagation first introduced by Courtillot (1982) in the Gulf of Aden is a major feature of divergent plate boundaries. Despite the well documented geological record of these changes of rate, and their implications for plate tectonics and kinematic reconstructions or the thermal regime of oblique margins, the dynamics of the propagation at long/geodynamic timescale remains poorly studied or understood.

Kinematic interpretations of V-shaped propagators are based on models established in the late 1970s and early 1980s, which are still in use today. The first model, called the rotational extension (Hey et al., 1980; Molnar et al., 2017; Mondy et al., 2018; Zwaan et al., 2020), assumes that seafloor spreading occurs near the Euler pole and is therefore described by very large variations in the opening rate in the direction of the ridge. In this case, the first magnetic anomaly forms parallel to the rifted margin then the linearly increasing rates of opening with distance from Euler pole produce a series concentric V's over time Figure 2c. This model infers that very rapid changes in tectonic regime must occur at the tip of the V-shaped ocean domain. The second kinematic model, published by Vink (1982), considers that the divergence remains constant in the direction of rift axis but it is variably partitioned between the oceanic accretion and the continental rifting (Figure 2b). This model applies only when magnetic anomalies indicate that the Euler Pole is far enough so that the rate of divergence is constant at regional scale. This is generally the case for a distance greater than 3000 km or 30° from the Euler pole. The Vink model allows the formation of V-shaped propagators when the propagation rate is similar to the oceanic accretion rate. For faster propagation, magnetic anomalies are parallel to the margin (Figure 2a). The Vink model is the basis for Courtillot's (1982) proposal (Figure 2d) who interpreted the change in angle between magnetic anomalies and the continental margins along the direction in the Gulf of Aden as the geological record of pulsating continental break-up propagation. It should be noted here that these two kinematic models are not contradictory but rather complement each other as two end-member cases. However, Vink's and Courtillot's models predictions in terms of magnetic anomalies and associated tectonic structures at the front of the propagator are different and therefore permit the description of different natural examples of V-shaped propagators. The Woodlark Basin only opens 500 km from the Euler Pole. It displays a strong segmentation, variations from extensional to strike-slip structure (Holm et al., 2016; Mondy et al., 2018) and only the youngest ridge ages at the tip which conform with characteristics of a rotational extension opening basin. While the South China Sea that opened at about 4500 km from its pole of rotation (Biais et al., 1993; Mazur et al., 2012; Le Pourhiet et al., 2018) with very little variations in the spreading rate along the strike would better illustrate the Vink kinematic model.

Dynamically speaking, on the one hand, the rotational opening model has been reproduced by a number of dynamic simulations based either on analogue materials (Molnar et al., 2017; Zwaan et al., 2020) or numerical methods (Mondy et al., 2018) while, on the other hand, almost no properly scaled dynamic simulations, including numerical (Allken et al., 2012; Beaussier et al., 2019; Liao & Gerya, 2015; Laetitia Le Pourhiet et al., 2017) or analogue models (Ding & Li, 2016; Zwaan et al., 2016), reproduces the kinematics proposed by Vink (1982) at lithospheric scale. With these boundary conditions and a realistic viscosity in the asthenospheric mantle, lithospheric necking should propagate approximately ten time faster than plates diverge (Allken et al., 2012; Laetitia Le Pourhiet et al., 2018) resulting invariably in the fast propagation kinematic displayed in Figure 2a. Therefore, to produce a kinematic solution, which resembles Vink's (1982) model (Figure 2b) and a number of geological observations (Figure 1), some additional ingredients must therefore be added to the dynamic simulations to stop or slow down the propagation in order to form a V-shaped oceanic propagator.

86 The obvious candidate to slow down the propagation of continental break-up is the presence of a
87 strong rheological heterogeneity, so called strong ribbon (Figure 2d). This hypothesis was first
88 proposed by Morgan & Parmentier (1985) based on the theory of crack propagation. It was supported
89 by small strain numerical simulations of the propagation of continental rifting published by Van Wijk
90 & Blackman (2005). The results of this first study are consistent with the results presented by Allken
91 et al. (2012): the propagation rate of a neck (continental rift) in a frictional layer (upper crust) lying
92 above a viscous layer (lower crust) decreases with the viscosity of the lower layer. Many similar
93 analogue or numerical experiments (Allken et al., 2012; Brune et al., 2017; Corti, 2012) have proved
94 that increasing the viscosity of the lower crust resulted in a slower lengthening of grabens in the upper
95 crust. However, at lithosphere scale there is nothing like a high viscosity asthenosphere to delay or
96 slow the propagation of continental break-up. At that scale, the strength of the lower crust has been
97 shown to enhance fast lithospheric necking and tends to increase the rate of propagation of continental
98 break-up (Benes & Scott, 1996; Le Pourhiet et al., 2018).

99 Oblique continental rifting controlled by the presence of pre-existing viscous weak zones oblique to
100 the extension direction creates en-échelon rift basins, which are separated by the transfer zones that
101 accommodate strain differences between segments (Ammann et al., 2017; Brune, 2014; Duclaux et al.,
102 2020; Mart & Dauteuil, 2000; McClay & White, 1995; Molnar et al., 2017). These isolated basins
103 form small segments of oceanic accretion before complete lithosphere continental plate break-up is
104 achieved. As they grow in width they also grow in length, propagating from the basin center towards
105 both the extremity. Yet their propagation is limited by the compressional stress build-up that arises
106 from the growth of nearby segment. Le Calvez & Vendeville, (2002) proposed a model set-up to study
107 in details the dynamic of linkage between en-échelon segments. This set-up consists of two offset
108 weak zones which grow laterally in map view and link or not depending on spacing, frictional layer
109 thickness and viscosity of the underlying layer. Allken et al. (2012) presented the result for a similar
110 set of numerical experiments and showed the importance of the rate of softening in the frictional layer
111 to produce a transform fault. More recently, Le Pourhiet et al. (2017) have extended this model to
112 study continental break-up at lithosphere scale (Figure 2e). At that scale, the linkage between
113 segments can be delayed by 10's of millions of years. Meanwhile, two asymmetric V-shaped
114 propagators form, as shown in Figure 2e. Full linkage occurs only after oblique necking of the mantle
115 lithosphere is reached showing similarities with the initial conditions of Ammann et al. (2017) study in
116 which an oblique weak zone is imposed. These asymmetric propagators better fit the observations in
117 the Gulf of Aden or in the equatorial Atlantic (Le Pourhiet et al., 2017) than the original propagator
118 models do (Figure 2a, b), but they do not apply to all geological examples of V-shaped propagators.

119 To produce oblique extension, pure shear boundary conditions in map view similar to the Alboran
120 Sea set up in Le Pourhiet et al., (2014) (i.e. stretching velocity in one horizontal direction and a
121 shortening velocity in the other horizontal direction) are a good alternative to the introduction of
122 oblique heterogeneities in cylindrical boundary conditions. Indeed, strain localisation causes the
123 formation of structures that can rotate internally within the pure shear far field strain and which
124 favours the emergence of instable 4 plates system that stabilizes into stable 3 plates systems (Gerya &
125 Burov, 2018). It has been demonstrated that pure shear conditions produce V-shaped propagators both
126 during the reorganisation of oceanic plates (Gerya & Burov, 2018) and during continental break-up
127 (Le Pourhiet et al., 2018). This results together with the analogue experiments of Ding & Li (2016),
128 who introduced a gradient in gravitational potential energy facing the propagation direction, suggests
129 that applying a small amount of shortening or compression out of the plane of the main driving
130 extension slows down continental rifting propagation rate. One should note that in 3D the transition
131 from transtensional to transpressional regime in pure shear map setting is related to change in surface

of the model and that as long as the surface remain in dilatational regime no shortening related structures form (Le Pourhiet et al., 2014). GPS velocities vectors plotted relative to fix Arabian plate in figure 1 clearly show frontal displacement towards actively propagating oceanic ridge exist in active propagating continental rifts of Aden and Red Sea. In the first case, one might suspect the influence of the Afar plume and in the second case the vicinity of the Mediterranean subduction zone. In both cases the frontal shortening is not associated to shortening tectonic structures which could “fossilize” its existence.

Here, we therefore intend to focus our attention in this still largely unexplored direction. We adopt the same boundary conditions as Le Pourhiet et al. (2018) (Figures 2f and 3) in order to analyse the sensitivity of continental break-up propagation rate to both the rate of ridge parallel shortening and lower crustal rheology. We first extract from the simulations synthetic propagation rates and expected magnetic anomalies in opening basins that corresponds to data that are widely available, relate them to large scale plate deformations. We then focus on the tectonic structures that accommodates the frontal shortening in the models in order to discuss what are key tectonic observations that might help to decipher past kinematics and lithospheric dynamics from seismic lines and thermal dataset.

2 Set Up of the thermo-mechanical simulations

2.1 Modelling approach

In order to model the long term deformation of the lithosphere we use pTatin3D (May et al., 2014, 2015), a highly scalable, massively parallel implementation of the finite element method which employs an Arbitrary Lagrangian-Eulerian (ALE) discretization together with the material point method to solve the conservation of momentum for an incompressible fluid:

$$\nabla \cdot (2\eta \dot{\epsilon}) - \nabla P = \rho \mathbf{g} \quad (1)$$

$$\nabla \cdot \mathbf{v} = 0 \quad (2)$$

where η is the non-linear effective viscosity, $\dot{\epsilon}$ the strain rate tensor, P the pressure, ρ the density, $\mathbf{g}$ the gravity acceleration vector and $\mathbf{v}$ the velocity vector of the fluid. This Stokes flow is coupled with the time dependent heat conservation equation:

$$\frac{\partial T}{\partial t} + \mathbf{v} \cdot \nabla T = \nabla \cdot (\kappa \nabla T) + \frac{H}{\rho C_p} \quad (3)$$

where T is the temperature, $\mathbf{v}$ the velocity vector of the fluid, κ the thermal diffusivity, H the heat source (here only radioactive heat sources are considered) and C_p is the heat capacity. The material density is considered to vary with temperature such as:

$$\rho = \rho_0 (1 - \alpha(T - T_0)) \quad (4)$$

where ρ_0 is the initial material density and α the thermal expansion coefficient.

The Stokes problem is solved using a Q_2 - P_1 discretization while the heat equation is discretized on Q_1 elements. pTatin3D uses a free surface boundary condition which can dynamically evolve due to deformation and surface processes that we model with a 2D diffusion equation discretized on Q_1 elements:

$$\frac{\partial h}{\partial t} = -\nabla \cdot (\kappa \nabla h) \quad (5)$$

where h is the altitude and k the diffusion coefficient. The equation is solved using a first order explicit time integration.

2.2 Rheological model

To simulate the long time behaviour of the lithosphere we use temperature and pressure dependent non-linear rheologies. Brittle parts of the lithosphere follow the Drucker-Prager pseudo-plastic yield criterion adapted to continuum mechanics:

$$\eta_p = \frac{C \cos(\phi) + P \sin(\phi)}{\dot{\epsilon}^{II}} \quad (6)$$

where C is the cohesion, ϕ the friction coefficient, P the pressure and $\dot{\epsilon}^{II}$ the second invariant of the strain rate tensor. To simulate the friction drop in deforming faults, we use a simple linear decrease of the friction angle from 30° to 5° with accumulated plastic strain from 0 to 1. However, laboratory experiments show that under high confining pressures (> 1 GPa) brittle behaviour changes to plastic behaviour (e.g. Kameyama et al., 1999; Precigout et al., 2007). We account for that change by limiting the Drucker-Prager yield stress to a maximum deviatoric stress of 400 MPa according to the calibration performed by Watremez et al. (2013).

Viscous parts of the lithosphere are simulated using an Arrhenius flow law for dislocation creep:

$$\eta_v = A^{-\frac{1}{n}} (\dot{\epsilon}^{II})^{\frac{1}{n-1}} \exp\left(\frac{Q+PV}{nRT}\right) \quad (7)$$

where A , n and Q are material dependant parameters (see Table 1), R is the gas constant and V is the activation volume.

2.3 Initial conditions

Our experiments represent a 1200 (in the x direction) $\times$ 600 (in the z direction) $\times$ 250 (in the y direction) km^3 domain discretized with $512 \times 256 \times 128$ elements respectively (Figure 3a). In the y direction, two third of the 128 elements are concentrated in the first third of the 250 km giving approximately 8 km^3 ($2 \times 2 \times 2 \text{ km}^3$) elements in the first 80 km. Four rheological layers subdivide the models domain to represent the upper and lower crust (each of 20 km thick), the lithosphere mantle (80 km thick) and the asthenosphere (170 km thick). The Moho is located at 40 km depth and the lithosphere-asthenosphere boundary is at 120 km depth (Figure 3a). Lithosphere mantle and asthenosphere are both simulated using a dry olivine flow law (Hirth & Kohlstedt, 2003) (Table 1).

Depending on the model, the crust rheology may vary. The rheology of the lower crust has been proven to be of first order importance for the strain localization and propagation in 3D (Allken et al., 2012; Le Pourhiet et al. 2014; Brune et al., 2017; Corti, 2012; Le Pourhiet et al., 2017; 2018). Therefore, in this study we use two different crustal lithologies denoted as weak and strong (Figure 3c). The weak crust corresponds to an upper and a lower crust modelled with a quartz rheology (Ranalli & Murphy, 1987) while the strong crust corresponds to an upper crust modelled with a quartz rheology and the lower crust with an anorthite rheology (Rybacki & Dresen, 2000). We modelled those two rheological layering with the same boundary and initial conditions in order to assess the role of the lower crust on the evolution of the rifting and continental break-up.

	Units	Quartz	Anorthite	Olivine
Reference		Ranalli and Murphy 1987	Rybacki and Dresen 2000	Hirth and Kohlstedt 2003
A	Mpa ⁻ⁿ .s ⁻¹	6.3x10 ⁻⁶	13.4637	1.1x10 ⁵
n		2.4	3	3.5
Q	KJ/mol	156	345	530
V	m ³ .mol ⁻¹	0	3.8x10 ⁻⁵	18x10 ⁻⁶

The initial temperature field is computed with a steady-state analytical solution (Turcotte & Schubert, 2002)

$$T_{init} = T_{y=0} + \frac{-yq_m}{k} + H \frac{y_p^2}{k} \left(1 - \exp\left(\frac{-y}{y_p}\right) \right) \quad (8)$$

for a $T_{y=0} = 0^\circ\text{C}$, an incoming mantle heat flux $q_m = 20 \text{ mW.m}^{-2}$, a radiogenic heat production $H = 1.2 \times 10^{-6} \text{ W.m}^{-3}$, a characteristic radiogenic layer of $y_p = 40 \text{ km}$ and a conductivity of $3.3 \text{ W.m}^{-1}.\text{K}^{-1}$. This analytical solution gives 610°C at 40 km (depth of the Moho, Figure 3c) and 1300°C at 120 km (depth of the LAB, Figure 3c). Then from 120 km to 250 km we compute a linear increase of temperature for an adiabatic gradient of 0.5°C/km (Figure 3c). Although this second part of the geotherm is not steady state, the cooling by diffusivity is very slow (less than 2°C/Myr for the maximum cooling rate) and it allows to keep reasonable conductivity values in the asthenosphere ($3.3 \text{ W.m}^{-1}.\text{K}^{-1}$).

The initial radiogenic heat production is set as an exponential decay of heat production with depth according to Turcotte & Schubert (2002) as follow:

$$H = H_0 \exp\left(\frac{-y}{y_p}\right) \quad (9)$$

for a surface production H_0 of $1.2 \times 10^{-6} \text{ W.m}^{-3}$.

In order to initiate localisation on one side of the modelling domain, we define a weak zone in which a random amount of initial plastic strain ranging from 0 to 1 is applied to the particles located in a cube of coordinates $x=550$ to $x=650$, $y=0$ to $y=100$ and $z=0$ to $z=100$ instead of 0.0 to 0.03 background value. Since we use plastic strain softening, the initial amount of plastic strain will decrease the initial friction angle in the concerned area as:

$$\phi = \phi_0 - \frac{\epsilon_p - \epsilon_{min}}{\epsilon_{max} - \epsilon_{min}} (\phi_0 - \phi_\infty)$$

Where ϕ is the friction angle, ϕ_0 the initial maximum friction angle (30°), ϕ_∞ the minimal friction angle reached after softening (5°), ϵ_{max} and ϵ_{min} are the bounds of maximum and minimum plastic strain amount between which softening is applied and ϵ_p the actual plastic strain computed from equation (6).

2.4 Boundary conditions

The boundary conditions to solve the heat conservation equation are $T=0^{\circ}\text{C}$ at the top boundary ($y=0$ km), $T=1440^{\circ}\text{C}$ at the bottom boundary ($y=250\text{km}$) and null heat fluxes on all vertical boundaries of the domain. To solve the conservation of momentum, we impose a free surface on the top boundary, a free slip boundary conditions on the face of normal z at coordinates $x, y, z=0$, and Dirichlet boundary conditions on the normal component of velocity on all other vertical sides. An extension velocity $v_x=5$ mm/yr is imposed on both faces of normal x and a shortening velocity v_z is imposed on the face of normal z at coordinates $x, y, z=600$ (Figure 3a, b). This shortening velocity may vary between models (from 1.2 mm/yr to 2.0 mm/yr). It introduces a part of constriction in the model domain that intrinsically modifies the ratio between normal stretching and thinning as compared to a cylindrical model. In this range of shortening rates, propagation rates are indeed of the same order of magnitude (less than 1.5 cm/yr) of the plates velocity imposed in the model (1 cm/yr) and it is for this range of velocities that we expect the highest dependence of deformation pattern to the rheology of the lithosphere (Le Pourhiet et al., 2018).

2.5 Post processing

In order to compare the numerical models results with natural examples and known geological and geophysical data, we computed an equivalent of seafloor magnetic anomalies. Magnetic anomalies measured on the seafloor are used to model seafloor spreading rates and oceanic crust ages in oceans (Müller et al., 2008). In our models, the formation of oceanic crust by partial melting of the mantle and magma crystallization is not implemented. Therefore, we compute an age of mantle exhumation below the 800°C isotherm which can be roughly approximated as the Curie Temperature of ferromagnetic material (Rushbrooke & Wood, 1958). This age corresponds to the time in the model at which the mantle cooled beneath 800°C . In the models we use 0 Myr as the beginning and time goes forward from this initial value while geological time is counted from present (0 Ma) to past. Therefore, the age of the mantle exhumed during the latest stages of the models is greater than the age of the mantle exhumed in the earlier stages. Those ages values need to be interpreted as the greater is also the younger.

Figures 8, 10 and 12 and supplementary figures 11 and 12 display the deformation regime computed from the stress tensor orientation also known as the Regime Stress Ratio (Brune, 2014; Brune & Autin, 2013 and reference therein). The colour intensity is related to the strain rate second invariant value.

Figures 4, 5, 7, 8 and 9 show the plastic and/or viscous strain deformation. Those values represent respectively the cumulated finite strain computed with the equations (6) and (7). Therefore, the plastic strain displays the deformations related to frictional (or brittle) parts of the lithosphere while the viscous strain shows the deformations related to the dislocation creep flow laws in the “ductile” parts of the lithosphere.

Supplementary Figure 1 shows the β factor of the crust computed as

$$\beta = \frac{h_c^{t=0}}{h_c^{t=n}} \quad (10)$$

where $h_c^{t=0}$ is the initial thickness of the crust at the beginning of the simulation ($h_c = 40$ km, $t = 0$ Myr) and $h_c^{t=n}$ is the thickness of the crust at a given time n . Thus, the β factor value represents the thinning ratio of the crust at a given time.

Figure 6a displays the position of the tip and the ocean propagation with respect to time. The position of the tip is measured as the higher z coordinates of the mantle-continent transition along the strike of the oceanic ridge. Therefore, it represents a particular area at the frontier between a plate boundary (the oceanic ridge) and an intra-continental deformation zone (the rift). As discuss in the numerical results section, this area may evolve with time in a triple junction or remain as a plate boundary to intra-continental deformation transition. The ridge propagation (Figure 6b) is the time derivative of this position computed as:

$$Vrp = \frac{\Delta z}{\Delta t} = \frac{z_{t=n} - z_{t=n-1}}{t_n - t_{n-1}} \quad (11)$$

where Vrp is the ridge propagation velocity, $z_{t=n}$ and $z_{t=n-1}$ are the z coordinates at time t_n and t_{n-1} respectively. Figure 6c shows the tectonic forces evolution through time. F_x and F_z represent the forces normal to the faces of normal x and z respectively. To compute forces we first compute the deviatoric stress tensor as:

$$\sigma' = 2\eta \dot{\epsilon} \quad (12)$$

where η is the viscosity and $\dot{\epsilon}$ the strain rate tensor. We then compute the traction along faces of normal x and z respectively:

$$\vec{T}_x = \iint \sigma' \cdot \vec{x} \, dzdy \quad (13a)$$

and:

$$\vec{T}_z = \iint \sigma' \cdot \vec{z} \, dx dy \quad (13b)$$

where $\vec{x}$ and $\vec{z}$ are the unit vectors in x and z directions respectively. We then compute the tectonic forces per linear meter as:

$$F_x = \frac{\|\vec{T}_x\|}{L_z} \quad (14a)$$

and:

$$F_z = \frac{\|\vec{T}_z\|}{L_x} \quad (14b)$$

where L_x and L_z are the length of the domain along x and z axis respectively.

3. Dynamic of continental break up propagation

Numerical models are aimed at investigating the role of both the crust rheology and the intensity of rift parallel shortening applied in our simulations. We show here that if we impose a sufficient velocity v_z (perpendicular to the extension direction) the propagation rate of the continental breakup is not instantaneous and does not exceed 1.5 cm/yr. Moreover, this is independent of the rheology of the crust. For the same amount of shortening velocity applied in the direction perpendicular to the extension, the main difference between strong and weak lower crust models resides in the propagation mode. Furthermore, for large shortening velocities, weak and strong lower crust models tend to show the same behaviour forming oblique rifts (Figures 4 and 5). In order to highlight the principal results of this study we first focus on the ridge propagation mode as a function of rift parallel inflow (obliquity).

3.1. Rheology of the crust and sensitivity to the shortening velocity

Our numerical modelling results show that the rift-parallel shortening velocity and the crust rheology strongly affect the propagation of the rift and its structure (Figures 4, 5 and 6). With a strong lower crust, the minimum shortening velocity necessary to form a strike slip conjugate and isolate a new micro-plate is 17% of the extension velocity ($v_z=1.7$ mm/yr, Figures 4e and 5e, model 3). With a weak lower crust, the rift propagation starts to be sensitive to the shortening velocity when the latter reaches 12% of the total extension velocity (Figure 4b and 5b, model 1). The influence of this third boundary condition becomes very important for 15% of the extension velocity (Figure 4d and 5d, model 2). For both lower crust rheologies (strong and weak), the shortening velocity represents a threshold, across which pure extension gives place to a transtensional and strike-slip deformation regime. A strong lower crust favours the formation of three tectonic plates joining at a Ridge-Ridge-Fault triple junction because it allows for a more localised strain than weaker crust rheologies (Figures 4e, g, h and 5e, g, h). Margins are narrow and few shear zones accommodate the crustal thinning (as shown by the beta factor varying from 1 to 5 in less than 50 km, Supplementary Figure 1). In contrast, a weak lower crust rheology favours a more distributed deformation (Figure 4b, d, f and 5b, d, f). Wide rifts are formed and many shear zones develop to accommodate the crustal thinning (beta factor of ~ 3 along 300 km from the exhumed mantle, Supplementary Figure 1). This distribution prevents the formation of three distinct plates and the effects of the out-of-plane boundary condition are visible in the rift evolution and propagation. Finally, for both weak and strong lower crusts a too large amount of shortening (20% for the strong case, Figures 4h and 5h and 17% for the weak case, Figures 4f and 5f) results in the formation of an oblique continental rift followed by oblique spreading.

3.2. Ridge propagation

The numerical models highlight three types of rifts propagation modes (Figure 4 and Figure 6a). The first mode is a very fast one (from 3 cm/yr to 1.5 cm/yr, Figure 6b) that results in an oceanic opening almost simultaneous along the rift axis. Models with a strong lower crust and a shortening velocity less than 17% of the total extension velocity (1.7 mm/yr) display this evolution (Figure 4 a, c and Figure 6).

The second propagation mode is characterized by a succession of fast propagation and stalling phases (Figure 6). This mode is principally visible in the weak lower crust models for shortening velocities less than 1.7 mm/yr (Figure 4b, d). Due to the weak lower crust, a long phase of continental rifting occurs (~ 40 Myr) before ridge propagation takes place (Figure 6d). The propagation velocity starts at about 1.5 cm/yr before slowing down in ~ 10 Myr to less than 5 mm/yr (Figure 6b). This propagation velocity drop can be related to crustal thickness variations and competition between extension velocity and shortening velocity. Indeed, the initial continental rift phase efficiently thins the crust along 200 km to 300 km (in z direction) where the velocity field is principally perpendicular to the rift. Therefore, once crustal breakup occurs, the propagation along this already thinned crust is fast. However, once the rift tip reaches a less thinned crust the propagation stops or slows down to allow the continental crust domain in front of the tip to thin enough before resuming the propagation (Supplementary Figures 4, 7 and 9). During the whole simulations time (~ 120 Myr) models were able to produce two or three stalling phases, each one lasting less than the previous one. This feature can be explained by the decreasing crustal thickness with time in front of the rift propagator.

Finally, a third propagation mode reveals a “jump” of the spreading centre from the tip of the stalling rift to a new basin. Models with strong and weak lower crust both present this propagation mode for shortening velocities larger than 1.7 mm/yr (Figure 4e, g, h and Figure 6a). As in the two previous propagation modes, the rift starts with a fast propagation (~ 1.5 cm/yr, Figure 6b) before stalling. But instead of propagating again from where it has stopped, a new spreading centre forms 200

km to 300 km farther (Figure 4e and Figure 6a). The main characteristic of this propagation mode resides in the deformation regime between the onset of the stalling phase and the formation of a new spreading centre. As soon as the along strike oceanic propagation stops, the intra-continental deformation at the front of the ridge switches from a dip-slip dominated regime to a strike-slip dominated. During this deformation regime, the rifting continues but localizes on an offset branch and leads to the formation of two offset oceanic segment and an ultra oblique margin (Figure 4e, g, h).

3.3. Tectonic forces evolution

Figure 6c shows the tectonic forces F_x (equation 14a) and F_z (equation 14b) evolution along boundaries of normal x and z respectively. Along the faces of normal x the F_x force corresponds to the extensional forces. Strong lower crust models display initial extensional forces (from 15 TN/m to 17 TN/m) greater than weak lower crust ones (from 10 TN/m to 12 TN/m) at the beginning of the experiments and through time. The F_x force display a small dependency to the rift parallel shortening magnitude at the beginning of the simulations. However, since the rift parallel shortening influences the strain localization, the force evolution differs between models. At first order, the F_x force decreases with the lithosphere thinning through time for all models. The force decrease is directly related to the thinning and propagation rates of the models: the faster the lithosphere thins, the faster the force decreases. Then, once the continental break-up occurred along the whole length of the modelled domain the force reaches or tend to approach a plateau representing the strength of the lithosphere.

Along the face of normal z a first order observation is that the F_z force is not dependent to the rift parallel shortening velocity but to the strength of the lithosphere. Indeed, the models with a strong lower crust display an initial state between 4 TN/m and 5 TN/m while the models with a weak lower crust show an initial state between at ~ 3 TN/m. During the deformation of the lithosphere, the F_z force shows few variations related to strain localization events in the models except for models with a strong lower crust and a rift parallel shortening of 1.2 mm/a to 1.5 mm/a. These two models display a fast force increase associated with the fast ridge propagation and a plateau between 8 TN/m and 9 TN/m when the ridge had propagated all along the domain. In absolute values, the F_z force is 2 to 5 times lower than the extensional force F_x .

4. Characteristic structures associated to propagation of break-up

In order to decipher the geological record of the different propagation modes and at the same time to gain insights on the transition from straight mode of propagation to partitioned mode we have chosen to details the evolution of models with rates of shortening 1.2 mm/a (model 1), 1.5 mm/a (model 2) and 1.7 mm/a (model 3).

4.1. Weak Lower crust, 12% of shortening velocity

The first model (model 1) is characterized by a weak lower crust and a shortening velocity $v_z=1.2$ mm/yr (12% of the total extension velocity). Continental rifting lasts 40 Myr before continental breakup occurs at the southern boundary of the model (Figure 7a, and Figure 8a). During this initial phase, normal faulting is perpendicular to the extension direction. The weak lower crust results in strain distribution over a wide continental rift (Figure 7a, b, cross sections 2 and 3 and Figure 8a, b). At 50 Myr, the continental breakup reaches $z=200$ km and a localized strain band replaces the wide strain distribution in the exhuming mantle between $z=0$ and $z=200$ km (Figures 7b, cross section 3 and Figure 8b), while in the rest of the model continental rifting is still active (Sections 1 and 2 of Figure 7b). At 60 Myr, the domain of exhumed mantle reaches $z=300$ km and the oceanic rift enters a second phase characterized by short episodes of propagation and stalling take place. Propagation first slows

down for 10 Myr while the stretching of the continental crust increases in a narrow domain and finally localized as crustal breakup occurs, allowing oceanic rift propagation to continue (Figure 7c, d). During this phase, a minor strike-slip crustal deformation initiates at the edge of the propagator tip (coordinates: $300 < z < 400$; $700 < x < 800$, Figure 7d, cross section 2 and Figure 8d) that accommodates the differential movement between the stalling ridge and ongoing crustal extension away from the ridge. Then, from 80 Myr to 100 Myr the ridge propagation resumes and as the mantle is exhumed, the active deformation in the crust progressively vanishes (Figure 7e-g, Figure 8e-g). The age map of mantle exhumation displays a typical pattern where the oldest ages are decreasing in the direction of propagation (Figure 7g). At the end of the model, the angle between the oceanic rift and the continental margins varies along strike. This angle variation developed during the slowing down of the propagation phase. The direction of normal shear zones changes in time and along the rift axis. The first faults formed during the continental rift phase have been rotated during the first stalling phase (between 50 Myr and 60 Myr, Figure 7c, d). Then, as the rift propagates the shear zones formed in the crust continue to rotate and align with the rifted margins.

4.2. Weak Lower crust, 15% of shortening velocity

The second model (model 2) also presents a weak lower crust but with a velocity $v_z=1.5$ mm/yr (15% of the total extension velocity). Here again continental rifting takes place during approximately 40 Myr before reaching the crustal breakup and mantle exhumation (Figure 8a and Supplementary Figures 5a and 6a). Like the previous model a wide rift develops but deformation appears to be more localized. At 50 Myr, the ridge is propagating northward (up to $z=200$) and normal faulting is reduced (Figure 9b and Figure 10b). At this stage, the propagation starts to slow down and the ridge propagation enters a stalling phase until 80 Myr. During these 30 Myr, the exhumed mantle domain widens, forming stripes of exhumation ages parallel to the direction of extension but perpendicular to the continental margins located at the edge of the propagator (Figure 9b, c, d). This stalling phase results in drastic modification of the rift architecture. The rifted margins located around the tip and normal faults rotate until they form an angle of 90° with the active ridge (Figure 9c, d, e and Figure 10c, d, e). As a result, the rifted margins located at the propagator tip are striking parallel to the extension direction. This rotation is accompanied with the development of a left-lateral strike-slip fault zone in the crust (Figures 9c, d, e and Figure 10c, d, e). Transcurrent deformation first develops at around 50 Myrs as ridge propagation stops (Figure 10b) and evolves as a highly localized crustal-scale strike-slip shear zone (Figure 10e). The rotated normal faults of the continental rift reach an obliquity of $\sim 45^\circ$ with respect to the stretching direction and ultimately form releasing bends aligned in the strike-slip fault direction. During this phase, the continental crust is thinned under a transtensional strain regime, exhuming the lower crust in elongated dome structures parallel to stretching direction as described on the field (e.g. Jolivet et al., 2004) and reproduced previous modelling studies (e.g. Le Pourhiet et al., 2012, 2014). Around 80 Myr, the propagation resumes and the exhumed mantle domain opens obliquely to the stretching direction along the former strike-slip zone (Figures 9e and Figure 10e). At 90 Myr, the ridge propagation slows down once again and a second left-lateral strike-slip structure develops (Figure 9f and Figure 10f). The most remarkable result is that the rift orientation change associated with the development of strike-slip deformation occurs spontaneously without involving changes in plate kinematics (Figure 9 and 10).

4.3. Strong Lower crust, 17% of shortening velocity

The third model (model 3) involves a strong lower crust and a velocity $v_z=1.7$ mm/yr (17% of the total extension velocity). In this model, the continental rifting phase stands for less than 30 Myr before crustal breakup occurs (Figure 6d). Crustal deformation is also more localized than in the previous models (Figure 11 and 12). At 30 Myr, the ridge reaches $z=200$ km (Figures 11a and Figure 12a) and

its propagation slows down. This propagation rate change coincides with the formation of two new deformation zones (Figure 12a): a right-lateral strike-slip structure to the left and a left-lateral transtensional shear zone to the right. At 40 Myr, the ridge propagation enters in a stalling phase and stops (Figure 11b). The two branches localize now all the intra-continental deformation at the front of the ridge and a ridge-rift-fault triple junction which rapidly evolves in ridge-ridge-fault forms at the tip of the propagator (Figure 12b). The strike-slip motion in both branches intensifies and becomes predominant (Figure 12b, c) until a new exhumed mantle domain appears at 60 Myr ($800 < x < 900$; $400 < z < 500$, Figures 11d, cross-section 1). Between the initial propagator and this new basin, a thin band of continental crust remains before the two exhumed mantle domains join at 70 Myr (Figures 11e, cross section 2). As soon as the new exhumed mantle domain forms, the deformation regime on the right branch changes once again from a strike-slip regime to transtensional extension (Figure 12e). As in the second model (weak lower crust, $v_z = 1.5$ mm/yr), the strike-slip deformation in the right branch accommodates the rotation of the continental margin previously located at the edge of the initial propagator. The rifted margin not only rotates but also thins during the transtensional deformation phase, before being torn apart when both exhumed mantle domains join (Figure 11e, cross-section 2). The finite rotation results in a rifted margin oriented parallel to the extension direction, while it was formed perpendicularly to it. This process may be responsible for the initiation of a transform margin. Here again, the lower crust exhumes preferentially in areas where strike-slip deformation is dominant. In the left branch, the right-lateral strike-slip deformation is active from 30 Myr to the end of the simulation (Figure 11 and Figure 12). This localized vertical shear zone progressively offsets the former rifted margin structures (Figure 11, cross section 3) and forms a vertical margin where the transition from continent to exhumed mantle is sharp (Figures 11e, f, cross section 4). Compared with the previous models, the lithosphere-scale strain partitioning is a main characteristic of model 3 and results in the formation of a triple junction and three distinct tectonic plates. As in the model 2, the change in deformation regime and rift orientation is provoked by the stalling phase while kinematic boundary conditions are kept constant.

5. Discussion

5.1. Geodynamic interpretation of rift parallel inflow

Lithospheric modelling aims at understanding the dynamics that lead to the emergence of new localised plate boundaries at regional scale. These plate boundaries are of 3 types (Fault, Ridge, Trench), which can be studied separately. But plates tectonics usually deal with triple junctions and the velocity field surrounding triple junction is not cylindrical in map, it is rather a combination of pure and simple shear components. The geodynamic pertinence of non cylindrical boundary conditions makes therefore no doubts if one wants to study the emergence of triple junction.

In term of modelling, Le Pourhiet et al. (2014) have shown that transtensional tectonic structures (A type domes) emerging from pure shear boundary conditions are similar to those produced by imposing simple shear boundary conditions. Yet, pure shear boundary conditions have the advantage over simple shear boundary conditions because they allow the formation of large strike slip structures without having to impose a step function in the boundary conditions of the models, that is, without prescribing the location of the strike slip faults nor their occurrence at all. They are therefore useful to study the emergence or not of localised strike slip plate boundary rather than oblique rift.

Finally, as for all models run with kinematic boundary conditions, one need to discuss the forces needed to deform the model. Our results and especially the force calculation show some unexpected results which need to be discuss.

For 3D deformation patterns to emerge within cylindrical boundary conditions one must include a form of inheritance in the initial conditions. Models having large area with random noise (Duclaux et al., 2020; Naliboff et al., 2020) demonstrates that the offset of oblique structures in cylindrical conditions is not very large without strong kinematic forcing (Ammann et al., 2017; Brune et al., 2012; Allken et al., 2012; Gerya, 2013; Liao & Gerya, 2015; Le Pourhiet et al., 2017) assuming that regional deformation is cylindrical. But also through boundary conditions such as oblique extension (Brune, 2014; Brune et al., 2012; Brune & Autin, 2013), pure shear (e.g. Gerya & Burov, 2018; Le Pourhiet et al., 2014) or combination between pure and simple shear (Le Pourhiet et al., 2018). Kinematic boundary conditions involving pure shear showed the ability to reproduce formation of more than two tectonic plates and triple junctions (Gerya & Burov, 2018).

In this study we used Le Pourhiet et al. (2018) kinematic boundary conditions to impose obliquity. As for pure shear boundary conditions we showed that accounting for sufficient rift parallel shortening leads to the formation of a triple junction between three tectonic plates in models with strong lower crust. The coupled rheology of the crust seems to be a first order constraint to produce large scale strike-slip fault in extensional contexts as also shown in analogue models (Zwaan et al., 2019).

In terms of tectonic forces, the models show that the forces are more dependants to the rheology of the lithosphere than to the rift parallel shortening magnitude. In a weak lower crust lithosphere, the force associated to the rift parallel shortening (F_z) is in the same order of magnitude than the force generated by gravitational potential energy (Bird et al., 2006). For a strong lower crust lithosphere, the gravitational potential energy may not be sufficient to explain such rift parallel shortening and an active tectonic process like a subduction or a collision may be required in the vicinity of the rift to allow rift parallel shortening. However, the force generated by this rift parallel shortening is ten times lower than the extensional force and likely represents the resistance of the strike-slip fault developing in strong lower crust models which is several hundreds of kilometres long. Such a force may be lowered by a more efficient strain softening in the brittle crust.

5.2. Ridge migration and comparison with previous modelling work

Models presented in this study show that a small amount (12% to 20%) of rift parallel shortening leads to alternation of ridge propagation phases and ridge stalling phases. These alternating phases are each associated with strain regime changes. During ridge propagation, the deformation in the continental crust located at the front of the propagator is dip-slip dominated, while during stalling phases the deformation in the continental crust is strike-slip dominated.

According to our models results, the ridge propagation occurs under three different mechanisms (Figure 6 and Figure 14). For low rift-parallel shortening velocities (12% to 15% of the extension rate, Figure 4a, b, c) the ridge propagation is fast (between 1.5 cm/yr and 3.5 cm/yr, Figure 6b) and straight. Therefore, in this case the ridge propagation velocity exceeds the total extension rate (1 cm/yr). This propagation mode has been documented in crustal scale models involving a low viscosity layer beneath a plastic layer (Allken et al., 2012) where the lower viscosities lead to faster propagation. The same behaviour is observed in lithospheric scale models with cylindrical boundary conditions (Le Pourhiet et al., 2018) where free slip boundary conditions prevent deformation in the rift parallel

direction (which is equivalent to a boundary opposing no resistance to propagation) but with the mantle acting as the viscous layer controlling the propagation rate.

Higher rift-parallel shortening velocities (15% to 17%, Figure 4d) lead to ridge propagation by pulses alternating between fast propagation (between 0.5 cm/yr and 1.5 cm/yr, Figure 6b) and stalling phases. This propagation mode obviously requires that the ridge propagation stalls at some point. Until now, this behaviour has been modelled only during soft linkage of en-échelon propagators (Ammann et al., 2017; Liao & Gerya, 2015; Le Pourhiet et al., 2017) and a rift parallel inflow boundary condition model (Le Pourhiet et al., 2018, and this study). The common feature between these models is that during the stalling phase the velocity field at the tip of propagator changes and leads to a strike-slip dominated deformation. This mode of propagation was also proposed by Courtillot (1982) in the Gulf of Aden to explain the obliquity between seafloor magnetic anomalies and the passive margin. In Courtillot's (1982) model, a hard node in the lithosphere structure would explain why the ridge propagation stopped. All these thermo-mechanical and conceptual models show that to enter in a stalling ridge phase, a non-cylindrical parameter is required. It can be rheological (Courtillot, 1982) or kinematic (Ammann et al., 2017; Liao & Gerya, 2015; Le Pourhiet et al., 2017, 2018).

Finally, for higher velocities (17% to 20%, Figure 4e, f, g) a ridge jump propagation mode occurs. This ridge propagation mode also requires a ridge stalling phase, but rather than resuming from the ridge tip, a new spreading centre localizes between 100 and 200 km away from the ridge (Figure 6a and Figure 11d). This propagation mode is particularly well observed in models evolving towards three distinct tectonic plates (Figure 4e, g, h), therefore for highly oblique kinematic conditions. A propagation by jump was also obtained in analogue models with rotational extension coupled with an oblique weak seed (Molnar et al., 2017) where only highly oblique weak zones lead to jumps of spreading centres. Effectively, a weak zone oblique to the ridge propagation direction has the same effect as a strong zone or an inflow in the rift parallel direction. In the first case it will guide the deformation, while in the second case it will force the deformation to propagate elsewhere.

These different propagation modes are highly dependent of the lithosphere ability to deform in the rift parallel direction, whether it is due to rheological heterogeneities (e.g. Courtillot, 1982; Molnar et al., 2017) or velocity field variations (e.g. Ammann et al., 2017; Le Pourhiet et al., 2017, 2018).

5.3. Comparison with natural cases

Although the four oceanic basins presented in Figure 1 are not at same time and space scales, they all share similar features between each other and with our models. All of them are well documented examples of ridge propagation characterized by pulses of alternating fast propagation along straight margins and stalling phases which often corresponds to the location of oblique and transform margins. Our numerical simulations show that during propagation phases the deformation occurs orthogonal to the rift axis whereas oblique extension and strike-slip deformation characterises stalling phases. Initiation of strike-slip deformation results in the reorientation of pre-existing rift normal faults. As stretching continues an oblique continental rift forms which may result in the formation of a transform margin or oblique continental break-up propagation (Figure 14). A first order observation is that as in the natural systems described in this study, none of the models involving rift parallel shortening produce shortening structures (as thrusts or folds). However, the non-cylindrical component of the velocity field is well expressed in the GPS velocities of active oblique rift systems (Gulf of Aden and Northern Red Sea basins).

Among the four oceanic basins in Figure 1, the South China Sea is probably the most compelling example of secondary oblique continental break-up propagation following a stalling phase. Figure 1b

576 outlines three V-shaped propagators which indicate that continental break-up propagated from east to
577 west by pulses of rapid propagation, separated by two periods of stalling. The longest period occurred
578 from 32 to 23 Ma where magnetic anomalies in the propagator region form at a 60° angle with the
579 continental margin indicating that the propagation rate is much lower than the spreading rate. Le
580 Pourhiet et al. (2018) proposed that rift parallel inflow could explain this first phase of stalling using a
581 model similar to the one with a weak crust and 1.2 mm/yr rift parallel inflow. They then posit that this
582 long stalling period ends when both the spreading and propagating direction of the mid-oceanic ridge
583 changed by 15° (Sibuet et al., 2016) in the South China Sea (Figure 1b). The origin of this rift parallel
584 inflow has been proposed as the expression of gravitational potential energy gradient (e.g. Ding & Li,
585 2016; Le Pourhiet et al., 2018) between the developing basin and the higher Indochina to the west.
586 Bird et al. (2006) showed that a topographic load can propagate vertical forces of 1 TN/m to 2 TN/m
587 over hundreds of kilometres. Here, with a more thorough exploration of the effect of shortening on the
588 propagation of continental break up, we show that increasing the shortening rate to 1.5 mm/yr
589 reproduces the 15° rotation of the spreading direction reported by Sibuet et al. (2016) and the
590 propagation of a younger oceanic propagator self consistently. Moreover, associated to the strike-slip
591 deformation, lower crust domes exhumed in the continental margin as in the South China Sea. This
592 finding questions whether magnetic anomalies do record a change in far field conditions or simply the
593 localization of the deformation on an oblique continental rift branch.

594 This result may have significant implications on plates reconstruction and geodynamic
595 interpretations. Indeed, changes of structural directions are common during continental rifting. These
596 phases coincide with a distribution of the deformation in the crust in front of the oceanic propagator as
597 in the Atlantic realm (Enachescu, 2006; Mohn et al., 2015; Nirrengarten et al., 2018; Sibuet et al.,
598 2007), in the Gulf of Aden (Fournier et al., 2010; Leroy et al., 2012) or in the northern Red Sea
599 (Almalki et al., 2016; Bartov et al., 1980; Bosworth, 2015; 2020) where major transform fault zones
600 are segmenting the oceanic basins at locations where stalling phases occurred.

601 Previous studies proposed that these stalling phases can be related to several processes acting as
602 an opposing force to the rift propagation in the undeformed continental lithosphere like the presence of
603 high strength continental domains or inherited structures in the lithosphere (Courtillot, 1982; Morgan
604 & Parmentier, 1985; Nirrengarten et al., 2018). As a result they are generally interpreted as indicating
605 varying kinematic conditions or inheritance. This hypothesis was already questioned by numerical
606 model of rift linkage at oblique continental margin which showed that with increasing strain
607 localisation the tectonic regime changes from initially dip-slip dominated to strike-slip dominated (Le
608 Pourhiet et al., 2017). Here, we show that strain partitioning and change in deformation regime can
609 occur at the tip of oceanic propagators under constant boundary kinematic conditions (Figures 9 and
610 11). This implies that a relationship seems to exist between major transform margins and the rift
611 stalling phase during ocean opening. In the North Atlantic Ocean we observe that the Newfoundland-
612 Azores-Gibraltar fault zone and the Charlie-Gibbs fault are the two major transform faults separate
613 segment of different spreading ages (Nirrengarten et al., 2018 and references therein). In the same
614 way, the Gulf of Aden (Figure 1a) opened almost synchronously between the Alula-Fartak fault zone
615 and the Shukra el Sheik fault zone but was stalling along the Shukra el Sheik fault zone for 9 Ma to 12
616 Ma before propagation restarted (Fournier et al., 2010; Leroy et al., 2012). However, in our models we
617 were not able to produce oceanic transform faults segmenting the ridge. Thermo-mechanical models
618 tend to show that magmatism and serpentinization are essential to produce oceanic transform faults
619 (e.g. Ammann et al., 2017; Gerya, 2012).

620 The South China Sea and the Gulf of Aden are both good examples to illustrate the rifting of a
621 weak continental lithosphere (Figure 4b, d and Figures 7 and 9) due to the geodynamic context in

which those two oceanic basins have formed. The Gulf of Aden is located near an important mantle thermal anomaly (the Afar hot spot, *e.g.* Fournier et al., 2010; Leroy et al., 2012) while the South China sea opening followed a long-term subduction associated with magmatism (Pubellier et al., 2003; Wang et al., 2013). The Gulf of Aden presents the advantage of being an active plate boundary monitored with GPS stations. The GPS velocities with respect to Arabia fixed (ArRajehi et al., 2010) display the Eastward motion (*i.e.* parallel to the ridge) of the continent at the tip of the oceanic propagator and an oblique North-eastward motion for Africa fixed. The oceanic basin and continental margins display stalling phases and propagation phases and the rifting phase is characterized as a wide oblique rift. Figure 13c shows the relative velocity field in the Model 1 (1.2 mm/yr rift parallel inflow) which display the oblique velocity at the tip of the propagator. In the Gulf of Aden, the force that triggers the rift parallel inflow as displayed by GPS velocities could come from the differential gravitational potential energy between the high topography above the Afar hot spot and the opening Gulf of Aden.

However, the strain localization and the evolution of the deformation in a stronger lithosphere (Figure 4e, g, h and Figure 11) results in the development of a localized strike-slip fault and the formation of a micro-plate. These models display similarities with the Northern Red Sea and its two branches, the Gulf of Aqaba to the East and the Gulf of Suez to the West (Almalki et al., 2016; Bartov et al., 1980; Bosworth, 2015; 2020). In the Northern Red Sea, the Sinai block behaves as a third tectonic micro-plate separated from Arabia by the Levant transform Fault and from Africa by an active oblique rift (with respect to the Red Sea). Although a continental rift is not a complete plate boundary it is the initiation. The GPS velocities with respect to Arabia display the southward motion of the Sinai block along the Levant Fault. Therefore, regarding the absence of shortening structures in the northern Red Sea, the GPS velocities showing the rift parallel displacement and the presence of the transform continental fault of the Levant, we propose that a rotational extension model (Molnar et al., 2017; Smith, 1993; Zwaan et al., 2020) may not be the best candidate to explain the dynamics of the Northern Red Sea. However, the formation of the triple junction between a continental transform fault, an active oblique rift and a ridge is well reproduced in our strong lower crust models when the rift parallel inflow is greater than 16% of extension velocity (Figure 14b, model 3). The geodynamic origin of this regional rift parallel inflow arise from the contrast between the fast motion of the Arabian plate driven by subduction in Zagros and Makran and the slow northward motion of Africa in collision with the European plate.

Moreover, the lithosphere-scale strain partitioning between a highly localized strike-slip fault and a very oblique rift leads to the development of two types of transform margins with extremely different kinematic, dynamics and thermal history.

The classical model for transform margins formation involves the development of transform faults during the intra-continental rifting phase (Basile, 2015; Francheteau & Le Pichon, 1972; Mascle & Blarez, 1987). The evolution from rifting to oceanic spreading should preserve the continental structures and develop oceanic transform faults along the previous continental ones. This model has major implications on development of the margin structure and temperature. It implies that oceanic ridge migrates along the transform margin as a triple junction at half the spreading rate (for large offset transform margin it may take several million years) heating the continental margin and its basins (Ruepke et al. 2010). It also implies that the strike-slip faults responsible for the steep structure of the transform margins (Mercier de Lépinay et al., 2016) are active at the beginning of the rifting phase and that the transform margin was born with characteristics observable today on the fossil transform margins. However, models of this study show different characteristic and evolution for transform margins.

On the one hand, models with high obliquity display the typical kinematic context of a transform margin (Basile, 2015; Francheteau & Le Pichon, 1972). In this condition one of the oblique continental rift branch (Figure 11, cross sections 1 and 2, Figure 12) reach continental break-up and the system evolves into two en-échellon ridges separated by a zone of soft linkage that evolves into an oblique rift with strike-slip deformation similarly to previous studies (Ammann et al., 2017; Liao & Gerya, 2015; Laetitia Le Pourhiet et al., 2017). However, structural properties of these modelled margins correspond to hyper-thinned continental crust but not steep margins along a localized transform fault. On the other hand, simulations with high obliquity and strong lower crust produce steep margins along a pure strike-slip fault (e.g. Figure 11f, corss section 4) that display the structural characteristics of a transform margin as a slope break, a marginal ridge close to the slope break and a strike-slip fault separating the continental crust from the exhumed mantle (or oceanic crust)(Mercier de Lépinay et al., 2016). However, these strike-slip margins form in atypical kinematic context for a transform margin. From an initially hyper thinned margin developed during the initial extension phase (until 30Myr), a steep margin develops (Figure 11e, f, cross section 4) as the deformation increases along the strike-slip fault. This strike-slip shear zone puts in contact a 50 km thick continental lithosphere with the asthenosphere. As a result, the formation these steep margins starts 10 Myr after the continental break-up meaning that the margin geometry is acquired after the intra-continental rifting phase. These results highlight that oblique and transform margins can acquire their final structure several million years after continental break-up due to the strike-slip deformation.

6. Conclusion

3D thermo-mechanical numerical models integrating a rift parallel inflow show that:

- The 3D boundary conditions can be interpreted as an absolute velocity field that can be converted to a relative velocity field comparable with natural cases of triple junctions involving continental transform faults.
- At least three propagation modes can exist. A fast propagation resulting in a straight oceanic basin (2 plates), an alternation of propagation and stalling phases (extension opposing velocity of 12% to 15% for weak lower crust) and propagation with a “jump” of the spreading centre (opposing extension velocity over 17% for strong and weak lower crust).
- .
- Continental break-up propagation slows down when rift parallel inflow is applied (Le Pourhiet et al., 2018) and we show that during these stalling phases the continental margins and tectonic structures rotate and the deformation regime changes from dip-slip dominated to strike-slip dominated while boundary kinematic conditions are kept constant.
- There might be a relationship between stalling phases, transform margin formation and the subsequent location of large oceanic transform fault zones due to margins rotation
- Two modes of transform margin can develop, steep margin associated with strike slip faults and hyper-extended ones associated with offset spreading segment similar to Le Pourhiet (2017)
- All oblique margins might not emerge from a single geodynamic process and that it could be worthwhile in the future to revisit the available data using the soft linkage oblique margin and the strike-slip oblique margin models as a classification

Acknowledgment

This study is part of the Orogen program, a tripartite research collaboration between CNRS, BRGM, and TOTAL SA. We thank TOTAL SA for the computing time on Pangea supercomputer. pTatin3D is an open source code, which is available publicly at <https://bitbucket.org/ptatin/ptatin3d/src/master/>. The version of the code used in this study, an example options file to reproduce the results and the models results are available publicly at <http://doi.org/10.5281/zenodo.3967184>. Maps from figure 1 were made with GeomapApp and post-processing with Paraview.

Figure captions

Figure 1: Topographic and bathymetric maps (GMRT, (Ryan et al., 2009) with oceanic crust ages of a) the Gulf of Aden (Fournier et al., 2010; Leroy et al., 2012), c) the South China Sea (Li et al., 2014; Le Pourhiet et al., 2018), e) the Red Sea (Almalki et al., 2016; Bosworth et al., 2020), g) the North Atlantic Ocean (Klitgord & Schouten, 1986). Main transform fault zones are reported with thick black lines. GPS velocities are from (ArRajehi et al., 2010; Reilinger & McClusky, 2011). Graphs with red curves represent the ridge propagation in b) the Gulf of Aden, d) the South China Sea, f) the Red Sea, h) the North Atlantic Ocean interpreted from the oceanic seafloor ages distribution along strike. Topographic and bathymetric maps made with GeoMapApp (www.geomapapp.org) / CC BY.

Figure 2: Kinematic models of ridge propagation and continental rift geometry. a) Fast ridge propagation (Vink, 1982), b) slow ridge propagation (Vink, 1982), c) Scissor opening (Hey et al., 1980). Dynamic models of the formation of oceanic V-shaped propagators and continental rifts d) strong ribbon in the continental crust slowing ridge propagation (Courtillot, 1982), e) offset propagators (Le Pourhiet et al., 2017), f) shortening perpendicular to extension direction (Le Pourhiet et al., 2018).

Figure 3: a) Numerical models setup. Black arrows display the initial velocity field in the models domain. b) Velocity boundary conditions to solve the conservation of momentum. c) Yield stress envelopes of the models domain computed for a constant strain rate of 10^{-14} s^{-1} and an initial geotherm represented by the red curve.

Figure 4: Map view of nine models with a strong and weak lower crust and varying v_z velocities displaying the plastic strain and the mantle exhumation age. The plastic strain represents all the deformations that occurred under the regime described by the equation 6. The mantle exhumation age is set when the mantle cools beneath the isotherm 800°C .

Figure 5: Serial cross-sections of nine models with a strong and weak lower crust and varying v_z velocities representing the plastic strain and the viscous strain. For every models, the cross-sections are regularly spaced every 50 km from $z=150 \text{ km}$ to $z=500 \text{ km}$. The plastic strain represents all the deformations that occurred under the regime described by the equation 6. The viscous strain represents all the deformations that occurred under the regime described by the equation 7.

Figure 6: a) Curves for every models representing the position of the tip along the z axis (in km) with respect to time (in Myr). b) Curves for every models representing the velocity of the rift propagation (in mm/yr) with respect to time (in Myr). The velocity of the rift propagation is computed as the time derivative of the position of the tip. c) Evolution of the tectonic forces through time. F_x (left panel) corresponds to the forces applied in the direction normal to the face of normal x (equation 14a) and F_z

corresponds to the forces applied in the direction normal to the face of normal z (equation 14b) d)
Diagram showing the rift evolution of every model in time for both strong and weak lower crust.

Figure 7: Numerical model with a weak lower crust and a v_z shortening velocity of 1.2 mm/yr (model 1). Left panels show a map view evolution of the plastic strain (deformation related to equation 6) in the crust and exhumation ages of the mantle below the 800°C isotherm. Red dashed lines shows the locations of cross-sections displayed in the right panels. On the cross-sections, background colours correspond to the simulated lithologies presented in Figure 3. The deformation drawing is interpreted from Supplementary Figures 2 and 3.

Figure 8: Numerical model with a weak lower crust and a v_z shortening velocity of 1.2 mm/yr (model 1). The background colours represent the lithologies detailed in Figure 3. The superimposed colours represent the stress inferred deformation regime (Brune, 2014; Brune & Autin, 2013). The intensity of the colourscale is related to the intensity of the strain rate tensor second invariant.

Figure 9: Numerical model with a weak lower crust and a v_z shortening velocity of 1.5 mm/yr (model 2). Left panels show a map view evolution of the plastic strain (deformation related to equation 6) in the crust and exhumation ages of the mantle below the 800°C isotherm. Red dashed lines shows the locations of cross-sections displayed in the right panels. On the cross-sections, background colours correspond to the simulated lithologies presented in Figure 3. The deformation drawing is interpreted from Supplementary Figures 5 and 6.

Figure 10: Numerical model with a weak lower crust and a v_z shortening velocity of 1.5 mm/yr (model 2). The background colours represent the lithologies detailed in Figure 3. The superimposed colours represent the stress inferred deformation regime (Brune, 2014; Brune & Autin, 2013). The intensity of the colourscale is related to the intensity of the strain rate tensor second invariant.

Figure 11: Numerical model with a strong lower crust and a v_z shortening velocity of 1.7 mm/yr (model 3). Left panels show a map view evolution of the plastic strain (deformation related to equation 6) in the crust and exhumation ages of the mantle below the 800°C isotherm. Red dashed lines shows the locations of cross-sections displayed in the right panels. On the cross-sections, background colours correspond to the simulated lithologies presented in Figure 3. The deformation drawing is interpreted from Supplementary Figures 8 and 9.

Figure 12: Numerical model with a strong lower crust and a v_z shortening velocity of 1.7 mm/yr (model 3). The background colours represent the lithologies detailed in Figure 3. The superimposed colours represent the stress inferred deformation regime (Brune, 2014; Brune & Autin, 2013). The intensity of the colourscale is related to the intensity of the strain rate tensor second invariant.

Figure 13: Map view of the horizontal velocity field extracted from model 3 ($v_z = 1.7$ mm/a, strong lower crust) at 50 Myr. a) Absolute horizontal velocity field. b) Relative horizontal velocity field (plate A fixed). c) Relative horizontal velocity field (plate C fixed). d) Relative horizontal velocity field (plate B fixed)

Figure 14: Synthetic sketch representing the three propagation modes highlighted by models of this study. Propagation modes are related to the degree of obliquity and strain partitioning. Relative velocities vector directions are extracted from the model with a strong lower crust and a shortening velocity $v_z=1.7$ mm/yr.

References

- Allken, V., Huismans, R. S., & Thieulot, C. (2012). Factors controlling the mode of rift interaction in brittle-ductile coupled systems: A 3D numerical study. *Geochemistry, Geophysics, Geosystems*, 13(5), 1–18. <https://doi.org/10.1029/2012GC004077>
- Almalki, K. A., Betts, P. G., & Ailleres, L. (2016). Incipient seafloor spreading segments: Insights from the Red Sea. *Geophysical Research Letters*, 43, 2709–2715. <https://doi.org/10.1002/2016GL068069>.Received
- Ammann, N., Liao, J., Gerya, T., & Ball, P. (2017). Oblique continental rifting and long transform fault formation based on 3D thermomechanical numerical modeling. *Tectonophysics*, (February), 1–16. <https://doi.org/10.1016/j.tecto.2017.08.015>
- ArRajehi, A., McClusky, S., Reilinger, R., Daoud, M., Alchalbi, A., Ergintav, S., ... Kogan, L. (2010). Geodetic constraints on present-day motion of the Arabian Plate: Implications for Red Sea and Gulf of Aden rifting. *Tectonics*, 29(3), 1–10. <https://doi.org/10.1029/2009TC002482>
- Bartov, Y., Steinitz, G., Eyal, M., & Eyal, Y. (1980). Sinistral movement along the Gulf of Aqaba - its age and relation to the opening of the Red Sea. *Nature*, 285, 220–222.
- Basile, C. (2015). Tectonophysics Transform continental margins — part 1 : Concepts and models. *Tectonophysics*, 661, 1–10. <https://doi.org/10.1016/j.tecto.2015.08.034>
- Beaussier, S. J., Gerya, T. V., & Burg, J. (2019). Near-ridge initiation of intraoceanic subduction : Effects of inheritance in 3D numerical models of the Wilson Cycle. *Tectonophysics*, 763(April), 1–13. <https://doi.org/10.1016/j.tecto.2019.04.011>
- Benes, V., & Scott, S. D. (1996). Oblique rifting in the havre trough and its propagation into the continental margin of New Zealand: Comparison with analogue experiments. *Marine Geophysical Research*, 18(2–4), 189–201. <https://doi.org/10.1007/BF00286077>
- Bird, P., Ben-Avraham, Z., Schubert, G., Andreoli, M., & Viola, G. (2006). Patterns of stress and strain rate in southern Africa. *Journal of Geophysical Research: Solid Earth*, 111(8), 1–14. <https://doi.org/10.1029/2005JB003882>
- Bosworth, W., Khalil, S. M., Ligi, M., Stockli, D. F., & McClay, K. R. (2020). Geology of Egypt: The Northern Red Sea. In Z. Hamimi, A. El-Barkooky, J. Martínez Frias, H. Fritz, & Y. A. El-Rahman (Eds.), *Geology of Egypt* (Springer, pp. 343–374). Cham, Switzerland: Springer Nature Switzerland AG. <https://doi.org/10.1007/978-3-030-15265-9>
- Bosworth, William. (2015). Geological Evolution of the Red Sea : Historical Background, Review, and Synthesis. In N. M. A. Rasul & I. C. F. Stewart (Eds.), *The Red Sea* (Springer E, pp. 45–78). Berlin Heidelberg: Springer. <https://doi.org/10.1007/978-3-662-45201-1>
- Briais, A., Patriat, P., & Tapponnier, P. (1993). Updated interpretation of magnetic anomalies and seafloor spreading stages in the south China Sea: Implications for the Tertiary tectonics of Southeast Asia. *Journal of Geophysical Research Solid Earth*, 98(B4), 6299– 6328. <https://doi.org/10.1029/92JB02280>
- Brune, S. (2014). Evolution of stress and fault patterns in oblique rift systems: 3-D numerical lithospheric-scale experiments from rift to breakup. *Geochemistry, Geophysics, Geosystems*, 15, 3392–3415. <https://doi.org/10.1002/2014GC005446>.Received
- Brune, S., & Autin, J. (2013). The rift to break-up evolution of the Gulf of Aden: Insights from 3D numerical lithospheric-scale modelling. *Tectonophysics*, 607, 65–79. <https://doi.org/10.1016/j.tecto.2013.06.029>

- 834 Brune, S., Corti, G., & Ranalli, G. (2017). Controls of inherited lithospheric heterogeneity on rift
835 linkage: Numerical and analog models of interaction between the Kenyan and Ethiopian rifts
836 across the Turkana depression. *Tectonics*, 36(9), 1767–1786.
837 <https://doi.org/10.1002/2017TC004739>
- 838 Brune, S., Popov, A. A., & Sobolev, S. V. (2012). Modeling suggests that oblique extension facilitates
839 rifting and continental break-up. *Journal of Geophysical Research*, 117(B08402), 1–16.
840 <https://doi.org/10.1029/2011JB008860>
- 841 Le Calvez, J., & Vendeville, B. (2002). Experimental designs to model along-strike fault interaction
842 fault interaction. *Journal of the Virtual Explorer*, 7, 1–17.
843 <https://doi.org/10.3809/jvirtex.2002.00043>
- 844 Corti, G. (2012). Tectonophysics Evolution and characteristics of continental rifting : Analog
845 modeling-inspired view and comparison with examples from the East African Rift System.
846 *Tectonophysics*, 522–523, 1–33. <https://doi.org/10.1016/j.tecto.2011.06.010>
- 847 Courtillot, V. (1982). Propagating rifts and continental breakup. *Tectonics*, 1(3), 239–250.
- 848 Ding, W., & Li, J. (2016). Propagated rifting in the Southwest Sub-basin , South China Sea : Insights
849 from analogue modelling. *Journal of Geodynamics*, 100, 71–86.
850 <https://doi.org/10.1016/j.jog.2016.02.004>
- 851 Duclaux, G., Huismans, R. S., & May, D. A. (2020). Rotation , narrowing , and preferential
852 reactivation of brittle structures during oblique rifting. *Earth and Planetary Science Letters*, 531.
853 <https://doi.org/10.1016/j.epsl.2019.115952>
- 854 Enachescu, M. (2006). Structural Setting and Petroleum Potential of the Orphan Basin , offshore
855 Newfoundland and Labrador. *Canadian Society Exploration Geophysics Recreation*, 31(2), 5–
856 13.
- 857 Fournier, M., Rooke, N. C., Petit, C., Huchon, P., Kathiri, A. Al, Audin, L., ... Merkouriev, S. (2010).
858 Arabia - Somalia plate kinematics , evolution of the Aden - Owen - Carlsberg triple junction ,
859 and opening of the Gulf of Aden. *Journal of Geophysical Research*, 115(B04102), 1–24.
860 <https://doi.org/10.1029/2008JB006257>
- 861 Francheteau, J., & Le Pichon, X. (1972). Marginal Fracture Zones as Structural Framework of
862 Continental Margins in South Atlantic Ocean. *AAPG Bulletin*, 56(6), 991–1007.
863 <https://doi.org/10.1306/819a40a8-16c5-11d7-8645000102c1865d>
- 864 Gerya, T. (2012). Origin and models of oceanic transform faults. *Tectonophysics*, 522–523, 34–54.
865 <https://doi.org/10.1016/j.tecto.2011.07.006>
- 866 Gerya, T., & Burov, E. (2018). Nucleation and evolution of ridge-ridge-ridge triple junctions :
867 Thermomechanical model and geometrical theory. *Tectonophysics*, 746, 83–105.
868 <https://doi.org/10.1016/j.tecto.2017.10.020>
- 869 Gerya, T. V. (2013). Three-dimensional thermomechanical modeling of oceanic spreading initiation
870 and evolution. *Physics of the Earth and Planetary Interiors*, 214, 35–52.
871 <https://doi.org/10.1016/j.pepi.2012.10.007>
- 872 Hey, R., Duennebier, F. K., & Morgan, W. J. (1980). Propagating rifts on midocean ridges. *Journal of*
873 *Geophysical Research*, 85(B7), 3647–3658.
- 874 Hirth, G., & Kohlstedt, D. L. (2003). Rheology of the Upper Mantle and the Mantle Wedge: A View
875 from the Experimentalists. *Geophysical Monograph*, 138, 83–105.
- 876 Holm, R. J., Rosenbaum, G., & Richards, S. W. (2016). Post 8 Ma reconstruction of Papua New

877 Guinea and Solomon Islands: Microplate tectonics in a convergent plate boundary setting. *Earth-*
878 *Science Reviews*, 156, 66–81. <https://doi.org/10.1016/j.earscirev.2016.03.005>

879 Jolivet, L., Famin, V., Mehl, C., Parra, T., Aubourg, C., Hébert, R., & Philippot, P. (2004). Strain
880 localization during crustal-scale boudinage to form extensional metamorphic domes in the
881 Aegean Sea. In D. L. Whitney, C. Teyssier, & C. S. Siddoway (Eds.), *Gneiss domes in orogeny*
882 (Geological, pp. 185–210). Boulder, Colorado: Geological Society of America.

883 Jourdon, A., Le Pourhiet, L., Mouthereau, F., & Masini, E. (2019). Role of rift maturity on the
884 architecture and shortening distribution in mountain belts. *Earth and Planetary Science Letters*,
885 512, 89–99. <https://doi.org/10.1016/j.epsl.2019.01.057>

886 Kameyama, M., Yuen, D. A., & Karato, S. (1999). Thermal-mechanical effects of low-temperature
887 plasticity (the Peierls mechanism) on the deformation of a viscoelastic shear zone. *Earth and*
888 *Planetary Science Letters*, 168, 159–172.

889 Klitgord, K., & Schouten, H. (1986). Plate kinematics of the central Atlantic. In P. R. Vogt & B. E.
890 Tucholke (Eds.), *The Western North Atlantic Region* (DNAG, pp. 351–378). Boulder, Colorado:
891 Geological Society of America.

892 Leroy, S., Razin, P., Autin, J., Bache, F., D’Acremont, E., Watremez, L., ... Al Lazki, A. (2012).
893 From rifting to oceanic spreading in the Gulf of Aden : A synthesis From rifting to oceanic
894 spreading in the Gulf of Aden : a synthesis. *Arabian Journal of Geosciences*, 5, 859–901.
895 <https://doi.org/10.1007/s12517-011-0475-4>

896 Li, C.-F., Xu, X., Lin, J., Sun, Z., Zhu, J., Yao, Y., ... Zhang, G.-L. (2014). Ages and magnetic
897 structures of the South China Sea constrained by deep tow magnetic surveys and IODP
898 Expedition 349. *Geochemistry, Geophysics, Geosystems*, 15, 4958–4983.
899 <https://doi.org/10.1002/2014GC005567>

900 Liao, J., & Gerya, T. (2015). From continental rifting to sea floor spreading : Insight from 3D thermo-
901 mechanical modeling. *Gondwana Research*, 28(4), 1329–1343.
902 <https://doi.org/10.1016/j.gr.2014.11.004>

903 Mart, Y., & Dauteuil, O. (2000). Analogue experiments of propagation of oblique rifts.
904 *Tectonophysics*, 316, 121–132.

905 Mascle, J., & Blarez, E. (1987). Evidence for transform margin evolution from the Ivory Coast–Ghana
906 continental margin. *Nature*, 326(6111), 378–381.

907 May, D. A., Brown, J., & Le Pourhiet, L. (2014). pTatin3D : High-Performance Methods for Long-
908 Term Lithospheric Dynamics. *Proceeding SC'14 Proceedings of the International Conference*
909 *for High Performance Computing, Networking, Storage and Analysis*, 274–284.

910 May, D. A., Brown, J., & Le Pourhiet, L. (2015). A scalable , matrix-free multigrid preconditioner for
911 finite element discretizations of heterogeneous Stokes flow. *Computer Methods in Applied*
912 *Mechanics and Engineering*, 290, 496–523. <https://doi.org/10.1016/j.cma.2015.03.014>

913 Mazur, S., Green, C., Stewart, M. G., Whittaker, J. M., Williams, S., & Bouatmani, R. (2012).
914 Displacement along the Red River Fault constrained by extension estimates and plate
915 reconstructions. *Tectonics*, 31(5), 1–22. <https://doi.org/10.1029/2012TC003174>

916 McClay, K. R., & White, M. J. (1995). Analogue modelling of orthogonal and oblique rifting. *Marine*
917 *and Petroleum Geology*, 12(2), 137–151.

918 Mercier de Lépinay, M., Loncke, L., Basile, C., Roest, W. R., Patriat, M., Maillard, A., & Clarens, P.
919 De. (2016). Transform continental margins – Part 2 : A worldwide review. *Tectonophysics*, 693,
920 96–115. <https://doi.org/10.1016/j.tecto.2016.05.038>

- 921 Mohn, G., Karner, G., Manatschal, G., & Johnson, C. (2015). Structural and stratigraphic evolution of
922 the Iberia and Newfoundland hyper-extended rifted margins: A quantitative modeling approach.
923 *Geological Society, London, Special Publications*, 16(June), 9156.
924 <https://doi.org/10.1144/SP413.9>
- 925 Molnar, N. E., Cruden, A. R., & Betts, P. G. (2017). Interactions between propagating rotational rifts
926 and linear rheological heterogeneities: Insights from three-dimensional laboratory experiments.
927 *Tectonics*, 36, 420–443. <https://doi.org/10.1002/2016TC004447>
- 928 Mondy, L. S., Rey, P. F., Duclaux, G., & Moresi, L. (2018). The role of asthenospheric flow during
929 rift propagation and breakup. *Geology*, 46(2), 103–106.
- 930 Morgan, J. P., & Parmentier, E. M. (1985). Causes and rate-limiting mechanisms of ridge propagation:
931 A fracture mechanics model. *Journal of Geophysical Research*, 90(B10), 8603–8612.
- 932 Müller, D. R., Sdrolias, M., Gaina, C., & Roest, W. R. (2008). Age, spreading rates, and spreading
933 asymmetry of the world's ocean crust. *Geochemistry, Geophysics, Geosystems*, 9(4), 1–19.
934 <https://doi.org/10.1029/2007GC001743>
- 935 Müller, R. D., Seton, M., Zahirovic, S., Williams, S. E., Matthews, K. J., Wright, N. M., ... Cannon, J.
936 (2016). Ocean Basin Evolution and Global-Scale Plate Reorganization Events Since Pangea
937 Breakup. *Annual Review of Earth and Planetary Sciences*, 44(1), 107–138.
938 <https://doi.org/10.1146/annurev-earth-060115-012211>
- 939 Naliboff, J. B., Glerum, A., Brune, S., Péron-Pinvidic, G., & Wrona, T. (2020). Development of 3-D
940 Rift Heterogeneity Through Fault Network Evolution. *Geophysical Research Letters*, 47(13), 0–
941 3. <https://doi.org/10.1029/2019GL086611>
- 942 Nirrengarten, M., Manatschal, G., Tugend, J., Kuszniir, N., & Sauter, D. (2018). Kinematic Evolution
943 of the Southern North Atlantic: Implications for the Formation of Hyperextended Rift Systems.
944 *Tectonics*, 37(1), 89–118. <https://doi.org/10.1002/2017TC004495>
- 945 Le Pourhiet, L., Huet, B., & Traoré, N. (2014). Links between long-term and short-term rheology of
946 the lithosphere: Insights from strike-slip fault modelling. *Tectonophysics*, 631(C), 146–159.
947 <https://doi.org/10.1016/j.tecto.2014.06.034>
- 948 Le Pourhiet, Laetitia, Huet, B., May, D. A., Labrousse, L., & Jolivet, L. (2012). Kinematic
949 interpretation of the 3D shapes of metamorphic core complexes. *Geochemistry, Geophysics,*
950 *Geosystems*, 13(9), 1–17. <https://doi.org/10.1029/2012GC004271>
- 951 Le Pourhiet, Laetitia, May, D. A., Huille, L., Watremez, L., & Leroy, S. (2017). A genetic link
952 between transform and hyper-extended margins. *Earth and Planetary Science Letters*, 465, 184–
953 192. <https://doi.org/10.1016/j.epsl.2017.02.043>
- 954 Le Pourhiet, Laetitia, Chamot-Rooke, N., Delescluse, M., May, D. A., Watremez, L., & Pubellier, M.
955 (2018). Continental break-up of the South China Sea stalled by far-field compression. *Nature*
956 *Geoscience*. <https://doi.org/10.1038/s41561-018-0178-5>
- 957 Precigout, J., Gueydan, F., Gapais, D., Garrido, C. J., & Essaifi, A. (2007). Strain localisation in the
958 subcontinental mantle — a ductile alternative to the brittle mantle. *Tectonophysics*, 445, 318–
959 336. <https://doi.org/10.1016/j.tecto.2007.09.002>
- 960 Pubellier, M., Ego, F., Chamot-Rooke, N., & Rangin, C. (2003). The building of pericratonic
961 mountain ranges : structural and kinematic constraints applied to GIS-based reconstructions of
962 SE Asia. *Bulletin de La Societe Geologique de France*, 174(6), 561–584.
963 <https://doi.org/10.2113/174.6.561>
- 964 Ranalli, G., & Murphy, D. C. (1987). Rheological stratification of the lithosphere. *Tectonophysics*,

- 965 132, 281–295.
- 966 Reilinger, R., & McClusky, S. (2011). Nubia-Arabia-Eurasia plate motions and the dynamics of
 967 Mediterranean and Middle East tectonics. *Geophysical Journal International*, 186(3), 971–979.
 968 <https://doi.org/10.1111/j.1365-246X.2011.05133.x>
- 969 Rushbrooke, G. S., & Wood, P. J. (1958). On the Curie points and high temperature susceptibilities of
 970 Heisenberg model ferromagnetics. *Molecular Physics: An International Journal at the Interface*
 971 *Between Chemistry and Physics*, 1:3, 257–283.
 972 <https://doi.org/http://dx.doi.org/10.1080/00268975800100321>
- 973 Ryan, W. B. F., Carbotte, S. M., Coplan, J., O'Hara, S., Melkonian, A., Arko, R., ... Zemsky, R.
 974 (2009). Global Multi-Resolution Topography (GMRT) synthesis data set. *Geochemistry,*
 975 *Geophysics, Geosystems*, 10. <https://doi.org/10.1029/2008GC002332>
- 976 Rybacki, E., & Dresen, G. (2000). Dislocation and diffusion creep of synthetic anorthite aggregates.
 977 *Journal of Geophysical Research: Solid Earth*, 105(B11), 26017–26036.
 978 <https://doi.org/10.1029/2000JB900223>
- 979 Seton, M., Müller, R. D., Zahirovic, S., Gaina, C., Torsvik, T., Shephard, G., ... Chandler, M. (2012).
 980 Global continental and ocean basin reconstructions since 200Ma. *Earth-Science Reviews*, 113(3–
 981 4), 212–270. <https://doi.org/10.1016/j.earscirev.2012.03.002>
- 982 Sibuet, J.-C., Srivastava, S. P., Enachescu, M., & Karner, G. D. (2007). Early Cretaceous motion of
 983 Flemish Cap with respect to North America : implications on the formation of Orphan Basin and
 984 SE Flemish Cap – Galicia Bank conjugate margins. In G. D. Karner, G. Manatschal, & L. M.
 985 Pinheiro (Eds.), *Imaging, Mapping and Modelling Continental Lithosphere Extension and*
 986 *Breakup* (Geological, pp. 63–76). Geological Society of London.
 987 <https://doi.org/10.1144/SP282.4>
- 988 Sibuet, Jean-Claude, Yeh, Y., & Lee, C. (2016). Geodynamics of the South China Sea.
 989 *Tectonophysics*, 692, 98–119.
- 990 Smith, J. V. (1993). Infinitesimal kinematics of rotational rifting with reference to en echelon marginal
 991 faults in the Red Sea region. *Tectonophysics*, 222(2), 227–235. [https://doi.org/10.1016/0040-](https://doi.org/10.1016/0040-1951(93)90050-T)
 992 [1951\(93\)90050-T](https://doi.org/10.1016/0040-1951(93)90050-T)
- 993 Turcotte, D. L., & Schubert, G. (2002). *Geodynamics. Cambridge University Press, Cambridge,*
 994 *Second Edition.* <https://doi.org/10.1007/s007690000247>
- 995 Vine, F. J., & Matthews, D. H. (1963). Magnetic Anomalies Over Oceanic Ridges. *Nature*, 4897, 947–
 996 949.
- 997 Vink, G. E. (1982). Continental rifting and the implications for plate tectonic reconstructions. *Journal*
 998 *of Geophysical Research*, 87(B13), 10677–10688.
- 999 Wang, Y., Fan, W., Zhang, G., & Zhang, Y. (2013). Phanerozoic tectonics of the South China Block:
 1000 Key observations and controversies. *Gondwana Research*, 23(4), 1273–1305.
 1001 <https://doi.org/10.1016/j.gr.2012.02.019>
- 1002 Watremez, L., Burov, E., D'Acremont, E., Leroy, S., Huet, B., Le Pourhiet, L., & Bellahsen, N.
 1003 (2013). Buoyancy and localizing properties of continental mantle lithosphere: Insights from
 1004 thermomechanical models of the eastern Gulf of Aden. *Geochemistry, Geophysics, Geosystems*,
 1005 14(8), 2800–2817. <https://doi.org/10.1002/ggge.20179>
- 1006 Van Wijk, J. W. ., & Blackman, D. K. (2005). Dynamics of continental rift propagation : the end-
 1007 member modes. *Earth and Planetary Science Letters*, 229, 247–258.
 1008 <https://doi.org/10.1016/j.epsl.2004.10.039>

1009 Zwaan, F., Schreurs, G., & Buiter, S. J. H. (2019). *A systematic comparison of experimental set-ups*
1010 *for modelling extensional tectonics. Solid Earth* (Vol. 10). [https://doi.org/10.5194/se-10-1063-](https://doi.org/10.5194/se-10-1063-2019)
1011 2019

1012 Zwaan, F., Schreurs, G., Naliboff, J., & Buiter, S. J. H. (2016). Insights into the effects of oblique
1013 extension on continental rift interaction from 3D analogue and numerical models.
1014 *Tectonophysics*, 693, 239–260. <https://doi.org/10.1016/j.tecto.2016.02.036>

1015 Zwaan, F., Schreurs, G., & Rosenau, M. (2020). Rift propagation in rotational versus orthogonal
1016 extension: Insights from 4D analogue models. *Journal of Structural Geology*, 135.
1017 <https://doi.org/10.1016/j.jsg.2019.103946>

1018

1019

1020

1021

Figure 1.

Gulf of Aden

South China Sea

Red Sea

North Atlantic Ocean

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.

Figure 7.

Weak Lower Crust
 $V_z = 1.2 \text{ mm/yr}$ (model 1)

Figure 8.

Figure 9.

Weak Lower Crust Vz = 1.5 mm/yr (model 2)

Figure 10.

Figure 11.

Strong Lower Crust
 $V_z = 1.7$ mm/yr (model 3)

Figure 12.

Strong Lower Crust
 $V_z = 1.7$ mm/yr (model 3)

Figure 13.

Figure 14.

Fast ridge propagation

Seafloor age

Pulses ridge propagation

Jump ridge propagation

Increasing obliquity and strain partitioning