



HAL
open science

First detection of *Francisella haliotica* in mussels *Mytilus* spp. experiencing mortalities in France

Maud Charles, Antonio Villalba, Gary Meyer, Suzanne Trancart, Coralie Lagy,
Ismaël Bernard, Maryline Houssin

► **To cite this version:**

Maud Charles, Antonio Villalba, Gary Meyer, Suzanne Trancart, Coralie Lagy, et al.. First detection of *Francisella haliotica* in mussels *Mytilus* spp. experiencing mortalities in France. *Diseases of Aquatic Organisms*, 2020, 140, pp.203-208. <10.3354/dao03505>. <hal-02987893>

HAL Id: hal-02987893

<https://hal.sorbonne-universite.fr/hal-02987893v1>

Submitted on 4 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

First detection of *Francisella haliotica* in mussels (*Mytilus* spp.) experiencing mortalities in France

Maud Charles^{1,2,1}, Antonio Villalba^{3,4,5}, Gary Meyer⁶, Suzanne Trancart², Coralie Lagy², Ismaël Bernard⁷, Maryline Houssin^{1,2}

¹ Normandie Université, Université de Caen Normandie, FRE BOREA, CNRS-2030, IRD-207, MNHN, UPMC, UCN, Esplanade de la Paix, 14032 Caen Cedex 4, France

² LABÉO Frank Duncombe, 1 Route de Rosel, 14053 Caen Cedex 4, France

³ Centro de Investigacións Mariñas (CIMA), Consellería do Mar, Xunta de Galicia, 36620 Vilanova de Arousa, Spain

⁴ Departamento de Ciencias de la Vida, Universidad de Alcalá, 28871 Alcalá de Henares, Spain

⁵ Research Centre for Experimental Marine Biology and Biotechnology (PIE), University of the Basque Country (UPV/EHU), 48620 Plentzia, Basque Country, Spain

⁶ Fisheries and Oceans Canada, Pacific Biological Station, Aquatic Animal Health Laboratory, 3190 Hammond Bay Road, Nanaimo, BC V9T 6N7

⁷ Eurêka Mer, 22740 Lézardrieux, France

¹Corresponding author: Maud Charles (maud.charles@outlook.fr)

2 **ABSTRACT**

3 This note describes the first detection of the bacteria *Francisella halioticida* in
4 mussels (*Mytilus* spp.) from locations in Normandy and northern Brittany (France)
5 experiencing high mussel mortalities, while it was not detected in Bay of St Brieuc (N
6 Brittany), which was not affected by abnormal mussel mortality. The distribution of
7 the bacteria in mussels seems to be restricted within inflammatory granulomas as
8 observed in Yesso scallops *Mizuhopecten yessoensis* from Canada and Japan.
9 *Francisella halioticida* has been identified as being involved in mass (>80%) mortality
10 of abalones *Haliotis gigantea* in Japan and high (up to 40%) mortality of Yesso scal-
11 lops *M. yessoensis* in Canada as well as in lesions reducing marketability of Yesso
12 scallops in Japan. The impact of this bacterium on the health of mussels needs to be
13 investigated in future research, especially since the cause of high mussel mortalities
14 that have been occurring in France for the past few years is still undetermined.

15 **KEY WORDS:**

16 *Francisella halioticida*, mussels, granulocytomas, polymerase chain reaction, *in situ*
17 hybridisation

1. INTRODUCTION

The gram-negative bacteria *Francisella haliotica* was originally described in farmed abalone (*Haliotis gigantea*) as the cause of mass mortality that occurred in Japan (Kamaishi et al. 2010, Brevik et al. 2011). Subsequently, this bacterium was identified in diseased Yesso scallops, *Mizuhopecten* (= *Patinopecten*) *yessoensis*, in conjunction with mortality events in Canada (Meyer et al. 2017) and Japan (Kawahara et al. 2018). In addition, Kawahara et al. (2019) demonstrated that this bacterium can be highly virulent in Yesso scallops. Thus, *F. haliotica* has shown pathogenic potential in both gastropod and bivalve species of molluscs.

Since 2014, unexplained mass mortalities of mussels (*Mytilus edulis*, *Mytilus galloprovincialis* and their hybrids) have occurred in France from the Atlantic to the English Channel coasts (Béchemin et al. 2014, 2015, Allain & Bernard 2016, Travers et al. 2016, Bernard & Allain 2017), with mortality rates ranging from 40 % to 70 % (Bernard & Allain 2017, Bernard et al. 2018, Charles et al. 2020). Several hypotheses have been proposed to explain these mortalities, including the involvement of pathogenic bacteria belonging to the *Vibrio splendidus* clade (Béchemin et al. 2014, 2015, François et al. 2015, Soletchnik & Robert 2016, Travers et al. 2016), possibility of genomic abnormalities and disseminated neoplasia (Benabdelmouna & Ledu 2016, Benabdelmouna et al. 2018,). Nevertheless, none of these proposals alone could explain such high mortality rates (Charles 2019). High mortality rates were recently observed among mussels in Northern Brittany and although *V. splendidus* strains and disseminated neoplasia were detected, the prevalence in mortality-affected versus non-affected areas did not correlate with mortality and no causative agent was identified (Bernard et al. 2018, Charles et al. 2020).

In a multi-parametric field study to identify potential causes of high mortality of mussels, Charles et al. (2020) reported the occurrence of heavy haemocyte infiltra-

tion and formation of granulomatous lesions in the connective tissue of various organs, however were not able to identify an aetiological agent. Similarities between the histological observations in mussels and the histopathology reported in Yesso scallops infected with *F. halioticida* (Meyer et al. 2017, Kawahara et al. 2018) is what prompted this investigation to test for the presence of *F. halioticida* in French mussel samples from previous studies.

To the best of our knowledge, *F. halioticida* has never been reported in mussels. The current study reveals the presence of *F. halioticida*, identified by polymerase chain reaction (PCR) and by *in situ* hybridisation (ISH), in cultured French mussels experiencing mortalities.

2. MATERIALS & METHODS

The 341 mussels analysed for detection of *Francisella halioticida* in the current study came from two overlapping multi-parametric studies in France: the first was conducted in 2017 with mussels taken from three sites in northern Brittany (Charles et al. 2020), two sites (Brest and Lannion) were affected by high mussel mortality while the third site (St Brieuc) did not experience mortalities. In this first study, mussel seed batches were periodically checked for mortality and sampled (live mussels randomly taken) for diagnosis. The second study was conducted in 2017 and 2018 by analysing mussels for obligatory reporting of increasing mortality (Council Directive 2006/88/EC) in Normandy and Hauts-de-France region (Charles 2019); in this second study, the sampling agents (from Administration and/or the Regional Shellfish Farming Committee) collected mostly moribund mussels and the information on mortality was obtained from Administration reports (Canier et al. 2018, Combette 2018). In the two previous studies, mussels were analysed using histological, bacteriological and molecular diagnostic procedures; and although a number of pathogens and

pathological conditions were detected none of them were believed to be the primary cause of high mussel mortality in northern France (Charles 2019, Charles et al. 2020).

The 17 mussel groups analysed in the current study are listed in Table 1 together with some data derived from the two previous studies, namely the sampling place, the corresponding farming method, the cumulative mortality at sampling and the condition of mussels in the samples (either alive or moribund). It is noteworthy that the mussels collected during the first previous study, designed in advance to monitor mortality, were alive (valves remained closed when out of seawater), while the mussels collected during the second previous study, based on sampling immediately after knowing mortality events, were primarily moribund specimens (valves were gaping and non-responsive) or a mixture of live and moribund. Two molecular diagnostic procedures were used targeting *F. halioticida* DNA, polymerase chain reaction (PCR) and *in situ* hybridisation (ISH). Information on how the mussels had been processed in the previous two studies was reported by Charles (2019) and Charles et al. (2020). Briefly, for histological analysis, an approximately 5 mm thick transverse section of mussel tissue containing mantle lobes, visceral mass (gut, digestive gland) and gills was excised, fixed in Davidson's solution for 48h, kept in 70% ethanol and then dehydrated through an ascending ethanol series and embedded in paraffin wax. From each mussel a pool of minced tissues from foot, adductor muscle, gills, mantle, and digestive gland were preserved in a microcentrifuge tube at -20 °C.

A total of 332 mussels were used for PCR assays, 20 mussels per group except for groups 11 and 15 in which 16 mussels were used. DNA was extracted from preserved tissues of individual sample mussels, using a QIAamp DNA minikit[®] (Qiagen, Courtaboeuf, France) following the manufacturer's protocol for blood or

body fluids (except elution was performed using 60 μ L Qiagen elution buffer AE). Following extraction, the DNA extracts of 5 mussels were pooled, except in groups 11 and 15 in which DNA pools derived from 4 mussels. Thus, 4 DNA pools were produced for each mussel group. PCR assays were conducted using the primer pair Fh-rpoB/F and Fh-rpoB/R (Brevik et al. 2011) designed to amplify the gene corresponding to bacterial DNA-directed RNA polymerase beta subunit (*rpoB*). For the conventional PCR, typical 25 μ L reaction mixtures contained 12.5 μ L of Premix Ex Taq[®] 2 X Takara[®] (Lonza, Verviers, Belgium), 0.5 μ L of each primer (20 μ M), 9.5 μ L of purified water and 2 μ L of DNA template (replaced with 2 μ L of purified water in the negative control). PCR amplifications were carried out in a T100[™] Thermal Cycler (Bio-Rad, France) using the following thermal cycling conditions: 10 s at 95 °C; 30 cycles for 10 s at 95 °C, 30 s at 55 °C, 40 s at 72 °C and a final extension of 3 min at 72 °C. All PCR products were purified and sequenced at LABÉO Frank Duncombe (Caen Cedex, France), and then analysed using the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST).

For ISH assays, only mussels from the first study could be retrospectively analysed; nine archived paraffin blocks of mussels showing different histological characteristics were selected: three healthy mussels (from groups 5, 6 and 7), two mussels with heavy haemocyte infiltration in the visceral mass (from groups 1 and 5), two mussels with granulomas (from group 5), one mussel with massive haemocyte diapedesis through stomach epithelium (from group 4) and one mussel with abundant degenerated digestive gland tubules filled with necrotic cells (from group 6). Samples analysed by ISH corresponded to mussels different from those analysed by PCR. The ISH assays were performed as described by Meyer et al. (2017) except that a 1 % Light Green counterstain was substituted for Bismark Brown to yield better

contrast.

ACCEPTED MANUSCRIPT

Table 1: Summary of mussel (*Mytilus* spp.) samples collected from France and test results for *Francisella haliotida* detection by conventional Polymerase Chain Reaction (PCR) and *in situ* hybridisation (ISH) assays. Each pool in the PCR assays resulted from pooling DNA from 5 mussels except in mussel groups 11 and 15 in which DNA from 4 mussels was pooled. Individual mussels assayed by PCR were different from those assayed by ISH. *Bouchot*: intertidal mussel stakes; Mix: mixture of alive and moribund mussels; *NA*: missing data; *NP*: not performed.

Group	Site	Region	Sampling date	Farming method	Cumulative mortality (%) at sampling time	Mussels condition	Number of pool of mussels analysed by PCR	PCR-results (positive pools/total pools)	ISH-results (positive mussels/total mussels analysed)
1	Lannion	Brittany	24/02/2017	Longlines	5 ^{(a) (b)}	Alive	4	0/4	0/1
2	Brest	Brittany	27/02/2017	<i>Bouchot</i>	12.5 ^{(a) (b)}	Alive	4	0/4	<i>NP</i>
3	St Brieuc	Brittany	27/02/2017	<i>Bouchot</i>	2.6 ^{(a) (b)}	Alive	4	0/4	<i>NP</i>
4	Brest	Brittany	24/04/2017	<i>Bouchot</i>	45 ^{(a) (b)}	Alive	4	0/4	0/1
5	Lannion	Brittany	17/05/2017	Longlines	20 ^{(a) (b)}	Alive	4	1/4	2/4
6	St Brieuc	Brittany	28/05/2017	<i>Bouchot</i>	4.5 ^{(a) (b)}	Alive	4	0/4	0/2
7	Brest	Brittany	30/05/2017	<i>Bouchot</i>	56.5 ^{(a) (b)}	Alive	4	2/4	0/1
8	Brest	Brittany	20/09/2017	<i>Bouchot</i>	66 ^{(a) (b)}	Alive	4	0/4	<i>NP</i>
9	Oye-Plage	Hauts-de-France	20/03/2017	<i>Bouchot</i>	25 ^(c)	Moribund	4	3/4	<i>NP</i>
10	Oye-Plage	Hauts-de-France	24/10/2017	<i>Bouchot</i>	10-50 ^(c)	Moribund	4	4/4	<i>NP</i>
11	Tardinghen	Hauts-de-France	14/03/2018	<i>Bouchot</i>	10-50 ^(d)	Moribund	4	3/4	<i>NP</i>
12	Donville-les-Bains	Normandy	19/06/2018	<i>Bouchot</i>	50 ^(d)	Mix	4	2/4	<i>NP</i>
13	Agon-Coutainville	Normandy	16/07/2018	<i>Bouchot</i>	<i>NA</i> ^(d)	Mix	4	2/4	<i>NP</i>
14	Chausey	Normandy	25/09/2018	<i>Bouchot</i>	10-50 ^(d)	Mix	4	0/4	<i>NP</i>
15	Wimereux	Hauts-de-France	15/10/2018	Wild	<i>NA</i> ^(d)	Moribund	4	4/4	<i>NP</i>
16	Tardinghen	Hauts-de-France	13/10/2018	<i>Bouchot</i>	<i>NA</i> ^(d)	Alive	4	2/4	<i>NP</i>
17	Dannes	Hauts-de-France	26/11/2018	<i>Bouchot</i>	25 ^(d)	Moribund	4	4/4	<i>NP</i>

^(a) Bernard *et al.* (2018); ^(b) Charles *et al.* (2020); ^(c) Canier *et al.* (2018); ^(d) Combette (2018)

3. RESULTS

The PCR results as well as the ISH results for detection of *Francisella halioticida* DNA in French *Mytilus* spp., are summarized in Table 1. BLAST results of all the PCR amplified fragments were a 99.9 % match to the reference *rpoB* sequences for *F. halioticida* (Brevik et al. 2011) (GenBank accession numbers JF290381 and JF290374). Regarding the first study conducted in northern Brittany during 2017 (groups 1 to 8), *F. halioticida* DNA was detected by PCR in mussels collected in May at Lannion and Brest which had mortality rates of 20 % and 56.5 % respectively, after 3 months of culture, but it was not detected in samples from St Brieuc, an area that was not affected by abnormal mortality (only 4.5 %). Concerning the groups collected following reports of abnormal mortality in Normandy and Hauts-de-France region in 2017 and 2018 (groups 9 to 17), PCR-positive results were obtained from every group except the one coming from Chausey's Island.

Although ISH assays were not conducted on all groups, positive reactions were observed within granuloma lesions from two specimens (Figure 1). These lesions were characterized by an intense accumulation of granulocytes surrounded by layers of flattened, epithelioid cells (Figure 1A & Figure 2A); and every tested mussel with granulomas showed positive reaction. Other types of pathology, such as heavy haemocyte infiltration and diapedesis, were negative via ISH; furthermore, mussels with no significant histopathology were also negative via ISH. Thus, infection with *F. halioticida* appears to be associated with the host's response and formation of granulomas (Figure 2), which is common in vertebrates infected with other *Francisella* species (Colquhoun & Duodu 2011).




Figure 1. *In situ* hybridisation (ISH) assay showing positive reaction (dark blue deposits). (A) Section through the digestive gland of a mussel showing a granuloma (*) with tiny dark blue deposits within them. (B) Higher magnification of the inside of a granuloma showing dark blue deposits (arrows) indicative of positive ISH reaction within cells.




Figure 2. Micrographs of a histological section consecutive to that of the Figure 1, stained with Harris' haematoxylin and eosin. (A) Large granuloma in the connective tissue of the digestive gland consisting of a mass of accumulated granulocytes (*) surrounded by layers of flattened, epithelioid cells (arrow). (B) Higher magnification of the inside of the granuloma showing numerous granulocytes with phagocytosed cells.

4. DISCUSSION

The occurrence of *Francisella halioticida* in French areas affected by high mussel mortality was revealed, whereas it was not detected in two areas, St Brieuc, which was not affected by abnormal mussel mortality, and Chausey's Island, where

mussels are known to suffer from intense predation by ducks, gulls and spider crabs, which were most likely contributing to the mortalities (CRC Normandie-Mer du Nord 2017, pers. com. 2019).. The findings from this study support a hypothesis proposed by Charles et al. (2020) concerning the possibility of missing an uncultivable bacterial pathogen with the classical medium used for bacterial culture. This could also be the case of the network for the monitoring of mollusc pathologies (REPAMO), coordinated by the French institute Ifremer, that monitors reports of mussel mortalities and conducts bacteriology using traditional culture media. Isolation and culture of *Francisella haliotica* requires a special culture medium (*Modified Eugon Agar*) and it can take several weeks for the bacteria to grow (Kamaishi et al. 2010, Kawahara et al. 2018). The limited sample size of this study prevents us from giving conclusive results; nevertheless, given that *F. haliotica* is known to be highly pathogenic in abalone and Yesso scallops, its detection in mussels from Normandy and northern Brittany areas affected by high mussel mortality highlights the need for further research concerning the association between mussel mortalities and infection with *F. haliotica*. Regarding the lack of geographical connection of the French mussel culture areas with the previous locations where *F. haliotica* was found, Canada and Japan, the importation of *M. yessoensis* in the beginning of the year 1987 in a hatchery of Brittany for several aquaculture trials (Buestel et al. 1988) could be considered as a potential hypothetical connection. It would be interesting to retrospectively test for the presence of *F. haliotica* DNA in samples of past mortalities, given that since 2015, several studies highlighted the presence of numerous inflammatory granulomas in French mussels affected by mortalities in the Atlantic coasts without any cause being identified (Robert et al. 2016, Travers et al. 2016, Pépin et al. 2017, 2018, Charles et al. 2020). In fact, some of these authors have even suggested a link

between the presence of the granuloma lesions and mortality. In addition, since 2014, mussels were not the only molluscs in France to be affected by unexplained mortalities: abalone (*Haliotis tuberculata*), cockles (*Cerastoderma edule*), clams (*Ruditapes philippinarum*) and scallops (*Pecten maximus*, *Chlamys varia*) were also affected (François et al. 2015, Lupo et al. 2016, Osta Amigo et al. 2018).

Considering the contagious and virulent nature of the *Francisella* spp. bacteria, and particularly *F. haliotida* in scallops, the detection of this bacterial species in French mussels opens up a new path to follow for further research concerning the cause of high mussel mortalities that have been occurring in northern France for the past few years. A field approach combined with laboratory challenges assessing the Koch's classic criteria would contribute to clarify the role of *F. haliotida* in mussel high mortality in N France.

5. ACKNOWLEDGEMENTS

The authors thank the Regional Shellfish Committee of North Brittany for supporting sampling in northern Brittany and the local mussel farmers for making this possible. The Regional Shellfish Committee of Normandy/ North Sea (CRC Normandie/Mer du Nord) and the Laboratory LABÉO financially supported this study. Maud Charles received cofunding from the Normandy region and from the laboratory LABÉO Frank Duncombe.

6. CONFLICT OF INTEREST

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

7. DATA AVAILABILITY STATEMENT

The authors confirm that one part of the data supporting the findings of this study is available within the article and the other part is available on request from the corresponding author.

8. LITERATURE CITED

Allain G, Bernard I (2016) Les mortalités de moules en 2014 et 2015 vues par les professionnels. Compte-rendu de la phase 1 : synthèse sur l'émergence, la propagation et l'installation des mortalités - Rapport technique. CRC Bretagne Nord

Béchemin C, Soletchnik P, Polsenaere P, Le Moine O, Pernet F, Protat M, Fuhrmann M, Quéré C, Goulitquer S, Corporeau C, Lapègue S, Travers M-A, Morga B, Garrigues M, Garcia C, Haffner P, Dubreuil C, Faury N, Baillon L, Baud J-P, Renaud T (2015) Épisodes de mortalité massive de moules bleues observés en 2014 dans les Pertuis charentais. Bull épidémiol santé anim aliment 67:6–9

Béchemin C, Soletchnik P, Polsenaere P, Le Moine O, Pernet F, Protat M, Fuhrmann M, Quéré C, Goulitquer S, Corporeau C, Renaud T, Lapègue S, Travers M-A, Morga B, Garrigues M, Garcia C, Haffner P, Dubreuil C, Faury N, Baillon L (2014) Surmortalités de la moule bleue *Mytilus edulis* dans les Pertuis Charentais (mars 2014) - Rapport d'expertise. Ifremer <http://archimer.ifremer.fr/doc/00229/34022/32387.pdf>

Benabdelmouna A, Ledu C (2016) The mass mortality of blue mussels (*Mytilus* spp.) from the Atlantic coast of France is associated with heavy genomic abnormalities as evidenced by flow cytometry. J Invertebr Pathol 138(0):30–38 <https://doi.org/10.1016/j.jip.2016.06.001>

Benabdelmouna A, Saunier A, Ledu C, Travers M-A, Morga B (2018) Genomic abnormalities affecting mussels (*Mytilus edulis-galloprovincialis*) in France are related to ongoing neoplastic processes, evidenced by dual flow cytometry and cell monolayer analyses. J Invertebr Pathol 157(0):45–52 <https://doi.org/10.1016/j.jip.2018.08.003>

Bernard I, Allain G (2017) Mortalités des moules en Bretagne nord : bilan des connaissances - Rapport technique. CRC Bretagne Nord

Bernard I, Charles M, Allain G, Burioli EAV, Villalba A, Le Foll F, Deydier Y, Houssin M (2018) Bilan de l'observatoire des mortalités de moules en Bretagne Nord pour la saison 2016-2017 et premiers éléments sur les organismes pathogènes présents - Rapport technique. CRC Bretagne Nord

Brevik ØJ, Ottem KF, Kamaishi, T, Watanabe K, Nylund A (2011) *Francisella halioticida* sp. nov., a pathogen of farmed giant abalone (*Haliotis gigantea*) in Japan. J Appl Microbiol 111(5): 1044–1056 <https://doi.org/10.1111/j.1365-2672.2011.05133.x>

Buestel D, Comps M, Moriceau J, Paquotte P (1988) Programme de recherche pour la mise au point d'une méthode d'élevage des coquilles Saint-Jacques en Méditerranée - Rapport d'avancement des travaux. Résultats obtenus en 1987. Ifremer

Canier L, Fleury E, Normand J, Garcia C, Geairon P (2018) REPAMO 2017 – Bulletins de la

surveillance - Janvier à Décembre 2017. Ifremer

Charles M. (2019). Étude des organismes pathogènes, des conditions physiologiques et pathologiques impliqués dans les mortalités anormales de moules (*Mytilus* sp.). PhD dissertation, University of Caen Normandy, Caen, France

Charles M, Bernard I, Villalba A, Oden E, Burioli EAV, Allain G, Trancart S, Bouchart V, Houssin M (2020) High mortality of mussels in northern Brittany – Evaluation of the involvement of pathogens, pathological conditions and pollutants. *J Invertebr Pathol* 170:107308 <https://doi.org/10.1016/j.jip.2019.107308>

Colquhoun DJ, Duodu S (2011) Francisella infections in farmed and wild aquatic organisms. *Vet Res* 42(1):47 <https://doi.org/10.1186/1297-9716-42-47>

Combette A (2018) REPAMO 2018 - Bulletins de la surveillance - Janvier à Décembre 2018. Ifremer

CRC Normandie-Mer du Nord (2017) Demande de dérogation pour la perturbation intentionnelle de goélands argentés sur l'archipel des îles Chausey http://www.normandie.developpement-durable.gouv.fr/IMG/pdf/dossier_dde_autorisation_chausey_2017.pdf

François C, Garcia C, Lupo C, Travers MA, Morga B, Tourbiez D, Fauray N, Haffner P, Serpin D, Dubreuil C, Chollet B, Baillon L, Lapègue S, Renault T (2015) Bilan 2014 du réseau Repamo - Réseau national de surveillance de la santé des mollusques marins. Ifremer <http://archimer.ifremer.fr/doc/00256/36691/>

Kamaishi T, Miwa S, Goto E, Matsuyama T, Oseko N (2010) Mass mortality of giant abalone *Haliotis gigantea* caused by a Francisella sp. bacterium. *Dis Aquat Org* 89:145–154 <https://doi.org/10.3354/dao02188>

Kawahara M, Meyer G, Lowe G, Kim E, Polinski M, Yoshinaga T, Itoh N (2019) Parallel studies confirm *Francisella halioticida* causes mortality in Yesso scallops *Patinopecten yessoensis*. *Dis Aquat Org* 135(2):127–134 <https://doi.org/10.3354/dao03383>

Kawahara M, Kanamori M, Meyer G, Yoshinaga T, Itoh N (2018) *Francisella halioticida*, Identified as the Most Probable Cause of Adductor Muscle Lesions in Yesso scallops *Patinopecten yessoensis* Cultured in Southern Hokkaido, Japan. *Fish Pathol* 53(2) :78–85 <https://doi.org/10.3147/jsfp.53.78>

Lupo C, Osta Amigo A, Fleury E, Robert S, Garcia C, Baillon L, Béchemin C, Canier L, Chollet B, Déchamps L, Dubreuil C, Fauray N, François C, Godfrin Y, Lapègue S, Morga B, Travers M-A, Tourbiez D, Masson J-C, Lamoureux, A (2016) Bilan 2015 du dispositif national de surveillance de la santé des mollusques marins. Ifremer

Meyer G, Lowe G, Gilmore S, Bower S (2017) Disease and mortality among Yesso scallops *Patinopecten yessoensis* putatively caused by infection with *Francisella halioticida*. *Dis Aquat Org* 125(1):79–84 <https://doi.org/10.3354/dao03130>

Osta Amigo A, Robert S, Fleury E, Lupo C, Garcia C, Geairon P, Canier L (2018) Repamo 2016 – bulletins de la surveillance janvier à juin et décembre 2016. Ifremer

Pépin J-F, Benabdelmouna A, Bierne N, Bouget J-F, Chabirand JM, Costes L, Dégremont L, Garcia C, Génauzeau S, Geairon P, Grizon J, Lamy J-B, Ledu C, Jolivet AL, Le Moine O,

Normand J, Palvadeau H, Polsenaere P, Robert S, Travers M-A (2018) Mortalités de moules bleues dans les secteurs mytilicoles : description et facteurs liés - MORBLEU. Ifremer

Pépin J-F, Benabdelmouna A, Degremont L, Guesdon S, Le Moine O, Morga B, Bierne N, Travers M-A, Robert S, Soletchnik P (2017) Mortalités de moules bleues dans les secteurs mytilicoles charentais et vendéens: description et facteurs liés – MORBLEU. Ifremer <http://archimer.ifremer.fr/doc/00391/50288/>

Robert S, Bouget J-F, Gabellec R, Louis W, Normand J, Blin J-L, Dagault F, Gerla D, Jolivet AL, Penot J, Chabirand J-M, Geairon P, Grizon J, Guesdon S, Le Moine O, Pépin J-F, Seugnet JL, Brunet A, Soletchnik P (2016) Réseau national d'observation de la moule bleue *Mytilus edulis*, Mytilobs, campagne 2015. Ifremer

Soletchnik P, Robert S (2016) Éléments de connaissance sur la mortalité et la reproduction de la moule bleue (*Mytilus edulis*) sur la façade atlantique. Ifremer <https://archimer.ifremer.fr/doc/00345/45634/45263.pdf>

Travers M-A, Pépin J-F, Soletchnik P, Guesdon S, Le Moine O (2016) Mortalités de moules bleues dans les Pertuis Charentais – MORBLEU. Ifremer <http://archimer.ifremer.fr/doc/00324/43539/>

ACCEPTED MANUSCRIPT