



HAL
open science

The Gut Microbiota at the Service of Immunometabolism

Chloé Michaudel, Harry Sokol

► **To cite this version:**

Chloé Michaudel, Harry Sokol. The Gut Microbiota at the Service of Immunometabolism. *Cell Metabolism*, 2020, 32 (4), pp.514-523. 10.1016/j.cmet.2020.09.004 . hal-02997641

HAL Id: hal-02997641

<https://hal.sorbonne-universite.fr/hal-02997641>

Submitted on 10 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18

Review

The gut microbiota at the service of immunometabolism

Chloé Michaudel^{1,2} & Harry Sokol^{1,2,3,*}

¹ INRA, UMR1319 Micalis & AgroParisTech, Jouy en Josas, France

² Paris Center for Microbiome Medicine (PaCeMM) FHU, Paris, France

³ Sorbonne Université, INSERM, Centre de Recherche Saint-Antoine, CRSA, AP-HP, Saint Antoine Hospital, Gastroenterology department, F-75012 Paris, France

*Correspondence:

Pr. Harry Sokol

Gastroenterology Department

Hôpital Saint-Antoine

184 rue du Faubourg Saint-Antoine, 75571 Paris CEDEX 12, France

Tel.: +33 1 49 28 31 71 Fax: +33 1 49 28 31 88

harry.sokol@aphp.fr

19 Summary

20 The gut microbiota is implicated in immune system functions. Regulation of the metabolic
21 processes occurring in immune cells is crucial for the maintenance of homeostasis and
22 immunopathogenesis. Emerging data demonstrate that the gut microbiota is an actor in
23 immunometabolism, notably through the effect of metabolites such as short-chain fatty acids,
24 bile acids, and tryptophan metabolites. In this review, we discuss the impact of the gut
25 microbiota on the intracellular metabolism of the different subtypes of immune cells, including
26 intestinal epithelial cells. Besides the effects on health, we discuss the potential consequences
27 in infection context and inflammatory bowel diseases.

28

29 *Keywords: Immunometabolism, Metabolism, Microbiota, Immunity*

30 Introduction

31 Metabolism involves cellular mechanisms to sustain life during physiological or pathological
32 processes. More generally, it is about energy; the utilization of metabolic substrates, notably
33 glucose, fatty acids (FAs), and amino acids (AA); and the balance between catabolism and
34 anabolism that maintain cellular homeostasis. Metabolism is impacted by lifestyles and dietary
35 habits, as illustrated by the increased rate of infection in malnourished populations (Blanton et
36 al., 2016; Hashimoto et al., 2012) and the metabolic syndrome-related disease outbreak in
37 overfed populations living in developed countries (Khare et al., 2020). In 2002,
38 immunometabolism, a new branch of metabolism, was brought to light with the discovery of
39 the link between CD28 activation and glycolysis in T cells (Frauwirth et al., 2002, p. 2). This
40 field notably aims to understand the impact of immune cells on metabolism and, conversely,
41 the metabolic needs of immune cells during homeostasis and pathological settings.

42 The microbiome is a major contributor to health, contributing to several development
43 processes, homeostatic states and responses to pathogenic situations. Although the human

44 microbiome is composed of several microbiotas colonizing different niches (e.g., lung, skin,
45 mouth, and vagina), the most studied is that in the gastrointestinal tract. It is composed of
46 diverse microbial communities, approximately 100 trillion microorganisms (Sarin et al., 2019;
47 Sender et al., 2016) and 150,000 microbial genomes (Pasolli et al., 2019). The gut microbiome
48 is composed of bacteria, fungi, viruses, and protists (Iliev and Leonardi, 2017; Richard and
49 Sokol, 2019; Shkoporov and Hill, 2019), and following millions of years of concomitant
50 evolution, it is in symbiosis with its host. The gut microbiome plays a role in the modulation of
51 both metabolism and immunity. Indeed, microbiome-derived molecules, either produced or
52 transformed by microorganisms, are major actors in the dialogue with immune cells (Bäckhed
53 et al., 2004; Cavallari et al., 2020; Lavelle and Sokol, 2020). Given the key role of the gut
54 microbiome in physiological processes, any alteration in its composition or function could
55 induce or participate in a disease (Pigneur and Sokol, 2016). The global role of the gut
56 microbiota in immunity has been extensively reviewed (Honda and Littman, 2016; Rooks and
57 Garrett, 2016). Here, we specifically discuss the effects of the gut microbiota on
58 immunometabolism, and more precisely, on the intracellular metabolism of immune cells, in
59 health and the potential consequences in diseases.

60 1- Immunometabolism: energy architecture to promote immunity

61 Immune system development/activation typically involves changes in the expression of large
62 numbers of genes and results in the acquisition of new functions, such as high production of
63 cytokines, lipid mediators, tissue-remodeling enzymes, ~~and toxic gases~~, and the ability to
64 migrate through tissues and/or undergo cellular division. Immune cells use the same pathways
65 as other cell types to generate energy and ensure their effective functioning. The main
66 metabolic pathways involved in immunometabolism are glycolysis, the TCA (tricarboxylic acid)
67 cycle, the pentose phosphate pathway (PPP), FA oxidation, FA synthesis, and AA metabolism.
68 Among the microbiome metabolism pathways impacting the metabolism of the immune cells,
69 we will notably discuss short-chain fatty acid (SCFA) production, tryptophan metabolism, lipid
70 metabolism and bile acid transformation. We present here the main actors we will discuss and

71 refer the reader to recent extensive reviews on this topic for more details (Bantug et al., 2018;
72 Goodpaster and Sparks, 2017; O'Neill et al., 2016).

73 **Glycolysis** is a relatively inefficient way to generate energy, as the breakdown of one unit of
74 glucose produces only 2 ATP molecules (Lunt and Vander Heiden, 2011). However, it is a
75 source of intermediate molecules for other pathways, including the PPP, AA and FA
76 metabolism pathways, and it can be swiftly activated, which is particularly relevant for
77 proliferating cells such as T cells.

78 The **TCA cycle** (or Krebs or citric acid cycle), which takes place in mitochondria in eukaryotes,
79 is a crucial engine in energy generation. Its primary substrate is acetyl-CoA produced either
80 from pyruvate by oxidative decarboxylation at the end of glycolysis or from FA oxidation. It is
81 estimated that the TCA cycle produces approximately 30 molecules of ATP from one molecule
82 of glucose, including the consumption of the NADH and FADH₂ molecules produced by
83 **oxidative phosphorylation** (OXPHOS) in mitochondria.

84 **FA synthesis** is required for the biosynthesis of the cell membrane, energy storage, and the
85 generation of signaling molecules. This pathway is tightly dependent on mTOR (mammalian
86 target of rapamycin) signaling and principally uses acetyl-CoA and other molecules provided
87 by glycolysis, the TCA cycle and the PPP. **Beta-oxidation** is the main metabolic pathway for
88 FA degradation. It leads to the production of acetyl-CoA, NADH and FADH₂ and then to a high
89 amount of energy through the TCA cycle and OXPHOS. **Cholesterol** is an essential precursor
90 of several biomolecules, including steroid hormones, vitamin D, oxysterols, and bile acids. **Bile**
91 **acids** (BAs) are produced through the oxidation of cholesterol. These molecules are a good
92 example of co-metabolism, as they are synthesized as primary and conjugated BAs by the
93 liver (Fiorucci et al., 2018); they reach the intestine through the bile duct and are converted by
94 gut microbiota enzymes into unconjugated secondary BAs. Most of the BAs are reabsorbed in
95 the terminal ileum and go back to the liver, completing their entero-hepatic cycle. Beyond their
96 role in lipid digestion, BAs are signaling molecules impacting many immune cell types through

97 several membranes and nuclear receptors, such as G protein-coupled bile acid receptor 5
98 (TGR5), farnesoid X receptor (FXR) and vitamin D receptor (VDR).

99 Besides their building blocks role for proteins, some **AAs** are also precursors of bioactive
100 molecules that contribute to the maintenance of signaling pathways and metabolism (Liu et al.,
101 2020). Glutamine and aspartate are involved in nucleotide synthesis (Cory and Cory, 2006;
102 Gots, 1971). Glutamine can also feed the TCA cycle to produce energy or be a substrate for
103 FA synthesis. Metabolites of other AAs, such as arginine and tryptophan, are involved in cell
104 proliferation and growth processes (Badawy, 2019; Milner, 1985). For example, tryptophan
105 can be metabolized into a myriad of active molecules through three major pathways: the
106 kynurenine pathway, the serotonin pathway, and the indole pathway. While the first two
107 pathways occur in mammalian cells, the last pathway takes place in the gut microbiota and
108 leads to the production of aryl hydrocarbon receptor (AhR) agonists that exhibit
109 immunomodulatory effects (Agus et al., 2018). The production of serotonin from
110 enterochromaffin cells in the gut is under the influence of the microbiome. It is well established
111 as a direct immunomodulatory factor, with seven receptor isoforms expressed on immune and
112 non-immune cell types (Shajib and Khan, 2015).

113 2- Key roles of the microbiota in immune cells metabolism

114 Several recent studies highlighted newly discovered mechanisms by which the gut microbiota
115 manipulates immunometabolism pathways in specific immune cell types (Figure 1).

116 **Epithelial cells**

117 The gastrointestinal epithelium is a highly relevant actor in host-microbiome interactions; it is
118 one of the first players in the immune response, and intestinal epithelial cells (IECs) are now
119 considered immune cells (Allaire et al., 2018). The energy metabolism of IECs, particularly in
120 the colon, is largely dependent on the gut microbiota. Early in life, before adaptive immune
121 system maturation, unidentified microbiota-derived molecules activate intraepithelial
122 lymphocytes (IELs) and ILC3 through STAT3 phosphorylation in an IL-23 and IL-22 dependent

123 manner. In the absence of adaptive immunity, the IL-23-ILC3-IL-22-IEC circuit allows to control
124 the gut microbiota, but the overactivated IL-22 production leads to an abnormal lipid
125 metabolism with reduced expression of key lipid transporters (e.g. CD36, Fabp1/2), and
126 reduction of triglycerides and free FA in serum (Mao et al., 2018). In germ-free mice,
127 colonocytes exhibit an energy-deprived state with decreased activity of enzymes of the TCA
128 cycle, β oxidation, and pyruvate dehydrogenase complex (Donohoe et al., 2011). Autophagy
129 is induced by the energetic stress to maintain homeostasis in colonocytes. The SCFA butyrate
130 produced by the gut microbiome in the colon is indeed the only source of carbon for
131 colonocytes. After being transformed into butyryl-CoA, it diffuses passively into the
132 mitochondria, undergoes β -oxidation, and feeds the TCA cycle and OXPHOS to produce
133 energy and dampen autophagy activation (Donohoe et al., 2011). IECs are massively exposed
134 to gut microbes and produce mucus and antimicrobial peptides to maintain a safety distance.
135 Butyrate also promotes intestinal homeostasis by downregulating IDO1 expression and the
136 kynurenine pathway in human IECs (Martin-Gallausiaux et al., 2018). The mechanisms involve
137 a reduction in signal transducer and activator of transcription (STAT) 1 expression and HDAC
138 (histone-deacetylase) inhibition.

139 Among the different IEC types, enterochromaffin (EC) cells are responsible for the production
140 of serotonin (5-HT), which has major effects on immune cells (see below). Serotonin production
141 in the colon is largely modulated by the gut microbiota and particularly spore-forming bacteria
142 metabolites. The mechanisms are not fully elucidated, but it has been shown that upregulation
143 of Tph1 expression, the rate-limiting enzyme in serotonin production, can be achieved by
144 SCFAs (butyrate and propionate) and some secondary bile acids, such as deoxycholate
145 produced by microbial biotransformation of cholate (Yano et al., 2015). Even if further
146 investigations are needed, this data suggest that modulating the gut microbiota composition or
147 directly administrating microbial metabolites could allow manipulating the production of
148 serotonin from a therapeutic perspective.

149

150 **Macrophages**

151 Macrophages are in the first line during the immune response but also sense and respond to
152 the microbiota to control it without initiating a detrimental inflammatory response. During the
153 pathogenic response, the metabolic profile of activated macrophages varies as a function of
154 the situation. In pro-inflammatory M1 macrophages, the TCA cycle is disrupted, leading to the
155 accumulation of itaconate and succinate and a shift to glycolysis (Rodríguez-Prados et al.,
156 2010; Tannahill et al., 2013). Itaconate is a major actor in immunometabolism that exhibits
157 immunomodulatory and antimicrobial effects. It is also involved in the accumulation of
158 succinate, as it directly inhibits its oxidation by blocking the activity of succinate dehydrogenase
159 (SDH) (Lampropoulou et al., 2016). Succinate exhibits a pro-inflammatory effect through its
160 oxidation that generates mitochondrial ROS (Reactive Oxygen Species) and leads to IL-1 β
161 production (Mills et al., 2016). Conversely, M2 macrophages have an intact TCA cycle and rely
162 mostly on OXPHOS (Huang et al., 2014; Vats et al., 2006). The gut microbiota modulates
163 these processes, notably through SCFAs. Butyrate, but not acetate or propionate, reprograms
164 macrophage metabolism toward OXPHOS and lipid metabolism leading to an anti-
165 inflammatory M2 phenotype (Scott et al., 2018). The detailed mechanisms are not identified
166 but involve the up-regulation of genes involved in OXPHOS (such as mitochondrial ATP
167 synthase and NADH dehydrogenase) and lipid metabolism (such as lipoprotein lipase)
168 pathways. As an illustration, the impaired production of butyrate induced by antibiotics
169 promotes the pro-inflammatory polarization of the intestinal macrophages, leading to a global
170 dysfunction of the immune response (Scott et al., 2018). This might play a role in the
171 association between antibiotics intake and the emergence of inflammatory and metabolic
172 diseases (Cox et al., 2014; Hviid et al., 2011).

173 **Innate lymphoid cells (ILCs)**

174 There are different types of innate lymphoid cells (ILCs) characterized by the expression of
175 specific membrane markers, transcription factors, and cytokine signatures. During their
176 activation, ILCs change their energy metabolism profoundly to fit their new functions (Rolot and

177 O'Sullivan, 2020). Transcriptomic analysis suggests that ILC1s use mTOR signaling, ILC2s
178 depend on sphingolipid and amino acid metabolism, and ILC3s rely on glycolysis (Gury-BenAri
179 et al., 2016). The gut microbiota profoundly impacts ILCs function as demonstrated by the
180 dramatic effects of antibiotics on the transcriptomic program of ILC1s, ILC2s and ILC3s (Gury-
181 BenAri et al., 2016). ILC3 is the main type of ILC present in the gastrointestinal tract. These
182 cells express ROR γ t, can produce IL-17 and IL-22, and are crucial regulators of inflammation,
183 infection, microbiota composition, and metabolism (Klose and Artis, 2016). ILC3 functions,
184 such as maintenance of the intestinal epithelium defense, depend on circadian signals
185 mediated by the circadian regulator ARNTL (Aryl Hydrocarbon Receptor Nuclear Translocator
186 Like). Light-dark cycles are key factors in this process, but the gut microbiota, which is known
187 to be an actor in diurnal rhythmicity (Thaiss et al., 2016), also had some impact (Godinho-Silva
188 et al., 2019). This signaling circuit connecting the gut microbiota, ILC3 and the intestinal
189 epithelial clock is also involved in the regulation of the local and systemic lipid metabolism
190 (Wang et al., 2017).

191 Gut microbiota-derived butyrate modulates ILC2 functions, inhibiting their uncontrolled
192 activation and, consequently, their negative role in lung inflammation and asthma. The
193 mechanism is not determined. Yet, the involvement of intracellular metabolism is supported by
194 the induction of changes in mROS production and glycolysis by butyrate (Lewis et al., 2019).
195 Moreover, the preferential use FAs over glucose by ILC2 to maintain their function in infection
196 or nutritional stress suggest that butyrate might directly fuel the TCA (Wilhelm et al., 2016).
197 Succinate, produced in the gut by protists and specific bacteria, stimulates the secretion of IL-
198 13 by ILC2, through an indirect action on Tuft cells and IL-25 (Schneider et al., 2018). The role
199 of succinate of other origin and its direct impact on ILC2 remains to be explored.

200

201 **T cells**

202 T cell metabolic plasticity is necessary to fit the permanently dynamic immune environment.
203 The gut microbiota actively participates in this programming via ROS, SCFA, and BA
204 production and REDOX signaling modification (Skelly et al., 2019). Effector and memory T
205 cells have very different functions and needs and thus exhibit different metabolism. It is
206 dominated by aerobic glycolysis in effector T cells and by FA oxidation and OXPHOS in
207 memory T cells. Mitochondrial dynamics are evidence of these differences, with fused
208 mitochondrial networks in memory T cells and punctate mitochondria in effector T cells (Buck
209 et al., 2016). In addition, mitochondria are a critical component of T cell activation, mainly
210 through ROS production (Sena et al., 2013). T cells stimulation via CD3 induces calcium influx
211 that stimulates the function of pyruvate dehydrogenase and TCA enzymes. TCA cycling
212 activates the mitochondrial electron transport chain and leads to the production of ROS, which
213 are required for T cell activation. ROS act in synergy with calcium influx to elicit IL-2 expression,
214 likely in an NF- κ B and AP-1 dependent manner (Kaminski et al., 2010).

215 Microbiota-derived SCFAs boost CD8⁺ T cell effector functions by modifying their cellular
216 metabolism (Trompette et al., 2018). SCFAs produced by the metabolism of dietary fibers by
217 the gut microbiota stimulate OXPHOS and mitochondrial mass in CD8⁺ T cells as well as their
218 glycolytic capacity. The mechanisms are not yet fully understood, but a part of these changes
219 depend on GPR41 activation. Besides, SCFAs can diffuse into the cytoplasm and serve as a
220 substrate for FAO, leading to the production of Acetyl-CoA that fuel TCA and then OXPHOS.
221 In activated CD8⁺ T cell, SCFAs, particularly butyrate, boosts the uptake and oxidation of FA,
222 leading to a disconnection of the TCA cycle from glycolytic input and favoring OXPHOS
223 through FA catabolism and glutamine utilization. This butyrate-induced cellular metabolism
224 adaptation is required for the differentiation to memory T cells (Bachem et al., 2019).

225 In stress situations, a massive amount of acetate is released into the extracellular space via
226 hydrolysis from acetyl-CoA. Acetate uptake by memory CD8⁺ T cells expands the acetyl-CoA
227 pool through TCA cycle and ATP citrate lyase activity and triggers the acetylation of GAPDH
228 (Glyceraldehyde 3-phosphate dehydrogenase), a key enzyme in glycolysis. The prompt
229 stimulation of glycolysis allows the rapid recall capacity of CD8⁺ memory T cells (Balmer et al.,

230 2016). Although these phenomena were described with host cell-derived acetate, they are
231 likely triggered, at least in the gut, by the massive amount of acetate produced by the gut
232 microbiota.

233 SCFAs also exhibit significant effects on CD4+ T cells, notably regarding the generation of T
234 helper (Th) 17, Th1 (Park et al., 2015), and regulatory T cells (Furusawa et al., 2013; Smith et
235 al., 2013). The mechanisms involve the inhibition of HDACs and regulation of the mTOR
236 pathway (a master regulator of cell growth and metabolism). This link has been recently shown
237 with pentanoate (also known as valerate), a subdominant microbiota-produced SCFA that can
238 stimulate the production of the anti-inflammatory cytokine IL-10 by providing additional acetyl-
239 CoA for histone acetyltransferases and enhancing glycolysis and mTOR activity (Luu et al.,
240 2019). Two mechanisms have been suggested regarding the activation of mTOR by SCFA
241 (Figure 2). Through their action on energy production pathways, SCFAs induce the production
242 of ATP and the depletion of AMP, which are inhibitor and activator of AMP-activated protein
243 kinase (AMPK), respectively. Consequently, the inhibitor activity of AMPK on mTOR is
244 repressed, thus leading to mTOR activation (Kim et al., 2016; Luu et al., 2019; Zhou et al.,
245 2018). The second potential mechanism involves the HDAC inhibition activity of SCFAs.
246 SCFAs, in association with P300/CBP (E1A binding protein p300/ CREB-binding protein),
247 promote acetylation of the Ribosomal protein S6 kinase beta-1 (S6K1), which is a downstream
248 target of mTOR, leading to more robust activation of the pathway (Park et al., 2015). Another
249 layer of complexity has been indicated recently by showing that the effects of SCFAs on T cell
250 metabolism are dependent on the inflammatory context (Trapecar et al., 2020).

251 BAs also have an essential impact on T cells. A derivative of lithocholic acid (LCA), 3-oxoLCA,
252 inhibits the differentiation of Th17 cells by directly interacting with the transcription factor
253 ROR γ t (Hang et al., 2019). Conversely, another derivative of LCA, isoalloLCA, promotes the
254 differentiation of Treg cells. The mechanism involves the stimulation of OXPHOS and the
255 production of mitochondrial ROS, which leads to the increased expression of FOXP3 by
256 increasing the levels of histone (H3K27) acetylation in the Foxp3 promoter (Hang et al., 2019).

257 In the colon specifically, BAs act through the bile acid receptor Breg to regulate the function of
258 ROR γ ⁺ Treg cells, which are significant players in the maintenance of colonic homeostasis
259 (Song et al., 2020).

260 **B cells**

261 B cell differentiation into plasma cells and the production of antibodies require a massive
262 amount of energy and a global change in cellular metabolism. Gut microbiota-derived SCFAs
263 contribute to fuel the cellular energy engine at different levels for these processes and to boost
264 antibody production. SCFAs are converted into acetyl-CoA that is integrated into the
265 mitochondrial TCA cycle leading to the production of ATP. SCFAs also stimulate glycolysis in
266 B cell via mTOR activation. SCFAs-derived acetyl-CoA is also a substrate in FA (particularly
267 palmitic acid) synthesis, which is crucial for plasma cell differentiation and stimulates antibody
268 production (Kim et al., 2016). Using an elegant strategy based on genetically engineered
269 *Clostridium sporogenes* in germ-free mice, it has recently been shown that branched SCFAs,
270 such as isobutyrate or isovalerate, can also modulate B cells functions. The absence of
271 branched SCFAs production in manipulated mice led to an increased frequency of IgA⁺ plasma
272 cells in the small intestine and increased levels of IgA bound to the surface of innate immune
273 cells such as neutrophils, macrophages and dendritic cells (Guo et al., 2019). The mechanisms
274 underlying these effects are not yet known.

275 B cells have a critical role in tolerance toward the gut microbiota through the production of
276 immunoglobulins and the action of IL-10-producing Bregs (regulatory B cells). In Bregs, Rosser
277 and colleagues recently showed that butyrate could divert tryptophan metabolism toward the
278 serotonin pathway and the production of 5-hydroxyindole-3-acetic acid (5-HIAA) (Rosser et al.,
279 2020). Surprisingly, 5-HIAA was shown to activate AhR in these cells, mediating the
280 suppressive effect of butyrate supplementation in a rheumatoid arthritis model *in vivo*.

281 3- Consequences for disease pathogenesis

282 Immunometabolism at steady-state promotes homeostasis. However, the energy requirement
283 of immune cells during inflammatory and infectious diseases is much higher, and their whole
284 metabolism is altered. These processes are involved in both the pathogenesis of nonseptic
285 inflammatory disorders and in the resolution of infection (Zmora et al., 2017). As seen above,
286 the gut microbiota modulates immunometabolism and thus can have positive or negative
287 effects on these pathological events (Figure 3).

288 **Infections**

289 Innate immune cells are the first bulwark against bacterial infection. TCR $\gamma\delta$ (T cell receptor)
290 IELs are key players in the initial response to intestinal pathogens. Their location within the
291 intestinal epithelium and their motility, which are dependent on the gut microbiota, allow
292 effective surveillance of the mucosal surface (Hoytema van Konijnenburg et al., 2017). Upon
293 infection with *Salmonella*, the change in $\gamma\delta$ IEL behavior is associated with the activation of
294 OXPHOS and anaerobic glycolysis. These metabolic changes are dependent on mTOR and
295 microbial cues in IECs. These data highlight a complex 3-partner system in which the gut
296 microbiota, through action on IECs, induces the metabolic reprogramming of $\gamma\delta$ IELs to boost
297 their mucosal surveillance capacity (Hoytema van Konijnenburg et al., 2017). Metabolic
298 changes are also observed in IECs in response to infection. In the early steps of infection with
299 the mouse pathogen *Citrobacter rodentium*, downregulation of the TCA cycle and OXPHOS is
300 observed in parallel with perturbations of cholesterol homeostasis. Cholesterol synthesis and
301 import are activated simultaneously with cholesterol efflux suggesting either an atypical
302 cholesterol metabolism regulation in IECs during stress or the manipulation of cholesterol
303 homeostasis by *C. rodentium* (Hopkins et al., 2019). Starting on the second week following
304 infection with *C. rodentium*, the Th17 cell response is activated and required to resolve the
305 infection. These pathogen-induced Th17 cells rely on anaerobic glycolysis and OXPHOS,
306 while commensal microbe-induced Th17 cells rely mostly on OXPHOS. These differences in
307 bioenergetic profiles are associated with different mitochondrial morphologies and a pro-
308 inflammatory phenotype in pathogen-induced Th17 cells (Omenetti et al., 2019). ILC3 are other

309 important actors in response to *C. rodentium*, notably through the production of IL-22 and IL-
310 17A, which occur in an mTOR dependant way. The activation of mTOR complex 1 (mTORC1)
311 leads to metabolic reprogramming of ILC3 characterized by enhanced glycolysis and mROS
312 production. Mechanistically, mTORC1 activates HIF1 α that supports ROR γ t and stimulates
313 glycolysis. The downstream produced mROS contribute to stabilize HIF1 α and to reprogram
314 ILC3 metabolism toward the response to bacterial pathogens (Di Luccia et al., 2019).

315 During the response to infection and sepsis, T cells generate NOX2 (NADPH oxidase 2)-
316 mediated ROS. Acetate can restore the oxidant-antioxidant imbalance in T cells during sepsis
317 independently of GPR43 (G-protein-coupled receptor) and likely through upregulation of HDAC
318 activity (Al-Harbi et al., 2018).

319 Mitochondrial FA metabolism in the intestinal epithelium is impaired in SIV-infected rhesus
320 macaques and HIV (human immunodeficiency viruses)-infected patients. The underlying
321 mechanisms involve altered PPAR α (peroxisome proliferator-activated receptor) signaling and
322 impaired FA β -oxidation of short- and medium-chain FAs, which correlate with an alteration in
323 the intestinal epithelial barrier. Interestingly, these phenomena are modulated by the gut
324 microbiome, as mitochondrial FA metabolism and intestinal barrier function can be rapidly
325 restored by the administration of the probiotic *Lactobacillus plantarum*, independent of any
326 effect on CD4+ T cells (Crakes et al., 2019).

327 **Inflammatory Bowel Disease**

328 The prominent role of the gut microbiota in the pathogenesis of inflammatory bowel disease
329 (IBD) has been demonstrated by both human and animal studies (Britton et al., 2019; Lavelle
330 and Sokol, 2020). The first actors in the interaction with the gut microbiota in IBD are epithelial
331 cells. Alterations in the metabolism and functions of IECs are involved in IBD and lead to an
332 impaired intestinal barrier and the translocation of microbial molecules, resulting in
333 overactivation of the gut immune system. Some studies are now linking the gut microbiota to
334 defective IEC metabolism in intestinal inflammation, notably through the Nod-like receptor
335 (NLR) family. NLRX1 (nucleotide-binding oligomerization domain, leucine-rich repeat

336 containing X1) is a mitochondria-associated NLR with potential anti-inflammatory effects in
337 colitis settings (Leber et al., 2018). NLRX1 is required to maintain balanced glutamine
338 metabolism and barrier functions in IECs. The mechanisms are not clearly demonstrated, but
339 it is suggested that NLRX1 may support the glutamine input into the TCA cycle through its
340 metabolism into glutamate and α -ketoglutarate. The impaired glutamine metabolism in IECs
341 leads to changes in AA availability for the gut microbiota, inducing changes in composition.
342 Interestingly, the altered gut microbiota exhibits a pro-inflammatory effect by itself, as
343 demonstrated by fecal microbiota transfer experiments (Leber et al., 2018). NLR-associated
344 inflammasomes are also involved. SCFAs induce the activation of NLRP3 (NOD-like receptor
345 family, pyrin domain containing 3) via their receptors GPR43 and GPR109a, inducing ion (K+
346 and Ca²⁺) efflux, and promoting epithelial repair in colitis setting through IL-18 maturation and
347 release (Macia et al., 2015). The impact of SCFAs on macrophage polarization is also relevant
348 in IBD. SCFAs depletion, for example, induced by antibiotics, favors an M1 hyperresponsive
349 phenotype leading to an overproduction of pro-inflammatory cytokines and to the promotion of
350 intestinal inflammation (Scott et al., 2018).

351 Previous studies have also shown a link between mitochondrial dysfunction and IBD. The
352 expression of prohibitin 1 (PHB1), an inner mitochondrial membrane component, is decreased
353 in colonic biopsies from IBD patients (Hsieh et al., 2006; Theiss et al., 2007). Moreover,
354 mitochondrial dysfunction in IECs and notably in Paneth cells can induce ileal inflammation in
355 mouse models (Jackson et al., 2020). Interestingly, Paneth cell abnormalities in patients with
356 Crohn's disease correlate with alterations in both microbiota composition and oxidative
357 phosphorylation in ileal tissue (Liu et al., 2016). Mechanistically, mitochondrial respiration
358 impairment forces IECs to acquire a dysfunctional Paneth cell phenotype, leading to metabolic
359 imbalance and inflammation (Khaloian et al., 2020). Moreover, mitochondrial impairment in
360 Crohn's patients also involves a decrease in H₂S detoxification, while the relative abundance
361 of H₂S-producing microbes is increased in the gut microbiota. The amount of *Atopobium*
362 *parvulum*, a keystone microbiota species for H₂S production, correlated with Crohn's disease

363 severity (Mottaweia et al., 2016). Overall, the net increase in H₂S due to increased microbiota
364 production and decreased mitochondrial detoxification is involved in intestinal inflammation
365 pathogenesis.

366 Conclusion

367 The effects of the gut microbiome on host immune cells is often examined with classical host-
368 microbes interaction concepts, relying on the recognition of conserved microbial motifs by
369 innate immunity sensors, or on the effect of microbial molecules on a host cell receptors.
370 Despite the crucial role of the cellular metabolism in the ability to mount an appropriate immune
371 response, the studies investigating how the gut microbiota directly affects it, remain scarce.
372 Yet, the gut microbiota has a special relationship with metabolism, notably via the mitochondria
373 due to their common origin. Mitochondria share a large part of their genome with bacteria, so
374 communication and regulation can be evoked between these entities, which are only separated
375 by the cell membrane (Lin and Wang, 2017). Host cell and gut microbiota are tightly connected
376 in an inter-kingdom metabolic network that allows the proper functioning of mammalian meta-
377 organisms. Each pathway is modulated by or depends on metabolites from others. It takes the
378 collapse of only one path to compromise the normal operation. These processes are even
379 more critical for immunometabolism, as immune cells need to react to stimuli rapidly and to
380 reprogram their metabolism to exercise their functions. Gut microbiota-derived metabolites are
381 genuinely represented in immunometabolism, with a particularly important role of SCFAs, BAs,
382 and AA metabolites. Deciphering all the ins and outs resulting from the action of the microbiota
383 on immunometabolism is highly challenging. Part of the complexity lies in the final effects of
384 the microbial products, which can be different depending on the context or the cell types. The
385 intrinsic diversity of the actors within the gut microbiota and the immune system brings an
386 additional level of difficulty in the exploration of these interactions.

387 The next step in the understanding of host-microbiota cross-talk is to decipher more precisely
388 the bidirectional impact of each metabolism on that of the partner in health and disease. This

389 effort is crucial to identify therapeutic targets that will be actionable through metabolic
390 modulation. These innovative treatments may take several forms. The modulation of the gut
391 microbiota to favor beneficial metabolite-producing bacteria is one possibility. However, an
392 even more attractive strategy is to precisely impact host-microbiota metabolism by accurately
393 supplementing a missing metabolite and/or inhibiting an overactivated pathway simultaneously
394 on both sides of the interkingdom cross-talk.

395 Acknowledgements

396 Harry Sokol received funding from the European Research Council (ERC) under the European
397 Union's Horizon 2020 Research and Innovation Programme (ERC-2016-StG-71577). We
398 thank Hugo Michaudel for the design of the figures.

399 Author Contributions

400 C.M. and H.S. wrote the paper.

401 Declaration of Interests

402 The authors declare no competing interests.

403

404 Figure legends

405 **Figure 1. Influence of the gut microbiota on immunometabolism**

406 In **epithelial cells**, after being transformed into butyrate-CoA, butyrate diffuses passively in
407 the mitochondria, undergoes β -oxidation, and feeds the TCA cycle and OXPHOS to produce
408 energy. Butyrate also repress IDO1 via HDAC inhibition. In enterochromaffin cells, the
409 metabolism of tryptophan into serotonin (5-HT) is stimulated by butyrate, propionate and bile
410 acids. The production of 5-HIAA from 5-HT is stimulated by butyrate, and 5-HIAA binds AhR
411 in Breg cells, inducing suppressive effects. SCFAs also increase glycolysis and mTOR activity
412 in **B cells**. SCFAs-derived acetyl-CoA are also substrate in FA synthesis and β -oxidation,
413 which is crucial for antibody production.. In **memory T cells**, butyrate activates β -oxidation,

414 while acetate-derived acetyl-CoA stimulates glycolysis through acetylation of GAPDH. In
415 **effector T cells**, the secondary bile acid isoalloLCA stimulates OXPHOS and the production
416 of mtROS, which leads to the upregulation of FOXP3 through histone acetylation in its
417 promoter region, resulting in Treg differentiation. Another secondary bile acid, 3-oxoLCA,
418 interacts directly with ROR γ t and inhibits the differentiation of Th17 cells. Pentanoate
419 stimulates glycolysis and mTOR activity and leads to the production of acetyl-CoA, which feeds
420 histone acetyltransferase activity and IL-10 production. SCFAs also boost CD8⁺ T cell effector
421 function via an increased glycolytic capacity, OXPHOS and mitochondrial mass. In
422 **macrophages**, butyrate promotes OXPHOS activation and the anti-inflammatory M2
423 phenotype. The impaired production of butyrate can be involved in the pro-inflammatory
424 polarization of the intestinal macrophages, leading to a global dysfunction of the immune
425 response. In M1 macrophages, the TCA cycle is broken, succinate and itaconate accumulate,
426 and glycolysis dominates energy production. The accumulation of succinate generates mtROS
427 and leads to IL-1 β production. Itaconate exhibits immunomodulatory and antimicrobial effects.
428 In **ILC3s**, the gut microbiota influences circadian gene expression, which favors STAT3
429 phosphorylation during development and modulates FA transport in IECs. Succinate, produced
430 in the gut by protists and specific bacteria, stimulates the secretion of IL-13 by **ILC2**, through
431 an indirect action on Tuft cells and IL-25.

432

433 **Figure 2. Mechanism of activation of the mTOR pathway by SCFA**

434 **A.** Through their action on energy production pathways, SCFAs induce the production of ATP
435 and the depletion of AMP, which are inhibitor and activator of AMP-activated protein kinase
436 (AMPK), respectively. Consequently, the inhibitor activity of AMPK on mTOR is repressed,
437 thus leading to mTOR activation. **B.** SCFAs, in association with P300/CBP, promote
438 acetylation of the Ribosomal protein S6 kinase beta-1 (S6K1), which is a downstream target
439 of mTOR, leading to more robust activation of the pathway with RPS6 (Ribosomal Protein S6)
440 phosphorylation and inhibition of 4EBP (Translation initiation factor 4E binding protein)
441 phosphorylation. ULK1: Unc-51 Like Autophagy Activating Kinase 1).

442

443 **Figure 3. Immunometabolism and the microbiota in diseases**

444 **Infections.** Upon infection with *Salmonella*, $\gamma\delta$ IEL behavior changes are associated with
445 activation of OXPHOS and anaerobic glycolysis and boost epithelial barrier protection. This
446 response is dependent on mTOR and microbiota signals. Metabolic changes are also
447 observed in IECs in response to *Citrobacter rodentium*, with downregulation of the TCA cycle
448 and OXPHOS. In parallel, IECs present a dysregulation of cholesterol homeostasis. During
449 this infection, pathogen-induced Th17 cells rely on glycolysis and OXPHOS, while commensal
450 microbe-induced Th17 cells rely mostly on OXPHOS. ILC3 are other important actors in
451 response to *C. rodentium*, notably through mTORC 1 activation (mTORC1) that leads to HIF1 α
452 activation that supports ROR γ t and stimulates glycolysis. In parallel, the downstream produced
453 mROS contribute to stabilizing HIF1 α . Metabolic reprogramming of ILC3 permits to produce
454 IL-22 and IL-17A to sustain immune response. These differences in bioenergetic profiles are
455 associated with different mitochondrial morphologies. During the response to infection and
456 sepsis, T cells generate NOX2-mediated ROS. Acetate can restore the oxidant-antioxidant
457 balance in T cells in this setting and likely through the upregulation of HDAC activity. During
458 pathogen infection, TLR engagement in macrophages induces the recruitment of TRAF6 to
459 mitochondria, leading to an increased production of ROS that is involved in response to
460 intracellular pathogens. In SIV- and HIV-infected individuals, the intestinal epithelium presents
461 altered PPAR α signaling and FA β -oxidation, which correlates with an alteration in the
462 intestinal epithelial barrier that can be restored by microbiota-derived factors.

463 **IBD.** The altered microbiota produces an insufficient amount of SCFAs, leading to defective
464 activation of NLRP3 and, subsequently, to inadequate production of IL-18 that normally
465 promotes epithelial repair. Defect in SCFAs promotes M1 polarization leading to the production
466 of pro-inflammatory cytokines promoting intestinal inflammation. An alteration in the
467 metabolism and functions of IECs, notably Paneth cells, is observed in IBD. It is linked to
468 altered microbiota signals leading to inhibition of OXPHOS. The accumulation of pro-

469 inflammatory H₂S is observed in IBD. It is connected to the increased H₂S production by the
470 abnormal microbiota associated with the impairment in mitochondrial detoxification. *Atopobium*
471 *parvulum* is a keystone microbiota species for the production of H₂S.

472

473 **References**

- 474 Agus, A., Planchais, J., Sokol, H., 2018. Gut Microbiota Regulation of Tryptophan
 475 Metabolism in Health and Disease. *Cell Host Microbe* 23, 716–724.
 476 <https://doi.org/10.1016/j.chom.2018.05.003>
- 477 Al-Harbi, N.O., Nadeem, A., Ahmad, S.F., Alotaibi, M.R., AlAsmari, A.F., Alanazi, W.A., Al-
 478 Harbi, M.M., El-Sherbeeney, A.M., Ibrahim, K.E., 2018. Short chain fatty acid, acetate
 479 ameliorates sepsis-induced acute kidney injury by inhibition of NADPH oxidase
 480 signaling in T cells. *Int. Immunopharmacol.* 58, 24–31.
 481 <https://doi.org/10.1016/j.intimp.2018.02.023>
- 482 Allaire, J.M., Crowley, S.M., Law, H.T., Chang, S.-Y., Ko, H.-J., Vallance, B.A., 2018. The
 483 Intestinal Epithelium: Central Coordinator of Mucosal Immunity. *Trends Immunol.* 39,
 484 677–696. <https://doi.org/10.1016/j.it.2018.04.002>
- 485 Bachem, A., Makhlof, C., Binger, K.J., de Souza, D.P., Tull, D., Hochheiser, K., Whitney,
 486 P.G., Fernandez-Ruiz, D., Dähling, S., Kastenmüller, W., Jönsson, J., Gressier, E.,
 487 Lew, A.M., Perdomo, C., Kupz, A., Figgett, W., Mackay, F., Oleshansky, M., Russ,
 488 B.E., Parish, I.A., Kallies, A., McConville, M.J., Turner, S.J., Gebhardt, T., Bedoui, S.,
 489 2019. Microbiota-Derived Short-Chain Fatty Acids Promote the Memory Potential of
 490 Antigen-Activated CD8+ T Cells. *Immunity* 51, 285-297.e5.
 491 <https://doi.org/10.1016/j.immuni.2019.06.002>
- 492 Bäckhed, F., Ding, H., Wang, T., Hooper, L.V., Koh, G.Y., Nagy, A., Semenkovich, C.F.,
 493 Gordon, J.I., 2004. The gut microbiota as an environmental factor that regulates fat
 494 storage. *Proc. Natl. Acad. Sci. U. S. A.* 101, 15718–15723.
 495 <https://doi.org/10.1073/pnas.0407076101>
- 496 Badawy, A.A.-B., 2019. Tryptophan Metabolism: A Versatile Area Providing Multiple Targets
 497 for Pharmacological Intervention. *Egypt. J. Basic Clin. Pharmacol.* 9.
 498 <https://doi.org/10.32527/2019/101415>
- 499 Balmer, M.L., Ma, E.H., Bantug, G.R., Grähler, J., Pfister, S., Glatter, T., Jauch, A., Dimeloe,
 500 S., Slack, E., Dehio, P., Krzyzaniak, M.A., King, C.G., Burgener, A.-V., Fischer, M.,
 501 Develioglu, L., Belle, R., Recher, M., Bonilla, W.V., Macpherson, A.J., Hapfelmeier,
 502 S., Jones, R.G., Hess, C., 2016. Memory CD8(+) T Cells Require Increased
 503 Concentrations of Acetate Induced by Stress for Optimal Function. *Immunity* 44,
 504 1312–1324. <https://doi.org/10.1016/j.immuni.2016.03.016>
- 505 Bantug, G.R., Galluzzi, L., Kroemer, G., Hess, C., 2018. The spectrum of T cell metabolism
 506 in health and disease. *Nat. Rev. Immunol.* 18, 19–34.
 507 <https://doi.org/10.1038/nri.2017.99>
- 508 Blanton, L.V., Barratt, M.J., Charbonneau, M.R., Ahmed, T., Gordon, J.I., 2016. Childhood
 509 undernutrition, the gut microbiota, and microbiota-directed therapeutics. *Science* 352,
 510 1533. <https://doi.org/10.1126/science.aad9359>
- 511 Britton, G.J., Contijoch, E.J., Mogno, I., Vennaro, O.H., Llewellyn, S.R., Ng, R., Li, Z.,
 512 Mortha, A., Merad, M., Das, A., Gevers, D., McGovern, D.P.B., Singh, N., Braun, J.,
 513 Jacobs, J.P., Clemente, J.C., Grinspan, A., Sands, B.E., Colombel, J.-F., Dubinsky,
 514 M.C., Faith, J.J., 2019. Microbiotas from Humans with Inflammatory Bowel Disease
 515 Alter the Balance of Gut Th17 and RORγt+ Regulatory T Cells and Exacerbate Colitis
 516 in Mice. *Immunity* 50, 212-224.e4. <https://doi.org/10.1016/j.immuni.2018.12.015>
- 517 Buck, M.D., O’Sullivan, D., Klein Geltink, R.I., Curtis, J.D., Chang, C.-H., Sanin, D.E., Qiu, J.,
 518 Kretz, O., Braas, D., van der Windt, G.J.W., Chen, Q., Huang, S.C.-C., O’Neill, C.M.,
 519 Edelson, B.T., Pearce, E.J., Sesaki, H., Huber, T.B., Rambold, A.S., Pearce, E.L.,
 520 2016. Mitochondrial Dynamics Controls T Cell Fate through Metabolic Programming.
 521 *Cell* 166, 63–76. <https://doi.org/10.1016/j.cell.2016.05.035>
- 522 Cavallari, J.F., Barra, N.G., Foley, K.P., Lee, A., Duggan, B.M., Henriksbo, B.D., Anhê, F.F.,
 523 Ashkar, A.A., Schertzer, J.D., 2020. Postbiotics for NOD2 require non-hematopoietic
 524 RIPK2 to improve blood glucose and metabolic inflammation in mice. *Am. J. Physiol.*
 525 *Endocrinol. Metab.* <https://doi.org/10.1152/ajpendo.00033.2020>

526 Cory, J.G., Cory, A.H., 2006. Critical Roles of Glutamine as Nitrogen Donors in Purine and
527 Pyrimidine Nucleotide Synthesis: Asparaginase Treatment in Childhood Acute
528 Lymphoblastic Leukemia. *In Vivo* 20, 587–589.

529 Cox, L.M., Yamanishi, S., Sohn, J., Alekseyenko, A.V., Leung, J.M., Cho, I., Kim, S.G., Li, H.,
530 Gao, Z., Mahana, D., Zárata Rodriguez, J.G., Rogers, A.B., Robine, N., Loke, P.,
531 Blaser, M.J., 2014. Altering the intestinal microbiota during a critical developmental
532 window has lasting metabolic consequences. *Cell* 158, 705–721.
533 <https://doi.org/10.1016/j.cell.2014.05.052>

534 Crakes, K.R., Santos Rocha, C., Grishina, I., Hirao, L.A., Napoli, E., Gaulke, C.A., Fenton,
535 A., Datta, S., Arredondo, J., Marco, M.L., Sankaran-Walters, S., Cortopassi, G.,
536 Giulivi, C., Dandekar, S., 2019. PPAR α -targeted mitochondrial bioenergetics mediate
537 repair of intestinal barriers at the host-microbe intersection during SIV infection. *Proc.*
538 *Natl. Acad. Sci. U. S. A.* 116, 24819–24829.
539 <https://doi.org/10.1073/pnas.1908977116>

540 Di Luccia, B., Gilfillan, S., Cella, M., Colonna, M., Huang, S.C.-C., 2019. ILC3s integrate
541 glycolysis and mitochondrial production of reactive oxygen species to fulfill activation
542 demands. *J. Exp. Med.* 216, 2231–2241. <https://doi.org/10.1084/jem.20180549>

543 Donohoe, D.R., Garge, N., Zhang, X., Sun, W., O’Connell, T.M., Bunger, M.K., Bultman, S.J.,
544 2011. The microbiome and butyrate regulate energy metabolism and autophagy in
545 the mammalian colon. *Cell Metab.* 13, 517–526.
546 <https://doi.org/10.1016/j.cmet.2011.02.018>

547 Fiorucci, S., Biagioli, M., Zampella, A., Distrutti, E., 2018. Bile Acids Activated Receptors
548 Regulate Innate Immunity. *Front. Immunol.* 9.
549 <https://doi.org/10.3389/fimmu.2018.01853>

550 Frauwirth, K.A., Riley, J.L., Harris, M.H., Parry, R.V., Rathmell, J.C., Plas, D.R., Elstrom,
551 R.L., June, C.H., Thompson, C.B., 2002. The CD28 signaling pathway regulates
552 glucose metabolism. *Immunity* 16, 769–777. [https://doi.org/10.1016/s1074-](https://doi.org/10.1016/s1074-7613(02)00323-0)
553 [7613\(02\)00323-0](https://doi.org/10.1016/s1074-7613(02)00323-0)

554 Furusawa, Y., Obata, Y., Fukuda, S., Endo, T.A., Nakato, G., Takahashi, D., Nakanishi, Y.,
555 Uetake, C., Kato, K., Kato, T., Takahashi, M., Fukuda, N.N., Murakami, S., Miyauchi,
556 E., Hino, S., Atarashi, K., Onawa, S., Fujimura, Y., Lockett, T., Clarke, J.M., Topping,
557 D.L., Tomita, M., Hori, S., Ohara, O., Morita, T., Koseki, H., Kikuchi, J., Honda, K.,
558 Hase, K., Ohno, H., 2013. Commensal microbe-derived butyrate induces the
559 differentiation of colonic regulatory T cells. *Nature* 504, 446–450.
560 <https://doi.org/10.1038/nature12721>

561 Godinho-Silva, C., Domingues, R.G., Rendas, M., Raposo, B., Ribeiro, H., da Silva, J.A.,
562 Vieira, A., Costa, R.M., Barbosa-Morais, N.L., Carvalho, T., Veiga-Fernandes, H.,
563 2019. Light-entrained and brain-tuned circadian circuits regulate ILC3s and gut
564 homeostasis. *Nature* 574, 254–258. <https://doi.org/10.1038/s41586-019-1579-3>

565 Goodpaster, B.H., Sparks, L.M., 2017. Metabolic Flexibility in Health and Disease. *Cell*
566 *Metab.* 25, 1027–1036. <https://doi.org/10.1016/j.cmet.2017.04.015>

567 Gots, J.S., 1971. CHAPTER 8 - Regulation of Purine and Pyrimidine Metabolism, in: Vogel,
568 H.J. (Ed.), *Metabolic Regulation (Third Edition)*. Academic Press, pp. 225–255.
569 <https://doi.org/10.1016/B978-0-12-299255-1.50015-8>

570 Guo, C.-J., Allen, B.M., Hiam, K.J., Dodd, D., Van Treuren, W., Higginbottom, S.,
571 Nagashima, K., Fischer, C.R., Sonnenburg, J.L., Spitzer, M.H., Fischbach, M.A.,
572 2019. Depletion of microbiome-derived molecules in the host using *Clostridium*
573 genetics. *Science* 366. <https://doi.org/10.1126/science.aav1282>

574 Gury-BenAri, M., Thaïss, C.A., Serafini, N., Winter, D.R., Giladi, A., Lara-Astiaso, D., Levy,
575 M., Salame, T.M., Weiner, A., David, E., Shapiro, H., Dori-Bachash, M., Pevsner-
576 Fischer, M., Lorenzo-Vivas, E., Keren-Shaul, H., Paul, F., Harmelin, A., Eberl, G.,
577 Itzkovitz, S., Tanay, A., Di Santo, J.P., Elinav, E., Amit, I., 2016. The Spectrum and
578 Regulatory Landscape of Intestinal Innate Lymphoid Cells Are Shaped by the
579 Microbiome. *Cell* 166, 1231-1246.e13. <https://doi.org/10.1016/j.cell.2016.07.043>

580 Hang, S., Paik, D., Yao, L., Kim, E., Trinath, J., Lu, J., Ha, S., Nelson, B.N., Kelly, S.P., Wu,
581 L., Zheng, Y., Longman, R.S., Rastinejad, F., Devlin, A.S., Krout, M.R., Fischbach,
582 M.A., Littman, D.R., Huh, J.R., 2019. Bile acid metabolites control TH17 and Treg cell
583 differentiation. *Nature* 576, 143–148. <https://doi.org/10.1038/s41586-019-1785-z>
584 Hashimoto, T., Perlot, T., Rehman, A., Trichereau, J., Ishiguro, H., Paolino, M., Sigl, V.,
585 Hanada, T., Hanada, R., Lipinski, S., Wild, B., Camargo, S.M.R., Singer, D., Richter,
586 A., Kuba, K., Fukamizu, A., Schreiber, S., Clevers, H., Verrey, F., Rosenstiel, P.,
587 Penninger, J.M., 2012. ACE2 links amino acid malnutrition to microbial ecology and
588 intestinal inflammation. *Nature* 487, 477–481. <https://doi.org/10.1038/nature11228>
589 Honda, K., Littman, D.R., 2016. The microbiota in adaptive immune homeostasis and
590 disease. *Nature* 535, 75–84. <https://doi.org/10.1038/nature18848>
591 Hopkins, E.G.D., Roumeliotis, T.I., Mullineaux-Sanders, C., Choudhary, J.S., Frankel, G.,
592 2019. Intestinal Epithelial Cells and the Microbiome Undergo Swift Reprogramming at
593 the Inception of Colonic *Citrobacter rodentium* Infection. *mBio* 10.
594 <https://doi.org/10.1128/mBio.00062-19>
595 Hoytema van Konijnenburg, D.P., Reis, B.S., Pedicord, V.A., Farache, J., Victora, G.D.,
596 Mucida, D., 2017. Intestinal epithelial and intraepithelial T cell cross-talk mediates a
597 dynamic response to infection. *Cell* 171, 783-794.e13.
598 <https://doi.org/10.1016/j.cell.2017.08.046>
599 Hsieh, S.-Y., Shih, T.-C., Yeh, C.-Y., Lin, C.-J., Chou, Y.-Y., Lee, Y.-S., 2006. Comparative
600 proteomic studies on the pathogenesis of human ulcerative colitis. *Proteomics* 6,
601 5322–5331. <https://doi.org/10.1002/pmic.200500541>
602 Huang, S.C.-C., Everts, B., Ivanova, Y., O’Sullivan, D., Nascimento, M., Smith, A.M., Beatty,
603 W., Love-Gregory, L., Lam, W.Y., O’Neill, C.M., Yan, C., Du, H., Abumrad, N.A.,
604 Urban, J.F., Artyomov, M.N., Pearce, E.L., Pearce, E.J., 2014. Cell-intrinsic lysosomal
605 lipolysis is essential for alternative activation of macrophages. *Nat. Immunol.* 15, 846–
606 855. <https://doi.org/10.1038/ni.2956>
607 Hviid, A., Svanström, H., Frisch, M., 2011. Antibiotic use and inflammatory bowel diseases in
608 childhood. *Gut* 60, 49–54. <https://doi.org/10.1136/gut.2010.219683>
609 Iliev, I.D., Leonardi, I., 2017. Fungal dysbiosis: immunity and interactions at mucosal barriers.
610 *Nat. Rev. Immunol.* 17, 635–646. <https://doi.org/10.1038/nri.2017.55>
611 Jackson, D.N., Panopoulos, M., Neumann, W.L., Turner, K., Cantarel, B.L., Thompson-
612 Snipes, L., Dassopoulos, T., Feagins, L.A., Souza, R.F., Mills, J.C., Blumberg, R.S.,
613 Venuprasad, K., Thompson, W.E., Theiss, A.L., 2020. Mitochondrial dysfunction
614 during loss of prohibitin 1 triggers Paneth cell defects and ileitis. *Gut*.
615 <https://doi.org/10.1136/gutjnl-2019-319523>
616 Kaminski, M.M., Sauer, S.W., Klemke, C.-D., Süß, D., Okun, J.G., Krammer, P.H., Gülow,
617 K., 2010. Mitochondrial reactive oxygen species control T cell activation by regulating
618 IL-2 and IL-4 expression: mechanism of ciprofloxacin-mediated immunosuppression.
619 *J. Immunol. Baltim. Md 1950* 184, 4827–4841.
620 <https://doi.org/10.4049/jimmunol.0901662>
621 Khaloian, S., Rath, E., Hammoudi, N., Gleisinger, E., Blutke, A., Giesbertz, P., Berger, E.,
622 Metwaly, A., Waldschmitt, N., Allez, M., Haller, D., 2020. Mitochondrial impairment
623 drives intestinal stem cell transition into dysfunctional Paneth cells predicting Crohn’s
624 disease recurrence. *Gut*. <https://doi.org/10.1136/gutjnl-2019-319514>
625 Kim, M., Qie, Y., Park, J., Kim, C.H., 2016. Gut Microbial Metabolites Fuel Host Antibody
626 Responses. *Cell Host Microbe* 20, 202–214.
627 <https://doi.org/10.1016/j.chom.2016.07.001>
628 Klose, C.S.N., Artis, D., 2016. Innate lymphoid cells as regulators of immunity, inflammation
629 and tissue homeostasis. *Nat. Immunol.* 17, 765–774. <https://doi.org/10.1038/ni.3489>
630 Lampropoulou, V., Sergushichev, A., Bambouskova, M., Nair, S., Vincent, E.E., Loginicheva,
631 E., Cervantes-Barragan, L., Ma, X., Huang, S.C.-C., Griss, T., Weinheimer, C.J.,
632 Khader, S., Randolph, G.J., Pearce, E.J., Jones, R.G., Diwan, A., Diamond, M.S.,
633 Artyomov, M.N., 2016. Itaconate Links Inhibition of Succinate Dehydrogenase with

634 Macrophage Metabolic Remodeling and Regulation of Inflammation. *Cell Metab.* 24,
635 158–166. <https://doi.org/10.1016/j.cmet.2016.06.004>

636 Lavelle, A., Sokol, H., 2020. Gut microbiota-derived metabolites as key actors in
637 inflammatory bowel disease. *Nat. Rev. Gastroenterol. Hepatol.*
638 <https://doi.org/10.1038/s41575-019-0258-z>

639 Leber, A., Hontecillas, R., Tubau-Juni, N., Zoccoli-Rodriguez, V., Abedi, V., Bassaganya-
640 Riera, J., 2018. NLRX1 Modulates Immunometabolic Mechanisms Controlling the
641 Host-Gut Microbiota Interactions during Inflammatory Bowel Disease. *Front. Immunol.*
642 9, 363. <https://doi.org/10.3389/fimmu.2018.00363>

643 Lewis, G., Wang, B., Shafiei Jahani, P., Hurrell, B.P., Banie, H., Aleman Muench, G.R.,
644 Maazi, H., Helou, D.G., Howard, E., Galle-Treger, L., Lo, R., Santosh, S., Baltus, A.,
645 Bongers, G., San-Mateo, L., Gilliland, F.D., Rehan, V.K., Soroosh, P., Akbari, O.,
646 2019. Dietary Fiber-Induced Microbial Short Chain Fatty Acids Suppress ILC2-
647 Dependent Airway Inflammation. *Front. Immunol.* 10.
648 <https://doi.org/10.3389/fimmu.2019.02051>

649 Lin, C.-C.J., Wang, M.C., 2017. Microbial metabolites regulate host lipid metabolism through
650 NR5A-Hedgehog signalling. *Nat. Cell Biol.* 19, 550–557.
651 <https://doi.org/10.1038/ncb3515>

652 Liu, T.-C., Gurram, B., Baldridge, M.T., Head, R., Lam, V., Luo, C., Cao, Y., Simpson, P.,
653 Hayward, M., Holtz, M.L., Bousounis, P., Noe, J., Lerner, D., Cabrera, J., Biank, V.,
654 Stephens, M., Huttenhower, C., McGovern, D.P., Xavier, R.J., Stappenbeck, T.S.,
655 Salzman, N.H., 2016. Paneth cell defects in Crohn’s disease patients promote
656 dysbiosis. *JCI Insight* 1, e86907. <https://doi.org/10.1172/jci.insight.86907>

657 Liu, Y., Hou, Y., Wang, G., Zheng, X., Hao, H., 2020. Gut Microbial Metabolites of Aromatic
658 Amino Acids as Signals in Host-Microbe Interplay. *Trends Endocrinol. Metab. TEM.*
659 <https://doi.org/10.1016/j.tem.2020.02.012>

660 Lunt, S.Y., Vander Heiden, M.G., 2011. Aerobic Glycolysis: Meeting the Metabolic
661 Requirements of Cell Proliferation. *Annu. Rev. Cell Dev. Biol.* 27, 441–464.
662 <https://doi.org/10.1146/annurev-cellbio-092910-154237>

663 Luu, M., Pautz, S., Kohl, V., Singh, R., Romero, R., Lucas, S., Hofmann, J., Raifer, H.,
664 Vachharajani, N., Carrascosa, L.C., Lamp, B., Nist, A., Stiewe, T., Shaul, Y.,
665 Adhikary, T., Zaiss, M.M., Lauth, M., Steinhoff, U., Visekruna, A., 2019. The short-
666 chain fatty acid pentanoate suppresses autoimmunity by modulating the metabolic-
667 epigenetic cross-talk in lymphocytes. *Nat. Commun.* 10, 1–12.
668 <https://doi.org/10.1038/s41467-019-08711-2>

669 Macia, L., Tan, J., Vieira, A.T., Leach, K., Stanley, D., Luong, S., Maruya, M., Ian McKenzie,
670 C., Hijikata, A., Wong, C., Binge, L., Thorburn, A.N., Chevalier, N., Ang, C., Marino,
671 E., Robert, R., Offermanns, S., Teixeira, M.M., Moore, R.J., Flavell, R.A., Fagarasan,
672 S., Mackay, C.R., 2015. Metabolite-sensing receptors GPR43 and GPR109A facilitate
673 dietary fibre-induced gut homeostasis through regulation of the inflammasome. *Nat.*
674 *Commun.* 6, 6734. <https://doi.org/10.1038/ncomms7734>

675 Mao, K., Baptista, A.P., Tamoutounour, S., Zhuang, L., Bouladoux, N., Martins, A.J., Huang,
676 Y., Gerner, M.Y., Belkaid, Y., Germain, R.N., 2018. Innate and adaptive lymphocytes
677 sequentially shape the gut microbiota and lipid metabolism. *Nature* 554, 255–259.
678 <https://doi.org/10.1038/nature25437>

679 Martin-Gallausiaux, C., Larraufie, P., Jarry, A., Béguet-Crespel, F., Marinelli, L., Ledue, F.,
680 Reimann, F., Blottière, H.M., Lapaque, N., 2018. Butyrate Produced by Commensal
681 Bacteria Down-Regulates Indolamine 2,3-Dioxygenase 1 (IDO-1) Expression via a
682 Dual Mechanism in Human Intestinal Epithelial Cells. *Front. Immunol.* 9.
683 <https://doi.org/10.3389/fimmu.2018.02838>

684 Mills, E.L., Kelly, B., Logan, A., Costa, A.S.H., Varma, M., Bryant, C.E., Tourlomousis, P.,
685 Däbritz, J.H.M., Gottlieb, E., Latorre, I., Corr, S.C., McManus, G., Ryan, D., Jacobs,
686 H.T., Szibor, M., Xavier, R.J., Braun, T., Frezza, C., Murphy, M.P., O’Neill, L.A., 2016.
687 Succinate Dehydrogenase Supports Metabolic Repurposing of Mitochondria to Drive

688 Inflammatory Macrophages. *Cell* 167, 457-470.e13.
689 <https://doi.org/10.1016/j.cell.2016.08.064>

690 Milner, J.A., 1985. Metabolic Aberrations Associated with Arginine Deficiency. *J. Nutr.* 115,
691 516–523. <https://doi.org/10.1093/jn/115.4.516>

692 Mottawea, W., Chiang, C.-K., Mühlbauer, M., Starr, A.E., Butcher, J., Abujamel, T., Deeke,
693 S.A., Brandel, A., Zhou, H., Shokralla, S., Hajibabaei, M., Singleton, R., Benchimol,
694 E.I., Jobin, C., Mack, D.R., Figeys, D., Stintzi, A., 2016. Altered intestinal microbiota–
695 host mitochondria cross-talk in new onset Crohn’s disease. *Nat. Commun.* 7, 1–14.
696 <https://doi.org/10.1038/ncomms13419>

697 Omenetti, S., Bussi, C., Metidji, A., Iseppon, A., Lee, S., Tolaini, M., Li, Y., Kelly, G.,
698 Chakravarty, P., Shoaie, S., Gutierrez, M.G., Stockinger, B., 2019. The Intestine
699 Harbors Functionally Distinct Homeostatic Tissue-Resident and Inflammatory Th17
700 Cells. *Immunity* 51, 77-89.e6. <https://doi.org/10.1016/j.immuni.2019.05.004>

701 O’Neill, L.A.J., Kishton, R.J., Rathmell, J., 2016. A guide to immunometabolism for
702 immunologists. *Nat. Rev. Immunol.* 16, 553–565. <https://doi.org/10.1038/nri.2016.70>

703 Park, J., Kim, M., Kang, S.G., Jannasch, A.H., Cooper, B., Patterson, J., Kim, C.H., 2015.
704 Short-chain fatty acids induce both effector and regulatory T cells by suppression of
705 histone deacetylases and regulation of the mTOR-S6K pathway. *Mucosal Immunol.*
706 8, 80–93. <https://doi.org/10.1038/mi.2014.44>

707 Pasolli, E., Asnicar, F., Manara, S., Zolfo, M., Karcher, N., Armanini, F., Beghini, F., Manghi,
708 P., Tett, A., Ghensi, P., Collado, M.C., Rice, B.L., DuLong, C., Morgan, X.C., Golden,
709 C.D., Quince, C., Huttenhower, C., Segata, N., 2019. Extensive Unexplored Human
710 Microbiome Diversity Revealed by Over 150,000 Genomes from Metagenomes
711 Spanning Age, Geography, and Lifestyle. *Cell* 176, 649-662.e20.
712 <https://doi.org/10.1016/j.cell.2019.01.001>

713 Pigneur, B., Sokol, H., 2016. Fecal microbiota transplantation in inflammatory bowel disease:
714 the quest for the holy grail. *Mucosal Immunol.* 9, 1360–1365.
715 <https://doi.org/10.1038/mi.2016.67>

716 Richard, M.L., Sokol, H., 2019. The gut mycobiota: insights into analysis, environmental
717 interactions and role in gastrointestinal diseases. *Nat. Rev. Gastroenterol. Hepatol.*
718 16, 331–345. <https://doi.org/10.1038/s41575-019-0121-2>

719 Rodríguez-Prados, J.-C., Través, P.G., Cuenca, J., Rico, D., Aragónés, J., Martín-Sanz, P.,
720 Cascante, M., Boscá, L., 2010. Substrate fate in activated macrophages: a
721 comparison between innate, classic, and alternative activation. *J. Immunol. Baltim.*
722 *Md* 1950 185, 605–614. <https://doi.org/10.4049/jimmunol.0901698>

723 Rolot, M., O’Sullivan, T.E., 2020. Living with Yourself: Innate Lymphoid Cell
724 Immunometabolism. *Cells* 9. <https://doi.org/10.3390/cells9020334>

725 Rooks, M.G., Garrett, W.S., 2016. Gut microbiota, metabolites and host immunity. *Nat. Rev.*
726 *Immunol.* 16, 341–352. <https://doi.org/10.1038/nri.2016.42>

727 Rosser, E.C., Piper, C.J.M., Matei, D.E., Blair, P.A., Rendeiro, A.F., Orford, M., Alber, D.G.,
728 Krausgruber, T., Catalan, D., Klein, N., Manson, J.J., Drozdov, I., Bock, C.,
729 Wedderburn, L.R., Eaton, S., Mauri, C., 2020. Microbiota-Derived Metabolites
730 Suppress Arthritis by Amplifying Aryl-Hydrocarbon Receptor Activation in Regulatory
731 B Cells. *Cell Metab.* <https://doi.org/10.1016/j.cmet.2020.03.003>

732 Sarin, S.K., Pande, A., Schnabl, B., 2019. Microbiome as a therapeutic target in alcohol-
733 related liver disease. *J. Hepatol.* 70, 260–272.
734 <https://doi.org/10.1016/j.jhep.2018.10.019>

735 Schneider, C., O’Leary, C.E., von Moltke, J., Liang, H.-E., Ang, Q.Y., Turnbaugh, P.J.,
736 Radhakrishnan, S., Pellizzon, M., Ma, A., Locksley, R.M., 2018. A Metabolite-
737 Triggered Tuft Cell-ILC2 Circuit Drives Small Intestinal Remodeling. *Cell* 174, 271-
738 284.e14. <https://doi.org/10.1016/j.cell.2018.05.014>

739 Scott, N.A., Andrusaite, A., Andersen, P., Lawson, M., Alcon-Giner, C., Leclaire, C., Caim,
740 S., Le Gall, G., Shaw, T., Connolly, J.P.R., Roe, A.J., Wessel, H., Bravo-Blas, A.,
741 Thomson, C.A., Kästele, V., Wang, P., Peterson, D.A., Bancroft, A., Li, X., Grecis,
742 R., Mowat, A.M., Hall, L.J., Travis, M.A., Milling, S.W.F., Mann, E.R., 2018. Antibiotics

743 induce sustained dysregulation of intestinal T cell immunity by perturbing
744 macrophage homeostasis. *Sci. Transl. Med.* 10.
745 <https://doi.org/10.1126/scitranslmed.aao4755>

746 Sena, L.A., Li, S., Jairaman, A., Prakriya, M., Ezponda, T., Hildeman, D.A., Wang, C.-R.,
747 Schumacker, P.T., Licht, J.D., Perlman, H., Bryce, P.J., Chandel, N.S., 2013.
748 Mitochondria are required for antigen-specific T cell activation through reactive
749 oxygen species signaling. *Immunity* 38, 225–236.
750 <https://doi.org/10.1016/j.immuni.2012.10.020>

751 Sender, R., Fuchs, S., Milo, R., 2016. Are We Really Vastly Outnumbered? Revisiting the
752 Ratio of Bacterial to Host Cells in Humans. *Cell* 164, 337–340.
753 <https://doi.org/10.1016/j.cell.2016.01.013>

754 Shajib, M.S., Khan, W.I., 2015. The role of serotonin and its receptors in activation of
755 immune responses and inflammation. *Acta Physiol. Oxf. Engl.* 213, 561–574.
756 <https://doi.org/10.1111/apha.12430>

757 Shkoporov, A.N., Hill, C., 2019. Bacteriophages of the Human Gut: The “Known Unknown” of
758 the Microbiome. *Cell Host Microbe* 25, 195–209.
759 <https://doi.org/10.1016/j.chom.2019.01.017>

760 Skelly, A.N., Sato, Y., Kearney, S., Honda, K., 2019. Mining the microbiota for microbial and
761 metabolite-based immunotherapies. *Nat. Rev. Immunol.* 19, 305–323.
762 <https://doi.org/10.1038/s41577-019-0144-5>

763 Smith, P.M., Howitt, M.R., Panikov, N., Michaud, M., Gallini, C.A., Bohlooly-Y, M., Glickman,
764 J.N., Garrett, W.S., 2013. The microbial metabolites, short-chain fatty acids, regulate
765 colonic Treg cell homeostasis. *Science* 341, 569–573.
766 <https://doi.org/10.1126/science.1241165>

767 Song, X., Sun, X., Oh, S.F., Wu, M., Zhang, Y., Zheng, W., Geva-Zatorsky, N., Jupp, R.,
768 Mathis, D., Benoist, C., Kasper, D.L., 2020. Microbial bile acid metabolites modulate
769 gut ROR γ ⁺ regulatory T cell homeostasis. *Nature* 577, 410–415.
770 <https://doi.org/10.1038/s41586-019-1865-0>

771 Tannahill, G.M., Curtis, A.M., Adamik, J., Palsson-McDermott, E.M., McGettrick, A.F., Goel,
772 G., Frezza, C., Bernard, N.J., Kelly, B., Foley, N.H., Zheng, L., Gardet, A., Tong, Z.,
773 Jany, S.S., Corr, S.C., Haneklaus, M., Caffrey, B.E., Pierce, K., Walmsley, S.,
774 Beasley, F.C., Cummins, E., Nizet, V., Whyte, M., Taylor, C.T., Lin, H., Masters, S.L.,
775 Gottlieb, E., Kelly, V.P., Clish, C., Auron, P.E., Xavier, R.J., O’Neill, L. a. J., 2013.
776 Succinate is an inflammatory signal that induces IL-1 β through HIF-1 α . *Nature* 496,
777 238–242. <https://doi.org/10.1038/nature11986>

778 Thaiss, C.A., Levy, M., Korem, T., Dohnalová, L., Shapiro, H., Jaitin, D.A., David, E., Winter,
779 D.R., Gury-BenAri, M., Tatirovsky, E., Tuganbaev, T., Federici, S., Zmora, N., Zeevi,
780 D., Dori-Bachash, M., Pevsner-Fischer, M., Kartvelishvily, E., Brandis, A., Harmelin,
781 A., Shibolet, O., Halpern, Z., Honda, K., Amit, I., Segal, E., Elinav, E., 2016.
782 Microbiota Diurnal Rhythmicity Programs Host Transcriptome Oscillations. *Cell* 167,
783 1495-1510.e12. <https://doi.org/10.1016/j.cell.2016.11.003>

784 Theiss, A.L., Idell, R.D., Srinivasan, S., Klapproth, J.-M., Jones, D.P., Merlin, D., Sitaraman,
785 S.V., 2007. Prohibitin protects against oxidative stress in intestinal epithelial cells.
786 *FASEB J. Off. Publ. Fed. Am. Soc. Exp. Biol.* 21, 197–206.
787 <https://doi.org/10.1096/fj.06-6801com>

788 Trapecar, M., Communal, C., Velazquez, J., Maass, C.A., Huang, Y.-J., Schneider, K.,
789 Wright, C.W., Butty, V., Eng, G., Yilmaz, O., Trumper, D., Griffith, L.G., 2020. Gut-
790 Liver Physiomics Reveal Paradoxical Modulation of IBD-Related Inflammation by
791 Short-Chain Fatty Acids. *Cell Syst.* 10, 223-239.e9.
792 <https://doi.org/10.1016/j.cels.2020.02.008>

793 Trompette, A., Gollwitzer, E.S., Pattaroni, C., Lopez-Mejia, I.C., Riva, E., Pernot, J., Ubags,
794 N., Fajas, L., Nicod, L.P., Marsland, B.J., 2018. Dietary Fiber Confers Protection
795 against Flu by Shaping Ly6c⁺ Patrolling Monocyte Hematopoiesis and CD8⁺ T Cell
796 Metabolism. *Immunity* 48, 992-1005.e8. <https://doi.org/10.1016/j.immuni.2018.04.022>

797 Vats, D., Mukundan, L., Odegaard, J.I., Zhang, L., Smith, K.L., Morel, C.R., Wagner, R.A.,
798 Greaves, D.R., Murray, P.J., Chawla, A., 2006. Oxidative metabolism and PGC-1beta
799 attenuate macrophage-mediated inflammation. *Cell Metab.* 4, 13–24.
800 <https://doi.org/10.1016/j.cmet.2006.05.011>

801 Wang, Y., Kuang, Z., Yu, X., Ruhn, K.A., Kubo, M., Hooper, L.V., 2017. The intestinal
802 microbiota regulates body composition through NFIL3 and the circadian clock.
803 *Science* 357, 912–916. <https://doi.org/10.1126/science.aan0677>

804 Wilhelm, C., Harrison, O.J., Schmitt, V., Pelletier, M., Spencer, S.P., Urban, J.F., Ploch, M.,
805 Ramalingam, T.R., Siegel, R.M., Belkaid, Y., 2016. Critical role of fatty acid
806 metabolism in ILC2-mediated barrier protection during malnutrition and helminth
807 infection. *J. Exp. Med.* 213, 1409–1418. <https://doi.org/10.1084/jem.20151448>

808 Yano, J.M., Yu, K., Donaldson, G.P., Shastri, G.G., Ann, P., Ma, L., Nagler, C.R., Ismagilov,
809 R.F., Mazmanian, S.K., Hsiao, E.Y., 2015. Indigenous bacteria from the gut
810 microbiota regulate host serotonin biosynthesis. *Cell* 161, 264–276.
811 <https://doi.org/10.1016/j.cell.2015.02.047>

812 Zhou, D., Chen, Y.-W., Zhao, Z.-H., Yang, R.-X., Xin, F.-Z., Liu, X.-L., Pan, Q., Zhou, H.,
813 Fan, J.-G., 2018. Sodium butyrate reduces high-fat diet-induced non-alcoholic
814 steatohepatitis through upregulation of hepatic GLP-1R expression. *Exp. Mol. Med.*
815 50, 1–12. <https://doi.org/10.1038/s12276-018-0183-1>

816 Zmora, N., Bashirdes, S., Levy, M., Elinav, E., 2017. The Role of the Immune System in
817 Metabolic Health and Disease. *Cell Metab.* 25, 506–521.
818 <https://doi.org/10.1016/j.cmet.2017.02.006>
819

Highlights and eTOC blurb

The gut microbiota is implicated in immune system functions. Regulation of the metabolic processes occurring in immune cells is crucial for the maintenance of homeostasis and immunopathogenesis. Michaudel & Sokol focused on emerging data demonstrating that the gut microbiota is an actor in immunometabolism, notably through the effect of metabolites.




Figure 1. Influence of the gut microbiota on immunometabolism




Figure 2. Mechanism of activation of the mTOR pathway by SCFA




Figure 3. Immunometabolism and the microbiota in diseases