



HAL
open science

Urban colonies are more resistant to a trace metal than their forest counterparts in the ant *Temnothorax nylanderi*

L Jacquier, C Doums, A Four-Chaboussant, R Peronnet, C Tirard, M Molet

► To cite this version:

L Jacquier, C Doums, A Four-Chaboussant, R Peronnet, C Tirard, et al.. Urban colonies are more resistant to a trace metal than their forest counterparts in the ant *Temnothorax nylanderi*. *Urban Ecosystems*, 2020. hal-03019372v1

HAL Id: hal-03019372

<https://hal.sorbonne-universite.fr/hal-03019372v1>

Submitted on 23 Nov 2020 (v1), last revised 4 Dec 2020 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Urban colonies are more resistant to a trace metal than their forest counterparts in the ant

2 *Temnothorax nylanderi*

3 L. Jacquier¹, C. Doums², A. Four-Chaboussant³, R. Peronnet¹, C. Tirard¹, M. Molet¹

4
5 ¹Institute of Ecology and Environmental Sciences of Paris, Sorbonne Université, Paris, France

6 ²EPHE, PSL Paris, France

7 ³Master EBE, Sorbonne Université, Paris, France

8
9 **Keywords:** Urbanization, common garden, cadmium resistance, social insects

10 **Abstract:** Cities can be used as open-sky replicates to assess the responses of organisms to
11 environmental changes, such as increased heat or pollution. In contrast with the numerous community-
12 level studies, population-level studies are rare, hugely biased towards birds, and rarely based on multiple
13 city replicates. They are also mostly based on solitary species, despite some social insects being
14 ecologically dominant and economically important. Our study aimed at assessing whether colonies of
15 the ant *Temnothorax nylanderi* from forest and urban populations exhibit distinct responses to pollution.
16 We collected colonies from four pairs of forest and urban habitats and reared them in the laboratory
17 under normal or cadmium-supplemented diets. We then measured the emergence rate and size of newly-
18 produced workers as well as the mortality rate of adult workers. We also assessed the effect of some
19 social factors such as colony size or size of field workers. In all four forest/city replicates, urban colonies
20 were less negatively affected by cadmium than forest colonies. We did not find any effect of social
21 factors on cadmium resistance, in contrast with what was found in other eusocial insects. We discuss
22 the potential adaptive nature of this differential response.

23 **Introduction**

24 United Nations predict that by 2030, more than 60% of the worldwide human population will live in
25 cities, resulting in a threefold increase of total urban areas (1). Urbanization is associated with many

26 environmental changes, both abiotic and biotic (e.g. modified land use, habitat fragmentation, decreased
27 soil permeability, increased temperatures known as Urban Heat Island effect, noise, light, and chemical
28 pollutions including trace metals, exotic species (2–4)). Because these changes are quite fast and recent
29 on the evolutionary timescale, cities can be seen as open sky laboratory replicates for understanding the
30 responses of species facing rapid environmental changes (5,6). Most studies investigating the impacts
31 of urbanization on wildlife have been conducted at the community level, with a taxonomic skew toward
32 birds, although fishes, lizards, amphibians, arthropods (7–14) and plants (15,16) have been occasionally
33 studied. Urbanization alters the composition of communities (higher or lower species abundance),
34 functional traits (e.g. reduced body size), and interaction networks (10–14,16–19). In contrast, species-
35 level studies are less common (see (6) for a good summary) Relative to non-urban populations, animals
36 from urban populations express altered behavior (e.g. songs with distinct pitches, loss of migratory
37 behavior (17,18)), changes in physiology (e.g. lower glucocorticoid secretion in response to stress,
38 higher oxidative stress (19)), phenology (earlier reproduction (20,21)), circadian rhythm (faster
39 circadian oscillations (22)), health (higher parasitism rate (23)) and/or morphology (7,24). These shifts
40 in life-history traits are often seen as a response, adaptive or not, to urban stressors. However, because
41 cities are extremely diverse and because most studies are based on a single city and not multiple city
42 replicates (but see (21,25)), it is difficult to make general conclusions and identify a robust syndrome of
43 urbanization for population-level traits. In addition, population studies are generally correlative
44 (17,24,26–34), and do not quantify the respective contributions of genetics and phenotypic plasticity.
45 So far, common garden experiments between urban and non-urban populations highlighted a genetic
46 basis for the lower dispersal ability of urban plants (15), earlier reproduction of urban birds and ants
47 (20,21) and for specific behaviors of urban birds (lower migratory behavior, increased boldness and
48 stress). Additional common garden experiments are thus needed to complement correlative studies and
49 assess causality and determination of various other life-history traits associated with urban life.

50 Pollution is one of the main chronic urban stressors. Urban habitats are usually more polluted because
51 of traffic and industrial activities (35). For example, traffic-generated air pollution is responsible of
52 increased oxidative stress leading to genetic damages, respiratory diseases and brain inflammation

53 (29,36,37) as well as higher susceptibility to parasites (23) in animals. Cities are also contaminated by
54 trace metals. For example, cadmium has a concentration of 0.30 mg.kg^{-1} in soil of rural woods and 2.45
55 mg.kg^{-1} in urban woods (35). Cadmium alters plant physiology by interfering with calcium uptake (38).
56 It also impacts various life history traits, causing longer development in fishes and arthropods, decreased
57 hatching rate in fishes, sea stars and arthropods (39–41), decreased body size in fishes (40) and
58 immunosuppression in rats (42).

59 Ants represent a large portion of the total earth animal biomass (43). They are found in a large diversity
60 of ecosystems including urban areas and can deeply affect the functioning of ecosystems (44–46). This
61 ecological success partly originates from their social lifestyle. Ants live in colonies made of few tens to
62 millions of individuals where labor is divided among individuals according to their caste. Eusociality
63 could enhance robustness against environmental changes. For example, eusocial species of bees are less
64 negatively impacted by neocotinoids than solitary species (47), and models suggest that, within species,
65 larger colonies resist better to pesticides (48). In ants, social life enhances resistance to parasites and
66 starvation (49,50). In spite of their ecological interest and their social life, few studies investigating the
67 effect of urbanization have been conducted on ants (5,51–53) and even fewer at the species level (54).
68 How sociality could help ants resist anthropic stressors is an open question.

69 The aim of this study was to test whether there is a general effect of urbanization on resistance to trace
70 metals (cadmium) in the small acorn ant *Temnothorax nylanderi*. Ants are a good model to study the
71 effects of trace metals because they live in long-lived colonies and they forage locally around their nest,
72 exposing themselves to bioaccumulation (55). We tested whether colonies of *T. nylanderi* from city and
73 forest populations exhibit a differential response to pollution by feeding them with food enriched with
74 cadmium or not, in a common garden laboratory setup, and by measuring several traits of the newly
75 produced workers. We expected urban colonies to resist cadmium better because of their long history
76 (several generations) of exposure to trace metal pollution relative to non-urban colonies. Moreover, this
77 species offers the opportunity to assess the effects of social life on cadmium resistance, as colony size
78 varies from less than ten to a few hundred individuals. The common-garden setup allowed to test for a

79 causal link between the stressor and the observed shift in traits (mortality of adult workers, and
80 emergence rate and size of newly produced workers).

81 **Material and methods**

82 *Colony collection and rearing*

83 *Temnothorax nylanderi* is a small ant with small colonies (up to few hundred workers) that nests in
84 acorns and twigs on the ground. Colonies are easy to collect and rear in the laboratory, and they produce
85 new workers in two months at 22-26°C cycle (56). Two hundred and thirty-eight colonies of *T. nylanderi*
86 were collected from eight sites at three locations in France (Paris, Bordeaux, Lyon). Each location had
87 one pair of collection sites: a central urban park (city habitat) and a forest located between 23 and 52
88 km from the city (forest habitat). Two pairs were collected in Paris location from two different parks
89 and forest sites and in two different years, 2017 and 2018 (see table 1). They are subsequently called
90 “Paris’17” for colonies collected in March 2017 and “Paris’18” for colonies collected in March 2018.
91 Colonies from Lyon and Bordeaux were collected in February and March 2018. After collection,
92 colonies were brought back to the laboratory and workers were counted. We discarded colonies with no
93 queen or more than one queen (72 colonies) as well as colonies parasitized by cestodes (9 colonies,
94 identified by the pale orange colour of workers). A total of 157 queenright colonies were used for the
95 experiment and reared in foraging plastic boxes (11.5 x 11.5 x 5.5 cm) containing an artificial nest made
96 of two microscope slides separated by a thin layer of plastic foam with hollow chambers. The top slide
97 was covered with dark plastic to protect the colony from the light. Plastic boxes were kept in
98 temperature-controlled chambers (reference CTS TP10/600) at 22°C from 7pm to 7am and 27°C from
99 7am to 7pm. These stressful temperatures were chosen to speed up development and to amplify the
100 effects of cadmium (57).

101 *Experimental design*

102 Our feeding treatment included two conditions: cadmium and control, applied to all locations and all
103 pairs of habitats. In order to avoid any bias in colony size between control and cadmium conditions, we
104 evenly assigned colonies to cadmium and control based on their colony size for each habitat within each

105 location. Colonies were randomly distributed between the two chambers independently of their location,
106 habitat and treatment. All original brood were removed from the colonies except for about 30 second
107 instar larvae. When colonies had less than 30 second instar larvae, we left all of them in the colony.
108 Newly laid eggs were removed every week to avoid the hatching of extra larvae. We fed colonies for 61
109 days every other day with a mixture of diluted honey and dried crickets containing 100µg/g cadmium
110 (cadmium condition) or not (control condition). This cadmium concentration corresponds to the LC₅₀
111 (concentration at which half of the treated individuals die) of *Spodoptera littoralis* (M.A. Pottier
112 personal communication). The mortality rate of the adult workers present in the colony from the
113 beginning of the experiment (i.e. workers born in the field, subsequently called field workers) was
114 computed based on the frequent removal of corpses and defined as the total number of dead field workers
115 at the end of the experiment divided by the total number of field workers at the start of the experiment.

116 The experiment was performed over two different years: 2017 for Paris'17, and 2018 for
117 Bordeaux, Lyon and Paris'18. For both years, colonies were acclimated for 14 days in laboratory before
118 the experiment started. The experiment lasted for 61 days, so any larva that would have escaped the egg
119 removal protocol and hatched could not have reached adulthood before the end of the experiment. The
120 emerged workers (subsequently called lab workers) were identified based on their pale color
121 (*Temnothorax* larvae do not spin a cocoon) and collected once they started to move. They were
122 anesthetized by cooling down and stored in ethanol for subsequent morphological measurements. We
123 computed the emergence rate of larvae as the ratio between the number of emerged adults (both workers
124 and sexuals) and the initial number of larvae. The date of emergence was recorded in order to compute
125 development duration for each lab worker, i.e. the duration from the second instar larva to the first
126 movements of the newly emerged lab worker. Nine queens died during the experiment (one from
127 Bordeaux-city-cadmium, one from Lyon-city-control, three from Paris'17-city-control, one from
128 Paris'17-city-cadmium, two from Paris'17-forest-control and one from Paris'18-forest-cadmium). We
129 excluded these colonies from the analyses, so 148 colonies were left in.

130

131 ***Morphological measurements***

132 At the end of the experiment, the heads of lab and field workers were stuck on double-sided tape and
133 photographed at a x5 magnification under a stereomicroscope (Zeiss Discovery.V12) connected to a
134 computer via a camera (Zeiss AxioCamICc). Head width was measured from eye to eye (just posterior
135 to the eyes) with ImageJ (<https://imagej.nih.gov/ij/>; (58) and used as a proxy of body size (56).

136

137 *Statistical analyses*

138 We used R v1.1.383 (R Core Team, 2015) to conduct our statistical analyses. We tested for the effect of
139 treatment (control, ie fed without cadmium, and cadmium, ie fed with cadmium), habitat (city or forest)
140 and location (Paris'17, Paris'18, Bordeaux, Lyon), and their two level interactions on three dependent
141 variables (mortality of field workers; emergence rate and size of lab workers). We considered the two
142 collections in Paris as two independent locations (Paris'17 and Paris'18) as they were carried out during
143 two different years and at two different pairs of sites. We also tested for the effect of two covariates
144 (initial colony size and mean size of field workers) on each dependent variable to investigate the effects
145 of these social parameters. To test for a buffering effect of colony size or field worker size, we included
146 in the models the interaction between treatment and colony size or field worker size. For the two colony-
147 level dependent variables (mortality rate of field workers and emergence rate of lab workers), we used
148 generalized linear models (GLMs). Because mortality and emergence rates were around 50%, using a
149 quasibinomial model was not better than using a Gaussian model with log-transformed data, and both
150 models provided the same qualitative results. We therefore presented data from the Gaussian model for
151 simplicity and because this allows to get AIC information. Regarding the individual-level dependent
152 variable (size of lab workers), we used a linear mixed-effect model (*lmer*) with the package *nlme* (59),
153 including colony as a random effect. We visually checked the homogeneity of variance and normality
154 of residuals. The minimal adequate model was chosen based on the AIC scores (*stepmodel*)(SI). The *P*-
155 values for the explanatory variables and interactions were obtained by removing or adding the variable
156 of interest from/to this minimal adequate model (60). *P*-values were obtained by comparing the model
157 fitted with maximum likelihood method with and without the variable of interest using a log likelihood

158 ratio test (*anova*). When necessary, we used *lsmeans* to compare means among the different levels of
159 the explanatory variables.

160 **Results**

161 The 148 colonies produced a total of 1269 lab workers (922 for the control condition and 347 for the
162 cadmium condition, Table 1), 215 males (176 for the control condition and 39 for the cadmium
163 condition, Table 1), and no gynes in 61 days. Because only few males (39) emerged under cadmium
164 condition, statistical power was lacking and we chose not to present results on male traits.

165

166 ***Field workers***

167 Only nine queens died during the experiment. There was no differential queen mortality between control
168 and cadmium conditions (Fisher's exact test, $P = 0.49$) nor between city and forest colonies (Fisher's
169 exact test, $P = 0.32$).

170 Colony size (the number of field workers by colony) was only affected by the location ($F = 4.90$, $P <$
171 0.001), with larger colonies found in Paris'18 (P -adjusted=0.041 for Paris'18-Bordeaux, P -
172 adjusted=0.014 for Paris'18-Lyon and P -adjusted = 0.002 for Paris'17-Paris'18). The size of field
173 workers also differed among locations ($F = 17.58$; $P < 0.001$) with the bigger field workers found in
174 Paris'17 and Paris'18 (P -adjusted Paris'17-Paris'18=0.95, P -adjusted Paris'17-Bordeaux and Paris'18-
175 Bordeaux $< 10^{-6}$, P -adjusted Paris'17- Lyon and Paris-18-Lyon $< 10^{-3}$), and it was positively correlated
176 with colony size ($F = 13.10$, $P < 0.001$; slope estimate = $1.285e-04$). However, there was no difference
177 in the size of field workers ($p = 0.187$, $F = 1.76$) and in colony size ($P = 0.87$, $F = 0.024$) between habitats.

178 At all locations, the mortality rate of field workers was significantly higher in cadmium condition than
179 in control condition ($F = 115$, $P < 10^{-16}$) but the magnitude of the effect varied among locations (Treatment-
180 Location interaction, $P = 0.00037$, $F = 6.5$, Figure 1, Table S2). Mortality rate significantly differed
181 among locations ($F = 2.9$, $P = 0.037$) with a higher mortality rate in Lyon ($P = 0.07$, $t = 1.824$, figure 1).

182 Under the hypothesis that urban colonies resist better to stressors than forest colonies, we were expecting
183 to find a differential effect of cadmium on mortality rate depending on habitat (Treatment: Habitat
184 interaction). However we did not find this effect (Treatment: Habitat interaction, $P = 0.91$, $F = 0.1$) and

185 there was no effect of habitat on mortality rate ($P=0.54$, $F=0.36$). Regarding social parameters, colony
186 size and size of field workers as well as their interactions with treatment did not affect mortality rates
187 (Table S2).

188

189

190 *Laboratory workers*

191 In all locations, the emergence rate was significantly lower in cadmium condition than in control
192 condition ($F=104$, $P<10^{-16}$, figure 2A) even though the magnitude of the effect differed significantly
193 among locations (Treatment: Location interaction, $F=2.7$, $P = 0.045$, Table S2). The emergence rate
194 significantly differed among locations ($F=21$, $P<10^{-10}$). Interestingly, when treated with cadmium, forest
195 colonies experienced a stronger decrease in emergence rate than city colonies for all location (Treatment:
196 Habitat interaction, $F=4.22$, $P = 0.04$, figure 2A, Table S2). This was associated with an increase in the
197 number of larvae that apparently paused their development at early stages (15% in control vs 30% in
198 cadmium condition, $t=3.76$, $df=134$, $P=0.00024$). Emergence rate in control condition was the same
199 between city and forest habitats at each location (Paris'17: $t=1.8$, $df=16$, $P=0.08$; Lyon: $t=-0.07$, $df=5.9$,
200 $P=0.9$; Paris'18: $t=-0.44$, $df=18.9$, $P= 0.65$) except for Bordeaux ($t = -2.99$, $df=10.3$, $P = 0.01$), so the
201 observed Treatment: Habitat interaction was not caused by an initial differential emergence rate between
202 forest and city colonies when exposed to a stressful temperature. The only social parameters affecting
203 the emergence rate was the mean size of field worker (Table S2), with colonies containing larger field
204 workers having lower emergence rate ($F=9.7$, $P = 0.0022$, slope estimate = -2.4) whatever the treatment
205 (no significant interaction treatment and size of field workers ($P=0.62$, $F=0.24$)).

206 Colonies treated with cadmium produced significantly smaller lab workers in all locations but as
207 previously, the effect significantly differed in magnitude among locations (Treatment: Location
208 interaction, $X_3=9.87$, $P = 0.0197$, Figure 2B, $P<10^{-2}$, Table S3). Interestingly, this decrease in the size
209 of lab workers was stronger in forest colonies than in city colonies for all location (Treatment: Habitat
210 interaction, $X_1=5.03$, $P=0.02485$, table S3, Figure 2B). The size of lab workers under control condition
211 did not differ between the two types of habitats for all location (Paris'17: $t=1$, $P=0.29$, Lyon: $t=0.8$,

212 $P=0.4$, Paris'18: $t=-1.7$, $P=0.09$) except for Bordeaux ($t = -3.2$, $P = 0.002$), so the observed Treatment:
213 Habitat interaction was not caused by an initial difference in the size of lab workers between forest and
214 city colonies when exposed to a stressful temperature. Regarding social parameters, only colony size
215 affected significantly and positively the size of lab workers ($X_1=12.52$, $P= 0.00040$, slope estimate
216 $=1.7e-04$) whatever the treatment (no significant interaction between treatment and colony size) (Table
217 S3 to see all statistical results).

218

219 **Discussion**

220 The aim of our study was to assess whether urban colonies of the small acorn ant *Temnothorax nylanderi*
221 are more resistant to a trace metal than forest colonies. We exposed them to food with a very high
222 cadmium concentration. As expected, this caused an increase in adult mortality rates, and a decrease in
223 size and emergence rate of lab workers relative to the control condition, confirming that cadmium
224 condition has a negative effect on life history traits in this species as found in other insects (61,62). More
225 interestingly, our experiment showed that city colonies were less affected by cadmium than the forest
226 ones and this was observed in the four city/forest pairs studied, strongly suggesting a general effect of
227 urbanization in term of cadmium resistance. Whether this effect of urbanization could be considered as
228 an adaptive response is discussed below.

229 Different hypotheses could explain the lower vulnerability of city colonies to cadmium. First, because
230 urban soils often contain more cadmium than non-urban soils (on average, 2.45mg/kg for urban wood
231 soil in Paris, and 0.3 mg/kg in rural wood soils of Ile-de-France (35)), we can hypothesize that urban
232 colonies have acquired a resistance to cadmium following exposure. Such adaptation has already been
233 documented in the collembola *Orchesella cincta*, where individuals from populations of heavily
234 cadmium-contaminated disused mining sites are more resistant to cadmium (63). Second, the better
235 resistance to cadmium of city colonies observed in our experiment could reflect a general resistance to
236 other trace elements such as arsenic, zinc, lead or mercury that could be found at a higher concentration
237 in urban leaf litter because of intense industrial activity and proximity with traffic. Accordingly, urban

238 ants may have evolved general detoxification systems that can cope with various trace elements. Third,
239 more generally, the lower sensitivity to cadmium of city colonies could reflect a better tolerance to
240 various stressors. Indeed, it is important to remind that our rearing temperature was stressful for ants
241 (22-27 °C as opposed to the typical 15-20 °C experienced in nature, (56)). Because of the Urban Heat
242 Island effect, populations living in urban centers frequently experience higher temperature than their
243 non-urban counterparts and could therefore be less stressed by heat (64) allowing them to resist better
244 to cadmium. City colonies did not perform better than forest colonies under control conditions, but this
245 may be due to the ability of forest colonies to buffer stressors. When exposed to several stressing factors
246 (heat + cadmium), forest colonies may however collapse whereas urban colonies would resist.
247 Interestingly, studies performed on European black birds reared under common garden experiments
248 showed that urban populations have a lower stress response to human handling than their rural
249 counterparts (19), suggesting that urban populations can evolve a general tolerance to stressors. These
250 conclusions should be confirmed by further experiments to assess whether the observed lower sensitivity
251 to cadmium of city colonies in *T. nylanderi* is due to a general resistance to stress or to a more efficient
252 detoxification system.

253 An unexpected effect of cadmium was the increased frequency of larvae stopping their development.
254 These larvae did not grow over the entire course of the experiment, but they still looked alive and were
255 kept in the nest with the rest of the brood and cared for by workers, unlike dead larvae that were typically
256 discarded outside. In insects, environmental factors such as cold temperature, can induce a pause in
257 development (65). In ants, diapause seems to be endogenous, even if external factors such as photoperiod
258 and temperature can secondarily affect its occurrence and duration (66). However diapause can be
259 induced by other non-seasonal (thus unpredictable) environmental factors, such as trace metals in adult
260 earthworms or hypoxia in *C. elegans* embryos (67,68). Diapausing individuals usually survive longer in
261 response to a stress, for example flesh-fly larvae exposed to anoxic conditions (69). Thus, diapause can
262 be interpreted as a stress-avoidance mechanism(67). The higher proportion of paused *T. nylanderi* larvae
263 under cadmium condition may be a general defense mechanism to face a wide range of stressors. It is
264 unknown whether paused larvae can resume development or are just barely surviving and are doomed

265 to death. This and the potential impact of paused development on the life history traits of workers that
266 would manage to resume development deserves further investigation.

267 The differential response to cadmium between urban and non-urban colonies of *T. nylanderi* could have
268 genetic or environmental determination. In some hexapodes, populations that are tolerant to heavy metal
269 have duplicated metallothionein genes (*Drosophila melanogaster*, (70)) or an increased expression of a
270 single metallothionein gene (*Orchesella cincta*, (71)). A recent genomic analysis (Rad-seq) based on
271 our *T. nylanderi* colonies revealed a lack of genetic differentiation between habitats (Khimoun et al,
272 submitted). However, the authors found that 19 loci are under divergent selection between urban and
273 non-urban populations, but none of them corresponding to metallothionein genes. Interestingly, two of
274 these loci (one from the Lyon pair of populations and one from the Bordeaux pair of populations) code
275 for kinase C proteins involved in signalization pathways (72). Two other loci (both from the Lyon pair
276 of populations) code for a protein involved in mRNA processing and for a membrane protein. These
277 results support the idea that, in spite of a high gene flow among populations, adaptive genetic
278 differentiation can occur between urban and forest populations. Similarly, Schrader et al (73) found that
279 the invasive ant *Cardiocondyla obscurior* could rapidly adapt to environmental changes because of
280 transposable elements (TE) that generate genetic diversity through transpositions. Given that stressors
281 could trigger transposition events (74), it would be interesting to test whether TE could trigger genetic
282 adaptation in urban populations exposed to various stressors in *T. nylanderi*. Phenotypic differences
283 found after a single laboratory generation could also result from maternal effects and/or epigenetic
284 effects. For example, queens experiencing stressful conditions like pollution may undergo epigenetic
285 modifications of their genome that would be transmitted and help workers to face stressors. In that case,
286 urban queens could resist better to cadmium; however we did not find differential queen mortality
287 between city and forest colonies. This may be due to a lack of statistical power as very few queens died
288 over the experiment. Moreover, there was no differential queen mortality between control and cadmium
289 colonies despite a strong differential worker mortality. Some studies suggested that workers act as a
290 filter, limiting queen exposure to toxic compounds such as cadmium (75). It is possible that queens are

291 exposed to pollution during colony foundation only, and not after, limiting the potential involvement of
292 maternal effects.

293 In social groups, the phenotype of an individual is also determined by the surrounding social
294 environment. The social environment includes colony size (number of individuals), the phenotype of
295 other individuals (caste, size, morphology, physiology, behaviour), the proportion of individual types,
296 and complex emerging group properties including communication. In this study, we tested for a potential
297 buffering effect of colony size and size of field workers on the response to cadmium. We found two
298 effects of social factors on the phenotype of lab workers but they were similar with and without
299 cadmium. 1/ Larger colonies produced larger lab workers, a typical phenomenon in ants (56,76). 2/
300 Colonies with larger field workers had a lower emergence rate, which may be due to the longer
301 development duration of large workers that were then exposed to cadmium for longer. In contrast, in
302 bees, larger colonies were found to be more resistant to cadmium (47,48). Even though we did not detect
303 any effect of the social environment on colony sensitivity to cadmium, workers from urban colonies
304 could still be more used to polluted food or stressors. Workers could change their behavior when facing
305 cadmium, either through direct induction by cadmium or indirectly through the higher mortality
306 triggered by cadmium. The relative contribution of intrinsic factors within larvae and social factors (i.e.
307 nestmates that limit or enhance environmental variations experienced by larvae) deserves further
308 investigation through cross-fostering experiments. Social life could deeply impact the ability of ants to
309 face environmental changes. On one hand, it may help species to cope with environmental changes to
310 an extent that cannot be achieved by solitary species. The social filtering of the environment can for
311 instance protect the brood from temperature fluctuations, drought, starvation or pathogens. On the other
312 hand, consequences of social life such as extreme task-specialization or obligatory collective behavior
313 may make social species more vulnerable to environmental changes (77,78). Because of the ecologic
314 and economic values of social species (46,75) it is crucial to understand how they react to environmental
315 changes.

316 **Conclusion**

317 Our study provides a unique example of replicated common garden experiment highlighting a better
 318 resistance to cadmium in urban colonies. The plastic response is convergent across all four urban/forest
 319 pairs tested. This differential resistance can be genetically determined, due to the social environment, or
 320 a combination of both. The social factors that we assessed had no effect on colony resistance to
 321 cadmium, in contrast with what was found in bees. To our knowledge, the role of sociality in the
 322 adaptation to environmental changes has barely been investigated. The relative contribution of
 323 individual and social factors to the final adult phenotype could be assessed through cross fostering
 324 experiments.

325 **Figures Legend**

326 **Figure 1: Boxplot of the mortality rate of field workers by habitat, location and treatment.**

327 Colonies fed with cadmium are in red, colonies fed with control are in blue. Boxes show median,
 328 quartiles and extreme values. Statistics are presented in Table S2.

329 **Figure 2: Boxplot of worker traits by habitat, location and treatment.** Colonies fed with cadmium

330 are in red, colonies fed with control are in blue. (A) Emergence rate of lab workers. (B) Head width of
 331 lab workers (mm). Boxes show median, quartiles and extreme values

332

333 **Tables**

	Control	Cadmium	Site	GPS	
Paris'17	City	13	14	Jardin	48°50'59.684"N
		247W, 21M	206W, 8M	Ecologique	2° 21' 40.385"E
	Forest	13	13	Fontainebleau	48° 26' 2.595"N
		213W, 6M	87W, 1M		2°43' 52.842"E
	City	7	7	Jardin public	44°51'16.9"N
		39W, 12M	12W, 1M		0°35'16.5"W

Bordeaux	Forest	12	11	Monfaucon	44°52'23.6"N
		130W, 29M	11W, 11M		0°08'14.7"W
Lyon	City	4	5	Tête d'Or	45°46'44.2"N
		18W, 22M	10W, 1M		4°52'54.3"E
	Forest	4	4	Arandon	45°43'07.5"N
		28W, 4M	7W, 3M		5°26'44.5"E
Paris'18	City	10	11	Buttes	48°52'47.1"N
		107W, 64M	7W, 7M	Chaumont	2°23'45.6"E
	Forest	10	10	Chantilly	49°10'59.8"N
		140W, 23M	7W, 2M		2°28'43.6"E

334 Table 1: Number of *Temnothorax nylanderi* colonies used and number of workers (W) and males (M)
 335 collected for each location, treatment and habitat type.

336

337 References

- 338 1. Seto KC, Güneralp B, Hutyra LR. Global forecasts of urban expansion to 2030 and direct
 339 impacts on biodiversity and carbon pools. *Proc Natl Acad Sci.* 2 oct 2012;109(40):16083-8.
- 340 2. Pickett STA, Cadenasso ML, Grove JM, Nilon CH, Pouyat RV, Zipperer WC, et al. Urban
 341 Ecological Systems: Linking Terrestrial Ecological, Physical, and Socioeconomic Components
 342 of Metropolitan Areas. *Annu Rev Ecol Syst.* 2001;32(1):127-57.
- 343 3. Bornstein RD. Observations of the Urban Heat Island Effect in New York City. *J Appl Meteorol.*
 344 1 août 1968;7(4):575-82.
- 345 4. McDonnell MJ, Pickett STA, Groffman P, Bohlen P, Pouyat RV, Zipperer WC, et al. Ecosystem
 346 processes along an urban-to-rural gradient. *Urban Ecosyst.* 1 mars 1997;1(1):21-36.
- 347 5. Buczkowski G, Richmond DS. The Effect of Urbanization on Ant Abundance and Diversity: A
 348 Temporal Examination of Factors Affecting Biodiversity. *PLOS ONE.* 2 août 2012;7(8):e41729.
- 349 6. McDonnell MJ, Hahs AK. Adaptation and Adaptedness of Organisms to Urban Environments.
 350 *Annu Rev Ecol Evol Syst.* 2015;46(1):261-80.

- 351 7. Weller B, Ganzhorn JU. Carabid beetle community composition, body size, and fluctuating
352 asymmetry along an urban-rural gradient. *Basic Appl Ecol.* janv 2004;5(2):193-201.
- 353 8. Niemelä J, Kotze DJ, Venn S, Penev L, Stoyanov I, Spence J, et al. Carabid beetle assemblages
354 (Coleoptera, Carabidae) across urban-rural gradients: an international comparison. *Landsc Ecol.*
355 1 juill 2002;17(5):387-401.
- 356 9. Banaszak-Cibicka W, Żmihorski M. Wild bees along an urban gradient: winners and losers. *J*
357 *Insect Conserv.* 1 juin 2012;16(3):331-43.
- 358 10. Blair RB. Birds and Butterflies Along Urban Gradients in Two Ecoregions of the United States:
359 Is Urbanization Creating a Homogeneous Fauna? In: Lockwood JL, McKinney ML, éditeurs.
360 Biotic Homogenization [Internet]. Boston, MA: Springer US; 2001 [cité 15 janv 2019]. p. 33-56.
361 Disponible sur: https://doi.org/10.1007/978-1-4615-1261-5_3
- 362 11. Clergeau P, Savard J-PL, Mennechez G, Falardeau G. Bird Abundance and Diversity along an
363 Urban-Rural Gradient: A Comparative Study between Two Cities on Different Continents. *The*
364 *Condor.* 1998;100(3):413-25.
- 365 12. Germaine SS, Wakeling BF. Lizard species distributions and habitat occupation along an urban
366 gradient in Tucson, Arizona, USA. *Biol Conserv.* 1 févr 2001;97(2):229-37.
- 367 13. Alexandre CV, Esteves KE, de Moura e Mello MAM. Analysis of fish communities along a
368 rural-urban gradient in a neotropical stream (Piracicaba River Basin, São Paulo, Brazil).
369 *Hydrobiologia.* 1 mars 2010;641(1):97-114.
- 370 14. Pillsbury FC, Miller JR. Habitat and Landscape Characteristics Underlying Anuran Community
371 Structure Along an Urban-Rural Gradient. *Ecol Appl.* 2008;18(5):1107-18.
- 372 15. Cheptou P-O, Carrue O, Rouifed S, Cantarel A. Rapid evolution of seed dispersal in an urban
373 environment in the weed *Crepis sancta*. *Proc Natl Acad Sci.* 11 mars 2008;105(10):3796-9.
- 374 16. McKinney ML. Effects of urbanization on species richness: A review of plants and animals.
375 *Urban Ecosyst.* 1 juin 2008;11(2):161-76.
- 376 17. Mendes S, Colino-Rabanal VJ, Peris SJ. Bird song variations along an urban gradient: The case
377 of the European blackbird (*Turdus merula*). *Landsc Urban Plan.* 30 janv 2011;99(1):51-7.
- 378 18. Evans KL, Newton J, Gaston KJ, Sharp SP, McGowan A, Hatchwell BJ. Colonisation of urban
379 environments is associated with reduced migratory behaviour, facilitating divergence from
380 ancestral populations. *Oikos.* 2012;121(4):634-40.
- 381 19. Partecke J, Schwabl I, Gwinner E. Stress and the City: Urbanization and Its Effects on the Stress
382 Physiology in European Blackbirds. *Ecology.* 2006;87(8):1945-52.
- 383 20. Partecke Jesko, Van't Hof Thomas, Gwinner Eberhard. Differences in the timing of reproduction
384 between urban and forest European blackbirds (*Turdus merula*): result of phenotypic flexibility
385 or genetic differences? *Proc R Soc Lond B Biol Sci.* 7 oct 2004;271(1552):1995-2001.
- 386 21. Chick LD, Strickler SA, Perez A, Martin RA, Diamond SE. Urban heat islands advance the
387 timing of reproduction in a social insect. *J Therm Biol.* 1 févr 2019;80:119-25.
- 388 22. Dominoni D. M., Helm B., Lehmann M., Dowse H. B., Partecke J. Clocks for the city: circadian
389 differences between forest and city songbirds. *Proc R Soc B Biol Sci.* 22 juill
390 2013;280(1763):20130593.

- 391 23. Bradley CA, Altizer S. Urbanization and the ecology of wildlife diseases. *Trends Ecol Evol.* 1
392 févr 2007;22(2):95-102.
- 393 24. Liker A, Papp Z, Bókony V, Lendvai ÁZ. Lean birds in the city: body size and condition of
394 house sparrows along the urbanization gradient. *J Anim Ecol.* 2008;77(4):789-95.
- 395 25. Johnson Marc T. J., Prashad Cindy M., Lavoignat Mélanie, Saini Hargurdeep S. Contrasting the
396 effects of natural selection, genetic drift and gene flow on urban evolution in white clover
397 (*Trifolium repens*). *Proc R Soc B Biol Sci.* 25 juill 2018;285(1883):20181019.
- 398 26. Lu P, Yu Q, Liu J, Lee X. Advance of tree-flowering dates in response to urban climate change.
399 *Agric For Meteorol.* 29 août 2006;138(1):120-31.
- 400 27. Chamberlain DE, Cannon AR, Toms MP, Leech DI, Hatchwell BJ, Gaston KJ. Avian
401 productivity in urban landscapes: a review and meta-analysis. *Ibis.* 2009;151(1):1-18.
- 402 28. Leong M, Ponisio LC, Kremen C, Thorp RW, Roderick GK. Temporal dynamics influenced by
403 global change: bee community phenology in urban, agricultural, and natural landscapes. *Glob*
404 *Change Biol.* 2016;22(3):1046-53.
- 405 29. Isaksson C. Pollution and Its Impact on Wild Animals: A Meta-Analysis on Oxidative Stress.
406 *EcoHealth.* 1 sept 2010;7(3):342-50.
- 407 30. Scales J, Hyman J, Hughes M. Behavioral Syndromes Break Down in Urban Song Sparrow
408 Populations. *Ethology.* 2011;117(10):887-95.
- 409 31. Hof AE van't, Campagne P, Rigden DJ, Yung CJ, Lingley J, Quail MA, et al. The industrial
410 melanism mutation in British peppered moths is a transposable element. *Nature.* juin
411 2016;534(7605):102-5.
- 412 32. Slabbekoorn H, den Boer-Visser A. Cities Change the Songs of Birds. *Curr Biol.* 5 déc
413 2006;16(23):2326-31.
- 414 33. Lazić MM, Carretero MA, Crnobrnja-Isailović J, Kaliontzopoulou A. Effects of Environmental
415 Disturbance on Phenotypic Variation: An Integrated Assessment of Canalization, Developmental
416 Stability, Modularity, and Allometry in Lizard Head Shape. *Am Nat.* 1 janv 2015;185(1):44-58.
- 417 34. Lowe EC, Wilder SM, Hochuli DF. Urbanisation at Multiple Scales Is Associated with Larger
418 Size and Higher Fecundity of an Orb-Weaving Spider. *PLOS ONE.* 20 août 2014;9(8):e105480.
- 419 35. Foti L, Dubs F, Gignoux J, Lata J-C, Lerch TZ, Mathieu J, et al. Trace element concentrations
420 along a gradient of urban pressure in forest and lawn soils of the Paris region (France). *Sci Total*
421 *Environ.* 15 nov 2017;598:938-48.
- 422 36. Ieradi LA, Cristaldi M, Mascanzoni D, Cardarelli E, Grossi R, Campanella L. Genetic damage in
423 urban mice exposed to traffic pollution. *Environ Pollut.* 1 janv 1996;92(3):323-8.
- 424 37. Guxens M, Sunyer J. A review of epidemiological studies on neuropsychological effects of air
425 pollution. *Swiss Med Wkly.* 2012;141:w13322.
- 426 38. Stroiński A. Some physiological and biochemical aspects of plant resistance to cadmium effect.
427 I. Antioxidative system. *Acta Physiol Plant.* 1 juin 1999;21(2):175-88.
- 428 39. Gomot A. Toxic Effects of Cadmium on Reproduction, Development, and Hatching in the
429 Freshwater Snail *Lymnaea stagnalis* for Water Quality Monitoring. *Ecotoxicol Environ Saf.* 1 nov
430 1998;41(3):288-97.

- 431 40. Cervera A, Maymó AC, Sendra M, Martínez-Pardo R, Garcerá MD. Cadmium effects on
432 development and reproduction of *Oncopeltus fasciatus* (Heteroptera: Lygaeidae). *J Insect*
433 *Physiol.* août 2004;50(8):737-49.
- 434 41. den Besten PJ, Herwig HJ, Zandee DI, Voogt PA. Effects of cadmium and PCBs on
435 reproduction of the sea star *Asterias rubens*: Aberrations in the early development. *Ecotoxicol*
436 *Environ Saf.* 1 oct 1989;18(2):173-80.
- 437 42. Demenesku J, Mirkov I, Ninkov M, Popov Aleksandrov A, Zolotarevski L, Kataranovski D, et
438 al. Acute cadmium administration to rats exerts both immunosuppressive and proinflammatory
439 effects in spleen. *Toxicology.* 4 déc 2014;326:96-108.
- 440 43. Schultz TR. In search of ant ancestors. *Proc Natl Acad Sci U S A.* 19 déc 2000;97(26):14028-9.
- 441 44. Farji-Brener AG, Lescano N, Ghermandi L. Ecological engineering by a native leaf-cutting ant
442 increases the performance of exotic plant species. *Oecologia.* 1 mai 2010;163(1):163-9.
- 443 45. Sanders D, Veen FJF van. Ecosystem engineering and predation: the multi-trophic impact of two
444 ant species. *J Anim Ecol.* 2011;80(3):569-76.
- 445 46. Philpott SM, Armbrecht I. Biodiversity in tropical agroforests and the ecological role of ants and
446 ant diversity in predatory function. *Ecol Entomol.* 2006;31(4):369-77.
- 447 47. Rundlöf M, Andersson GKS, Bommarco R, Fries I, Hederström V, Herbertsson L, et al. Seed
448 coating with a neonicotinoid insecticide negatively affects wild bees. *Nature.* 7 mai
449 2015;521(7550):77-80.
- 450 48. Crall JD, de Bivort BL, Dey B, Ford Versypt AN. Social Buffering of Pesticides in Bumblebees:
451 Agent-Based Modeling of the Effects of Colony Size and Neonicotinoid Exposure on Behavior
452 Within Nests. *Front Ecol Evol* [Internet]. 4 mars 2019 [cité 15 mai 2019];7. Disponible sur:
453 <https://www.frontiersin.org/article/10.3389/fevo.2019.00051/full>
- 454 49. Modlmeier AP, Foitzik S, Scharf I. Starvation endurance in the ant *Temnothorax nylanderi*
455 depends on group size, body size and access to larvae. *Physiol Entomol.* 2013;38(1):89-94.
- 456 50. Scharf I, Modlmeier AP, Beros S, Foitzik S. Ant Societies Buffer Individual-Level Effects of
457 Parasite Infections. *Am Nat.* 1 nov 2012;180(5):671-83.
- 458 51. Lessard J-P, Buddle CM. The effects of urbanization on ant assemblages (Hymenoptera:
459 Formicidae) associated with the Molson Nature Reserve, Quebec. *Can Entomol. avr*
460 2005;137(2):215-25.
- 461 52. Thompson B, McLachlan S. The effects of urbanization on ant communities and myrmecochory
462 in Manitoba, Canada. *Urban Ecosyst.* 1 mars 2007;10(1):43-52.
- 463 53. McIntyre NE. Ecology of Urban Arthropods: a Review and a Call to Action. *Ann Entomol Soc*
464 *Am.* 1 juill 2000;93(4):825-35.
- 465 54. Buczkowski G. Extreme life history plasticity and the evolution of invasive characteristics in a
466 native ant. *Biol Invasions.* 1 sept 2010;12(9):3343-9.
- 467 55. Grześ IM. Ants and heavy metal pollution – A review. *Eur J Soil Biol.* 1 nov 2010;46(6):350-5.
- 468 56. Molet M, Péronnet R, Couette S, Canovas C, Doums C. Effect of temperature and social
469 environment on worker size in the ant *Temnothorax nylanderi*. *J Therm Biol.* 1 juill
470 2017;67:22-9.

- 471 57. Abdel-Lateif HM, Donker MH, Straalen NMV. Interaction between temperature and cadmium
472 toxicity in the isopod *Porcellio scaber*. *Funct Ecol*. 1998;12(4):521-7.
- 473 58. Abràmoff MD, Magalhães PJ, Ram SJ. Image processing with ImageJ. *Biophotonics Int*.
474 2004;11(7):36-42.
- 475 59. Pinheiro J, Bates D, DebRoy S, Sarkar D, Team RC. *nlme: Linear and Nonlinear Mixed Effects*
476 *Models*, 2015. R Package Version. 2016;3:103.
- 477 60. Zuur A, Ieno EN, Walker N, Saveliev AA, Smith GM. *Mixed effects models and extensions in*
478 *ecology with R*. Springer Science & Business Media; 2009.
- 479 61. Malakar C, Ganguly A, Haldar P. Influence of cadmium on growth, survival and clutch size of a
480 common Indian short horned grasshopper, *Oxya fuscovittata*. *Am J Toxicol Sci*. 2009;1:32-6.
- 481 62. Mirčić D, Janković-Tomanić M, Nenadović V, Franeta F, Lazarević J. The effects of cadmium
482 on the life history traits of *Lymantria dispar* L. *Arch Biol Sci*. 2010;62(4):1013-20.
- 483 63. Roelofs D, Janssens TKS, Timmermans MJTN, Nota B, Mariën J, Bochdanovits Z, et al.
484 Adaptive differences in gene expression associated with heavy metal tolerance in the soil
485 arthropod *Orchesella cincta*. *Mol Ecol*. 2009;18(15):3227-39.
- 486 64. Diamond SE, Chick L, Perez A, Strickler SA, Martin RA. Rapid evolution of ant thermal
487 tolerance across an urban-rural temperature cline. *Biol J Linn Soc*. 1 juin 2017;121(2):248-57.
- 488 65. Andrewartha HG. Diapause in Relation to the Ecology of Insects. *Biol Rev*. 1952;27(1):50-107.
- 489 66. Kipyatkov VE, Lopatina EB. Temperature and photoperiodic control of diapause induction in the
490 ant *Lepisiota semenovi* (Hymenoptera, formicidae) from Turkmenistan. *J Evol Biochem Physiol*.
491 1 avr 2009;45(2):238-45.
- 492 67. Morgan JE, Morgan AJ. Seasonal changes in the tissue-metal (Cd, Zn and Pb) concentrations in
493 two ecophysiologically dissimilar earthworm species: pollution-monitoring implications.
494 *Environ Pollut*. 1 janv 1993;82(1):1-7.
- 495 68. Miller DL, Roth MB. *C. Elegans* Are Protected from Lethal Hypoxia by an Embryonic
496 Diapause. *Curr Biol*. 28 juill 2009;19(14):1233-7.
- 497 69. Kukul O, Denlinger DL, Lee RE. Developmental and metabolic changes induced by anoxia in
498 diapausing and non-diapausing flesh fly pupae. *J Comp Physiol B*. 1 nov 1991;160(6):683-9.
- 499 70. Maroni G, Wise J, Young JE, Otto E. Metallothionein Gene Duplications and Metal Tolerance in
500 Natural Populations of *Drosophila melanogaster*. *Genetics*. 1 déc 1987;117(4):739-44.
- 501 71. Roelofs D, Overhein L, Boer ME de, Janssens TKS, Straalen NM van. Additive genetic variation
502 of transcriptional regulation: metallothionein expression in the soil insect *Orchesella cincta*.
503 *Heredity*. janv 2006;96(1):85.
- 504 72. Thévenod F. Cadmium and cellular signaling cascades: To be or not to be? *Toxicol Appl*
505 *Pharmacol*. 1 août 2009;238(3):221-39.
- 506 73. Schrader L, Kim JW, Ence D, Zimin A, Klein A, Wyschetzki K, et al. Transposable element
507 islands facilitate adaptation to novel environments in an invasive species. *Nat Commun*. 16 déc
508 2014;5:5495.

- 509 74. Mirouze M, Paszkowski J. Epigenetic contribution to stress adaptation in plants. *Curr Opin Plant*
510 *Biol.* 1 juin 2011;14(3):267-74.
- 511 75. Klatt Björn K., Holzschuh Andrea, Westphal Catrin, Clough Yann, Smit Inga, Pawelzik Elke, et
512 al. Bee pollination improves crop quality, shelf life and commercial value. *Proc R Soc B Biol*
513 *Sci.* 22 janv 2014;281(1775):20132440.
- 514 76. Porter SD, Tschinkel WR. Fire Ant Polymorphism (Hymenoptera: Formicidae): Factors
515 Affecting Worker Size. *Ann Entomol Soc Am.* 1 mai 1985;78(3):381-6.
- 516 77. Chapman RE, Bourke AFG. The influence of sociality on the conservation biology of social
517 insects. *Ecol Lett.* 2001;4(6):650-62.
- 518 78. Luque GM, Giraud T, Courchamp F. Allee effects in ants. *J Anim Ecol.* 2013;82(5):956-65.
519