



HAL
open science

Post-translational regulation and conformational plasticity of PTEN

Larissa Kotelevets, Barbara Trifault, Eric Chastre, Mark Scott

► **To cite this version:**

Larissa Kotelevets, Barbara Trifault, Eric Chastre, Mark Scott. Post-translational regulation and conformational plasticity of PTEN. Cold Spring Harbor Perspectives in Medicine, 2020, 10 (11). hal-03020718

HAL Id: hal-03020718

<https://hal.sorbonne-universite.fr/hal-03020718v1>

Submitted on 24 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Post-translational regulation and conformational plasticity of PTEN

Larissa Kotelevets^{1,2}, Barbara Trifault^{3,4,5}, Eric Chastre^{1,2¶} and Mark G.H. Scott^{3,4,5¶}

¹Institut National de la Santé et de la Recherche Médicale, UMR S 938, Centre de Recherche Saint-Antoine, Paris, France;

²Sorbonne Université, UFR de Médecine, Site Saint-Antoine, Paris, France

³Université de Paris, Paris, France;

⁴Institut Cochin, Inserm, U1016, Paris, France;

⁵CNRS, UMR8104, Paris, France;

¶Corresponding authors:

mark.scott@inserm.fr

Address:

Institut Cochin, INSERM U1016, CNRS UMR8104, Université de Paris

27, Rue du Faubourg Saint Jacques, 75014 Paris, France

eric.chastre@inserm.fr

Address:

Centre de Recherche Saint-Antoine, Inserm UMR S 938, Hôpital Saint-Antoine,

Bâtiment Kourilsky, 34 rue Crozatier, 75012 Paris, France

Short title: Post-translational regulation of PTEN

Abstract

PTEN is a tumour suppressor that is frequently downmodulated in human cancer. PTEN inhibits the PI3K/AKT pathway through its lipid phosphatase activity. Multiple PI3K/AKT-independent actions of PTEN, protein-phosphatase activities and functions within the nucleus have also been described. PTEN therefore regulates many cellular processes including cell proliferation, survival, genomic integrity, polarity, migration and invasion. Even a modest decrease in the functional dose of PTEN may promote cancer development. Understanding the molecular and cellular mechanisms that regulate PTEN protein levels and function, and how these may go awry in cancer contexts, is therefore key to fully understanding the role of PTEN in tumourigenesis. Here, we discuss current knowledge on post-translational control and conformational plasticity of PTEN, as well as therapeutic possibilities towards re-establishment of PTEN tumour suppressor activity in cancer.

Introduction

PTEN (phosphatase and tensin homolog deleted on chromosome ten) is a ubiquitously expressed tumour suppressor that is frequently deregulated in human cancers. Three different groups initially identified the *PTEN* gene in 1997 (Li and Sun 1997; Li et al. 1997; Steck et al. 1997). The gene is located on chromosome 10q23, a region that displays loss of heterozygosity (LOH) in several cancers (Teng et al. 1997). Following on from this, *PTEN* mutations were documented in the germline of patients with a group of rare autosomal dominant syndromes, collectively termed as the PTEN hamartoma tumour syndromes (PHTS) (Yehia et al. 2019). These disorders are characterized by the presence of multiple hamartomas and increased cancer predisposition, therefore providing direct evidence for the deregulation of PTEN in cancer. *In vivo* studies in mice have also analysed the importance of Pten in cancer development. Whereas homozygous deletion of *Pten* provokes early embryonic death, *Pten*^{+/-} mice develop various cancers demonstrating that Pten is haploinsufficient for suppressing cancer (Di Cristofano et al. 1998; Stambolic et al. 1998; Suzuki et al. 1998; Podsypanina et al. 1999). Furthermore, studies using a series of mice engineered to carry different combinations of wild-type, null and hypomorphic *Pten* alleles, have shown that even subtle reductions in Pten levels are sufficient to promote cancer susceptibility (Trotman et al. 2003; Alimonti et al. 2010). These findings suggest a “continuum model” of PTEN tumour suppression, whereby modest alterations in the functional dose of PTEN may promote tumourigenesis without mutation or loss of even one allele. In light of these studies, understanding the molecular and cellular mechanisms that regulate PTEN protein levels and function, and how these may go awry in cancer settings, is key to fully understanding the role of PTEN in tumourigenesis.

The PTEN protein comprises 403 amino acids (Fig. 1a). The crystal structure of human PTEN, using PTEN truncated at its termini and an internal loop, revealed the existence of an N-terminal phosphatase domain (amino acids 7-185) and a C-terminal C2 domain involved in membrane binding (amino acids 186-351) (Lee et al. 1999). PTEN also contains a short N-terminal PIP2-binding motif (amino acids 6-15) involved in membrane targeting, and a regulatory C-terminal tail (amino acids 352-403) that contains two phosphorylation site clusters and a PDZ binding motif. In addition to this 403 amino acid form of PTEN, several N-terminally extended isoforms expressed at lower levels, translated from upstream alternative start codons, have recently been

described: PTEN-O, PTEN-N, PTEN-M/PTEN- β and PTEN-L/PTEN- α (Malaney et al. 2017). The N-terminal extensions affect their subcellular localizations and the first discovered isoform PTEN-L, with a 173 amino acid N-terminal extension including a signal peptide, is secreted and enters into neighbouring cells (Hopkins et al. 2013).

PTEN is a phosphatase that can dephosphorylate both protein and lipid substrates. It can dephosphorylate protein substrates on serine/threonine and tyrosine residues. The protein phosphatase activity of PTEN is however weak, and determining what the physiological protein substrates of PTEN are remains an important area of investigation. Several potential protein substrates of PTEN have been proposed and include FAK (Gu et al. 1998), IRS1 (Shi et al. 2014) and Dvl2 (Shnitsar et al. 2015). PTEN can also autodephosphorylate threonine residues contained within its regulatory carboxy-terminal region (Raftopoulou et al. 2004; Tibarewal et al. 2012). The lipid phosphatase function of PTEN serves to negatively regulate the proliferative PI3K/AKT pathway by dephosphorylating the lipid signalling intermediate PIP3 that is generated by PI3Ks (Maehama and Dixon 1999) (Fig. 1b). PIP3 is considered the major physiological substrate for PTEN. PIP3 serves to recruit proteins, such as AKT, containing pleckstrin homology domains, to the membrane. By counterbalancing the PI3K/AKT pathway the lipid phosphatase activity of PTEN therefore serves to limit cell proliferation, survival, migration and invasiveness (Stambolic et al. 1998; Chalhoub and Baker 2009). When PTEN function is impaired or lost, PIP3 levels are not properly regulated leading to hyperactivation of the PI3K/AKT pathway, cell transformation and tumourigenesis (Suzuki et al. 1998; Podsypanina et al. 1999; Di Cristofano et al. 1998; Leslie and Downes 2004).

A number of phosphatase-independent biological actions of PTEN have also been described in the cytoplasm and nucleus that rely on protein-protein interactions. For example, PTEN directly binds tumour suppressor p53 in the nucleus, increasing its stability and transcriptional activity (Freeman et al. 2003). Nuclear PTEN also interacts with the APC/C (anaphase-promoting complex) and promotes its association with CDH1 (CDC20 homolog 1) to enhance the tumour-suppressive activity of the APC-CDH1 complex in a phosphatase-independent fashion (Song et al. 2011). The isolated membrane-targeting PTEN C2 domain is also able to mimic effects of full-length PTEN on inhibition of glioma cell migration (Raftopoulou et al. 2004; Lima-Fernandes et al. 2011) and glandular morphogenesis in 3D colorectal cancer cell systems (Javadi et al. 2017). Mechanisms controlling PTEN subcellular

compartmentalization are therefore key in spatially controlling both phosphatase-dependent and -independent functions.

PTEN levels and function are subject to tight regulation at the transcriptional, post-transcriptional and post-translational levels (Salmena et al. 2008; Leslie and Foti 2011; Song et al. 2012; Lee et al. 2018) (Fig. 2). Initially believed to be a constitutively active phosphatase, pleiotropic post-translational modifications and protein-protein interactions are now known to calibrate PTEN function (Keniry and Parsons 2008; Salmena et al. 2008; Wang and Jiang 2008; Leslie and Foti 2011; Song et al. 2012; Kotelevets et al. 2018; Lee et al. 2018). These post-translational mechanisms control PTEN conformation, phosphatase activity and association with protein complexes, drive subcellular compartmentalization and impact protein stability. Studying how these mechanisms contribute to the loss of PTEN function in cancer is therefore essential. With this knowledge in hand it may be possible to develop strategies with the goal of restoring or enhancing PTEN tumour suppressor activity in cancer contexts. Here, we discuss post-translational control and conformational plasticity of PTEN, and implications in human cancer.

PTEN Regulation

PTEN expression and function are tightly regulated at the transcriptional, post-transcriptional and post-translational levels (Salmena et al. 2008; Leslie and Foti 2011; Song et al. 2012; Lee et al. 2018) (Fig. 2). In addition to genetic loss or somatic mutations in human cancers, disorder in these regulatory molecular mechanisms controlling PTEN expression and function can lead to a spectrum of different loss-of-function PTEN protein categories and/or levels, which contribute to tumorigenesis in diverse manners (Lee et al. 2018). Epigenetic and transcriptional silencing as well as dysregulation of miRNAs and competitive endogenous RNA (ceRNA) systems have all been shown to suppress PTEN expression (Fig. 2). Regulation of PTEN at the post-translational level via dynamic post-translational modifications (PTMs) and protein-protein interactions calibrates PTEN function. Dysregulation of these non-genomic mechanisms can again lead to repression of PTEN function (Leslie and Foti 2011). Mechanisms of post-translational control of PTEN are discussed below.

Post-translational modifications of PTEN

PTEN function is calibrated by multiple PTMs summarized in Fig. 3a and Table 1. Aberrant regulation of mechanisms controlling these PTMs can lead to inhibited PTEN function. This point of control on PTEN also provides therapeutic avenues to explore with the aim of re-establishing or enhancing PTEN function.

Phosphorylation

A key PTM regulating PTEN function is phosphorylation. Phosphorylation controls PTEN conformation, activity, stability and subcellular targeting. PTEN is phosphorylated on serine/threonine clusters contained in its regulatory C-terminal tail (Fig. 3a: Ser362, Thr366, Ser370, Ser380, Thr382, Thr383 and Ser385). Ser380, Thr382, Thr383 and Ser385 are phosphorylated to high stoichiometry by casein kinase 2 (CK2) (Torres and Pulido 2001). CK2 also phosphorylates Ser370 and this promotes subsequent phosphorylation of Thr366 and Ser362 by glycogen synthase kinase 3 (GSK3) (Al-Khoury et al. 2005; Maccario et al. 2007). Phosphorylation of the Ser380, Thr382, Thr383, Ser385 cluster promotes a "closed", less active but stable form of PTEN that has increased conformational compaction, reduced interaction with membrane-anchored PDZ-domain-containing proteins, such as MAGI-2 and MAGI-3, and decreased plasma membrane targeting (Vazquez et al. 2000; Torres and Pulido

2001; Vazquez et al. 2001; Das et al. 2003; Bolduc et al. 2013) (Fig. 3b). This "closed" conformation of PTEN is more stable due to it displaying reduced access to E3 ubiquitin ligases and therefore being less targeted for proteasomal-mediated degradation (Maccario et al. 2010). An intramolecular interaction of the phosphorylated C-terminal tail with basic residues located in the N-terminal PIP2-binding motif, the catalytic and C2 domains, maintains PTEN in its "closed" form, blocking the active site (Odriozola et al. 2007; Rahdar et al. 2009). Mutation of these C-terminal phosphorylation sites disrupts the intramolecular interaction leading to an "open" conformation of PTEN, which displays enhanced plasma membrane targeting and increased activity (Fig. 3b). In human T cell acute lymphoblastic leukaemia (T-ALL) cells PTEN is inactivated by hyperphosphorylation on C-terminal phosphorylation sites resulting from CK2 overexpression and hyperactivation (Silva et al. 2008). This leads to decreased PTEN lipid phosphatase activity and constitutive hyperactivation of the PI3K/AKT pathway. CK2 inhibition using selective CK2 inhibitors, re-established PTEN function to impair PI3K/AKT signalling in T-ALL cells.

Another mechanism leading to enhanced PTEN C-terminal tail phosphorylation, is via loss of protein phosphatase 2A (PP2A)-mediated dephosphorylation of PTEN (Nakahata et al. 2014). The adaptor protein N-myc downstream-regulated gene (NDGR2) is a PTEN-binding protein that recruits PP2A to PTEN (Nakahata et al. 2014). NDGR2 is frequently downregulated in T-ALL, leading to enhanced PTEN phosphorylation and increased PI3K/AKT signalling. PTEN is also thought to be capable of auto-dephosphorylation of Thr366 and Thr383 with implications for glioma cell migration/invasion (Raftopoulou et al. 2004; Tibarewal et al. 2012). Phosphorylation of PTEN in the C-terminal tail region can also lead to changes in cytonuclear distribution of PTEN. For example, phosphorylation of PTEN at Thr398 by the kinase ataxia telangiectasia mutated (ATM) promotes its nuclear export and sensitizes cells to DNA damage (Bassi et al. 2013). Polo-like kinase 1 (PLK1), which regulates cell cycle-related processes, phosphorylates PTEN on Ser380 enhances PTEN accumulation on chromatin, and normal mitotic progression (Choi et al. 2014). A phospho-deficient PTEN Ser380Ala mutant promotes enhanced mitotic exit. Interestingly, PTEN dephosphorylates and destabilizes PLK1 (Zhang et al. 2016). Outwith the regulatory C-terminal tail of PTEN, Ser229/Thr232 and Thr319/Thr321 in the C2 domain are phosphorylated by Rho-associated kinase 1 (ROCK1), promoting enhanced lipid phosphatase activity, polarized distribution of PTEN in neutrophils and

chemotaxis (Li et al. 2005). Genotoxic agents activate ATM, which besides Thr398, can also phosphorylate PTEN at Ser113 to trigger nuclear translocation and autophagy (Chen et al. 2015).

PTEN is also phosphorylated by tyrosine kinases. PTEN function is inhibited by phosphorylation on tyrosine residues by Src family kinases (Koul et al. 2002; Lu et al. 2003; Fenton et al. 2012). The oncoprotein Src phosphorylates PTEN in the C2 domain at Tyr240 and Tyr315, leading to decreased stability and also likely decreased PTEN binding to cellular membranes. Interestingly, trastuzumab, used in the treatment of ERBB2-overexpressing breast cancers, induces PTEN membrane targeting and phosphatase activity by reducing PTEN tyrosine phosphorylation via Src inhibition (Nagata et al. 2004). FGFR2 and FGFR3, but not EGFR or PDGFR, also phosphorylate PTEN on Tyr240. Tyr240 phosphorylation is associated with resistance to EGFR inhibitors in patients with glioblastoma multiforme and a shortened overall survival (Fenton et al. 2012). This process might be independent of PTEN phosphatase activity but related to its targeting to specific subcellular compartments. Accordingly, in response to DNA damage, pTyr240-PTEN binds to chromatin through interaction with Ki-67 and facilitates the recruitment of RAD51 to promote DNA repair and thus resistance to ionizing radiation therapy (Ma et al. 2019). Tyr336 in the PTEN C2 domain is phosphorylated by FAK (Tzenaki et al. 2015) and Rak (Yim et al. 2009), leading to increased phosphatase activity and protein stability of PTEN. Increased PTEN stability in the presence of Rak is due to decreased binding of the E3 ubiquitin ligase NEDD4-1 to PTEN. In terms of tyrosine phosphatases, SHP-1 has been shown to dephosphorylate PTEN in Src transfected cells, restoring PTEN stability (Lu et al. 2003).

Ubiquitination

Several ubiquitin E3 ligases are implicated in the ubiquitination of PTEN, which has implications for PTEN stability and subcellular localization. Polyubiquitination leads to PTEN degradation via the ubiquitin-proteasome system, whereas monoubiquitination on Lys13 and Ly289 promotes its nuclear import. The HECT-domain ubiquitin E3 ligase NEDD4-1 was the first ubiquitin E3 ligase identified for PTEN (Wang et al. 2007). NEDD4-1 is implicated in both polyubiquitination and monoubiquitination of PTEN (Trotman et al. 2007; Wang et al. 2007). Depletion of Nedd4-1 in mice, however, found no change in Pten protein stability *in vivo*, suggesting

that there is some redundancy and other E3 ubiquitin ligases are likely also implicated in the regulation of PTEN proteosomal degradation (Fouladkou et al. 2008). In line with this, several other E3 ubiquitin ligases have been implicated in PTEN ubiquitination and increased PTEN turnover. These include, WWP2 the NEDD4-like protein family E3 ligase (Maddika et al. 2011), the RING-domain X-linked inhibitor of apoptosis protein (XIAP) (Van Themsche et al. 2009), the chaperone-assisted E3 ligase C-terminus of Hsc70-interacting protein (CHIP) (Ahmed et al. 2012), and RING finger protein 146 (RNF146) (Li et al. 2015). Non-degradative polyubiquitination by RFP, negatively regulates PTEN activity, without affecting PTEN stability (Lee et al. 2013). Very interestingly, non-degradative Lys27-linked polyubiquitination of PTEN by the E3 ubiquitin ligase WWP1 was recently shown to suppress PTEN oligomerization (see protein partners), plasma membrane recruitment, and tumour-suppressive functions both *in vitro* and *in vivo* (Lee et al. 2019). Genetic deletion or pharmacological inhibition of WWP1 provoked PTEN reactivation and tumour suppressive activity. This study therefore indicates a potential therapeutic strategy to reactivate PTEN for cancer treatment.

Several deubiquitinases (DUBs) have also been identified for PTEN. These include ubiquitin-specific proteases-10 (Sun et al. 2018), -11 (Park et al. 2019) and -13 (Zhang et al. 2013) (USP-10, USP-11 and USP-13), as well as OTU-deubiquitinase (OTUD3) (Yuan et al. 2015) that are all implicated in PTEN deubiquitination and stabilization. In addition, ubiquitin-specific protease-7 (USP7 or HAUSP) is involved in the reversal of PTEN monoubiquitination, leading to nuclear exclusion of PTEN (Song et al. 2008).

SUMOylation

Small ubiquitin-like modifier (SUMO) protein can be conjugated to Lys254 and Lys266 in the C2 domain of PTEN (Huang et al. 2012; Bassi et al. 2013; Wang et al. 2014). SUMOylation on Lys266 in the CBR3 loop of PTEN promotes plasma membrane binding via electrostatic interactions (Huang et al. 2012). This enhanced membrane targeting of SUMOylated PTEN leads to decreased PI3K/AKT signalling, suppression of anchorage-independent cell growth and tumour growth *in vivo*. PTEN SUMOylation on Lys254, enhances its nuclear retention (Bassi et al. 2013). This promotes the nuclear function of PTEN in DNA repair mechanisms. Following genotoxic stress, SUMO-PTEN is rapidly excluded from the nucleus in an ATM kinase-

dependent manner (phosphorylation of Thr398 in the C-terminal tail of PTEN) (Bassi et al. 2013). Cells lacking nuclear PTEN were found to be hypersensitive to DNA damage. Finally, cross-talk between SUMOylation and ubiquitination exists on PTEN. PTEN SUMOylation is increased by the SUMO E3 ligase PIASx α , resulting in decreased polyubiquitination, increased PTEN stability, leading to inhibition of the PI3K/AKT pathway, inhibition of cell proliferation and tumour suppression (Wang et al. 2014).

Acetylation

Following growth factor receptor stimulation, the histone acetylase p300/CBP-associated factor (PCAF) acetylates PTEN on Lys125 and Lys128, which are both located within the catalytic pocket (Okumura et al. 2006). Acetylation on these residues negatively regulates PTEN lipid phosphatase activity leading to enhanced AKT signalling. PTEN can also be acetylated by p300-CREB-binding protein (CBP) on Lys402, located in its C-terminal PDZ domain-binding motif (Ikenoue et al. 2008). Acetylation here does not impact catalytic activity but instead affects PTEN interaction with PDZ-domain containing proteins. The deacetylase sirtuin 1 (SIRT1) is mainly responsible for PTEN deacetylation (Ikenoue et al. 2008). In Sirt1-knockout cells, PTEN is hyperacetylated and excluded from the nucleus, indicating that acetylation modulates PTEN subcellular localization. It was also recently demonstrated that PTEN acetylation at Lys163, following inhibition of Histone Deacetylase 6 (HDAC6), switched PTEN into an "open" conformation that was targeted to the plasma membrane (Meng et al. 2016). This resulted in the inhibition of cell proliferation, migration and invasion, as well as xenograft tumour growth. This suggests that HDAC inhibitors may be clinically relevant in tumours expressing wild-type PTEN.

Oxidation

As with other protein tyrosine phosphatases PTEN contains a catalytic site cysteine nucleophile that is susceptible to oxidation. Reactive oxygen species (ROS) promote the oxidation of Cys124, which leads to the formation of an intramolecular disulphide bond with Cys71, suppressing PTEN phosphatase activity (Lee et al. 2002). The oxidation of PTEN is reversible. Endogenous ROS produced in activated cells promotes transient oxidation of PTEN leading to inactivation of a fraction of cellular PTEN, an increase in PIP3 levels, and activation of AKT (Leslie et al. 2003; Kwon et

al. 2004). This suggests localized control of a subcellular pool of PTEN by oxidation. Thioredoxin-interacting protein (TXNIP) regulates reversible PTEN oxidation by maintaining sufficient thioredoxin NADPH activity to reductively reactivate oxidized PTEN and counterbalance PI3K/AKT signalling (Hui et al. 2008). Oxidation of PTEN-binding partners also influences PTEN activity. For example, the oncoprotein DJ-1 binds PTEN and inhibits its catalytic activity (Kim et al. 2009). Oxidation of DJ-1 increases its affinity for binding PTEN, leading to a more profound decrease in PTEN activity. In contrast two other PTEN-binding partners peroxidase peroxiredoxin 1 (Prdx1) (Cao et al. 2009) and apoptosis-inducing factor (AIF) (Shen et al. 2015) protect PTEN from oxidation-induced inactivation via direct interaction.

S-Nitrosylation

Low concentrations of nitric oxide (NO) lead to S-nitrosylation of PTEN (SNO-PTEN) at Cys83 (Numajiri et al. 2011). This event results in inhibition of PTEN catalytic activity and enhanced AKT signalling, demonstrating that Cys83 is an important site for redox regulation of PTEN. In addition to the effect on lipid phosphatase activity, S-nitrosylation of PTEN also promotes PTEN protein degradation via the ubiquitin-proteasome system through NEDD4-1-mediated ubiquitination (Kwak et al. 2010). Depletion of the E3 ubiquitin ligase Parkin contributes to AMPK-mediated activation of endothelial nitric oxide synthase (eNOS), increased levels of reactive oxygen species, and a concomitant increase in oxidized NO levels (Gupta et al. 2017). This promotes S-nitrosylation of PTEN and its ubiquitination. Together, these studies highlight the influence of PTEN S-nitrosylation in supporting AKT signalling, cell survival and proliferation.

Methylation

PTEN is methylated on Lys313 by the oncogenic protein methyltransferase SMYD2, which has been proposed to lead to negative regulation of PTEN activity and increased PI3K/AKT signalling (Nakakido et al. 2015). DNA double-strand breaks (DSBs) also increase NSD2-mediated di-methylation of PTEN at Lys349, which is recognized by the tudor domain of 53BP to recruit PTEN to DNA damage sites, allowing for efficient repair of DSBs (Zhang et al. 2019). Interestingly, inhibiting NSD2-mediated PTEN methylation sensitizes cancer cells to combination treatment with PI3K inhibitor and DNA-damaging agents.

Ribosylation

PTEN is also a substrate for tankyrases: tankyrase1 (TNKS1) and tankyrase2 (TNKS2) ribosylate PTEN at Glu40/Glu150 in the phosphatase domain and Asp326 in the C2 domain (Li et al. 2015). PTEN ribosylation promotes the recognition of PTEN by the E3 ubiquitin ligase, RNF146, leading to subsequent PTEN ubiquitination and degradation. TNKS levels were found to negatively correlate with PTEN levels in human colon carcinomas. Simultaneous knockdown of TNKS1/2 in colorectal cancer cells resulted in inhibition of tumour growth in PTEN-expressing but not PTEN-depleted cells, indicating that targeting TNKS may only be effective in wild-type PTEN contexts. Taken together, these data support the idea to explore the development of tankyrase inhibitors to restore PTEN function.

Protein-protein interactions

PTEN interacts with a number of protein partners that influence its subcellular localization, activity and stability. These PTEN partners include integral plasma membrane proteins, adaptor proteins, transport proteins, and proteins that influence PTEN PTMs. PTEN monomers can also oligomerize, a requirement for full PTEN activation at the plasma membrane. A list of some of these PTEN partners, where they bind on PTEN, and their effects on PTEN function are shown in [Table 2](#), and discussed below.

Negative regulators of PTEN function include PIP3-dependent Rac exchanger protein (PREX2) (Fine et al. 2009; Hodakoski et al. 2014), shank-interacting protein-like 1 (SIPL1) (He et al. 2010; De Melo et al. 2014), α -mannosidase 2C1 (MAN2C1) (He et al. 2011), DJ1 (Kim et al. 2009; Das et al. 2011), BMI (Fan et al. 2009), and Protein Phosphatase 1 Regulatory subunit 10 (PNUTS) (Kavela et al. 2013). These negative regulators all inhibit PTEN function through diverse mechanisms leading to enhanced PI3K/AKT signalling in cells, and therefore potentially represent therapeutic targets.

A number of partners affect PTEN stability by protecting PTEN from ubiquitination by E3 ligases. For example, both the regulatory subunit of PI3K, p85 α , in a homodimeric form (Cheung et al. 2015), and the G protein-coupled receptor (GPCR) Melanocortin receptor 1 (MC1R) (Cao et al. 2013) protect PTEN from proteasomal degradation by E3 ubiquitin ligase WWP2. Interestingly, cancer-

associated mutants of MC1R or p85 α lead to decreased PTEN binding and increased PI3K-AKT signalling. The Rak tyrosine kinase ([Table 1](#)) interacts and phosphorylates PTEN on Tyr336, and this decreases binding of PTEN to the E3 ubiquitin ligase NEDD4-1 (Yim et al. 2009). Discs Large MAGUK Scaffold 1 (DLG1) a PDZ-domain containing scaffold binds the PDZ-binding motif of PTEN to promote PTEN stability and decreased PI3K/AKT signalling (Adey et al. 2000; Valiente et al. 2005). DLG1 also interacts with APC and therefore may serve as a platform to bring PTEN and APC in proximity, providing a functional network of tumour suppressor activities (Sotelo et al. 2012).

Protein interaction partners of PTEN can also regulate its subcellular localization. Plasma membrane targeting of PTEN by these partners localizes PTEN to membrane-localized PIP3, resulting in decreased PI3K/AKT signalling. A number of PDZ-domain containing proteins bind the PDZ-binding motif of PTEN and target it to the plasma membrane/junctional complexes. These include MAGI-1, MAGI-2, MAGI-3, MAST1, MAST2, MAST3, PAR-3 and NHERF1, with functional consequences for PTEN recruitment into high molecular weight complexes, inhibition of PI3K/AKT signalling, regulation of epithelial cell polarity, stabilization of adherens junctions and suppression of invasiveness (Wu et al. 2000; Tolkacheva et al. 2001; Kotelevets et al. 2005; Valiente et al. 2005; Takahashi et al. 2006; Wu et al. 2007; Chastre et al. 2009). Neural endopeptidase (NEP) (Sumitomo et al. 2004), the GPCR Sphingosine-1-Phosphate Receptor 2 (S1PR2) (Sanchez et al. 2005), the Cystic Fibrosis Transmembrane Conductance Regulator (CFTR) (Riquelme et al. 2017), and the motor protein Myosin V (van Diepen et al. 2009) also all target PTEN to the plasma membrane towards membrane-localized PIP3. The RhoA effector, RhoA associated protein kinase (ROCK), interacts with and phosphorylates PTEN ([Table 1](#)), promoting its membrane recruitment and activation (Li et al. 2005). ROCK activation also promotes the association of the multifunctional adaptor proteins β -arrestin1 (β -arr1) and β -arrestin2 (β -arr2) with PTEN (Lima-Fernandes et al. 2011). β -arrestins recruit PTEN to the plasma membrane following GPCR/Rho/ROCK stimulation and enhance its lipid phosphatase activity. However, during glioma cell migration β -arrestins bind the C2 domain of PTEN to inhibit its lipid-phosphatase-independent anti-migratory function. β -arr1 also binds the PTEN C2 domain as part of a plasma membrane-associated regulatory complex incorporating the Cdc42 GTPase-activating protein ARHGAP21 and Cdc42

(Javadi et al. 2017). This complex regulates Cdc42-dependent mitotic spindle formation and lumen formation in 3D cultures of colorectal cancer cells. Disruption of this protein network provokes mitotic spindle misorientation and abnormal multilumen that are evocative of colorectal cancer.

PTEN also has function in the nucleus, including enhancing both tumour suppressor p53 function (Freeman et al. 2003) and the tumour-suppressive activity of the APC-CDH1 complex in a phosphatase-independent fashion (Song et al. 2011). Nuclear translocation of PTEN occurs through various mechanisms including interaction with the Major Vault Protein (Chung et al. 2005; Minaguchi et al. 2006) and the nuclear transport receptor Importin-11 (Chen et al. 2017). Importin-11 protects PTEN from cytoplasmic degradation by NEDD4-1, and there is correlative loss of both Importin-11 and PTEN in lung tumours.

Finally, PTEN monomers can oligomerize at the plasma membrane, adopting an active conformation with full lipid phosphatase activity against PIP3 (Papa et al. 2014). Catalytically inactive cancer-associated PTEN mutants can heterodimerize with wild-type PTEN and act in a dominant-negative fashion to inhibit wild-type catalytic activity. A subsequent study demonstrated that PTEN can form homodimers *in vitro* and furnished a possible structural model for the complex (Heinrich et al. 2015). The C-terminal tail of PTEN functions to stabilize the homodimer and C-tail phosphorylation interferes with this stabilization.

Probing PTEN conformation

The combination of PTEN post-translational modifications and protein partner interactions occurring at a given moment in time will give rise to a particular conformation of PTEN. To rapidly probe changes in PTEN conformation dynamics in live cells, we recently developed an intramolecular bioluminescence energy transfer (BRET)-based biosensor of PTEN (Lima-Fernandes et al. 2014). For this, PTEN is sandwiched between the energy donor Renilla luciferase (Rluc) and the energy acceptor yellow fluorescent protein (YFP) (Fig. 4a). The energy transfer between donor and acceptor can be measured in live cells following the addition of coelenterazine, the substrate for Rluc (Fig. 4a and b). In cells expressing the biosensor a strong basal BRET signal was detected indicating molecular proximity of the donor and acceptor molecules (Fig. 4c). As the biosensor is unimolecular, BRET signals remain constant over a wide range of Rluc concentrations (Lima-Fernandes et al. 2014; Misticone et al.

2016) (Fig. 4c). This is particularly useful in situations where, for example, a mutant of PTEN is less stable, as a change in BRET will only reflect a conformational change, allowing conformational readout comparisons of cancer-associated PTEN mutants that may be less stable than wild-type PTEN. BRET from the biosensor can also be detected at “endogenous expression” levels of PTEN (Lima-Fernandes et al. 2014).

The biosensor can be used in different modes to probe PTEN conformational change occurring at the cellular and molecular levels (Fig. 4d). In a structure-function mode we demonstrated that combined mutation of key C-terminal tail phosphorylation sites (Ser380, Thr382, Thr383 and Ser385) implicated in PTEN intramolecular interaction and control of PTEN plasma membrane targeting/cellular activity, results in a decreased BRET signal (Fig. 4e), compared to baseline BRET, indicative of conformational rearrangement (Lima-Fernandes et al. 2014). This change is consistent with the "opening" of the PTEN molecule upon dephosphorylation of its C-terminal tail with increased membrane targeting and indicates that PTEN conformational changes can be followed in live cells using the biosensor. These carboxy-terminal regulatory sites are phosphorylated by the serine/threonine kinase CK2. CK2 overexpression has been documented in several types of cancer, leading to phospho-dependent PTEN inhibition and enhanced PI3K/AKT pathway activation (Silva et al. 2008). CK2 inhibitors have been shown to re-establish PTEN signalling and restrain the PI3K pathway (Silva et al. 2008). Negative BRET shifts (Fig. 4e) in the biosensor were also obtained following pharmacological inhibition of CK2 using either TBB or CX-4945, which is in development for clinical use (Chon et al. 2015). The BRET shifts obtained using CX-4945 paralleled decreased phosphorylation of the regulatory C-terminal residues (Misticone et al. 2016). In contrast to the results obtained with the CK2 inhibitors, overexpression of CK2 provoked a significant increase above baseline BRET of the biosensor (Lima-Fernandes et al. 2014) (Fig. 4e). This indicates that both positive and negative modification of a signalling pathway impacting PTEN function can be detected using the biosensor. Use of the biosensor, in live cells, therefore validates the model of a PTEN switch between a "closed/less active" cytoplasmic state and an "open/active" conformation that is targeted to the plasma membrane. The significance of PTEN membrane localization is underscored by studies demonstrating that certain cancer-associated mutants display normal catalytic activity *in vitro*, but display defective plasma membrane targeting (Nguyen et al. 2015). This impaired

plasma membrane localization leads to a PTEN-null phenotype of these mutants in cells.

The biosensor can also be used to probe conformational change that occurs in cancer-associated mutants: Lys13 is mutated to Glu (K13E) in spontaneous cancer (Duerr et al. 1998) and Lys289 to Glu (K289E) is associated with Cowden syndrome (Trotman et al. 2007). These amino acids are both located on unstructured regions of PTEN and are major monoubiquitination sites that are important for PTEN nuclear import. Introduction of the K13E/K289E mutations, which renders PTEN defective in nuclear import, provoked a BRET change signifying conformational change. The biosensor can therefore detect conformational changes associated with changes in different post-translational modifications of PTEN (i.e. phosphorylation and ubiquitination).

PTEN-protein partner interactions and the effect of upstream intracellular signalling proteins on PTEN conformation can also be detected using the biosensor. Co-expression of the biosensor with active RhoA and the molecular scaffold β -arrestin2, both proteins known to activate PTEN, promoted changes in PTEN conformation (Lima-Fernandes et al. 2014; Misticone et al. 2016). Previously characterized or candidate PTEN protein partners that may form part of the PTEN interactome, as well as signalling proteins that may impinge on PTEN function can therefore be rapidly assessed for their effects on PTEN conformation in live cells by coexpression with the biosensor.

The biosensor can also be used to report real-time changes in PTEN conformational change occurring in kinetic experiments in live cells, following incubation with either physiological ligands or therapeutic agents that target cell surface receptors. We found that the lipid sphingosine 1-phosphate (S1P) changed the BRET signal of the biosensor in cells expressing the GPCR S1PR2, but not in cells expressing the S1PR1, in agreement with previous studies that S1PR2, but not S1PR1, activates PTEN (Sanchez et al. 2005). Using this kinetic approach we also identified several other GPCRs that elicited PTEN conformational rearrangement upon ligand incubation. The ERBB2-targeting antibody, trastuzumab, used in the treatment of ERBB2-overexpressing breast cancers, induces PTEN membrane targeting and lipid phosphatase activity by provoking a decrease in PTEN tyrosine phosphorylation via Src inhibition (Nagata et al. 2004). We showed that trastuzumab elicited a time-dependent change in BRET that coincided with enhanced PTEN lipid phosphatase

activity and decreased PI3K/AKT signalling. This series of experiments therefore provides proof of principle that the biosensor can detect real-time changes in PTEN conformation downstream of cell surface receptor targeting and therapeutic agents.

The different BRET signatures obtained using the biosensor indicate that PTEN can adopt a spectrum of different conformations, depending on changes in post-translational regulation mechanisms, and that the biosensor provides a means of identifying additional signalling contexts that influence PTEN function. As the biosensor can be used as a rapid, direct and sensitive reporter to detect potential changes in PTEN localization and activity, it could be used as readout for drug discovery screening of small molecule libraries to identify chemicals that enhance or restore PTEN-dependent pathways. The identification of such chemical agents would help pave the way towards reactivating PTEN tumour suppressor as a potential cancer treatment. Indeed, proof of principle supporting this concept comes from a recent study demonstrating that pharmacological inhibition of WWP1 provokes PTEN reactivation and tumour suppressive activity, uncovering a potential therapeutic strategy to reactivate PTEN for cancer treatment (Lee et al. 2019).

Concluding remarks

PTEN acts as a tumour suppressor via its lipid phosphatase activity to inhibit the PI3K/AKT oncogenic pathway, and also via PI3K/AKT-independent functions. Even a modest reduction in the functional dose of PTEN may promote tumourigenesis, indicating that re-establishing or enhancing PTEN function may provide a therapeutic strategy in a wide range of human cancers. Indeed, in favour of this, an enhanced engineered PTEN (ePTEN) has been developed that displays increased ability to inhibit PIP3 signalling (Nguyen et al. 2014), and pharmacological inhibition of WWP1, which suppresses PTEN function, triggers PTEN reactivation and tumour suppressive activity (Lee et al. 2019). In light of these studies, fully understanding the molecular and cellular mechanisms that regulate PTEN function, and how these go awry in cancer, is key to defining PTEN dysregulation in tumourigenesis, in addition to providing therapeutic avenues for exploration to restore or enhance PTEN function as anti-cancer therapy.

Acknowledgements

This work was supported by the CNRS and Inserm. Work in the group of MGHS is also supported by the Ligue Contre le Cancer (comité de L'Oise), La Fondation ARC, Université de Paris, and the Who am I? laboratory of excellence (grant ANR-11-LABX-0071) funded by the "Investments for the Future" program operated by The French National Research Agency (grant ANR-11-IDEX-0005-01).

Table 1: Partial list of proteins implicated in PTEN post translational modifications (PTMs) and functional consequences

Gene symbol	Full name	Activity, function	PTEN PTMs	Impact on PTEN	References
Kinases, phosphatases					
ATM	ATM Serine/Threonine Kinase, Ataxia Telangiectasia Mutated	Cell cycle checkpoint Ser/Thr kinase	Following DNA double-strand breaks ATM Phosphorylates PTEN at Ser113 and Thr398.	Phosphorylation of Ser113 triggers PTEN nuclear translocation and induction of autophagy via p-JUN-SESN2-AMPK pathway. Following DNA-damaging chemotherapeutic agents, SUMO conjugated PTEN is excluded from the nucleus in an ATM protein kinase manner (phosphorylation of Thr398)	Bassi et al. 2013; Chen et al. 2015
CSNK2, CK2	Casein Kinase 2	Ser/Thr protein kinase. Tetrameric holoenzyme : 2 catalytic subunits (alpha1, alpha2), 2 beta regulatory subunits	Two independent cascades of ordered phosphorylations target 1) Ser385, Ser380, Thr383, Thr382; and 2) Ser370, and Thr361, Ser363 (following phosphorylation of Thr366 and Ser362 by GSK3)	Induces PTEN closed conformation. Increased PTEN stability, decreased activity	Vasquez et al. 2000; Torres and Pulido. 2001; Al-Khouri et al. 2005; Maccario et al. 2007; Cordier et al. 2012
GSK3-beta	Glycogen Synthase Kinase 3 beta	Ser/Thr protein kinase	Phosphorylation of Ser362 and Thr366. Phosphorylation of Ser370 by CK2 primes phosphorylation at Thr366 by GSK3.	Decreased PTEN stability and/or activity (according to cell types)	Al-Khouri et al. 2005; Maccario et al. 2007; Cordier et al. 2012
PLK1	Polo Like Kinase 1	Ser/Thr protein kinase. Regulator of cell cycle progression	Phosphorylation of Ser380	PTEN phosphorylation impairs its interaction with Cdh1 and stabilizes its association with chromatin. PTEN inhibits PLK1	Song et al. 2011; Choi et al. 2014; Li et al. 2014; Zhang et al. 2016
ROCK1	RhoA-dependent kinase	Ser/Thr protein kinase, activated when bound to GTP-bound Rho	Phosphorylation of Ser229/Thr232 and Thr319/Thr321	Increased PTEN phosphatase activity, increased PTEN/ β -arrestin interaction, decreased Akt activity and cell proliferation, polarized PTEN distribution at plasma membrane	Li et al. 2005; Lima-Fernandes et al. 2011
STK11, LKB1	Serine/Threonine Kinase 11	Ser/Thr protein kinase. Regulates cell polarity. Tumor suppressor	Phosphorylation of Ser385, then Ser380/Thr382/Thr383	Unknown	Mehenni et al. 2005
TOPK, PBK	PDZ Binding Kinase	Ser/Thr protein kinase	Phosphorylation of Ser380	Decreased PTEN activity and accumulation, Akt activation, G2/M progression	Shih et al. 2012; Shinde et al. 2013
FGFR2	Fibroblast Growth Factor Receptor 2	Receptor Tyr protein kinase	Phosphorylation of Tyr240	PTEN lipid phosphatase activity unaffected, inhibition of PI3K signaling preserved. pY240-PTEN binds to chromatin in response to irradiation and facilitates the recruitment of RAD51 to promote DNA repair. Tyr240 phosphorylation associated with resistance to EGFR inhibitor therapy	Fenton et al. 2012; Ma et al. 2019
FGFR3	Fibroblast Growth Factor Receptor 3	Receptor Tyr protein kinase	Phosphorylation of Tyr240	PTEN lipid phosphatase activity unaffected, inhibition of PI3K signaling preserved	Fenton et al. 2012
FRK, RAK	Fyn Related Src Family Tyrosine Kinase	Tyr protein kinase	Phosphorylation of Tyr336	Decreased polyubiquitination by NEDD4, PTEN stabilisation	Yim et al. 2009
Lck	LCK Proto-Oncogene, Non-Receptor Tyrosine Kinase	Src Family Tyr protein kinase	Phosphorylation of Tyr 240, 315	Altered interaction with plasma membrane, phosphatase activity unchanged, decreased stability. Increased Akt activity, anchorage-independent cell growth	Lu et al. 2003; Koul et al. 2002
Lyn	Lyn proto-oncogene, Non-receptor Tyrosine kinase	Src Family Tyr protein kinase	Phosphorylation of Tyr240	Unknown	Fenton et al. 2012

PTK2, FAK1	Protein Tyrosine Kinase 2	Cytoplasmic Tyr protein kinase concentrated at focal adhesions	Phosphorylation of Tyr336	Increased phosphatase activity, protein–lipid interaction, and protein stability. FAK is positively regulated by Rock	Tzenaki et al. 2015
SRC	SRC proto-oncogene, Non-receptor Tyrosine kinase	Tyr protein kinase	Phosphorylation of Tyr240 and Tyr315 (C2 domain), Tyr68 and Tyr155 (catalytic domain), and Tyr377 (tail)	Phosphorylation of Tyr240 and Tyr315: altered interaction with plasma membrane, phosphatase activity unaffected, decreased stability; Phosphorylation of Tyr68 and Tyr155: PTEN destabilization. Increased Akt activity, anchorage-independent cell growth	Koul et al. 2002; Lu et al. 2003; Fenton et al. 2012
PP2A	Protein Phosphatase 2A	Ser/Thr-protein Phosphatase	Dephosphorylation of PTEN at the Ser380, Thr382 and Thr383 cluster within the C-terminal tail	NDRG2 recruits protein phosphatase 2A (PP2A) to PTEN	Nakahata et al. 2014
PTEN	Phosphatase and tensin homolog deleted on chromosome 10	Protein / lipid phosphatase; Tumor suppressor	Probable auto-dephosphorylation of Thr366 and Thr383	PTEN protein phosphatase activity together with lipid phosphatase activity inhibits invasion of U87 glioblastoma cells	Raftopoulou et al. 2004; Tibarewal et al. 2012
PTPN6, SHP1	Protein Tyrosine Phosphatase Non-receptor type 6	Tyr protein phosphatase	Tyr dephosphorylation, reversion of Src/Lck effects	Decreased Akt activity	Lu et al. 2003

Ubiquitination/Deubiquitination/SUMOylation

Ndfip1 and Ndfip2	Nedd4 Family Interacting Protein 1 and protein 2	Endosomal membrane proteins that bind to and activate members of the Nedd4 family of E3 ubiquitin ligases	PTEN ubiquitination	Induces polyubiquitination of PTEN by either Nedd4 or Itch. PTEN destabilization. Neuronal progenitors : Ndfip1 induces nuclear PTEN targeting and decreases cell proliferation (downregulation of cyclinD1, PLK1)	Mund and Pelham 2010; Howitt et al. 2012; Howitt et al. 2015
NEDD4-1	Neural Precursor Cell Expressed, Developmentally Down-Regulated 4, E3 Ubiquitin Protein Ligase	E3 Ubiquitin Protein Ligase	PTEN (mono/poly) ubiquitination	PTEN polyubiquitination by NEDD4-1, leads to PTEN degradation. NEDD4-1 can also induce PTEN monoubiquitination and nuclear translocation	Wang et al. 2007; Trotman et al. 2007
SHARPIN, SIPL1	Shank-interacting protein-like 1	Component of the LUBAC complex which conjugates linear polyubiquitin chains in a head-to-tail manner to substrates	Promotes PTEN polyubiquitination	Inhibits PTEN lipid phosphatase activity, does not induce PTEN degradation	He et al. 2010; De Melo et al. 2014
TRIM27, RFP	Tripartite Motif Containing 27	Transcription repressor. RING-Type E3 Ubiquitin transferase	Polyubiquitination	RFP-mediated ubiquitination negatively regulates PTEN phosphatase activity, resulting in increase in AKT activity and inhibition of TRAIL-mediated apoptosis. PTEN stability or localization unaffected	Lee et al. 2013
WWP1	WW domain-containing ubiquitin E3 ligase 1	Ubiquitin E3 ligase	Non-degradative Lys27-linked polyubiquitination of PTEN	Suppresses PTEN oligomerization, plasma membrane recruitment, and tumour suppressor functions in vitro and in vivo	Lee et al. 2019
WWP2, AIP-2	WW domain-containing E3 Ligase 2	Ubiquitin E3 ligase	PTEN polyubiquitination and degradation	Increased Akt activity, Decreased apoptosis. PTEN phosphorylation on Tyr155 prevents interaction with WWP2	Maddika et al. 2011

XIAP, BIRC4	X-Linked Inhibitor Of Apoptosis	E3 Ubiquitin Protein Ligase, apoptotic suppressor protein,	PTEN mono and poly-ubiquitination	Regulates PTEN accumulation and compartmentalization	van Themsche et al. 2009
STUB1, CHIP	STIP1 Homology And U-Box Containing Protein 1, RING-Type E3 Ubiquitin Transferase	STIP1 Homology And U-Box Containing Protein 1	PTEN polyubiquitination and degradation	HSP70 and HSP90 favor PTEN interaction with STUB1	Ahmed et al. 2012
OTUD3	OTU Deubiquitinase 3	Deubiquitinating enzyme that hydrolyzes Lys-6- and Lys-11-linked polyubiquitin	Deubiquitination: removes Lys6, Lys11, Lys27 and Lys48 types of ubiquitin chains on PTEN. Does not remove monoubiquitin on PTEN	PTEN Stabilization, inhibition of Akt signaling. Depletion of OTUD3 leads to the activation of Akt signalling, induction of cellular transformation and cancer metastasis.	Yuan et al. 2015
USP7 (HAUSP)	Ubiquitin Specific Peptidase 7	Deubiquitinates target proteins	Nuclear PTEN exclusion through decreased monoubiquitination (Lys13, Lys289)	USP7 regulates PTEN subcellular localization but not its stability. High levels of USP7 is associated with tumor aggressiveness. Nuclear/monoubiquitinated PTEN possesses greater apoptotic potential, nuclear exclusion is observed in more aggressive cancers.	Song et al. 2008
USP10	Ubiquitin Specific Peptidase 10	Cleaves ubiquitin from ubiquitin-conjugated protein substrates	Deubiquitination and stabilization of PTEN	Knockdown of USP10 promotes tumor growth and invasion of Lewis carcinoma cells	Sun et al. 2018
USP11	Ubiquitin Specific Peptidase 11	Deubiquitinating enzyme (cysteine protease). X-linked tumor suppressor gene	Reverses PTEN polyubiquitination in cytoplasm and nucleus	PTEN stabilization	Park et al. 2019
USP-13	Ubiquitin Specific Peptidase 13	Deubiquitinates target proteins	Deubiquitination : removes Lys6, Lys29 and Lys63 types of ubiquitin chains on PTEN. Does not remove monoubiquitination	PTEN stabilization, inhibition of Akt signaling. Subcellular localization unaffected (cytoplasm). Overexpression of USP13 suppresses tumorigenesis and glycolysis in PTEN-positive but not PTEN-null breast cancer cells	Zhang et al. 2013
UBE2I	SUMO-conjugating enzyme Ubc9	SUMO E2 Conjugase involved in SUMOylation of targets impacting their subcellular localization, transcriptional regulation, and protein stability	Sumoylation of Lys266 targets PTEN to the plasma membrane. Sumoylation of Lys254 controls PTEN nuclear localization.	PTEN SUMOylated on Lys 266 decreases Akt activity, suppresses anchorage independent cell proliferation and tumor growth in vivo. PTEN sumoylated on Lys254 involved in DNA damage repair pathways	Huang et al. 2012; Bassi et al. 2013
PIASX α	SUMO E3 ligase PIAS3	SUMO E3 ligase that facilitates SUMOylation	Facilitates PTEN SUMOylation on Lys254 and Lys266	Leads to protection from polyubiquitination, increased PTEN stability, inhibition of cell proliferation and tumour suppression	Wang et al. 2014
Acetylation/Deacetylation					
PCAF KAT2B	Lysine Acetyltransferase 2B	Acetylates histone and non-histone proteins	Acetylation of Lys125 and Lys128 in response to growth factors. Decreased PTEN lipid Ptase activity	Decreased PTEN-induced G1 cell cycle arrest	Okumura et al. 2006
HDAC6	Histone Deacetylase 6	Deacetylation of lysine residues, closely related to sirtuins	Deacetylation of Lys163	Lys163 acetylation inhibited the interaction of the PTEN C-tail with the remaining part of PTEN. Deacetylation of Lys163 decreases PTEN translocation to plasma membrane	Meng et al. 2016
CREBBP, CBP	CREB Binding Protein	Acetyltransferase	PTEN acetylation on Lys402	Increased interaction with PDZ domain (hDLG, MAGI-2), PTEN phosphatase activity not directly affected	Ikenoue et al. 2008
SIRT1	Sirtuin 1	NAD-dependent protein deacetylase	Deacetylation of Lys402	Antagonizes CBP-induced PTEN acetylation	Ikenoue et al. 2008
Oxidation/Reduction					

PRDX1	Peroxiredoxin 1	Antioxidant enzyme which reduces hydrogen peroxide and alkyl hydroperoxides	Prdx1 interaction with PTEN protects oxidation-induced inactivation and promotes PTEN lipid phosphatase activity under oxidative stress	Prdx1 tumour suppression of Ras- or ErbB-2-induced transformation mainly via PTEN	Cao et al. 2009
TXNIP	Thioredoxin-interacting protein	Negative regulator of thioredoxin-NADPH-dependent reduction of disulphide bonds in proteins	Reactivation of oxidized PTEN	H ₂ O ₂ produced under pathological conditions or via activation of cell surface receptors inactivates PTEN through formation of a disulfide bridge between Cys124 and Cys71. TXNIP acts to reactivate oxidized PTEN	Lee et al. 2002; Kwon et al. 2004; Hui et al. 2008
Nitrosylation					
eNOS	Endothelial nitric oxide synthase	Synthesizes NO	S-nitrosylation of PTEN	S-nitrosylation of PTEN promotes its ubiquitination	Gupta et al. 2017
Ribosylation					
TNKS1 and TNKS2	Tankyrase 1 and Tankyrase 2	Poly-ADP-ribosyltransferase (PARP)	PTEN ADP-rybosylation (Glu40, Glu150, Asp326)	PTEN ribosylation promotes recognition by PAR-binding E3 ubiquitin ligase RNF146, leading to PTEN ubiquitination and degradation. Knockdown of tankyrases stabilizes PTEN, down-regulates AKT and suppresses cell proliferation and glycolysis in vitro and tumor growth in vivo	Li et al. 2015
Methylation					
SMYD2	SET And MYND Domain Containing 2	Protein-lysine N-methyltransferase	PTEN methylation at lysine 313 favors PTEN phosphorylation at Ser380	PTEN inactivation. Increased Akt activity	Nakakido et al. 2015
NSD2	Histone-lysine N-methyltransferase	Histone methyltransferase	PTEN methylation at Lys 349	Recruitment of PTEN to DNA damage sites	Zhang et al. 2019

Table 2: Partial list of PTEN-interacting partners

Molecular Partner	Full Name	Function	PTEN Interacting region	Subcellular localization of interaction	Functional effect on PTEN	References
PTEN	Phosphatase and tensin homolog deleted on chromosome 10	Protein / lipid phosphatase; Tumor suppressor	Multiple interfaces, including catalytic domain and C-terminal tail	Plasma membrane	Wild-type PTEN oligomerization results in increased lipid phosphatase activity. Mutant/inactive PTEN exerts dominant negative activity on wild-type PTEN via hetero-oligomerization	Papa et al. 2014; Heinrich et al. 2015
CFTR	Cystic Fibrosis Transmembrane Conductance Regulator	Chloride channel	NH2 terminus and central core of PTEN (loop and C2 domain)	Plasma membrane	Positions PTEN at the membrane. Independent of Cl-CFTR channel function. Downregulation of PI3K/Akt pathway, enhanced TLR4-TIRAP signaling and host immunity against <i>Pseudomonas aeruginosa</i> infection	Riquelme et al. 2017
MME, NEP	Membrane Metallo-endopeptidase	Neutral endopeptidase	Tail	Plasma membrane	Recruits PTEN to plasma membrane, enhances stability and phosphatase activity. Decreased Akt activity	Sumitomo et al. 2004
S1PR2	Sphingosine-1-Phosphate Receptor 2	G protein-coupled receptor	ND	Plasma membrane	Recruits PTEN to plasma membrane and enhances lipid phosphatase activity; increases tyrosine phosphorylation; promotes PTEN conformational change upon ligand activation. Inhibition of HUVEC and MEF migration	Sanchez et al. 2005; Lima-Fernandes et al. 2014
MC1R	Melanocortin 1 Receptor	G protein-coupled receptor for melanocyte-stimulating hormone (MSH). Controls melanogenesis	C2 domain	Plasma membrane	UVB exposure combined with α -MSH induces MC1R-PTEN interaction. MC1R protects PTEN from degradation by competing with WWP2 for binding PTEN. Decreased AKT activity	Cao et al. 2013
MYO5A	Myosin VA	Actin-based motor protein involved in cytoplasmic vesicle transport and anchorage, spindle-pole alignment and mRNA translocation.	Tail (aa 354–399)	Plasma membrane	Increased PTEN activity. Decreased neuronal soma size. PTEN phosphorylation by GSK3/CK2 increases its interaction with MYO5A	van Diepen et al. 2009
PIK3R1 (p85 α)	Phosphoinositide-3-Kinase Regulatory Subunit 1	Regulatory subunit of PI3K, adapter required for stabilization and localization of the p110-PI3K catalytic unit	Catalytic domain	Plasma membrane	Increases PTEN membrane association, lipid phosphatase activity and impairs PTEN degradation by competing with the E3 ligase WWP2. Binds preferentially PTEN in its unphosphorylated form (residues Ser380/Thr382/Thr383)	Cheung et al. 2015
β -arr1	β -arrestin 1	Regulatory scaffolding protein	C2 domain	Cytoplasm, Plasma membrane	Increased PTEN membrane association and lipid phosphatase activity. Decreased Akt activity, control of cell proliferation, also blocks the inhibitory effect of PTEN on glioma cell migration and involved in 3D multicellular assembly, pointing to differential regulation of multiple PTEN functions	Lima-Fernandes et al. 2011; Javadi et al. 2017
β -arr2	β -arrestin 2	Regulatory scaffolding protein	C2 domain	Cytoplasm, Plasma membrane	Increased PTEN membrane association and lipid phosphatase activity. Decreased Akt activity, control of cell proliferation, pointing to differential regulation of multiple PTEN functions; Also can alter PTEN conformation.	Lima-Fernandes et al. 2011; Lima-Fernandes et al. 2014
DLG1, SAP-97	Discs Large MAGUK Scaffold Protein 1	Scaffolding and/or regulatory molecules	PDZ domain-binding motif	Plasma membrane	Increased PTEN stability, decreased Akt activity. DLG1 may serve as a platform to bring in proximity APC and PTEN tumor suppressor activities (APC binds to the three PDZ domains of DLG1, PTEN binds PDZ2).	Adey et al. 2000; Valiente et al. 2005; Sotelo et al. 2012
MAGI-1, BAP1	Membrane Associated Guanylate Kinase, WW And PDZ Domain Containing 1	Scaffolding and/or regulatory molecules	PDZ domain-binding motif	Plasma membrane, junctional complexes	Recruitment to E-cadherin junctional complexes, stabilization of adherens junctions and suppression of invasiveness. Decreased Akt activity	Kotelevets et al. 2005; Chastre et al. 2009

MAGI-2, SSCAM, and MAGI-3	Membrane Associated Guanylate Kinase, WW and PDZ Domain Containing 2; and PDZ Domain Containing 3	Scaffolding and/or regulatory molecules	PDZ domain-binding motif, decreased interaction with PTEN phosphorylated at Thr382	Plasma membrane, junctional complexes	Increased PTEN protein stability; favors PTEN recruitment into high molecular weight molecular complexes and enhances PTEN-mediated down-regulation of PI3K/Akt pathway	Wu et al. 2000; Tolkacheva et al. 2001; Valiente et al. 2005
MAST1, 2 and 3	Microtubule Associated Serine Threonine Kinases 1, 2 and 3	Scaffolding and/or regulatory molecules, Ser/Thr protein kinases	PDZ domain-binding motif	Plasma membrane, junctional complexes	Binding of PTEN to the PDZ domain of MAST2 (MAST205) facilitates its phosphorylation by this kinase	Valiente et al. 2005.
PAR3, PAR-3	Par-3 Family Cell Polarity Regulator	Scaffolding molecule involved in asymmetrical cell division, tight junctions and cell polarization	PDZ domain-binding motif	Plasma membrane, junctional complexes	The 2nd PDZ domain of Par-3 binds to phosphatidylinositol (PI) lipid membranes, the 3rd binds PTEN. Epithelial cell polarization	Wu et al. 2007
SLC9A3R1, NHERF1, EBP50	SLC9A3 Regulator 1	Na ⁺ /H ⁺ exchanger regulatory cofactor, interacts with and regulates various proteins including CFTR and G-protein coupled receptors	PDZ domain-binding motif, decreased interaction with PTEN phosphorylated at Ser380/Thr382/Thr383	Plasma membrane	Assembly of a ternary complex between PTEN, NHERFs and PDGFR. Enhances PTEN stability, restricts activation of PI3K by PDGFR; inhibits Akt pathway.	Takahashi et al. 2006
MAN2C1	Cytosolic α -mannosidase 2C1	Catabolic enzyme for the breakdown of free oligosaccharides	ND	Cytoplasm	Prevents PTEN recruitment to the cell membrane, Inhibits lipid phosphatase activity	He et al. 2011
PARK7, DJ1	Parkinsonism Associated Deglycase	Peptidase	ND	Cytoplasm	Oxidized DJ1 Inhibits PTEN activity. High glucose increases PTEN/DJ1 interaction	Kim et al. 2009; Das et al. 2011
BMI1	Proto-Oncogene, Polycomb Ring Finger	Component of polycomb group complex 1 (PRC1) that modulates gene expression through epigenetic/ chromatin remodeling.	C2 domain (186-286)	Nucleus	Sequestering PTEN in the nucleus decreases PTEN's ability to inhibit the PI3K-AKT pathway	Fan et al. 2009
IPO11	Importin-11	Nucleocytoplasmic transport of protein and RNA cargoes	Monoubiquitinated PTEN	Nuclear translocation	Decreased degradation: nuclear translocation protects PTEN from cytoplasmic proteins that cause its degradation	Chen et al. 2017
MVP	Major Vault Protein	Component of a multi-subunit ribonucleoprotein structures, might be involved in nucleocytoplasmic transport	C2 domain, Ca ²⁺ dependent	Nuclear translocation	Nuclear transport of PTEN, increased PI3K/AKT pathway activity	Chung et al. 2005; Minaguchi et al. 2006
PPP1R10, PNU10	Protein Phosphatase 1 Regulatory Subunit 10	Protein phosphatase 1 binding protein	C2-domain	Nucleus	Sequesters PTEN in the nucleus in an inactive state (no induction of rad51 and p53 expression), decreases PTEN's ability to inhibit the PI3K-AKT pathway	Kavela et al. 2013
PREX2	Phosphatidylinositol-3,4,5-Trisphosphate Dependent Rac Exchange Factor 2	Guanine-nucleotide exchange factors for Rac family small G proteins	C-term tail (phosphorylated), Catalytic/ C2 domains	ND	Inhibits PTEN lipid phosphatase activity. Stimulation of cell growth, decrease of glucose uptake and insulin resistance	Fine et al. 2009; Hodakoski et al. 2014

Figure legends

Fig. 1. PTEN structure and function.

(a) The PTEN protein contains five functional domains: a short N-terminal PIP2-binding domain (PIP2-BD), the catalytic phosphatase domain, a C2 membrane-targeting/scaffold domain, a regulatory C-terminal tail containing multiple phosphorylation sites and several PEST sequences, and a PDZ-binding motif (PDZ-BM). **(b)** Schematic representation of the PTEN/PI3K/AKT pathway. PI3Ks are activated by a wide range of stimuli, including receptor tyrosine kinase receptors (RTKs) and G protein-coupled receptors (GPCRs). PI(3,4,5)P3 is produced from PI(4,5)P2 through the action of PI3Ks. PI(3,4,5)P3 acts as a lipid signalling intermediate to promote the recruitment and activation of a subset of pleckstrin homology (PH) domain-containing proteins, including AKT. AKT subsequently phosphorylates downstream protein substrates, leading to increased cell survival, proliferation, migration and invasion. PTEN acts as a brake on the PI3K/AKT pathway via its lipid phosphatase activity, which removes the D3 phosphate from PI(3,4,5)P3 to produce PI(4,5)P2.

Fig. 2. Molecular mechanisms controlling PTEN expression and activity.

The diagram depicts the different molecular mechanisms involved in control of PTEN expression and function. Methylation of the promoter results in *PTEN* silencing and hypermethylation of the promoter provokes reduced *PTEN* expression in different cancers. *PTEN* transcription is also under the control of a variety of transcription factors that bind the *PTEN* promoter to positively (e.g., p53, Egr-1) or negatively (e.g., NF- κ B, c-Jun) regulate its expression. At the post-transcriptional level microRNAs (miRNA) negatively impact PTEN expression. By acting as decoys for PTEN-targeting miRNAs, competitive endogenous RNAs (ceRNAs) that present partial sequence homology to PTEN as well as PTEN pseudogene (*PTENP1*) mRNAs, act to enhance PTEN expression. At the post-translational level the PTEN protein is subject to regulation by post-translational modifications (PTMs) and partner proteins, which impact PTEN conformation, localization, stability and activity. The culmination of all these regulatory events results in the "functional dose" of PTEN in the cell and corresponding functional outputs.

Fig. 3. Post-translational modifications mediating PTEN regulation.

(a) The diagram shows post-translational modifications (PTMs) implicated in PTEN regulation. ATM: ataxia telangiectasia kinase, CBP: CREB-binding protein, CK2: casein kinase 2, FAK: Focal adhesion kinase, GSK3 β : glycogen synthase 3 β , HDAC6: histone deacetylase 6, NEDD4: neural precursor cell expressed developmentally down-regulated protein 4, PCAF: p300/CBP-associated factor, histone acetyltransferase, Rak: Fyn Related Src family kinase, ROCK: RhoA-associated, coiled-coil containing protein kinase, SMYD2: SET and MYND domain containing 2, lysine methyltransferase, Src: protein tyrosine kinase, TNKS1/2: tankyrase1/2, poly(ADP-ribose) polymerase, Ubc9: SUMO E2 Conjugase. **(b)** The diagram shows the PTEN "phosphorylation switch" conformational model between "closed" and "open" states. Phosphorylation of the C-terminal tail residues Ser380, Thr382, Thr383 and Ser385 promotes a PTEN intramolecular interaction, resulting in the "closed" conformation. Dephosphorylation of the C-terminal tail phosphorylation cluster results in the loss of this intramolecular interaction and PTEN switches to an "open"

conformation that is targeted to the plasma membrane where its substrate PI(3,4,5)P3 resides.

Fig. 4. A PTEN biosensor to monitor conformational change in live cells.

(a) The schematic shows the Rluc-PTEN-YFP biosensor with PTEN sandwiched between the energy donor Rluc and the energy acceptor YFP. The diagram illustrates how PTEN conformational rearrangement may lead to changes in BRET measurements, although the actual orientations of donor and acceptor proteins are not known. **(b)** The BRET ratio is calculated by the ratio of light emitted by YFP at 530nm and the light emitted by Rluc at 480nm (YFP/Rluc) in the presence of the Rluc substrate coelenterazine. **(c)** As the biosensor is unimolecular, BRET signals remain constant over a wide range of Rluc concentrations (theoretical values are presented here based on previous experimental data). This is particularly useful in situations where, for example, a mutant of PTEN is less stable, as a change in BRET will only reflect a conformational change. Background BRET is determined using a Rluc-PTEN fusion, lacking YFP acceptor. Specific "net" BRET is then calculated by subtracting this value (background "bystander" BRET) from that obtained with the biosensor Rluc-PTEN-YFP (shaded grey area). **(d)** A list of potential experimental modes in which the biosensor can be used. **(e)** Specific BRET values multiplied by 1000 to generate mBRET (milliBRET units). Wild-type PTEN biosensor establishes the basal BRET signal (black bar and dotted line). Conformational changes are detected by changes in BRET signal that can be either negative (Condition X) or positive (Condition Y) compared to basal BRET.

References

- Adey NB, Huang L, Ormonde PA, Baumgard ML, Pero R, Byreddy DV, Tavtigian SV, Bartel PL. 2000. Threonine phosphorylation of the MMAC1/PTEN PDZ binding domain both inhibits and stimulates PDZ binding. *Cancer Res* **60**: 35-37.
- Ahmed SF, Deb S, Paul I, Chatterjee A, Mandal T, Chatterjee U, Ghosh MK. 2012. The chaperone-assisted E3 ligase C terminus of Hsc70-interacting protein (CHIP) targets PTEN for proteasomal degradation. *J Biol Chem* **287**: 15996-16006.
- Al-Khouri AM, Ma Y, Togo SH, Williams S, Mustelin T. 2005. Cooperative phosphorylation of the tumor suppressor phosphatase and tensin homologue (PTEN) by casein kinases and glycogen synthase kinase 3beta. *J Biol Chem* **280**: 35195-35202.
- Alimonti A, Carracedo A, Clohessy JG, Trotman LC, Nardella C, Egia A, Salmena L, Sampieri K, Haveman WJ, Brogi E et al. 2010. Subtle variations in Pten dose determine cancer susceptibility. *Nature genetics* **42**: 454-458.
- Bassi C, Ho J, Srikumar T, Dowling RJ, Gorrini C, Miller SJ, Mak TW, Neel BG, Raught B, Stambolic V. 2013. Nuclear PTEN controls DNA repair and sensitivity to genotoxic stress. *Science* **341**: 395-399.
- Bolduc D, Rahdar M, Tu-Sekine B, Sivakumaren SC, Raben D, Amzel LM, Devreotes P, Gabelli SB, Cole P. 2013. Phosphorylation-mediated PTEN conformational closure and deactivation revealed with protein semisynthesis. *Elife* **2**: e00691.
- Cao J, Schulte J, Knight A, Leslie NR, Zagozdzon A, Bronson R, Manevich Y, Beeson C, Neumann CA. 2009. Prdx1 inhibits tumorigenesis via regulating PTEN/AKT activity. *Embo J* **28**: 1505-1517.
- Cao J, Wan L, Hacker E, Dai X, Lenna S, Jimenez-Cervantes C, Wang Y, Leslie NR, Xu GX, Widlund HR et al. 2013. MC1R is a potent regulator of PTEN after UV exposure in melanocytes. *Mol Cell* **51**: 409-422.
- Chalhoub N, Baker SJ. 2009. PTEN and the PI3-kinase pathway in cancer. *Annu Rev Pathol* **4**: 127-150.
- Chastre E, Abdessamad M, Kruglov A, Bruyneel E, Bracke M, Di Gioia Y, Beckerle MC, van Roy F, Kotelevets L. 2009. TRIP6, a novel molecular partner of the MAGI-1 scaffolding molecule, promotes invasiveness. *Faseb J* **23**: 916-928.
- Chen JH, Zhang P, Chen WD, Li DD, Wu XQ, Deng R, Jiao L, Li X, Ji J, Feng GK et al. 2015. ATM-mediated PTEN phosphorylation promotes PTEN nuclear translocation and autophagy in response to DNA-damaging agents in cancer cells. *Autophagy* **11**: 239-252.
- Chen M, Nowak DG, Narula N, Robinson B, Watrud K, Ambrico A, Herzka TM, Zeeman ME, Minderer M, Zheng W et al. 2017. The nuclear transport receptor Importin-11 is a tumor suppressor that maintains PTEN protein. *J Cell Biol* **216**: 641-656.
- Cheung LW, Walkiewicz KW, Besong TM, Guo H, Hawke DH, Arold ST, Mills GB. 2015. Regulation of the PI3K pathway through a p85alpha monomer-homodimer equilibrium. *Elife* **4**: e06866.
- Choi BH, Pagano M, Dai W. 2014. Plk1 protein phosphorylates phosphatase and tensin homolog (PTEN) and regulates its mitotic activity during the cell cycle. *J Biol Chem* **289**: 14066-14074.
- Chon HJ, Bae KJ, Lee Y, Kim J. 2015. The casein kinase 2 inhibitor, CX-4945, as an anti-cancer drug in treatment of human hematological malignancies. *Frontiers in pharmacology* **6**: 70.

- Chung JH, Ginn-Pease ME, Eng C. 2005. Phosphatase and tensin homologue deleted on chromosome 10 (PTEN) has nuclear localization signal-like sequences for nuclear import mediated by major vault protein. *Cancer Res* **65**: 4108-4116.
- Cordier F, Chaffotte A, Terrien E, Préhaud C, Theillet F-X, Delepierre M, Lafon M, Buc H, Wolff N. 2012. Ordered phosphorylation events in two independent cascades of the PTEN C-tail revealed by NMR. *J Am Chem Soc* **134**: 20533–20543.
- Das S, Dixon JE, Cho W. 2003. Membrane-binding and activation mechanism of PTEN. *Proc Natl Acad Sci U S A* **100**: 7491-7496.
- Das F, Dey N, Venkatesan B, Kasinath BS, Ghosh-Choudhury N, Choudhury GG. 2011. High glucose upregulation of early-onset Parkinson's disease protein DJ-1 integrates the PRAS40/TORC1 axis to mesangial cell hypertrophy. *Cell Signal* **23**: 1311-1319.
- De Melo J, Lin X, He L, Wei F, Major P, Tang D. 2014. SIPL1-facilitated PTEN ubiquitination contributes to its association with PTEN. *Cell Signal* **26**: 2749-2756.
- Di Cristofano A, Pesce B, Cordon-Cardo C, Pandolfi PP. 1998. Pten is essential for embryonic development and tumour suppression. *Nat Genet* **19**: 348-355.
- Duerr EM, Rollbrocker B, Hayashi Y, Peters N, Meyer-Puttlitz B, Louis DN, Schramm J, Wiestler OD, Parsons R, Eng C et al. 1998. PTEN mutations in gliomas and glioneuronal tumors. *Oncogene* **16**: 2259-2264.
- Fan C, He L, Kapoor A, Rybak AP, De Melo J, Cutz JC, Tang D. 2009. PTEN inhibits BMI1 function independently of its phosphatase activity. *Mol Cancer* **8**: 98.
- Fenton TR, Nathanson D, Ponte de Albuquerque C, Kuga D, Iwanami A, Dang J, Yang H, Tanaka K, Oba-Shinjo SM, Uno M et al. 2012. Resistance to EGF receptor inhibitors in glioblastoma mediated by phosphorylation of the PTEN tumor suppressor at tyrosine 240. *Proc Natl Acad Sci U S A* **109**: 14164-14169.
- Fine B, Hodakoski C, Koujak S, Su T, Saal LH, Maurer M, Hopkins B, Keniry M, Sulis ML, Mense S et al. 2009. Activation of the PI3K pathway in cancer through inhibition of PTEN by exchange factor P-REX2a. *Science* **325**: 1261-1265.
- Fouladkou F, Landry T, Kawabe H, Neeb A, Lu C, Brose N, Stambolic V, Rotin D. 2008. The ubiquitin ligase Nedd4-1 is dispensable for the regulation of PTEN stability and localization. *Proc Natl Acad Sci U S A* **105**: 8585-8590.
- Freeman DJ, Li AG, Wei G, Li HH, Kertesz N, Lesche R, Whale AD, Martinez-Diaz H, Rozengurt N, Cardiff RD et al. 2003. PTEN tumor suppressor regulates p53 protein levels and activity through phosphatase-dependent and -independent mechanisms. *Cancer Cell* **3**: 117-130.
- Gu J, Tamura M, Yamada KM. 1998. Tumor suppressor PTEN inhibits integrin- and growth factor-mediated mitogen-activated protein (MAP) kinase signaling pathways. *J Cell Biol* **143**: 1375-1383.
- Gupta A, Anjomani-Virmouni S, Koundouros N, Dimitriadi M, Choo-Wing R, Valle A, Zheng Y, Chiu YH, Agnihotri S, Zadeh G et al. 2017. PARK2 Depletion Connects Energy and Oxidative Stress to PI3K/Akt Activation via PTEN S-Nitrosylation. *Mol Cell* **65**: 999-1013 e1017.
- He L, Ingram A, Rybak AP, Tang D. 2010. Shank-interacting protein-like 1 promotes tumorigenesis via PTEN inhibition in human tumor cells. *J Clin Invest* **120**: 2094-2108.
- He L, Fan C, Kapoor A, Ingram AJ, Rybak AP, Austin RC, Dickhout J, Cutz JC, Scholey J, Tang D. 2011. alpha-Mannosidase 2C1 attenuates PTEN function in prostate cancer cells. *Nat Commun* **2**: 307.
- Heinrich F, Chakravarthy S, Nanda H, Papa A, Pandolfi PP, Ross AH, Harishchandra RK, Gericke A, Losche M. 2015. The

- PTEN tumor suppressor forms homodimers in solution. *Structure* **23**: 1952-1957.
- Hodakoski C, Hopkins BD, Barrows D, Mense SM, Keniry M, Anderson KE, Kern PA, Hawkins PT, Stephens LR, Parsons R. 2014. Regulation of PTEN inhibition by the pleckstrin homology domain of P-REX2 during insulin signaling and glucose homeostasis. *Proc Natl Acad Sci U S A* **111**: 155-160.
- Hopkins BD, Fine B, Steinbach N, Dendy M, Rapp Z, Shaw J, Pappas K, Yu JS, Hodakoski C, Mense S et al. 2013. A secreted PTEN phosphatase that enters cells to alter signaling and survival. *Science* **341**: 399-402.
- Howitt J, Lackovic J, Low LH, Naguib A, Macintyre A, Goh CP, Callaway JK, Hammond V, Thomas T, Dixon M et al. 2012. Ndfip1 regulates nuclear Pten import in vivo to promote neuronal survival following cerebral ischemia. *J Cell Biol* **196**: 29-36.
- Howitt J, Low LH, Putz U, Doan A, Lackovic J, Goh CP, Gunnensen J, Silke J, Tan SS. 2015. Ndfip1 represses cell proliferation by controlling Pten localization and signaling specificity. *J Mol Cell Biol* **7**: 119-131.
- Huang J, Yan J, Zhang J, Zhu S, Wang Y, Shi T, Zhu C, Chen C, Liu X, Cheng J et al. 2012. SUMO1 modification of PTEN regulates tumorigenesis by controlling its association with the plasma membrane. *Nat Commun* **3**: 911.
- Hui ST, Andres AM, Miller AK, Spann NJ, Potter DW, Post NM, Chen AZ, Sachithanatham S, Jung DY, Kim JK et al. 2008. Txnip balances metabolic and growth signaling via PTEN disulfide reduction. *Proc Natl Acad Sci* **105**: 3921-3926.
- Ikenoue T, Inoki K, Zhao B, Guan KL. 2008. PTEN acetylation modulates its interaction with PDZ domain. *Cancer Res* **68**: 6908-6912.
- Javadi A, Deevi RK, Evergren E, Blondel-Tepaz E, Baillie GS, Scott MG, Campbell FC. 2017. PTEN controls glandular morphogenesis through a juxtamembrane beta-Arrestin1/ARHGAP21 scaffolding complex. *Elife* **6**.
- Kavela S, Shinde SR, Ratheesh R, Viswakalyan K, Bashyam MD, Gowrishankar S, Vamsy M, Pattnaik S, Rao S, Sastry RA et al. 2013. PNUTS functions as a proto-oncogene by sequestering PTEN. *Cancer Res* **73**: 205-214.
- Keniry M, Parsons R. 2008. The role of PTEN signaling perturbations in cancer and in targeted therapy. *Oncogene* **27**: 5477-5485.
- Kim YC, Kitaura H, Taira T, Iguchi-Arigo SM, Ariga H. 2009. Oxidation of DJ-1-dependent cell transformation through direct binding of DJ-1 to PTEN. *Int J Oncol* **35**: 1331-1341.
- Kotelevets L, van Hengel J, Bruyneel E, Mareel M, van Roy F, Chastre E. 2005. Implication of the MAGI-1b/PTEN signalosome in stabilization of adherens junctions and suppression of invasiveness. *Faseb J* **19**: 115-117.
- Kotelevets L, Scott MGH, Chastre E. 2018. Targeting PTEN in Colorectal Cancers. *Adv Exp Med Biol* **1110**: 55-73.
- Koul D, Jasser SA, Lu Y, Davies MA, Shen R, Shi Y, Mills GB, Yung WK. 2002. Motif analysis of the tumor suppressor gene MMAC/PTEN identifies tyrosines critical for tumor suppression and lipid phosphatase activity. *Oncogene* **21**: 2357-2364.
- Kwak YD, Ma T, Diao S, Zhang X, Chen Y, Hsu J, Lipton SA, Masliah E, Xu H, Liao FF. 2010. NO signaling and S-nitrosylation regulate PTEN inhibition in neurodegeneration. *Mol Neurodegener* **5**: 49.
- Kwon J, Lee SR, Yang KS, Ahn Y, Kim YJ, Stadtman ER, Rhee SG. 2004. Reversible oxidation and inactivation of the tumor suppressor PTEN in cells stimulated with peptide growth factors. *Proc Natl Acad Sci U S A* **101**: 16419-16424.

- Lee JO, Yang H, Georgescu MM, Di Cristofano A, Maehama T, Shi Y, Dixon JE, Pandolfi P, Pavletich NP. 1999. Crystal structure of the PTEN tumor suppressor: implications for its phosphoinositide phosphatase activity and membrane association. *Cell* **99**: 323-334.
- Lee SR, Yang KS, Kwon J, Lee C, Jeong W, Rhee SG. 2002. Reversible inactivation of the tumor suppressor PTEN by H₂O₂. *J Biol Chem* **277**: 20336-20342.
- Lee JT, Shan J, Zhong J, Li M, Zhou B, Zhou A, Parsons R, Gu W. 2013. RFP-mediated ubiquitination of PTEN modulates its effect on AKT activation. *Cell Res* **23**: 552-564.
- Lee YR, Chen M, Pandolfi PP. 2018. The functions and regulation of the PTEN tumour suppressor: new modes and prospects. *Nat Rev Mol Cell Biol*.
- Lee YR, Chen M, Lee JD, Zhang J, Lin SY, Fu TM, Chen H, Ishikawa T, Chiang SY, Katon J et al. 2019. Reactivation of PTEN tumor suppressor for cancer treatment through inhibition of a MYC-WWP1 inhibitory pathway. *Science* **364**.
- Leslie NR, Bennett D, Lindsay YE, Stewart H, Gray A, Downes CP. 2003. Redox regulation of PI 3-kinase signalling via inactivation of PTEN. *Embo J* **22**: 5501-5510.
- Leslie NR, Downes CP. 2004. PTEN function: how normal cells control it and tumour cells lose it. *Biochem J* **382**: 1-11.
- Leslie NR, Foti M. 2011. Non-genomic loss of PTEN function in cancer: not in my genes. *Trends Pharmacol Sci* **32**: 131-140.
- Li DM, Sun H. 1997. TEP1, encoded by a candidate tumor suppressor locus, is a novel protein tyrosine phosphatase regulated by transforming growth factor beta. *Cancer Res* **57**: 2124-2129.
- Li J, Yen C, Liaw D, Podsypanina K, Bose S, Wang SI, Puc J, Miliareis C, Rodgers L, McCombie R et al. 1997. PTEN, a putative protein tyrosine phosphatase gene mutated in human brain, breast, and prostate cancer. *Science* **275**: 1943-1947.
- Li Z, Dong X, Wang Z, Liu W, Deng N, Ding Y, Tang L, Hla T, Zeng R, Li L et al. 2005. Regulation of PTEN by Rho small GTPases. *Nat Cell Biol* **7**: 399-404.
- Li Z, Li J, Bi P, Lu Y, Burcham G, Elzey BD, Ratliff T, Konieczny SF, Ahmad N, Kuang S, et al. 2014. Plk1 phosphorylation of PTEN causes a tumor-promoting metabolic state. *Mol Cell Biol* **34**: 3642–3661.
- Li N, Zhang Y, Han X, Liang K, Wang J, Feng L, Wang W, Songyang Z, Lin C, Yang L et al. 2015. Poly-ADP ribosylation of PTEN by tankyrases promotes PTEN degradation and tumor growth. *Genes & development* **29**: 157-170.
- Lima-Fernandes E, Enslin H, Camand E, Kotelevets L, Boularan C, Achour L, Benmerah A, Gibson LC, Baillie GS, Pitcher JA et al. 2011. Distinct functional outputs of PTEN signalling are controlled by dynamic association with beta-arrestins. *Embo J* **30**: 2557-2568.
- Lima-Fernandes E, Misticone S, Boularan C, Paradis JS, Enslin H, Roux PP, Bouvier M, Baillie GS, Marullo S, Scott MG. 2014. A biosensor to monitor dynamic regulation and function of tumour suppressor PTEN in living cells. *Nat Commun* **5**: 4431.
- Lu Y, Yu Q, Liu JH, Zhang J, Wang H, Koul D, McMurray JS, Fang X, Yung WK, Siminovitch KA et al. 2003. Src family protein-tyrosine kinases alter the function of PTEN to regulate phosphatidylinositol 3-kinase/AKT cascades. *J Biol Chem* **278**: 40057-40066.
- Ma J, Benitez JA, Li J, Miki S, Ponte de Albuquerque C, Galatro T, Orellana L, Zanca C, Reed R, Boyer A et al. 2019. Inhibition of Nuclear PTEN Tyrosine

- Phosphorylation Enhances Glioma Radiation Sensitivity through Attenuated DNA Repair. *Cancer Cell* **35**: 504-518 e507.
- Maccario H, Perera NM, Davidson L, Downes CP, Leslie NR. 2007. PTEN is destabilized by phosphorylation on Thr366. *Biochem J* **405**: 439-444.
- Maccario H, Perera NM, Gray A, Downes CP, Leslie NR. 2010. Ubiquitination of PTEN (phosphatase and tensin homolog) inhibits phosphatase activity and is enhanced by membrane targeting and hyperosmotic stress. *J Biol Chem* **285**: 12620-12628.
- Maddika S, Kavela S, Rani N, Palicharla VR, Pokorny JL, Sarkaria JN, Chen J. 2011. WWP2 is an E3 ubiquitin ligase for PTEN. *Nat Cell Biol* **13**: 730-735.
- Maehama T, Dixon JE. 1999. PTEN: a tumour suppressor that functions as a phospholipid phosphatase. *Trends Cell Biol* **9**: 125-128.
- Malaney P, Uversky VN, Dave V. 2017. PTEN proteoforms in biology and disease. *Cell Mol Life Sci* **74**: 2783-2794.
- Mehenni H, Lin-Marq N, Buchet-Poyau K, Reymond A, Collart MA, Picard D, Antonarakis SE. 2005. LKB1 interacts with and phosphorylates PTEN: a functional link between two proteins involved in cancer predisposing syndromes. *Hum Mol Genet* **14**: 2209–2219.
- Meng Z, Jia LF, Gan YH. 2016. PTEN activation through K163 acetylation by inhibiting HDAC6 contributes to tumour inhibition. *Oncogene* **35**: 2333-2344.
- Minaguchi T, Waite KA, Eng C. 2006. Nuclear localization of PTEN is regulated by Ca(2+) through a tyrosil phosphorylation-independent conformational modification in major vault protein. *Cancer Res* **66**: 11677-11682.
- Misticone S, Lima-Fernandes E, Scott MG. 2016. Rapid Detection of Dynamic PTEN Regulation in Living Cells Using Intramolecular BRET. *Methods Mol Biol* **1388**: 95-110.
- Mund T, Pelham HR. 2010. Regulation of PTEN/Akt and MAP kinase signaling pathways by the ubiquitin ligase activators Ndfip1 and Ndfip2. *Proc Natl Acad Sci U S A* **107**: 11429-11434.
- Nagata Y, Lan KH, Zhou X, Tan M, Esteva FJ, Sahin AA, Klos KS, Li P, Monia BP, Nguyen NT et al. 2004. PTEN activation contributes to tumor inhibition by trastuzumab, and loss of PTEN predicts trastuzumab resistance in patients. *Cancer cell* **6**: 117-127.
- Nakahata S, Ichikawa T, Maneesaay P, Saito Y, Nagai K, Tamura T, Manachai N, Yamakawa N, Hamasaki M, Kitabayashi I et al. 2014. Loss of NDRG2 expression activates PI3K-AKT signalling via PTEN phosphorylation in ATLL and other cancers. *Nat Commun* **5**: 3393.
- Nakakido M, Deng Z, Suzuki T, Dohmae N, Nakamura Y, Hamamoto R. 2015. Dysregulation of AKT Pathway by SMYD2-Mediated Lysine Methylation on PTEN. *Neoplasia* **17**: 367-373.
- Nguyen HN, Yang JM, Afkari Y, Park BH, Sesaki H, Devreotes PN, Iijima M. 2014. Engineering ePTEN, an enhanced PTEN with increased tumor suppressor activities. *Proc Natl Acad Sci U S A* **111**: E2684-2693.
- Nguyen HN, Yang JM, Jr., Rahdar M, Keniry M, Swaney KF, Parsons R, Park BH, Sesaki H, Devreotes PN, Iijima M. 2015. A new class of cancer-associated PTEN mutations defined by membrane translocation defects. *Oncogene* **34**: 3737-3743.
- Numajiri N, Takasawa K, Nishiya T, Tanaka H, Ohno K, Hayakawa W, Asada M, Matsuda H, Azumi K, Kamata H et al. 2011. On-off system for PI3-kinase-Akt

- signaling through S-nitrosylation of phosphatase with sequence homology to tensin (PTEN). *Proc Natl Acad Sci U S A* **108**: 10349-10354.
- Odrizola L, Singh G, Hoang T, Chan AM. 2007. Regulation of PTEN activity by its carboxyl-terminal autoinhibitory domain. *J Biol Chem* **282**: 23306-23315.
- Okumura K, Mendoza M, Bachoo RM, DePinho RA, Cavenee WK, Furnari FB. 2006. PCAF modulates PTEN activity. *J Biol Chem* **281**: 26562-26568.
- Papa A, Wan L, Bonora M, Salmena L, Song MS, Hobbs RM, Lunardi A, Webster K, Ng C, Newton RH et al. 2014. Cancer-associated PTEN mutants act in a dominant-negative manner to suppress PTEN protein function. *Cell* **157**: 595-610.
- Park MK, Yao Y, Xia W, Setijono SR, Kim JH, Vila IK, Chiu HH, Wu Y, Billalabeitia EG, Lee MG et al. 2019. PTEN self-regulates through USP11 via the PI3K-FOXO pathway to stabilize tumor suppression. *Nat Commun* **10**: 636.
- Podszypanina K, Ellenson LH, Nemes A, Gu J, Tamura M, Yamada KM, Cordon-Cardo C, Catoretti G, Fisher PE, Parsons R. 1999. Mutation of Pten/Mmac1 in mice causes neoplasia in multiple organ systems. *Proc Natl Acad Sci U S A* **96**: 1563-1568.
- Raftopoulou M, Etienne-Manneville S, Self A, Nicholls S, Hall A. 2004. Regulation of cell migration by the C2 domain of the tumor suppressor PTEN. *Science* **303**: 1179-1181.
- Rahdar M, Inoue T, Meyer T, Zhang J, Vazquez F, Devreotes PN. 2009. A phosphorylation-dependent intramolecular interaction regulates the membrane association and activity of the tumor suppressor PTEN. *Proc Natl Acad Sci U S A* **106**: 480-485.
- Riquelme SA, Hopkins BD, Wolfe AL, DiMango E, Kitur K, Parsons R, Prince A. 2017. Cystic Fibrosis Transmembrane Conductance Regulator Attaches Tumor Suppressor PTEN to the Membrane and Promotes Anti *Pseudomonas aeruginosa* Immunity. *Immunity* **47**: 1169-1181 e1167.
- Salmena L, Carracedo A, Pandolfi PP. 2008. Tenets of PTEN tumor suppression. *Cell* **133**: 403-414.
- Sanchez T, Thangada S, Wu MT, Kontos CD, Wu D, Wu H, Hla T. 2005. PTEN as an effector in the signaling of antimigratory G protein-coupled receptor. *Proc Natl Acad Sci U S A* **102**: 4312-4317.
- Shen SM, Guo M, Xiong Z, Yu Y, Zhao XY, Zhang FF, Chen GQ. 2015. AIF inhibits tumor metastasis by protecting PTEN from oxidation. *EMBO Rep* **16**: 1563-1580.
- Shih M-C, Chen J-Y, Wu Y-C, Jan Y-H, Yang B-M, Lu P-J, Cheng H-C, Huang M-S, Yang C-J, Hsiao M, et al. 2012. TOPK/PBK promotes cell migration via modulation of the PI3K/PTEN/AKT pathway and is associated with poor prognosis in lung cancer. *Oncogene* **31**: 2389–2400.
- Shi Y, Wang J, Chandrapaty S, Cross J, Thompson C, Rosen N, Jiang X. 2014. PTEN is a protein tyrosine phosphatase for IRS1. *Nat Struct Mol Biol* **21**: 522-527.
- Shinde SR, Gangula NR, Kavela S, Pandey V, Maddika S. 2013. TOPK and PTEN participate in CHFR mediated mitotic checkpoint. *Cell Signal* **25**: 2511–2517.
- Shnitsar I, Bashkurov M, Masson GR, Ogunjimi AA, Mosessian S, Cabeza EA, Hirsch CL, Trcka D, Gish G, Jiao J et al. 2015. PTEN regulates cilia through Dishevelled. *Nat Commun* **6**: 8388.
- Silva A, Yunes JA, Cardoso BA, Martins LR, Jotta PY, Abecasis M, Nowill AE, Leslie NR, Cardoso AA, Barata JT. 2008. PTEN posttranslational inactivation and

- hyperactivation of the PI3K/Akt pathway sustain primary T cell leukemia viability. *J Clin Invest* **118**: 3762-3774.
- Song MS, Salmena L, Carracedo A, Egia A, Lo-Coco F, Teruya-Feldstein J, Pandolfi PP. 2008. The deubiquitylation and localization of PTEN are regulated by a HAUSP-PML network. *Nature* **455**: 813-817.
- Song MS, Carracedo A, Salmena L, Song SJ, Egia A, Malumbres M, Pandolfi PP. 2011. Nuclear PTEN regulates the APC-CDH1 tumor-suppressive complex in a phosphatase-independent manner. *Cell* **144**: 187-199.
- Song MS, Salmena L, Pandolfi PP. 2012. The functions and regulation of the PTEN tumour suppressor. *Nat Rev Mol Cell Biol* **13**: 283-296.
- Sotelo NS, Valiente M, Gil A, Pulido R. 2012. A functional network of the tumor suppressors APC, hDlg, and PTEN, that relies on recognition of specific PDZ-domains. *J Cell Biochem* **113**: 2661-2670.
- Stambolic V, Suzuki A, de la Pompa JL, Brothers GM, Mirtsos C, Sasaki T, Ruland J, Penninger JM, Siderovski DP, Mak TW. 1998. Negative regulation of PKB/Akt-dependent cell survival by the tumor suppressor PTEN. *Cell* **95**: 29-39.
- Steck PA, Pershouse MA, Jasser SA, Yung WK, Lin H, Ligon AH, Langford LA, Baumgard ML, Hattier T, Davis T et al. 1997. Identification of a candidate tumour suppressor gene, MMAC1, at chromosome 10q23.3 that is mutated in multiple advanced cancers. *Nat Genet* **15**: 356-362.
- Sumitomo M, Iwase A, Zheng R, Navarro D, Kaminetzky D, Shen R, Georgescu MM, Nanus DM. 2004. Synergy in tumor suppression by direct interaction of neutral endopeptidase with PTEN. *Cancer Cell* **5**: 67-78.
- Sun J, Li T, Zhao Y, Huang L, Sun H, Wu H, Jiang X. 2018. USP10 inhibits lung cancer cell growth and invasion by upregulating PTEN. *Mol Cell Biochem* **441**: 1-7.
- Suzuki A, de la Pompa JL, Stambolic V, Elia AJ, Sasaki T, del Barco Barrantes I, Ho A, Wakeham A, Itie A, Khoo W et al. 1998. High cancer susceptibility and embryonic lethality associated with mutation of the PTEN tumor suppressor gene in mice. *Curr Biol* **8**: 1169-1178.
- Takahashi Y, Morales FC, Kreimann EL, Georgescu MM. 2006. PTEN tumor suppressor associates with NHERF proteins to attenuate PDGF receptor signaling. *Embo J* **25**: 910-920.
- Teng DH, Hu R, Lin H, Davis T, Iliev D, Frye C, Swedlund B, Hansen KL, Vinson VL, Gumpfer KL et al. 1997. MMAC1/PTEN mutations in primary tumor specimens and tumor cell lines. *Cancer Res* **57**: 5221-5225.
- Tibarewal P, Zilidis G, Spinelli L, Schurch N, Maccario H, Gray A, Perera NM, Davidson L, Barton GJ, Leslie NR. 2012. PTEN protein phosphatase activity correlates with control of gene expression and invasion, a tumor-suppressing phenotype, but not with AKT activity. *Sci Signal* **5**: ra18.
- Tolkacheva T, Boddapati M, Sanfiz A, Tsuchida K, Kimmelman AC, Chan AM. 2001. Regulation of PTEN binding to MAGI-2 by two putative phosphorylation sites at threonine 382 and 383. *Cancer Res* **61**: 4985-4989.
- Torres J, Pulido R. 2001. The tumor suppressor PTEN is phosphorylated by the protein kinase CK2 at its C terminus. Implications for PTEN stability to proteasome-mediated degradation. *J Biol Chem* **276**: 993-998.
- Trotman LC, Niki M, Dotan ZA, Koutcher JA, Di Cristofano A, Xiao A, Khoo AS, Roy-Burman P, Greenberg NM, Van Dyke T et al. 2003. Pten dose dictates cancer progression in the prostate. *PLoS biology* **1**: E59.

- Trotman LC, Wang X, Alimonti A, Chen Z, Teruya-Feldstein J, Yang H, Pavletich NP, Carver BS, Cordon-Cardo C, Erdjument-Bromage H et al. 2007. Ubiquitination regulates PTEN nuclear import and tumor suppression. *Cell* **128**: 141-156.
- Tzenaki N, Aivaliotis M, Papakonstanti EA. 2015. Focal adhesion kinase phosphorylates the phosphatase and tensin homolog deleted on chromosome 10 under the control of p110delta phosphoinositide-3 kinase. *Faseb J* **29**: 4840-4852.
- Valiente M, Andres-Pons A, Gomar B, Torres J, Gil A, Tapparel C, Antonarakis SE, Pulido R. 2005. Binding of PTEN to specific PDZ domains contributes to PTEN protein stability and phosphorylation by microtubule-associated serine/threonine kinases. *J Biol Chem* **280**: 28936-28943.
- van Diepen MT, Parsons M, Downes CP, Leslie NR, Hindges R, Eickholt BJ. 2009. MyosinV controls PTEN function and neuronal cell size. *Nat Cell Biol* **11**: 1191-1196.
- Van Themsche C, Leblanc V, Parent S, Asselin E. 2009. X-linked inhibitor of apoptosis protein (XIAP) regulates PTEN ubiquitination, content, and compartmentalization. *J Biol Chem* **284**: 20462-20466.
- Vazquez F, Ramaswamy S, Nakamura N, Sellers WR. 2000. Phosphorylation of the PTEN tail regulates protein stability and function. *Mol Cell Biol* **20**: 5010-5018.
- Vazquez F, Grossman SR, Takahashi Y, Rokas MV, Nakamura N, Sellers WR. 2001. Phosphorylation of the PTEN tail acts as an inhibitory switch by preventing its recruitment into a protein complex. *J Biol Chem* **276**: 48627-48630.
- Wang X, Trotman LC, Koppie T, Alimonti A, Chen Z, Gao Z, Wang J, Erdjument-Bromage H, Tempst P, Cordon-Cardo C et al. 2007. NEDD4-1 is a proto-oncogenic ubiquitin ligase for PTEN. *Cell* **128**: 129-139.
- Wang X, Jiang X. 2008. Post-translational regulation of PTEN. *Oncogene* **27**: 5454-5463.
- Wang W, Chen Y, Wang S, Hu N, Cao Z, Tong T, Zhang X. 2014. PIAS α ligase enhances SUMO1 modification of PTEN protein as a SUMO E3 ligase. *J Biol Chem* **289**: 3217-3230.
- Wu X, Hepner K, Castelino-Prabhu S, Do D, Kaye MB, Yuan XJ, Wood J, Ross C, Sawyers CL, Whang YE. 2000. Evidence for regulation of the PTEN tumor suppressor by a membrane-localized multi-PDZ domain containing scaffold protein MAGI-2. *Proc Natl Acad Sci U S A* **97**: 4233-4238.
- Wu H, Feng W, Chen J, Chan LN, Huang S, Zhang M. 2007. PDZ domains of Par-3 as potential phosphoinositide signaling integrators. *Mol Cell* **28**: 886-898.
- Yehia L, Ngeow J, Eng C. 2019. PTEN-opathies: from biological insights to evidence-based precision medicine. *J Clin Invest* **129**: 452-464.
- Yim EK, Peng G, Dai H, Hu R, Li K, Lu Y, Mills GB, Meric-Bernstam F, Hennessey BT, Craven RJ et al. 2009. Rak functions as a tumor suppressor by regulating PTEN protein stability and function. *Cancer Cell* **15**: 304-314.
- Yuan L, Lv Y, Li H, Gao H, Song S, Zhang Y, Xing G, Kong X, Wang L, Li Y et al. 2015. Deubiquitylase OTUD3 regulates PTEN stability and suppresses tumorigenesis. *Nat Cell Biol* **17**: 1169-1181.
- Zhang J, Zhang P, Wei Y, Piao HL, Wang W, Maddika S, Wang M, Chen D, Sun Y, Hung MC et al. 2013. Deubiquitylation and stabilization of PTEN by USP13. *Nat Cell Biol* **15**: 1486-1494.
- Zhang Z, Hou SQ, He J, Gu T, Yin Y, Shen WH. 2016. PTEN regulates PLK1 and controls chromosomal stability during cell division. *Cell Cycle* **15**: 2476-2485.

Zhang J, Lee YR, Dang F, Gan W, Menon AV, Katon JM, Hsu CH, Asara JM, Tibarewal P, Leslie NR et al. 2019. PTEN Methylation by NSD2 Controls Cellular Sensitivity to DNA Damage. *Cancer Discov.*