



HAL
open science

BEHAVIOR, FEEDING AND GROWTH OF YOUNG LOLIGO FORBESI (CEPHALOPODA : MYOPSIDA) REARED IN THE LABORATORY

R T Hanlon, R F Hixon, P E Turk, P G Lee, W T Yang

► **To cite this version:**

R T Hanlon, R F Hixon, P E Turk, P G Lee, W T Yang. BEHAVIOR, FEEDING AND GROWTH OF YOUNG LOLIGO FORBESI (CEPHALOPODA : MYOPSIDA) REARED IN THE LABORATORY. Vie et Milieu / Life & Environment, 1985, pp.247. hal-03022189

HAL Id: hal-03022189

<https://hal.sorbonne-universite.fr/hal-03022189v1>

Submitted on 24 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

BEHAVIOR, FEEDING AND GROWTH OF YOUNG *LOLIGO FORBESI* (CEPHALOPODA : MYOPSIDA) REARED IN THE LABORATORY

R.T. HANLON, R.F. HIXON, P.E. TURK, P.G. LEE and W.T. YANG

The Marine Biomedical Institute
The University of Texas Medical Branch
Galveston, Texas 77550-2772, U.S.A.

CEPHALOPODA, *LOLIGO FORBESI*, REARING, JUVENILE BEHAVIOUR, GROWTH

Loligo forbesi is the largest known loliginid squid, an important biomedical research animal and a commercially fished species in the NE Atlantic. We reared this species from hatching to sexual maturity (140 mm ML) in 369 days at a mean temperature of 15 °C (s.d. 1.62). Eggs were obtained from England and France, air-shipped to Galveston, and hatched successfully (> 70 %) in both natural and artificial seawater in recirculating systems. The hatchlings were very large (3.2-4.0 mm ML) and were fed size-sorted zooplankton (mostly copepods) for the first 50 days. Mysidacean shrimps (1.3-5.3 mm) were mixed in with the zooplankton within 10 days and were the primary food from day 20 through 100; *Palaemonetes* sp. shrimp larvae (2.1-2.4 mm) were also eaten readily from day 10 to 80. One-week-old squids could capture and ingest prey of a wide size range, from less than 1 mm to 5.3 mm. Growth was rapid, exponential during the first two months. In 1983 (mean temperature 14 °C), the growth equation for live wet weight was :

$$\text{Weight} = 8.108e^{0.0541t} \quad (r^2 = 0.98)$$

and for live mantle length it was :

$$\text{Length} = 4.409e^{0.0182t} \quad (r^2 = 0.98)$$

This corresponds to approximately 5.4 % and 1.8 % increase in body weight and length per day, respectively. Slightly lower rates, 3.6 % daily weight increase and 1.3 % daily length increase, were obtained in a 1985 experiment (mean temperature 13.2 °C). In both experiments mortality was high : 94 % in 1983 and 97 % in 1985 by day 60, despite active feeding. By the time squids were approximately 12 mm ML and 0.2 g (approximately 30 — 60 days posthatching) they were capable of schooling together and maintaining their position in a current (2 cm/second). By comparison, *L. opalescens* in culture schooled at a size of 10 to 15 mm ML (Yang *et al.*, 1983, *Aquaculture* 31 : 77-88). These results indicate that during the first 2 months of life, wild *L. forbesi* are subject to distribution by currents and can feed upon a wide variety and size of prey.

Funding is acknowledged from NIH grant RR 01024 to R.T.H. and W.T.Y.

Reçu le 18 octobre 1985; received October 18, 1985
Accepté le 2 novembre 1985; accepted November 2, 1985