



Three-Dimensional Evolution of Mesoscale Anticyclones in the Lee of Crete

Artemis Ioannou, Alexandre Stegner, Franck Dumas, Briac Le Vu

► To cite this version:

Artemis Ioannou, Alexandre Stegner, Franck Dumas, Briac Le Vu. Three-Dimensional Evolution of Mesoscale Anticyclones in the Lee of Crete. *Frontiers in Marine Science*, 2020, 7, 10.3389/fmars.2020.609156 . hal-03028693

HAL Id: hal-03028693

<https://hal.sorbonne-universite.fr/hal-03028693>

Submitted on 27 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Three-dimensional evolution of mesoscale anticyclones in the lee of Crete

Artemis Ioannou^{1,*}, Alexandre Stegner¹, Franck Dumas² and Briac Le Vu¹

¹*Laboratoire de Météorologie Dynamique, CNRS, Ecole Polytechnique, Palaiseau, 91128, France*

²*Service Hydrographique et Océanographique de la Marine (SHOM), 13 rue du Chatellier, 29200 Brest, France*

Correspondence*:

Artemis Ioannou

innartemis@gmail.com

2 ABSTRACT

Motivated by the recurrent formation of mesoscale anticyclones in the southeast of Crete, we investigated with a high resolution model the response of the ocean to orographic wind jets driven by the Cretean mountain range. As shown in the dynamical process study of Ioannou et al. (2020) which uses a simplified shallow-water model, we confirm here, using the CROCO (Coastal and Regional Ocean COmmunity) model, that the main oceanic response to the Etesian wind forcing is the formation of mesoscale anticyclones. Moreover, we found that the intensity of the wind-induced Ekman pumping acting on the eddies, once they are formed, modulate their intensity. Among the various coastal anticyclones formed during summer and fall 2015, only one of them will correspond to a long lived structure which is similar to the Ierapetra Eddy detected in 2015 (IE2015) on the AVISO/DUACS products. Thanks, to the DYNED-Atlas data base we were able to perform a quantitative comparison of the vertical structure of such long-lived anticyclone between the numerical model and the in-situ measurements of the various Argo profilers trapped inside the eddy core. Even without assimilation or any nudging, the numerical model was able to reproduce correctly the formation period, the seasonal evolution and the vertical structure of the IE2015. The main discrepancy between the model and the altimetry observations is the dynamical intensity of the anticyclone. The characteristic eddy velocity, derived from the AVISO/DUACS product for the IE2015 is much lower than in the numerical model. This is probably due to the spatio temporal interpolation of the AVISO/DUACS altimetry products. More surprisingly, several coastal anticyclones were also formed in the model in the lee of Crete area during summer 2015 when the Etesian winds reach strong values. However, these coastal anticyclones respond differently to the wind forcing since they remain close to the coast, in shallow-waters, unlike the IE2015 which propagates offshore in deep water. The impact of the bottom friction or the coastal dissipation seems to limit the wind amplification of these coastal anticyclones.

Keywords: coastal eddies, wind-forced anticyclones, orographic wind forcing, Ierapetra eddies, Ekman pumping, island wakes

1 INTRODUCTION

Even if the generation of coastal eddies induced by orographic winds have been documented in several studies, it is still difficult to identify what are the main mechanisms that drive their dynamical characteristics (size and intensity) and their vertical extend. The simultaneous combination of several extra processes

(coastal currents, bottom friction, tides. . .) is often a source of complexity for the analysis of real wind-induced eddies. Such eddies could be very intense and/or long-lived, therefore they have a strong impact on the export of coastal nutrients or biogeochemical species into the open sea or the ocean.

The formation of both cyclonic and anticyclonic eddies was frequently observed in the lee of oceanic mountainous islands (Yoshida et al., 2010; Jia et al., 2011; Caldeira et al., 2014; Couvelard et al., 2012; Piedeleu et al., 2009; Kersalé et al., 2011; Barton et al., 2000; Jiménez et al., 2008; Caldeira and Marchesiello, 2002; Caldeira and Sangrà, 2012). The Hawaiian archipelago was one of the first case studies that required the use of high-resolution numerical models. The interaction between the North-Equatorial Current and the archipelago is enough to generate eddies, but the use of higher spatial ($1/4^\circ$ degrees instead of $1/2^\circ$) and temporal (daily instead of monthly) resolution of wind forcing for the regional models was shown to capture eddy intensities in agreement with the observations (Calil et al., 2008; Kersalé et al., 2011; Jia et al., 2011). For Madeira Island, both numerical simulations (Couvelard et al., 2012) and oceanic observations (Caldeira et al., 2014) indicate that the wind wake, induced by the mountain orography, could be the dominant mechanism of coastal eddy generation. For larger mountain chains, gaps or valleys could locally amplify the upstream synoptic winds and lead to strong wind-jets on the sea. The numerical study of Pullen et al. (2008) has shown that intensified wind jets in the lee of Mindoro and Luzon Islands induce the generation and the migration of a pair of counter-rotating oceanic eddies. In a similar way, the complex orography of Crete island acts as an obstacle for the wind propagation inducing channeling and deflection of the Etesian winds that impact the regional circulation in the south Aegean Sea and the Levantine basin. This study focuses on this specific area where intense coastal anticyclones are formed recurrently during the summer months (Larnicol et al., 1995; Matteoda and Glenn, 1996; Hamad et al., 2005, 2006; Taupier-Letage, 2008; Amitai et al., 2010; Menna et al., 2012; Mkhinini et al., 2014; Ioannou et al., 2017).

Kotroni et al. (2001) performed simulations with and without Crete and they concluded that, the Crete mountain ranges (three mountains in the row with height around 2000 m in Figure 1(a)) modify the Etesian intensity and pathways. The work of Bakun and Agostini (2001) extracts and computes the composite mean wind stress estimates for each one-half degree latitude-longitude quadrangle for the long-term mean seasonal cycle. This observational data-set confirms that the wind-stress curl drives an intense oceanic downwelling at the southeast tip of Crete. Miglietta et al. (2013) simulated the influence of the orography in the same area, capturing the lee waves patterns in the wakes of the Crete, Karpathos, Kasos and Rhodes islands. The statistical analysis of the monthly surface wind of the ERA-Interim reanalysis (at grid resolution of $1/12^\circ$) performed by Mkhinini et al. (2014) exhibits a seasonal correlation between strong negative wind stress curl and the formation of long-lived anticyclones in the eastern Mediterranean Sea. However, correlation does not imply causation and the recent work of Ioannou et al. (2020) provides a dynamical understanding for the formation of long-lived mesoscale anticyclones induced by a seasonal wind-jet that mimics the Etesian winds deflected by Crete island. This study shows that the oceanic response to a symmetric wind jet could be a symmetric dipole or a strongly asymmetric structure dominated by an intense and robust anticyclone. Since, the anticyclonic wind shear, for the mean summer Etesian wind jet, is two times larger than the cyclonic one, the asymmetry of the oceanic response is enhanced and the formation of large mesoscale anticyclones is expected to be favored in this area. Nevertheless, the reduce-gravity rotating shallow-water model used by Ioannou et al. (2020) might be too simple to reproduce the complexity of the oceanic response to the Etesian winds in the southeast of Crete. In order to better understand the different mechanisms involved in the formation of the real wind-induced anticyclones we performed a high-resolution numerical modelling, of the Mediterranean circulation, using the CROCO model forced by realistic winds from August 2012 to December 2016. The main advantage of such

high-resolution simulation is to describe the rapid dynamics of meso- and sub-mesoscale vortex structures and to have a precise view of their vertical structure. We focus especially on summer and fall 2015 when several Argo floats were present in this area and allowed for a quantitative comparison between the regional model and the in-situ observations. Our main goal is to investigate how the variability of the wind forcing, the complex bathymetry of the shelf or the local outflow impacts on the dynamical characteristics and the vertical extend of these coastal anticyclones.

The paper is organized as follows. In Section 2, we describe the various data set-used in this study, the ARPEGE winds and the DYNED-Atlas eddy data base, and the CROCO ocean model used for our realistic numerical simulations of the Mediterranean Sea in 2015 and 2016. Section 3 presents the dynamical characteristics and the vertical structure of the robust coastal anticyclones which formed at the southeast tip of Crete during summer and fall 2015. Throughout comparisons are carried out between the regional model and remote sensing or in-situ observations. We then discuss, in Section 4, the impact of various forcing on the vertical extend of these coastal anticyclones. Finally, we sum up our results and conclude in Section 5.

2 DATA & METHODS

2.1 Regional Wind forcing

We used the ARPEGE data-set to provide the most realistic wind-forcing for our regional simulations of the Mediterranean Sea. This data-set is based on 4-D variational assimilation of wind observations into the Meteo-France system of Forecast and Analysis ARPEGE. This reanalysis provides the atmospheric fields at high spatial ($1/10^\circ$) and temporal (hourly 1 h) resolution. To test the accuracy of the ARPEGE data-set in the Crete area and especially in the Kasos strait we collected regional wind speed data from three Meteorological stations of the Hellenic National Meteorological Service (HNMS) located on the islands of Crete, Kasos and Karpathos at heights 15 m, 17 m and 114 m respectively. We first build the time series of the mean wind speed in the Kasos strait (inside the black circle of Figure 1(a)) and compare the temporal variability with the in-situ data of (HNMS). We found that the synoptic variability of the ARPEGE data in 2015 and 2016 is in good agreement with the local observations (see Figure 1 for summer 2015). However, if we compare the wind intensities we could find some local discrepancies. There is a correct agreement with Sitia weather station, which is located at the southeast of Crete, but a slight overestimation is found with the Karpathos station and an underestimation with the Kasos station. Hence, if the main components of the synoptic wind variability in the Kasos area are accurate in the ARPEGE data-set the local intensities of the surface winds, which are strongly impacted by the complex orography of Crete, should always be taken with care. Nevertheless, as far as we know, this is the best wind data-set available at high resolution for this specific area in 2015 and 2016. The ALADIN data-set used by Mkhinini et al. (2014) have an slightly higher resolution but it ends in 2012.

2.2 CROCO Ocean Model

We use outputs of realistic numerical simulations that were carried out for the Mediterranean Sea using the CROCO numerical model (<http://www.croco-ocean.org>). We refer to (Shchepetkin and McWilliams, 2005), Debreu et al. (2012) as well as to (Auclair et al., 2018) for details regarding the CROCO inherited numerics from ROMS, its barotropic time-stepping set-up and its solver. The simulation under investigation, CROCO-MED60v40-2015, was forced at the ocean top with ARPEGE wind forcing, thanks to the classical bulk COARE formula (Fairall et al., 2003) that takes into account the wind stress acting on the ocean surface as

$$\tau(x, y, t) = \rho_{air} C_d |\underline{U}| \underline{U} \quad (1)$$

where $\rho_{air} = 1.25 \text{ kg m}^{-3}$ is the air density, C_d the drag coefficient that varies based on the exchanges between the atmosphere-ocean turbulent surface heat fluxes and $\underline{U}$ the surface wind. The model configuration solves the classical primitive equations in an horizontal resolution of $1/60^\circ$ in both longitudinal and latitudinal directions, a well fitted resolution to capture the dynamics of interest. The vertical coordinate used is a generalised terrain following one. It is a stretched coordinate that allows to keep flat levels near the surface whatever the bathymetry gradient. Forty unevenly distributed vertical levels discretized the water column. They are closer one from each other next to the surface and more spaced by the bottom where the vertical gradients of hydrology parameters (temperature or salinity) are weak. This distribution was designed in order to properly catch the intense surface dynamics. Moreover, the bathymetry has been produced at SHOM for modeling purposes (10.12770/50b46a9f-0c4c-4168-9d1c-da33cf7ee188) and was built up from DTM at 100m and 500m resolution, that was optimally interpolated at first and then smoothed to control the pressure gradient truncation error associated with the terrain following coordinate system (Shchepetkin and McWilliams (2003)). The initial and boundary conditions were built from CMEMS global system analysis optimally interpolated on the computational grid. CROCO-MED60v40-2015 is a result of a free run simulation (no nudging nor assimilation of any kind) that started on the 1st of August 2012 when the water column stability is at its maximum to avoid static instability in the spinning up phase of three years. It ran till the end of December 2016. For the purposes of this paper, we extracted oceanic numerical fields for the year 2015. To track and quantify full trajectories of mesoscale eddies reproduced in the model, we used AMEDA eddy detection algorithm (Le Vu et al., 2018). Adapted to CROCO $1/60^\circ$ numerical fields, AMEDA can identify the eddy characteristics from the daily mean surface geostrophic velocities derived from Sea Surface Height of the model averaged during 24h.

2.3 Eddy database DYNEDAtlas

In order to compare the mesoscale eddies formed in the southeast of Crete in the regional simulation CROCO-MED60v40-2015 with both remote sensing and in-situ observations, we used the dynamical eddy data-base DYNED-Atlas (<https://www.lmd.polytechnique.fr/dyned/>). This recent data-base provides 17 years (2000-2017) of eddy detection and tracking in the Mediterranean Sea along with the co-localisation of Argo floats for each detected eddy (<https://doi.org/10.14768/2019130201.2>). The dynamical characteristics of the eddies contained in the DYNED-Atlas database were computed by the AMEDA eddy detection algorithm (Le Vu et al., 2018) applied on daily surface velocity fields. The latter were derived from the Absolute Dynamic Topography (ADT) maps produced by Salto/Duacs and distributed by CMEMS with a spatial resolution of $1/8^\circ$ which is much coarser than the spatial resolution of the numerical simulations. Hence, we will compare in this study only the characteristics of mesoscale eddies having a characteristic radius R_{max} (i.e. the radius where the azimuthal velocity V_{max} is maximal) higher than 15 km. In order to estimate the vertical structures of the detected eddies, DYNED-Atlas uses all the Argo profiles available since 2000 in the Mediterranean Sea. Once all the detected eddies are identified during the 2000-2017 period, we can separate the Argo profiles in two groups: the ones that are located inside an eddy (i.e. inside the last closed streamline) and the ones which are outside of all the detected eddies. With the second group we can build unperturbed climatological profiles (T, S and ρ) around a given position and a given date. We consider here all the Argo profiles (out of eddies) located at less than 150 km around the selected position and at ± 30 days from the target day during the 17 years. Such climatological

profiles (plotted in black) give a reference for the T, S and ρ profiles associated to an unperturbed ocean (i.e. without coherent eddies). Hence, the difference between this climatological density profile with the Argo profile taken inside an eddy allows us to compute the profile of the density anomaly $\sigma_A(z)$ ($kg\ m^{-3}$) and estimates its vertical extend as shown in Figure 2. We use the depth of the maximal density anomaly Z_{max} to quantify the vertical extension of the eddy. A similar methodology was used to estimate the depth of the coastal anticyclones in the regional simulation CROCO-MED60v40-2015 and perform quantitative comparisons with the DYNED-Atlas data. Since all the physical fields are available in the numerical model, the core eddy density profile corresponds to an average of all profiles located at less than 10 km from the eddy center. The background profile corresponds to an average of all the vertical profiles located along the last closed streamline.

3 RESULTS

3.1 Etesian wind-forcing and formation of coastal anticyclones

The Etesian winds blowing across the complex orography of Crete induce strong wind jets in its wake. As shown by Ioannou et al. (2020) such orographic winds could lead to the formation of long-lived mesoscale anticyclones in this area. We show in Figure 3 the seasonal variations of both the surface wind stress and the wind stress curl of the ARPEGE wind reanalysis for the year 2015. We note that the intensity of the negative wind stress curl in the Kasos strait (area inside the circle of Figure 3) is not strictly correlated to the wind intensity (Figure 3E,F). For this specific year, the maximum wind intensity occurs in February while intense negative wind stress curls occur in July. During the summer months, strong wind jets occur with a large area of negative Ekman pumping (deep blue area in Figure 3C) that extends a hundred of kilometers away from the Kasos strait and tends to favor the formation of coastal anticyclones. It can therefore be expected that the long-lived Ierapetra anticyclone (IE2015) will form in July or early August this year. However, this was not the case. Indeed, a long-lived anticyclone that survives more than six months was formed in late September in the numerical simulation CROCO-MED60v40-2015 while a similar eddy was detected in early October in the DYNED-Atlas database. Such long-lived and robust anticyclone, which is formed in the Southeast of Crete is usually called a Ierapetra anticyclone and will be labeled IE15 in what follows. Nevertheless, according to CROCO-MED60v40-2015, several other anticyclones were formed in the same area during summer 2015. These coastal anticyclones (labeled AE1, AE2 and AE3) were formed the 9 of June, the 14 of August and the 12 of September respectively (Figure 4). The lifetime of these robust eddies exceeds two months, but not three. This is still low compared to the IE15, which survives more than 15 months. Hence, the realistic simulation CROCO-MED60v40-2015 reveals that several coastal anticyclones are formed in the Kasos strait area when intense wind-jets, driven by the Etesian winds, occur. Among all these robust coastal anticyclones, only one will survive more than six months and will have the dynamical characteristics of an Ierapetra eddy.

3.2 Comparison between the CROCO model and the DYNED-Atlas data-base

A regional model that runs without assimilation, such as the CROCO-MED60v40-2015 is very unlikely to reproduce the exact dynamics and trajectory of mesoscale eddies. However, if the wind forcing is correct in the Kasos strait (as shown in the Figure 1), the wind-induced coastal anticyclones should have similar characteristics both in the model and the observations. Therefore, a systematic comparison is made between the dynamical eddy characteristics of the CROCO-MED60v40-2015 numerical model and the observations compiled in the Mediterranean eddy data-base: DYNED-Atlas. Besides, such analysis will help to quantify the dynamical differences between the numerous coastal anticyclones, which are formed during summer

months, and the long-lived Ierapetra Eddy (IE2015).

3.2.1 Dynamical characteristics and trajectories

The Figure 5 compares the temporal evolution of the characteristic radius (R_{max}) and the intensity (V_{max}) of the long-lived IE15 formed in CROCO-MED60v40-2015 with the IE2015 detected in DYNED-Atlas. These two mesoscale anticyclones were formed mid-fall at the end of September or early October, respectively. Since the spatial resolution of the numerical model ($1/60^\circ$) is seven times greater than that of merged altimetry products ($1/8^\circ$), it makes sense that the initial formation of such coastal eddy is better detected in the regional simulation CROCO-MED60v40-2015. If we assume that the model simulates correctly the IE formation, the AMEDA algorithm will detect it earlier in the regional model than in the coarse AVISO/CMEMS data set. In both cases, the radius of the IE15 and the IE2015 exceeds the local deformation radius ($R_d = 10 - 12 \text{ km}$) by at least a factor three (Figure 5 B). Such large radius is in good agreement with previous observations of Ierapetra Eddies (Hamad et al., 2006; Matteoda and Glenn, 1996; Taupier-Letage, 2008; Mkhinini et al., 2014; Ioannou et al., 2017, 2019).

However, the eddy intensity seems to reach higher values in the numerical model than in the eddy database. The maximal azimuthal velocity V_{max} could reach up to 70 cm/s in the CROCO-MED60v40-2015 while it never exceeds 40 cm/s in the DYNED-Atlas data base (Figure 5A). The underestimation of the IE's intensity in AVISO/CMEMS products, in comparison with in-situ measurements, was previously documented in Ioannou et al. (2017) and typical velocity values of 60 cm/s were confirmed by VMADCP measurements for IE eddies (Ioannou et al., 2017, 2019). The trajectory of the simulated IE15 and the observed one also differs even if both of them quickly propagate offshore 60 km south of the Kasos strait (Figure 5C).

As in the numerical model, a shorter-lived coastal anticyclone, that remained close to the shore in the southeast of Crete, was also detected the 29 of July according to the DYNED Atlas data-base. Such coastal anticyclone was detected from altimetry despite its decreased accuracy near the coast. The formation and the location of the short-lived anticyclone was also confirmed by a careful analysis of SST images. Hence, both remote sensing data sets, visible images and altimetry maps, show that coastal anticyclones could be formed in this area earlier during the summer months. We compare in the Figure 6, the dynamical characteristics of this coastal anticyclone detected in the DYNED-Atlas with the three structures formed by the regional simulation in June, August and early September. These anticyclones are smaller and weaker than the IE2015, their characteristic radius R_{max} does not exceed 25 km while the maximal azimuthal velocities V_{max} remain in the range $20 - 40 \text{ cm/s}$. More striking, they all seem to follow the same type of trajectory. The centers of these eddies remain attached to the coastline of Crete and the anticyclone stays above shelf even if they propagate westward, far away from their formation area (Figure 6C). Hence, the dynamical characteristics of these wind-induced eddies differ significantly from the long-lived Ierapetra anticyclone.

3.2.2 Comparison of Vertical eddy characteristics

The growing number of Argo floats deployed in the Mediterranean Sea in recent years makes it possible to characterize more precisely the three-dimensional evolution of long-lived eddies. Fortunately, the Ierapetra Eddy was sampled by several Argo profiles in the autumn of 2015 just after its formation and later on during winter 2016. The Figure 7 shows the temporal evolution of the density anomaly in the core of the Ierapetra anticyclone according to the Argo profiles taken in November 2015, in January 2016 and in February 2016. We select here only the profiles that were located at a distance of less than 35 km from

the eddy center (Figure 7A). The maximal density anomaly induced by the eddy on the climatological density background that contains no eddy signature was then estimated. As expected for an anticyclonic eddy, the density anomaly is negative. Moreover, we compute the depth of the maximal density anomaly Z_{max} (black dots in Figure 7(B,C,D)) to quantify the vertical extend of the IE2015.

According to Figure 7B, one month after its first detection, the density anomaly is confined between 50 m and 125 m, with a maximum anomaly of $\sigma_A = -1 \text{ kg m}^{-3}$ located at -100 m . Few months later, in January and February 2016, the maximal density anomaly propagated in depth, down to $Z_{max} = -150 \text{ m}$ and $Z_{max} = -225 \text{ m}$ respectively, but decreased in amplitude. The significant deepening of the IE2015, during the winter months, coincides with the seasonal deepening of the mixed layer depth in the Mediterranean Sea (Moschos et al., 2020)). During winter months, when the air-sea interactions are strong, the mixed layer could reach deeper value in the anticyclonic eddy core in comparison with the surrounding (Kouketsu et al., 2011; Dufois et al., 2017). We found that the mixed layer could go down to 200 m inside the Ierapetra eddy in February 2016. It is then very simple to quantify the vertical extend of the IE15 that is formed in the CROCO-MED60v40-2015 and compare them with the in-situ observations. Since we can track the eddy center with a high accuracy in the model, we can easily follow the temporal evolution of the density anomaly within the eddy core. The Figure 8 presents the monthly average of this anomaly for IE15 in comparison with the Argo profiles in November 2015, January 2016 and February 2016. The model is in correct agreement with the in-situ observations (Figure 8 B-D) and exhibits the same trend: a significant deepening of the IE15 during winter months. However, the maximal density anomaly reaches deeper values, down to $Z_{max} = -220 \text{ m}$ and $Z_{max} = -240 \text{ m}$ in January and February 2016, in CROCO-MED60v40-2015. The main advantage of such a realistic regional model is that it is possible to follow the three-dimensional evolution of all eddies and to compare them with each other. We could then check how the vertical structure of the coastal anticyclones, that are formed during summer months, differs from the long-lived Ierapetra anticyclone. The Figure 9 shows the temporal evolution of the size, the intensity and the vertical core density anomaly of one short-lived coastal anticyclone (AE3) in comparison with the IE15. We observe that during the initial stage of formation (the month that follows the first detection) these two types of anticyclones exhibit the same vertical structure and a moderate value of the radius R_{max} around 20 km. It is about a month later (in November 2015) that the Ierapetra anticyclone changes its structure: it increases in size and intensity as it expands in depth. Hence, it appears that the deepening of this long-lived anticyclone is induced by a dynamical process which is independent from its initial generation. The initial structure and the dynamical characteristics of the Ierapetra eddy, few weeks after its formation, does not differ significantly from the coastal anticyclones that are generated in summer by the wind-jet channelized by the Kasos strait. It is later on, that another mechanism leads to a drastic change in the vertical and the horizontal extend of the IE15.

4 DISCUSSION

Several coastal anticyclones were formed during summer and fall 2015 at the southeast tip of Crete, but only one of them will evolve into a large, deep and long-lived Ierapetra Eddy. Distinct physical processes could lead to this dynamical evolution. On one hand, the orographic wind-jets that occur in the wake of Crete induce strong and localized Ekman pumping. These upwelling or downwelling could then re-intensify or attenuate some coastal eddies. The intensification of a pre-existing mesoscale anticyclone was confirmed by in-situ observations (Ioannou et al., 2017) and idealized numerical simulations (Ioannou et al., 2020). If all these coastal anticyclones seem to be wind driven, their lifetime does not seem to be correlated with the wind-jet intensity in the Kasos strait and probably some more complex mechanisms should be considered to explain the robustness and the lifetime of the IE15. On the other hand, the Aegean outflow through the

287 Kasos strait (Kontoyiannis et al., 1999, 2005) may also contribute to the formation of coastal eddies or
 288 interact with them in this area and therefore modify their intensity and their vertical extend. Both processes
 289 are discussed in what follows.

290

291 4.1 Wind-eddy interactions

292 We first investigate the impact of local winds on coastal anticyclones once they are formed. We track
 293 these eddies with the AMEDA algorithm and compute for each of them the evolution of the, daily averaged,
 294 surface wind-stress inside the eddy contour. The local wind-stress curl will drive horizontal divergence
 295 and convergence of the Ekman transport and induce a mean vertical Ekman pumping inside the eddy
 296 (Ekman, 1905; Stern, 1965). The cumulative effect of this local Ekman pumping could lead to a significant
 297 isopycnal displacement. In order to take into account the core vorticity of the coastal anticyclones we use
 298 the non-linear relation derived by Stern (1965). Assuming a quasi-steady response (i.e. neglecting inertial
 299 waves generation) the additional isopycnal displacement $\Delta\eta$ induced by the cumulative wind-forcing is
 300 given by the following relation:

$$\Delta\eta = \int_{t_0}^t W_E dt = \int_{t_0}^t \frac{1}{A} \iint -\frac{1}{\rho} \nabla \times \left(\frac{\underline{\tau}}{f + \zeta} \right) dA dt \quad (2)$$

301 where $t = t_0$ is the beginning of the eddy detection, ρ the density of water, f the Coriolis parameter, ζ the
 302 vorticity within the eddy core and A the area enclosed by a radial distance of 1.5 of the maximum eddy
 303 contour R_{max} . The surface wind stress $\underline{\tau}$ is estimated by the bulk formula:

$$\underline{\tau} = \rho C_D V_{wind} \underline{V_{wind}} \quad (3)$$

304 where the drag coefficient C_d is set constant $C_d = 1.6 \cdot 10^{-3}$ and V_{wind} is the 10 m wind speed. We plot in
 305 Figure 10A the temporal evolution of the cumulative isopycnal displacement induced in the core of the
 306 three coastal anticyclones (AE1, AE2 and AE3) in comparison with the Ierapetra anticyclone IE15. For
 307 all the anticyclones, the vortex intensity V_{max} follows the temporal evolution of the cumulative Ekman
 308 pumping (Figure 10A,B). Indeed, the azimuthal velocity V_{max} of AE1, AE3 and IE2015 reached their
 309 highest values, respectively in July, November and December 2015 when the wind-induced isopycnal
 310 displacement reaches its maximum value for each eddy. For AE2, the wind-stress curl, in the eddy core, is
 311 zero or negative and therefore, unlike the other ones, the intensity of AE2 stays roughly constant in August
 312 and starts to decay in September. However, if the short-lived anticyclones AE1 and AE3 experienced a
 313 similar Ekman pumping than the long-lived IE2015, their intensity and their vertical extends differ strongly
 314 from Ierapetra 2015. Thus, for the same wind-stress curl amplitude, the dynamic response can be very
 315 different from one anticyclone to another. It seems that the intensity and vertical extent of the Ierapetra
 316 anticyclone is more strongly intensified by local wind forcing than of the other eddies.

317 One of the main differences between IE15 and other coastal anticyclones lies in their trajectories. Quite
 318 rapidly, after its formation, the Ierapetra eddy escapes from the shore and propagates into deep water
 319 unlike other eddies that travel along the Crete coast. The seabed under the eddies AE1, AE2 and AE3
 320 is between -600 and -1000 m, when the wind forcing is strong, while, in November-December 2015,
 321 when Ierapetra anticyclone intensifies, the seabed stays below -2000 m and may reaches -3000 m depth
 322 (Figure 10C). Hence, the bottom friction could be a possible explanation of the limitation of the intensity

and the isopycnal downwelling of these short-lived coastal eddies. Moreover, according to the Figure 10D, the characteristic eddy contours of AE1 and AE3 tangent the Crete coastline in July and November 2015 when the cumulative Ekman pumping is maximum for these eddies. The alongshore dissipation could also attenuate the wind induced intensification of these coastal eddies.

The temporal evolution of the vertical density structure and the cumulative isopycnal displacement $\eta = Z_{max}(t_0) + \Delta\eta$, induced by the local Ekman pumping, are shown in Figure 11 for AE1, AE2, AE3 and IE15 respectively. We find that the depth of maximal density anomaly Z_{max} follows roughly the evolution of η . Even if the numerical values are not strictly equal, these two characteristic depths are very close in the first months of the eddy lifetime. This correct agreement between the temporal evolution of $\Delta\eta$ (given by the Equation 2) and Z_{max} confirms that the local wind-stress curl drives the vertical structure of these anticyclones few months after their formation. However, we note for the Ierapetra eddy that during winter months (December, January and February) the density anomaly deepens while the isopycnal downwelling, induced by the wind-stress curl (i.e. $\Delta\eta$), does not increase. We also notice, during this period, that the intensity of the maximal density anomaly weakens from $\sigma_A = -1 \text{ kgm}^{-3}$ mid-November to $\sigma_A = -0.3 \text{ kgm}^{-3}$ mid-February. Such an evolution is probably due to air-sea fluxes at the surface that tend to extract a significant amount of heat from the mixed layer that deepens into the anticyclonic core. Such heat fluxes are not taken into account in the Equation 2. Hence, in addition to the local wind-shear, the air-sea fluxes could have a significant impact, especially during winter months, on the vertical structure of long-lived mesoscale anticyclones.

4.2 Kasos strait outflow

Another forcing mechanism that could generate strong anticyclonic eddies is the density bulge induced by river or strait outflows. A well-know example, at the entrance of the Mediterranean Sea, is the intense Alboran gyre which is forced by the fresh Atlantic water which enters through the Gibraltar strait. Such anticyclone is mainly driven by the amplitude of the inflow rather than the local wind forcing. It is therefore questionable whether the flow out of the Kasos Strait can control the formation of the long-lived Ierapetra anticyclones. Some snapshots of the surface circulation show that the jet corresponding to the Kasos strait outflow, seems to be connected with the periphery of the Ierapetra anticyclone (Figure 12A). Therefore, we first quantify the Kasos strait outflow and its variability in CROCO-MED60v40-2015 (Figure 12C) both in the surface ($0 - 200 \text{ m}$) and the subsurface layer (below -200 m). This outflow of lighter water coming from the Aegean Sea could be quite significant with a total flow rate that could exceed 2.4 Sv during few days, in July, August or November 2015. We then compare the variability of this outflow with the intensity of the coastal anticyclones (AE1, AE2, AE3) and the IE15 that stays in the vicinity of the Kasos strait (i.e. the area delimited by the black box in Figure 12A). We find that the intensification of the AE1 in July and the IE15 in November seems to be both correlated to the outflow intensity according to the Figure 12C,D. We should note that the eddy intensification is always associated to an amplification of the density anomaly (Figure 12E). However, the outflow is relatively strong during the whole period and we can also observe the intensification of AE3 without a significant change in the outflow of the Kasos strait. Hence, there is no systematic correlation between the variations of the outflow and the intensification (of V_{max} or σ_A) of pre-existing anticyclones in this area.

5 SUMMARY AND CONCLUSIONS

Using high-resolution ($1/60^\circ$) regional model CROCO-MED60v40, we analyzed the formation of a coastal eddy in the southeastern wake of Crete island. It is in this area that an intense, large -scale and long-lived

anticyclone is formed almost every year, commonly known as the Ierapetra eddy (Hamad et al., 2005, 2006; Taupier-Letage, 2008; Amitai et al., 2010; Menna et al., 2012; Mkhinini et al., 2014; Ioannou et al., 2017). Our previous studies (Mkhinini et al., 2014; Ioannou et al., 2020) have confirmed that the intensity of the summer wind jets (i.e. Etesian winds) blowing through the Kasos strait is one of the main reasons for the formation of robust anticyclones in this area. Motivated by the fact that the regional model CROCO-MED60v40 was driven by the ARPEGE hourly wind reanalysis (at $1/10^\circ$), which are the most accurate wind dataset in this region, we study the relation between the wind forcing and the dynamical characteristics of the wind-induced eddies. More specifically, we focus on the year 2015 where an Ierapetra eddy was formed both in the observational as well as in the numerical fields. During that year, Argo profilers were trapped for several months (October 2015 - April 2016) in the core of the Ierapetra anticyclone, allowing us to compare the evolution of the vertical structure of the anticyclone from the in-situ data with the numerical outputs.

Even without in-situ data assimilation, the numerical model was able to reproduce the formation and the dynamical evolution of a long-lived and robust anticyclone (IE15) similar to the Ierapetra eddy (IE2015) that was detected this specific year. The IE15 was formed at the end of September while the IE2015 was detected in early November according to the DYNED-Atlas eddy data-base. The temporal evolution of the characteristic radius and the vertical extend of the IE15 are also very close to the observations. However, the trajectories of these two eddies diverge after a few weeks and the eddy intensity reach higher values in the numerical model than in the eddy database. The latter could be due to a systematic under-evaluation of the eddy amplitude when we use the sea surface height or the surface velocity field derived from altimetry data-set. Indeed, similar under-evaluation in comparison with local VMADCP measurements were found by (Ioannou et al., 2017, 2019).

More surprisingly, according to CROCO-MED60v40-2015, several other anticyclones were formed in the same area during summer 2015 when the Etesian winds reach strong values. Three of them survive more than two months but their trajectories differ from the IE15. These coastal anticyclones travel along the Crete shelf while the Ierapetra eddy propagates offshore after its formation. A careful analysis of the DYNED-Atlas eddy data base reveals that a similar coastal anticyclone was also detected, during August and September 2015, on the standard AVISO/CMEMS Mediterranean altimetry data. Hence, such coastal anticyclones that propagate along the south coast of Crete are both present in the model and the observations. Thus, even if it does not exactly reproduce the observed ocean circulation (since there is no data assimilation), the regional simulation CROCO-MED60v40-2015 seems to provide a realistic description of the formation and the evolution of coastal eddies in the south of Crete island. Therefore, we can rely on this high-resolution model to study the impact of the orographic wind forcing on the formation and the subsequent evolution of realistic coastal anticyclones.

We do found that the intensity of the wind-induced Ekman pumping acting on the eddies, once they are formed, modulate their intensity. However, these coastal anticyclones respond differently to the wind forcing if they remain close to the coast, in shallow-waters, or if they propagate offshore in deep water. The impact of the bottom friction or the coastal dissipation seems to limit the wind amplification of coastal eddies. Among all the coastal anticyclones, which are formed by the summer intensification of the wind jet, only the one that escapes from the shelf will lead to a deep and long-lived eddy. Hence, a strong surface wind-jet is not enough to form an Ierapetra anticyclone and several others factors play a role. The wind-induced Ekman pumping should occur when the eddy is in deep waters and presumably the outflow from the Kasos strait reinforces this mechanism. However, these assumptions must be confirmed by a

409 longer numerical simulation that will allow to investigate the formation of different Ierapetra anticyclones
410 several years in a row.

ACKNOWLEDGMENTS

411 This work and especially A. Ioannou and B.Le Vu were funded by the Shom with research contract Catoobs
412 18CP01. The Hellenic National Meteorological Service (HNMS) is kindly acknowledged for providing the
413 wind observations used in this study.

DATA AVAILABILITY STATEMENT

414 Wind datasets used in the current study are available by <http://www.meteo.fr/> for the ARPEGE
415 dataset and by <http://www.hnms.gr> for the meteorological stations. The DYNED-Atlas data base for
416 the Mediterranean Sea (<https://doi.org/10.14768/2019130201.2>) is available at <https://www.lmd.polytechnique.fr/dyned/>.
417

AUTHOR CONTRIBUTIONS

418 A.Ioannou and A.Stegner designed the study, performed the data analysis and contributed to the
419 writing. F.Dumas performed the numerical simulation CROCO-MED60v40 and provided guidance in
420 the interpretation of the results of the numerical simulation. B.Le Vu, adapted the AMEDA algorithm to
421 perform the automatic eddy detection on the numerical simulations.

FIGURE CAPTIONS




Figure 1. A) Elevation map (m) of Crete island from ETOPO2 dataset. The locations of the meteorological stations (HNMS) of the Karpathos (A), Kasos (B) and Sitia (C) islands are displayed with the blue diamond points. The selected area for analyzing the ARPEGE wind forcing climatology is illustrated with the black circle of $R = 60$ km. Time-series of wind speed (m/s), as extracted from the 3 meteorological stations (Station A,B and C), are shown against the mean wind speed variations from ARPEGE data in panels B) and C).




Figure 2. A) Position of the IE2015 in November 2015. The eddy characteristic and last contour are illustrated with the blue solid and black dashed lines. The position of an Argo float trapped inside the eddy is shown with the magenta diamond. Vertical B) density σ and C) density anomaly σ_A profile of the Argo float trapped inside the IE2015 (magenta color). The black line shows the mean climatological profile as computed by Argo profiles that are detected outside of eddies.




Figure 3. Climatological wind stress $\langle \tau \rangle$ (vectors) and wind stress curl $\langle \nabla \times \tau \rangle$ (colors) during 2015 for the winter (A), spring (B) summer (C) and autumn (D) months based on ARPEGE wind data. The selected area for analyzing the wind forcing climatology is illustrated with the black circle in panel (B). The mean monthly climatological variations of wind stress $\langle \tau \rangle$ and wind stress curl $\langle \nabla \times \tau \rangle$ are shown in panels (E) and (F) respectively.




Figure 4. Time-series of A) wind speed (m/s) and B) negative wind stress curl $\langle \nabla \times \tau \rangle$ are shown with the orange color as extracted from ARPEGE wind data. The selected area for analyzing the wind forcing climatology is illustrated with the orange circle in panel (C). Trajectories of long-lived (> 2 weeks) eddies (gray colors) detected in CROCO model in the southeastern lee of Crete island during the summer & autumn of 2015. The first points of the eddy detection are plotted with the black square points while the points of last detection are illustrated with the open white circles. The longest-lived detected anticyclone is illustrated with the thick black line.




Figure 5. A) Temporal evolution of long-lived Ierapetra anticyclones as detected in 2015 with AMEDA algorithm from DYNED-Atlas eddy database and from CROCO-MED60v40-2015 simulation. The dynamical characteristics of the eddy velocity V_{max} (m/s) and eddy radius R_{max} (km) are shown in panels A) and B) respectively while their trajectories are illustrated in panel C).




Figure 6. A) Temporal evolution of short-lived coastal anticyclones detected in 2015 with AMEDA algorithm from DYNED-Atlas eddy database and from CROCO-MED60v40-2015 simulation. The dynamical characteristics of the eddy, its velocity V_{max} (m/s) and radius R_{max} (km) are shown in panels A) and B) respectively while their trajectories are illustrated in panel C). The evolution and the trajectory of the anticyclones AE1, AE2 and AE3 tracked in the regional model are plotted with blue lines while the observed anticyclone (DYNED-Atlas data-base) is plotted with a black line.




Figure 7. A) Temporal evolution of the IE2015 eddy characteristic contour (blue line) and the last contour (black dashed line) as extracted from the DYNED-Atlas eddy database. The squared points illustrate the position of the Argo profiles as a function of their distance from the eddy center. Vertical profiles of density anomaly $\sigma_A (\text{kgm}^{-3})$ are shown for the months of November (B), January (C) and February (D) as obtained from the Argo float profiles that were trapped in the IE2015.




Figure 8. A) Temporal evolution of the IE15 eddy characteristic contour (solid line) and the last contour (black dashed line) computed with AMEDA algorithm applied on CROCO geostrophic fields. Vertical profiles of density anomaly σ_A (kgm^{-3}) for the months of November (B), January (C) and February (D) as obtained from the Argo float profiles that were trapped in the IE2015 (shown with magenta color) and as obtained from the IE15 in CROCO-MED60v40-2015 simulation for the same period (shown with the blue color).




Figure 9. Temporal evolution of dynamical characteristics of A) velocity V_{max} (m/s) and B) radius R_{max} (km) for the coastal eddies AE3 and IE15 as computed with AMEDA algorithm applied on CROCO geostrophic fields. Comparison between the vertical profiles of density anomaly σ_A ($kg m^{-3}$) are shown for the months of September (C), October (D) and November (D) for the two anticyclones AE3 and IE15 from the CROCO-MED60v40-2015 simulation.




Figure 10. Comparison between the temporal evolution of the anticyclones AE1, AE2, AE3 and IE15. The isopycnal displacements after generation $\Delta\eta$ (m), associated with the wind stress curl $\langle \nabla \times \tau \rangle$ extracted above the eddies (in an area of 30 km from the eddy center) are illustrated in panel A). The temporal evolution of the eddy characteristic velocities V_{max} and the mean depth along the eddy trajectories are shown in panels B) and C) for the different anticyclones. The eddy trajectories of the different anticyclones are shown in panel D). The background map corresponds to the CROCO model bathymetry.




Figure 11. Comparison between the vertical profiles of density anomaly σ_A (kg m^{-3}) for the AE1, AE2, AE3 and IE15 anticyclones generated during summer and fall period in the CROCO-MED60v40-2015 simulation southeast of Crete. The temporal evolution of the minimum density anomaly Z_{max} and the isopycnal displacement $\Delta\eta$ associated with the wind stress curl above the eddies are illustrated with the black and blue lines respectively.




Figure 12. A) Surface vorticity fields ζ/f and B) vertical section of the density σ (kg/m^{-3}) in the Kasos Strait for the 11 November 2015. Temporal evolution of the Kasos Strait outflow Q separated in surface ($0 - 200\text{ m}$) and subsurface outflow ($200 - 700\text{ m}$) in panel C). The dates shown with the vertical black lines indicate the first detection of long-lived anticyclones during the same period. The temporal evolution of the eddy characteristic velocity V_{max} and the density anomaly σ_A is shown for the different anticyclones AE1, AE2, AE3 and IE15 in panels D) and E).

REFERENCES

- Amitai, Y., Lehahn, Y., Lazar, A., and Heifetz, E. (2010). Surface circulation of the eastern mediterranean levantine basin: insights from analyzing 14 years of satellite altimetry data. *Journal of Geophysical Research* 115, C10058. doi:10.1029/2010JC006147
- Auclair, F., Bordoio, L., Dossmann, Y., Duhaut, T., Paci, A., Ulses, C., et al. (2018). A non-hydrostatic non-boussinesq algorithm for free-surface ocean modelling. *Ocean Modelling* 132, 12–29. doi:https://doi.org/10.1016/j.ocemod.2018.07.011
- Bakun, A. and Agostini, V. N. (2001). Seasonal patterns of wind-induced upwelling/downwelling in the mediterranean sea. *SCIENTIA MARINA* 65, 243–257. doi:10.3989/scimar.2001.65n3243
- Barton, E., Basterretxea, G., Flament, P., Mitchelson-Jacob, E., B.Jones, Aristegui, J., et al. (2000). Lee region of gran ca-naria. *Journal of Geophysical Research* 105, 17173–17193
- Caldeira, R. and Sangrà, P. (2012). Complex geophysical wake flows madeira archipelago case study. *Journal of Ocean Dynamics* 62, 683–700. doi:10.1007/s10236-012-0528-6
- Caldeira, R. M. A. and Marchesiello, P. (2002). Ocean response to wind sheltering in the southern california bight. *Journal of Geophysical Research Letters* 29, 1–4. doi:10.1029/2001GL014563
- Caldeira, R. M. A., Stegner, A., Couvelard, X., Araújo, I. B., Testor, P., and Lorenzo, A. (2014). Evolution of an oceanic anticyclone in the lee of madeira island: In situ and remote sensing survey. *Journal of Geophysical Research: Oceans* 119, 1195–1216. doi:10.1002/2013JC009493
- Calil, P. H., Richards, K. J., Jia, Y., and Bidigare, R. R. (2008). Eddy activity in the lee of the hawaiian islands. *Deep-Sea Research II* 55, 1179–1194. doi:10.1016/j.dsr2.2008.01.008
- Couvelard, X., Caldeira, R., Araújo, I., and Tomé, R. (2012). Wind mediated vorticity-generation and eddy-confinement, leeward of the madeira island: 2008 numerical case study. *Journal of Atmospheres and Oceans* 58, 128–149
- Debreu, L., Marchesiello, P., Penven, P., and Cambon, G. (2012). Two-way nesting in split-explicit ocean models: Algorithms, implementation and validation. *Ocean Modelling* 49–50, 1–21. doi:10.1016/j.ocemod.2012.03.003
- Dufois, F., Hardman-Mountford, N. J., Fernandes, M., Wojtasiewicz, B., Shenoy, D., Slawinski, D., et al. (2017). Observational insights into chlorophyll distributions of subtropical south indian ocean eddies. *Geophysical Research Letters* 44, 3255–3264. doi:10.1002/2016GL072371
- Ekman, W. (1905). On the influence of the earth's rotation on ocean-currents. *Arkiv Fur Matematik Astronomi Och Fysik*. 11, 355–367
- Fairall, C. W., Bradley, E. F., Hare, J. E., Grachev, A. A., and Edson, J. B. (2003). Bulk parameterization of air-sea fluxes: Updates and verification for the coare algorithm. *Journal of Climate* 16, 571–591
- Hamad, N., Millot, C., and Taupier-Letage, I. (2005). A new hypothesis about the surface circulation in the eastern basin of the mediterranean sea. *Progress in Oceanography* 66, 287–298. doi:10.1016/j.pocean.2005.04.002
- Hamad, N., Millot, C., and Taupier-Letage, I. (2006). The surface circulation in the eastern basin of the mediterranean sea. *SCIENTIA MARINA* 70, 457–503
- Ioannou, A., Stegner, A., Dubos, T., Le Vu, B., and Speich, S. (2020). Generation and intensification of mesoscale anticyclones by orographic wind jets: The case of ierapetra eddies forced by the etesians. *Journal of Geophysical Research: Oceans* 125. doi:10.1029/2019JC015810
- Ioannou, A., Stegner, A., Le Vu, B., Taupier-Letage, I., and Speich, S. (2017). Dynamical evolution of intense ierapetra eddies on a 22 year long period. *Journal of Geophysical Research: Oceans* 122, 9276–9298. doi:10.1002/2017jc013158

- Ioannou, A., Stegner, A., Tuel, A., Le Vu, B., Dumas, F., and Speich, S. (2019). Cyclostrophic corrections of AVISO/DUACS surface velocities and its application to mesoscale eddies in the mediterranean sea. *Journal of Geophysical Research: Oceans* 124, 8913–8932. doi:10.1029/2019jc015031
- Jia, Y., Calil, P. H. R., Chassignet, E. P., Metzger, E. J., Potemra, J. T., Richards, K. J., et al. (2011). Generation of mesoscale eddies in the lee of the hawaiian islands. *Journal of Geophysical Research* 116. doi:10.1029/2011JC007305
- Jiménez, B., Sangà, P., and Mason, E. (2008). A numerical study of the relative importance of wind and topographic forcing on oceanic eddy shedding by tall, deep water islands. *Journal of Ocean Modelling* 22, 146–157. doi:10.1016/j.ocemod.2008.02.004
- Kersalé, M., Doglioli, A. M., and Petrenko, A. A. (2011). Sensitivity study of the generation of mesoscale eddies in a numerical model of hawaii islands. *Ocean Science* 7, 227–291. doi:10.5194/os-7-277-2011
- Kontoyiannis, H., Balopoulos, E., Gotsis-Skretas, O., Pavlidou, A., Assimakopoulou, G., and Papageorgiou, E. (2005). The hydrology and biochemistry of the cretan straits (antikithira and kassos straits) revisited in the period june 1997–may 1998. *Journal of Marine Systems* 53, 37–57. doi:10.1016/j.jmarsys.2004.06.007
- Kontoyiannis, H., Theocharis, A., Balopoulos, E., Kioroglou, S., Papadopoulos, V., Collins, M., et al. (1999). Water fluxes through the cretan arc straits, eastern mediterranean sea: March 1994 to june 1995. *Progress in Oceanography* 44, 511 – 529. doi:https://doi.org/10.1016/S0079-6611(99)00044-0
- Kotroni, V., Lagouvardos, K., and Lalas, D. (2001). The effect of the island of crete on the etesian winds over the aegean sea. *Journal of the Royal Meteorological Society* 127, 1917–1937. doi:10.1002/qj.49712757604
- Kouketsu, S., Tomita, H., Oka, E., Hosoda, S., Kobayashi, T., and Sato, K. (2011). The role of meso-scale eddies in mixed layer deepening and mode water formation in the western north pacific. *Journal of Oceanography* 68, 63–77. doi:10.1007/s10872-011-0049-9
- Larnicol, G., Traon, P.-Y. L., Ayoub, N., and Mey, P. D. (1995). Mean sea level and surface circulation variability of the mediterranean sea from 2 years of TOPEX/POSEIDON altimetry. *Journal of Geophysical Research: Oceans* 100, 25163–25177. doi:10.1029/95JC01961
- Le Vu, B., Stegner, A., and Arsouze, T. (2018). Angular momentum eddy detection and tracking algorithm (ameda) and its application to coastal eddy formation. *J. Atmos. Oceanic Technol* 35, 739–762. doi:10.1175/JTECH-D-17-0010.1
- Matteoda, A. and Glenn, S. (1996). Observations of recurrent mesoscale eddies in the eastern mediterranean. *Journal of Geophysical Research* 101, 20687–20709. doi:10.1029/96JC01111
- Menna, M., Poulain, P., Zodiatis, G., and Gertman, I. (2012). On the surface circulation of the levantine sub-basin derived from lagrangian drifters and satellite altimetry data. *Deep-Sea Research* 65, 46–58. doi:10.1016/j.dsr.2012.02.008
- Miglietta, M. M., Zecchetto, S., and Biasio, F. D. (2013). A comparison of wrf model simulations with sar wind data in two case studies of orographic lee waves over the eastern mediterranean sea. *Journal of Atmospheric Research* 120–121, 127–146
- Mkhinini, N., Coimbra, A. L. S., Stegner, A., Arsouze, T., Taupier-Letage, I., and Béranger, K. (2014). Long-lived mesoscale eddies in the eastern mediterranean sea: analysis of 20 years of AVISO geostrophic velocities. *Journal of Geophysical Research: Oceans* 119, 8603–8626. doi:10.1002/2014JC010176
- Moschos, E., Stegner, A., Schwander, O., and Gallinari, P. (2020). Classification of eddy sea surface temperature signatures under cloud coverage. *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing* 13, 3437–3447. doi:10.1109/JSTARS.2020.3001830

- 509 Piedeleu, M., Sangrà, P., Sánchez-Vidal, A., Fabrès, J., Gordo, C., and Calafat, A. (2009). An observational
510 study of oceanic eddy generation mechanisms by tall deep-water islands (gran canaria). *Geophysical*
511 *Research Letters* 36, 1–5. doi:10.1029/2008GL037010
- 512 Pullen, J., Doyle, J. D., May, P., Chavanne, C., Flament, P., and Arnone, R. A. (2008). Monsoon
513 surges trigger oceanic eddy formation and propagation in the lee of the philippine islands. *Journal of*
514 *Geophysical Research Letters* 35, L07604. doi:10.1029/2007GL033109
- 515 Shchepetkin, A. F. and McWilliams, J. C. (2003). A method for computing horizontal pressure-gradient
516 force in an oceanic model with a nonaligned vertical coordinate. *Journal of Geophysical Research* 108.
517 doi:10.1029/2001jc001047
- 518 Shchepetkin, A. F. and McWilliams, J. C. (2005). The regional oceanic modeling system (ROMS): a split-
519 explicit, free-surface, topography-following-coordinate oceanic model. *Ocean Modelling* 9, 347–404.
520 doi:10.1016/j.ocemod.2004.08.002
- 521 Stern, M. E. (1965). Interaction of a uniform wind stress with a geostrophic vortex. *Deep Sea Research*
522 *and Oceanographic Abstracts* 12, 355–367
- 523 Taupier-Letage, I. (2008). On the use of thermal images for circulation studies: Applications to the eastern
524 mediterranean basin. In *Remote Sensing of the European Seas*, eds. V. Barale and M. Gade (Springer
525 Netherlands). 153–164. doi:10.1007/978-1-4020-6772-3_12
- 526 Yoshida, S., Qiu, B., and Hacker, P. (2010). Wind generated eddy characteristics in the lee of the island of
527 hawaii. *Journal of Geophysical Research: Oceans* 115, C03019. doi:10.1029/2009JC005417