



# Viral infection of prokaryotic plankton during early formation of the North West Atlantic Deep Water

Markus G Weinbauer, Christian Griebler, Hendrik van Aken, Gerhard J Herndl

## ► To cite this version:

Markus G Weinbauer, Christian Griebler, Hendrik van Aken, Gerhard J Herndl. Viral infection of prokaryotic plankton during early formation of the North West Atlantic Deep Water. *Aquatic Microbial Ecology*, 2020, 84, pp.175-189. 10.3354/ame01934 . hal-03033957

**HAL Id: hal-03033957**

**<https://hal.sorbonne-universite.fr/hal-03033957>**

Submitted on 1 Dec 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copy-edited by LD Todd

## **Viral infection of prokaryotic plankton during early formation of the North West Atlantic Deep Water**

Markus G. Weinbauer<sup>1,\*</sup>, Christian Griebler<sup>2</sup>, Hendrik M. van Aken<sup>3</sup>, Gerhard J. Herndl<sup>4,5</sup>

<sup>1</sup>Sorbonne Universités, UPMC Univ Paris 06, CNRS, Laboratoire d’Océanographie de Villefranche (LOV), 181 Chemin du Lazaret, 06230 Villefranche-sur-Mer, France

<sup>2</sup>Department of Limnology & Bio-Oceanography, University of Vienna, Althanstrasse 14, 1090 Vienna, Austria

<sup>3</sup>Department of Physical Oceanography, Royal Netherlands Institute for Sea Research (NIOZ), 1790 AB Den Burg, The Netherlands

<sup>4</sup>Department of Marine Biology, University of Vienna, Althanstrasse 14, 1090 Vienna, Austria

<sup>5</sup>Department of Biological Oceanography, Royal Netherlands Institute for Sea Research (NIOZ), 1790 AB Den Burg, The Netherlands

\*Corresponding author: [wein@obs-vlfr.fr](mailto:wein@obs-vlfr.fr)

Running head: Weinbauer et al.: Viruses in the North Atlantic Deep Water

**ABSTRACT:** Viral abundance was assessed in different water masses of the NW Atlantic, and the development of viral abundance, lytic viral infection and lysogeny was followed for the first ca. 5000 km (corresponding to ca. 50 yr in the oceanic conveyor belt) of the western branch of the North Atlantic Deep Water (NADW). Viral abundance was significantly higher in the 100 m layer than in the NADW (2400–2700 m depth) and the Denmark Strait Overflow Water (2400–3600 m depth). The virus-to-prokaryote ratio (VPR) increased with depth, ranging from 32–43 for different water masses of the bathypelagic ocean, thus corroborating the enigma of high viral abundance in the dark ocean. The O<sub>2</sub>-minimum layer (250–600 m) also showed high viral abundance and VPRs. Viral abundance, a viral subgroup and VPRs decreased in a non-linear form with distance from the NADW origin. Viral production (range: 0.2–2.4 × 10<sup>7</sup> viruses l<sup>-1</sup>) and the fraction of lytically

infected cells (range: 1–22%) decreased with increasing distance from the formation site of the NADW. Conservative estimations of virus-mediated mortality of prokaryotes in the NADW averaged  $20 \pm 12\%$ . The fraction of the prokaryotic community with lysogens (i.e. harboring a functional viral DNA) in the NADW averaged  $21 \pm 14\%$ . Hence, we conclude that (1) viral abundance and subgroups differ between water masses, (2) virus-mediated mortality of prokaryotes as well as lysogeny are significant in the dark ocean and (3) the lysogenic life strategy becomes more important than the lytic life style during the early formation of the NADW.

KEY WORDS: NADW · Thermohaline circulation · Dark ocean · Viral production · Lysogeny · Microorganisms

## 1. INTRODUCTION

It is now well accepted that viruses represent the most abundant ‘life forms’ in the ocean and that viral lysis is a major mortality factor for marine microorganisms in the upper ocean (e.g. (Wommack & Colwell 2000, Weinbauer 2004, Winter et al. 2010, Zimmerman et al. 2020)). Lysis not only causes cell death but also releases the cell contents as dissolved organic matter (DOM) and converts cell walls into small detritus; this viral shunt plays an important role in the cycling of carbon and nutrients (Gobler et al. 1997, Wilhelm & Suttle 1999, Brussaard et al. 2008, Weinbauer et al. 2009b) and in carbon export by the biological pump (Weinbauer et al. 2009a, Yamada et al. 2018).

During the last 1–2 decades, viral ecology of the dark ocean has been increasingly studied. Data sets are available on viral abundance (VA) and production (VP) in the water column of the dark realm of the Atlantic Ocean (Parada et al. 2007, De Corte et al. 2010, 2012, 2016, Muck et al. 2014, Winter et al. 2018), Pacific and Southern Ocean (Li et al. 2014, Yang et al. 2014), Mediterranean Sea (Winter et al. 2009, Umani et al. 2010) and from a circumnavigation expedition (Lara et al. 2017). Some of these studies reported that prokaryotic abundance (PA) decreased faster with depth than VA, thus resulting in a high virus-to-prokaryote ratio (VPR) in the bathypelagic and abyssopelagic ocean; however, there are also exceptions (Winter et al. 2009, Lara et al. 2017). This presents an enigma, since host abundances are 1–2 orders of magnitude lower in deep than in surface waters, which should—according to theory—result in a reduced contact rate between viruses and hosts and thus in reduced viral infection and production (Murray & Jackson 1992). However, it has been demonstrated that lytic VP is a major source of prokaryotic mortality in the dark ocean. Data on viral production and virus-mediated mortality in the dark ocean are available for the Chukchi Sea (Steward et al. 1996), Mediterranean Sea (Weinbauer et al. 2003, Winter et al.

2009, Umani et al. 2010), Atlantic Ocean (Parada et al. 2007, De Corte et al. 2010, 2012, Muck et al. 2014) and Pacific Ocean (Li et al. 2014, Yang et al. 2014).

Lysogeny, i.e. the phenomenon that prokaryotic cells can harbour a provirus (viral DNA) which can be induced by specific environmental conditions, has also been studied in the dark ocean (Weinbauer et al. 2003, De Corte et al. 2010, 2012). It is believed that lysogeny is established when the encounter rate between phages and host cells is low or when viral decay rates are high, i.e. in conditions unfavourable to hosts (e.g. Stewart & Levin 1984, Weinbauer 2004). Using isolates, it has been argued that a high host density destabilizes the interaction between a lytic phage and its host (Bohannan & Lenski 1997). Therefore, development of lysogeny would stabilize this interaction and support the survival of the phage line (Williams 1994). In this case, lysogeny might also be beneficial in high host-density environments (Weinbauer 2004). Indeed, it has been suggested that lysogeny can dominate when host abundance and production is high, i.e. in conditions favourable for hosts ('piggyback-the-winner' model; Knowles et al. 2016). Metagenomic studies suggest that provirus gene induction, and thus lysogeny, is the predominant life strategy in the deep sea (Williamson et al. 2008, Mizuno et al. 2016). Some studies have reported that lysogeny and inducible lysogenic VP typically increase with water depth compared to lytic VP; however, other studies have also reported the inverse trend. The inducing agents of lysogenic bacteria are not known for the dark ocean but it has been suggested that mixing of water masses can cause provirus induction (Winter et al. 2018).

Depth-related variations in prokaryotic biomass and production have been summarized (e.g. Nagata et al. 2000, Arístegui et al. 2009). In contrast to such depth trends, the functioning of the global ocean is primarily explained by the lateral transport of physically distinct water masses (thermohaline circulation) (Broecker 1997). Thermohaline circulation begins in the Greenland–Iceland–Norwegian (GIN) Sea. Differences in prokaryotic activity and community composition were investigated in distinct water masses of this North Atlantic Deep Water (NADW) during several cruises (Reinthal et al. 2006, Teira et al. 2006, Agogué et al. 2011). Here, we investigated VA and distinct viral groups as assessed by flow cytometry in the different water masses and followed the development of VA, VP, lytic viral infection and lysogeny in the western branch of the NADW during the TRNSAT-II cruise. The water mass was sampled from close to its formation for about 5500 km, thus covering approximately the first 50 yr of the NADW in the oceanic conveyor belt system.

## 2. MATERIALS AND METHODS

### 2.1. Study site and sampling

96 The western branch of the NADW was followed with the R/V 'Pelagia' from near its source  
97 of origin in the GIN Sea for over ca. 5000 km (Fig. 1). The TRANSAT-II cruise (May 2003)  
98 followed a track from 62.5° N, 30.3° W to 37.7° N, 69.7° W in the western basin of the North  
99 Atlantic, covering 34 stations (Fig. 1). The distance of the stations from the origin of the NADW  
100 was calculated using Ocean Data View (<http://odv.awi.de>). Water was collected with a CTD rosette  
101 sampler holding twenty four 12 l no oxygen exchange (NOEX) bottles. Samples were taken from  
102 100 m depth (subsurface layer [SSL]), the oxygen minimum zone and the main deep water masses  
103 encountered during the cruise. The main water masses sampled were the Labrador Sea Water  
104 (LSW), the NADW and the Denmark Strait Overflow Water (DSOW). These specific water masses  
105 were identified based on their temperature and salinity characteristics (see Table 1) and their  
106 oxygen concentrations, using a Seabird SBE43 oxygen sensor mounted on the CTD frame. For  
107 more details of the sampling and water mass characterization, see Reinthaler et al. (2006) and Teira  
108 et al. (2006). From these water masses, seawater samples were collected for physical-chemical  
109 parameters, PA, VA and VP and to estimate lytic and lysogenic infection.

## 110 2.2. Methods for data used from previous publications

111 The following data were obtained from previous publications on the TRANSAT-II cruise.  
112 For details and references see Reinthaler et al. (2006) and Teira et al. (2006). Briefly, apparent  
113 oxygen utilization (AOU) was calculated as the difference between the saturation oxygen  
114 concentration and the observed oxygen concentration. The concentrations of inorganic nutrients  
115 (NH<sub>4</sub>, NO<sub>3</sub>, NO<sub>2</sub> and PO<sub>4</sub>) were determined immediately after sample collection and gentle  
116 filtration through 0.2 mm filters (Acrodisc; Gelman Science) in a TRAACS autoanalyzer system.  
117 NH<sub>4</sub> was detected with the indo-phenol blue method (pH 10.5) at 630 nm. NO<sub>2</sub> was determined  
118 after diazotation with sulfanilamide and N-(1-naphtyl)-ethylene diammonium-dichloride as the  
119 reddish-purple dye complex at 540 nm. NO<sub>3</sub> was reduced in a copper cadmium coil to NO<sub>2</sub> (with  
120 imidazole as a buffer) and then measured as NO<sub>2</sub>. PO<sub>4</sub> was determined via the molybdenum blue  
121 complex at 880 nm.

122 For enumeration of heterotrophic prokaryotes, samples (1 ml) of unfiltered seawater were  
123 fixed with 37% 0.2 mm filtered (Acrodisc; Gelman) formaldehyde (2% final concentration), stained  
124 with 0.5 ml of SYBR Green I (Molecular Probes) at room temperature in the dark for 15 min and  
125 subsequently analyzed on a FACSCalibur flow cytometer (BD Biosciences). Counts were  
126 performed with an argon laser at 488 nm set at an energy output of 15 mW. Prokaryotic cells were  
127 enumerated according to their right-angle light scatter and green fluorescence measured at 530 nm.

128 Prokaryotic heterotrophic production (PHP) in seawater was measured by  $^3\text{H}$ -leucine  
129 incorporation (specific activity: 595.7 3 1010 Bq mmol<sup>-1</sup>; final concentration: 10 nmol l<sup>-1</sup>). Two  
130 10–40 ml samples and 1 blank were incubated in the dark. The blank was fixed with concentrated  
131 0.2 mm filtered formaldehyde (4% final concentration, v/v) 10 min prior to adding the tracer. After  
132 incubating the samples and the blank at *in situ* temperature for 4–12 h, depending on the expected  
133 activity, the samples were fixed with formaldehyde (4% final concentration), filtered onto 0.2 mm  
134 nitrocellulose filters (Millipore HA; 25 mm diameter) and rinsed twice with 5 ml ice-cold 5%  
135 trichloroacetic acid (Sigma Chemicals) for 5 min. The filters were dissolved in 1 ml ethylacetate,  
136 and after 10 min, 8 ml of scintillation cocktail (Insta-Gel Plus; Canberra Packard) was added. The  
137 radioactivity incorporated into cells was counted in a liquid scintillation counter (Model 1212; LKB  
138 Wallac). Leucine incorporated into prokaryotic biomass was converted to carbon production using  
139 the theoretical conversion factor of 3.1 kg C mol<sup>-1</sup> Leu, assuming a two-fold isotope dilution.

### 140 2.3. Enumeration of viruses

141 Water samples for viral enumeration were preserved with glutaraldehyde (0.5% final  
142 concentration) at 4°C for 30 min, then flash-frozen in liquid nitrogen and stored at –80°C until  
143 analysis. Virus samples were diluted 20-fold in autoclaved and 0.2 µm prefiltered TE buffer (10  
144 mmol l<sup>-1</sup> Tris, 1 mmol l<sup>-1</sup> EDTA, pH 8.0) and stained with SYBR Green I (Molecular Probes) (at a  
145 20000-fold dilution of the stock solution) in an 80°C water bath for 10 min before counting. Viruses  
146 were detected by their signatures in a side-scatter-versus-green-fluorescence (530 nm wavelength,  
147 fluorescence channel 1 of the instrument) plot and counted by flow cytometry (FACSCalibur; BD  
148 Biosciences) following the protocol of Brussaard et al. (2010). Data analysis was performed using  
149 BD Cell Quest Pro software version 4.0.2 (BD Biosciences). Viral subgroups V1, V2 and V3 were  
150 distinguished by increasing fluorescence intensity with settings for the different subgroups that were  
151 identical for all analyzed samples. The difference between replicates was typically better than 10%.

### 152 2.4. Burst size

153 To assess *in situ* burst size (BS; i.e. the number of viruses released upon cell lysis), 50 ml of  
154 the prokaryotic concentrates (see below) were preserved in glutaraldehyde (0.5% final  
155 concentration), kept briefly at 4°C and then stored at –80°C until analysis. Prokaryotic cells in  
156 thawed samples were collected by centrifugation onto formvar-coated transmission electron  
157 microscope (TEM) grids (copper, 400 mesh size) and stained with uranyl acetate (Weinbauer &  
158 Suttle 1999). Duplicate grids were used for each sample. The minimum BS (BS<sub>min</sub>) was estimated  
159 as the average from >20 visibly infected cells grid<sup>-1</sup>. This number is a conservative estimate

160 because viruses could still be assembled in the cells; therefore, a conversion was used to calculate  
161 maximum BS ( $BS_{\max}$ ):  $BS_{\max} = 1.41 \times BS_{\min} + 0.87$  (Parada et al. 2006).

## 162 2.5. VP and infection of prokaryotic plankton

163 VP, the fraction of infected cells (FIC) and the fraction of lysogenic cells (FLC) were  
164 estimated with a dilution technique (Wilhelm et al. 2002) using a modification described elsewhere  
165 (virus-reduction approach [VRA]: Weinbauer et al. 2002, 2010). Large water samples (150–200 l)  
166 were filtered through 0.8 mm pore-size polycarbonate filters (142 mm diameter; Millipore) and  
167 prokaryotes were concentrated using a Pellicon (Millipore) tangential flow filtration system  
168 equipped with a 0.2  $\mu\text{m}$  filter cartridge (Durapore; Millipore) as described in (Weinbauer et al.  
169 2009b). The first 20 l of the 0.2  $\mu\text{m}$  filtrate were processed with a 100 kDa cutoff polysulfone  
170 cartridge (Prep-Scale/TFF; Millipore: 0.23  $\text{m}^2$  nominal filter area, operated by a peristaltic pump at  
171 150000 Pa) to produce virus-free water. Aliquots of the prokaryote concentrate were added to virus-  
172 free water to obtain roughly *in situ* abundance assuming (based on previous findings) that half of  
173 the prokaryotes were lost during the prefiltration and ultrafiltration steps. This procedure reduces  
174 contact rates between viruses and hosts and thus new infection. Incubations were performed in the  
175 dark at *in situ* temperature (3.0°C) in duplicate 50 ml sterile conical tubes for 24 h. Samples were  
176 taken at incubation times ( $t$ ) 0, 6, 12, 18 and 24 h. VP was calculated as:

$$177 \quad VP = (VA_2 - VA_1) / (t_2 - t_1) \quad (1)$$

178 where  $VA_1$  and  $VA_2$  are the viral abundances at incubation times  $t_1$  and  $t_2$ , respectively. Note that  
179 individual incubations were treated separately and values at the start of incubations were not always  
180 used for calculations. Rather, the lowest viral abundance served as  $VA_1$  (Weinbauer et al. 2009b).  
181 Thus,  $VA_1$  and  $VA_2$  are the minimum and maximum of viral abundance in the incubation. VP was  
182 corrected for the changes in PA at the start of the experiment compared to *in situ* abundances.

183 Dividing the number of produced viruses by the BS yields the number of lysed cells and  
184 thus gives an estimate of FIC (Weinbauer et al. 2002), which was calculated by:

$$185 \quad FIC = 100(VA_2 - VA_1) / BS / PA \quad (2)$$

186 where PA is the prokaryotic abundance at the start of the experiment ( $t_1$ ). Virus-mediated mortality  
187 of prokaryotes (VMMP) was either calculated as:

$$188 \quad VMMP_{VP} = 100(VP / BS / PHP) \quad (3)$$

189 or using FIC values and the model of Binder (1999) ( $VMMP_{FIC}$ ).

190 The FIC treatment also served as a control in the lysogeny bioassays. To induce the lytic  
191 cycle in lysogenic cells (containing a prophage), samples were treated with mitomycin C (Sigma



192 Chemicals; final concentration: 0.5 µg ml<sup>-1</sup>; Paul & Weinbauer 2010). The difference in VA  
193 between this treatment and the control is the number of induced viruses, which is divided by the BS  
194 to estimate the number of induced cells and thus the FLC. FLC was calculated as percentage by:

195 
$$FLC = 100(V_{AMC} - V_{AC}) / BS / PA \quad (4)$$

196 where V<sub>AMC</sub> and V<sub>AC</sub> are the maximum difference in viral abundance at corresponding time points  
197 in mitomycin C and control treatments, respectively. Induced VP (VP<sub>i</sub>) was calculated analogous to  
198 VP after by subtracting V<sub>AC</sub> from V<sub>AMC</sub>.

## 199 **2.6. Statistics**

200 Spearman rank correlations were used to assess the covariation of parameters, since some  
201 variables did not comply with normality even after logarithmic transformation. The non-parametric  
202 Kruskal-Wallis and Mann-Whitney tests were used for comparing specific parameters obtained in  
203 different water masses, since normality was not always attained; p < 0.05 (after applying a  
204 Bonferroni correction) was considered significant. To test a potential change of viral parameters  
205 with distance from the GIN Sea, regressions with linear, logarithmic, exponential and power  
206 functions were calculated; p < 0.05 was considered significant. Statistics were performed with  
207 Aabel\_3.

## 208 **3. RESULTS**

### 209 **3.1. Characterization of water masses**

210 Some basic physical–chemical characteristics of the main water masses sampled during the  
211 study are given in Table 1. More details can be found elsewhere (Teira et al. 2006). The LSW,  
212 characterized by low salinity, was clearly identifiable at depths between 700 and 2100 m, except  
213 between 40 and 45° N. The NADW (2000–3000 m) was identifiable by its salinity maximum  
214 (34.90–34.95) south of 60° N. The DSOW underlying the NADW, with seawater temperature  
215 between 0.8 and 2.4°C and salinity < 34.90, was detected at all stations between 45 and 65° N. A  
216 local moderate oxygen minimum (ca. 30% less than in overlaying and underlaying water) was  
217 found between ca. 200 and 700 m depth in the southern part of the transect (from 40–50° N). Data  
218 were only used when these water masses could be clearly identified.

### 219 **3.2. VA and PA in different water masses; depth relationships**

220 The distribution of PA is shown in Fig. 2. PA decreased with depth by ca. an order of  
221 magnitude (Table 2), and this decrease was exponential (Reinthal et al. 2006). The various deep-  
222 water masses exhibited some differences in PA. PA was highest in the SSL (>100 m depth),



intermediate in the oxygen minimum zone and lowest in the bathypelagic water masses. These differences were significant (Kruskal-Wallis and Mann-Whitney tests,  $p < 0.05$ ; Table 2).

VA (Fig. 2) showed a local maximum in the oxygen minimum zone at ca.  $54^{\circ}$  N,  $47^{\circ}$  W (ca. 500 m water depth) and was slightly higher close to the GIN Sea. VA decreased significantly with depth (Table 3) and was, on average, highest in the SSL ( $3.1 \pm 1.8 \times 10^9 \text{ l}^{-1}$ ) and oxygen minimum zone ( $2.7 \pm 2.9 \times 10^9 \text{ l}^{-1}$ ) and lowest in the NADW ( $1.2 \pm 0.9 \times 10^9 \text{ l}^{-1}$ ) (Table 2). Significant differences were found between the SSL and the NADW/DSOW (Kruskal-Wallis and Mann-Whitney tests,  $p < 0.05$ ); however, differences between depth layers were less pronounced than for PA and were a maximum of 3.3-fold. The VPR also showed a local maximum in the oxygen minimum zone at  $47.6^{\circ}$  W,  $53.5^{\circ}$  W and was also highest towards the GIN Sea. In addition, VPR increased significantly with depth (Table 3). VPR was lowest in the SSL and highest in the DSOW; VPR was significantly lower in the SSL than in bathypelagic waters (Kruskal-Wallis and Mann-Whitney tests,  $p < 0.05$ ; Table 2).

There was a tendency that %V3 (the viral subgroup with the highest fluorescence) was highest in surface water, whereas %V1 was lower and %V2 was higher in the bathypelagic ocean than in surface water (Fig. 3, Table 2). A local maximum of %V1 and a local minimum of %V2 and %V3 was found at the oxygen minimum at  $47.6^{\circ}$  W,  $53.5^{\circ}$  W. Significant differences between some water masses were found for all 3 viral flow cytometer groups (Kruskal-Wallis and Mann-Whitney tests,  $p < 0.05$ ; Table 2). Significant differences with depth were only found for %V1 and %V2 (Table 3).

Co-variation of physico-chemical and biological parameters was assessed across water masses (Table 3). PA and VA increased with temperature and decreased with AOU,  $\text{PO}_4$  and  $\text{NO}_3$ . There were positive correlations between PA, VA and PHP.

### 3.3. Viral and microbial parameters in the NADW

In order to assess potential changes of parameters with the formation and ageing of the NADW, the distance of stations from the GIN Sea was calculated as a proxy for the length of the NADW (Fig. 1). Temperature and salinity increased significantly with distance from the GIN Sea.  $\text{PO}_4$  and  $\text{NO}_3$  concentrations increased also with distance. Oxygen concentrations decreased significantly with distance, whereas AOU increased (Table 4). VA and VPR decreased significantly with distance, but in a non-linear way (see also Fig. 3). The %V1 increased (from the 2 stations closest to the GIN Sea) and V2 decreased significantly with distance (Table 4, Fig. 3).

BS was assessed at Stns 1, 13 and 31, averaging  $28 \pm 5$  (range: 24–33 for the 3 samples) after correction to  $\text{BS}_{\text{max}}$ . For VP measurements, VA in the incubations was reduced by  $77 \pm 13\%$

256 compared to *in situ* VA, whereas PA was  $69 \pm 16\%$  of *in situ* abundance. VP averaged  $0.95 \pm 0.78$   
257  $\times 10^4$  viruses  $l^{-1} d^{-1}$  (Table 5). FIC averaged  $14 \pm 7\%$ , mean VMMP<sub>VP</sub> was  $59 \pm 52\%$  and VMMP<sub>FIC</sub>  
258  $20 \pm 12\%$ . VP<sub>i</sub> averaged  $1.7 \pm 1.4 \times 10^7$  viruses  $l^{-1} d^{-1}$ , while FLC averaged  $21 \pm 14\%$ . VP and FIC  
259 decreased with distance from the GIN Sea, whereas for VP<sub>i</sub> and FLC no significant relationship was  
260 found (Fig. 4).

261 Co-variation of parameters was also assessed within the NADW (Table 4). PA was  
262 positively related to oxygen concentration and negatively related to salinity and AOU. VA was  
263 negatively related to AOU, PO<sub>4</sub> and NO<sub>3</sub> concentrations and positively to oxygen.

## 264 4. DISCUSSION

265 The data presented here indicate that different water masses maintain specific viral  
266 characteristics during their early lateral flow in the oceanic conveyor belt. Nevertheless, for the  
267 NADW, successional changes with distance from the origin could be observed for VA, subgroups,  
268 infection and lytic production.

### 269 4.1. Depth distribution of viruses

270 The decrease of VA with depth during early formation of the NADW was much less  
271 pronounced than for PA (Fig. 1, Table 2). Similar trends were found for the Atlantic (Parada et al.  
272 2007, De Corte et al. 2010, 2012, 2016, Muck et al. 2014, Winter et al. 2018) and the Pacific (Li et  
273 al. 2014, Yang et al. 2014). Vertical transport of sinking particles is probably not always  
274 responsible for the high VPRs in the dark ocean as previously suggested (Hara et al. 1996), since  
275 very high VPRs were found in bathypelagic areas where sinking particle fluxes are generally low  
276 (Yang et al. 2014). However, since viruses can enhance aggregate formation and export into the  
277 dark ocean (i.e. viral shuttle; Peduzzi & Weinbauer 1993, Sullivan et al. 2017, Yamada et al. 2018,  
278 Boeuf et al. 2019), viral lysis can also contribute to the sinking of viruses attached to aggregates.

279 Among the main causes for viral decay in the absence of sunlight are high temperatures,  
280 high molecular weight DOM and microscopic (inorganic) particles (Suttle & Chen 1992, Cottrell &  
281 Suttle 1995, Noble & Fuhrman 1997). One of the obvious reasons for the high abundances of  
282 viruses in bathypelagic waters could be low decay due to low temperatures (Parada et al. 2007).  
283 This is supported by the finding that the VPR was lower and water temperature ca. 13°C higher in  
284 the bathypelagic zone of the Mediterranean Sea (Magagnini et al. 2007, Winter et al. 2009) than in  
285 the Atlantic Ocean.

286 While the low DOM concentrations in the dark ocean (e.g. Aristegui et al. 2009) likely mean  
287 reduced decay, large microscopic particles are plentiful (Bochdansky et al. 2010, 2016, Boeuf et al.

288 2019) and could thus be a significant cause of viral decay. However, since there is evidence that  
289 organic particles (marine snow) are viral factories rather than viral traps (Weinbauer et al. 2009a,  
290 Bettarel et al. 2016), the particles in the dark ocean could protect viruses against decay and even  
291 foster VP. If the emerging notion holds that deep-sea prokaryotes are preferentially particle-  
292 attached (e.g. DeLong et al. 2006, Baltar et al. 2009, Swan et al. 2011), these particles might be  
293 hotspots of viral infection by increasing contact rates and, hence, abundance (De Corte et al. 2012).

294 It has often been assumed that there is a trade-off for prokaryotes between competition for  
295 nutrients and resistance against viral infection (Thingstad 2000, Winter et al. 2010). From a fitness  
296 perspective (Thingstad et al. 2014, Thingstad & Våge 2019), one could argue that—as the supply  
297 with organic material is low in the deep-sea (Arístegui et al. 2009)—favouring DOM uptake  
298 abilities should occur at the expense of defence against viral infection. This should result in higher  
299 VP, especially when contact rates remain high in cases where the viruses and microbes are mainly  
300 particle-attached (see paragraph above), and thus in a high VPR in the deep-sea.

301 There is no simple relationship between fluorescence intensity and genome size of viruses.  
302 However, as viruses do not have their own metabolism, the staining intensity with dyes such as  
303 SYBR Green does not vary for specific types of viruses. Therefore, changes in the relative  
304 proportion of viral subgroups indicate changes of viral community composition (Brussaard et al.  
305 2010). Consequently, the variation of the viral subgroups between depth layers (Table 2) suggests  
306 differences in viral community composition. Such differences between viral subgroups have also  
307 been found in other studies (De Corte et al. 2010, Muck et al. 2014) and were confirmed by using  
308 pulsed-field gel electrophoresis (Parada et al. 2007), randomly amplified polymorphic DNA-PCR  
309 (RAPD-PCR; De Corte et al. 2010, Winter & Weinbauer 2010, Muck et al. 2014) and  
310 metagenomics (Mizuno et al. 2016, Winter et al. 2018, Gregory et al. 2019, Liang et al. 2019).  
311 Using viral subgroups, the strongest differences were found between the SSL and bathypelagic  
312 water masses, suggesting that specific viral communities are inhabiting these environments. A  
313 likely reason for the differences in viral community composition is differences in host activity and  
314 community structure diversity in these water masses as assessed during the same cruise (Teira et al.  
315 2004, Reinthaler et al. 2006).

## 316 4.2. Lysogeny in the NADW

317 FLC in the NADW data ranged from 4.5–40.1%, which is similar to another study from the  
318 deep ocean using the VRA approach (10.1–27.3%; Muck et al. 2014). Using a whole seawater  
319 approach, FLC values were found to be highest in the bathypelagic zone of the Mediterranean Sea

320 (73.2%; Weinbauer et al. 2003). Using the VRA approach, FLC values from the NADW were  
321 higher than FIC values (Fig. 4, Table 5).

322 FLC and lysogenic VP did not vary significantly with distance from the GIN Sea (Fig. 4b,d,  
323 Table 5). Also, FLC and lysogenic VP did not vary significantly with PHP (Spearman rank  
324 correlation,  $\rho < 0.65$ ,  $p > 0.15$ ). In other studies of the Atlantic Ocean, inducible VP did not change  
325 (or changed little) with depth from the epipelagic to the abyssopelagic zone, whereas PA, PHP and  
326 lytic VP decreased strongly with depth (De Corte et al. 2010, 2012, Muck et al. 2014).  
327 Metagenomic data indicate that lysogeny is the predominant life strategy in the deep ocean  
328 (Williamson et al. 2008, Mizuno et al. 2016). Thus, these data support the general idea that  
329 lysogeny dominates when the encounter rate between phages and host cells is low (Stewart & Levin  
330 1984, Weinbauer et al. 2003, Weinbauer 2004). However, in the Malaspina Circumnavigation  
331 Experiment, Lara et al. (2017) found that lysogeny dominated in surface water and lytic VP became  
332 more important in deep water, hence supporting both hypotheses, i.e. promotion of lysogeny by low  
333 growth rates and low nutrient conditions ('classic' explanation) and promotion at high host  
334 abundance ('piggyback-the-winner' model) (Knowles et al. 2016). This indicates that the lytic–  
335 lysogenic switch is likely more complex than previously thought, which calls for further studies  
336 (Lara et al. 2017). Independent of the lytic–lysogeny switch argument, all studies show that  
337 lysogeny is significant in the deep sea. It has been hypothesized that marine prophages directly  
338 contribute to host survival in unfavourable environments by suppressing superfluous metabolic  
339 activities (Paul 2008), which could be particularly important in the carbon-limited dark ocean.

### 340 4.3. VA and infection during the formation of the NADW

341 VA, VPR, lytic VP and FIC decreased with distance from the GIN sea. Among the factors  
342 which could have influenced these parameters during the early formation of the NADW are (1)  
343 physical factors (temperature or mixing of the NADW with adjacent water masses), (2) sinking of  
344 particles from surface water into the interior of the ocean (export of viruses or stimulation of host  
345 activity) and (3) successional (i.e. internal temporal changes such as changes in community  
346 composition or resistance).

347 VA decreased in the NADW and temperature increased with distance from its formation  
348 (Fig. 3, Table 4). It is possible that the ca. 50 yr transport into warming water resulted in an  
349 increased decay and contributed to losses of viruses. The increase in temperature and salinity in the  
350 NADW with distance from its formation (Table 4) is caused by mixing with adjacent water masses  
351 such as the overlying LSW, which is warmer and lower in salinity than the NADW (van Aken  
352 2000a,b). Since VA was higher in the LSW than in the NADW, one would expect VA to increase

353 with distance in the NADW; however, they decreased, instead indicating that other factors were  
354 more important than direct mixing processes. Mixing of deep sea water masses can cause an  
355 increase in prokaryotic production and hence VP (Muck et al. 2014) e.g. by induction of lysogens  
356 (Winter et al. 2018). Our data set did not allow us to evaluate this possibility for the NADW.

357 General distribution patterns of chlorophyll *a* (chl *a*) during the TRANSAT-II cruise suggest  
358 a potentially decreasing sinking particle flux and thus a potentially decreasing viral export flux with  
359 distance from the GIN Sea (Teira et al. 2004, Reinthaler et al. 2006). If sinking particles are a  
360 source of viruses for the dark ocean, such a pattern could contribute to the finding of a decrease in  
361 VA and VPR with age of the NADW (but see discussion on transport of viruses on particles in  
362 Section 4.1). Release of viruses from sinking particles could for example explain local maxima of  
363 viral parameters as detected in this study (Fig. 2). The decrease of VA, VPR, VP and FIC with  
364 distance from the GIN Sea was not accompanied by a change in PA and PHP (Table 4). Thus, there  
365 is no support for the hypothesis that sinking aggregates (transporting viruses and prokaryotes) or the  
366 supply of organic matter into the NADW caused the observed viral (and prokaryotic) patterns with  
367 distance from the GIN Sea.

368 Assuming a conservative prokaryotic turnover time of ca. 1 mo in the NADW (Reinthaler et  
369 al. 2006) and an investigated time frame for the formation of the NADW of 50 yr, prokaryotes  
370 produced ca. 600 generations of offspring. Correlation analysis indicates that (micro)organisms  
371 consume oxygen and remineralize PO<sub>4</sub> and NO<sub>3</sub> during this transport (Table 4). It is possible that  
372 prokaryotic community composition changed during these 600 generations. Changes in community  
373 composition of hosts can affect the community composition of virioplankton (e.g. Winter et al.  
374 2010). Such a mechanism could explain the finding that viral community composition, as indicated  
375 by the relative abundance of viral groups, changed significantly with age of the NADW. Since the  
376 BS of viruses is quite variable (Børsheim 1993), changes in the community composition to virus–  
377 host systems with lower BSs could contribute to the decrease of VA and VPR with distance from  
378 the GIN Sea. The few data on BS support this idea (Table 5).

379 In a 2 yr pressure incubation (corresponding to pressure at 3000 m water depth) with various  
380 phage isolates and a natural virus community, it was found that small and low fluorescence viruses  
381 decayed slower than larger or high fluorescence viruses (Tian et al 2020). The finding that the  
382 proportion of the V1 group (low fluorescence) became more important with distance from the GIN  
383 Sea compared to V2 and V3 groups (higher fluorescence) (Fig. 3b,c) could therefore be explained  
384 by lower decay rates. Thus, variable decay rates between different types of viruses could change the  
385 virus community and hence infection patterns.

386 Resistance against infection is a well known phenomenon from studies with isolates (Avrani  
387 et al. 2012). Among the more recently detected resistance mechanisms is CRISPR (Barrangou et al.  
388 2007). CRISPR is a sort of immune system for prokaryotes in the sense that it confers resistance to  
389 bacterial and archaea cells against mobile genetic elements such as viruses (e.g. Barrangou et al.  
390 2007, Vestergaard et al. 2008). Exposure of prokaryotes to viruses for 600 generations and  
391 development of resistance is therefore another possible cause for the decrease of VA and lytic  
392 infection with ageing NADW. If the resistance hypothesis holds, viral types belonging to the V2  
393 group would be the loser in the arms-race with prokaryotic hosts compared to type V1, which  
394 decrease in relative abundance with distance from the GIN Sea.

395 Applying the reasoning of a fitness penalty in a low-nutrient environment (see Section 4.1),  
396 it can be argued that the fitness costs of resistance should increase with distance from the GIN Sea,  
397 i.e. with the ageing of NADW. Hence, a strategy towards competitive traits with high susceptibility  
398 to viral infection could be anticipated. However, there is no evidence that viral infection or VPR  
399 increased; on the contrary, these parameters decreased with distance from the GIN Sea. Overall,  
400 VPR was very high at the origin of the GIN Sea, and despite the decline with ageing of the NADW  
401 it remained higher than in surface water. It is possible that the high VPR values in origin water  
402 masked fitness-related trends. In this context, it is important to mention that temperature increased  
403 in the NADW with distance from the GIN Sea. Alternatively, the origin water could have been  
404 already characterized by conditions favouring competitive over defence traits.

#### 405 **4.4. VP, mortality and carbon release in the NADW**

406 Lytic VP ranged from  $2.2\text{--}2.5 \times 10^7 \text{ l}^{-1} \text{ d}^{-1}$ . These values are slightly lower than the lytic VP  
407 estimated in other bathy- and abyssopelagic environments ( $3.6 \times 10^7\text{--}8.4 \times 10^9 \text{ l}^{-1} \text{ d}^{-1}$  (De Corte et  
408 al. 2010 2012, Umani et al. 2010, Li et al. 2014, Muck et al. 2016, Lara et al. 2017, Winter et al.  
409 2018). The FIC (0.6–22.0%) was at the lower range compared to other studies (15–143%; Muck et  
410 al. 2014, Winter et al. 2018).

411 Using 2 different methods for estimating the role of viral lysis for prokaryotic mortality,  
412 estimates based on  $\text{VMMP}_{\text{VP}}$  were on average 2.7 higher than estimates based on  $\text{VMMP}_{\text{FIC}}$  (Table  
413 5). It is a well known but poorly understood phenomenon that different methods for estimating  
414 VMMP are not always fully congruent (Winter et al. 2004, Helton et al. 2005, Winget et al. 2005,  
415 Weinbauer et al. 2009b). Nevertheless, data from both methods suggest significant mortality due to  
416 viral lysis in the NADW (on average 20% by  $\text{VMMP}_{\text{FIC}}$  and 59% by  $\text{VMMP}_{\text{VP}}$ , respectively).  
417 Finally, viruses have been detected within cells (Weinbauer et al. 2003, this study). This finding  
418 supports that of Li et al. (2014) that there is an autochthonous active virus community in the deep



419 sea. Also, endemic deep-sea virus communities have been documented by metagenomics (Winter et  
420 al. 2014, 2018, Mizuno et al. 2016, Gregory et al. 2019, Liang et al. 2019).

421 Using a conversion factor of  $12.4 \text{ fg cell}^{-1}$  for the dark ocean (Fukuda et al. 1998) and data  
422 from Table 5, the carbon release by viral lysis of prokaryotes would be  $8.6 \pm 5.5 \text{ ng C l}^{-1} \text{ d}^{-1}$  using  
423 VMMP<sub>VP</sub> and  $2.9 \pm 1.4 \text{ ng C l}^{-1} \text{ d}^{-1}$  using VMMP<sub>FIC</sub>. This is lower than the  $0.03\text{--}0.69 \text{ } \mu\text{g C l}^{-1} \text{ d}^{-1}$   
424 estimated by Li et al. (2014). The majority of DOM in the deep sea is characterized by low turnover  
425 times and is either too recalcitrant or too diluted to be used by prokaryotes (Jiao et al. 2011, Arrieta  
426 et al. 2015). Viral lysis products consist of cell contents and wall cell debris such as DNA, RNA,  
427 carbohydrates, amino acids, glucosamine and diaminopimelic acid (Weinbauer et al. 1993,  
428 Weinbauer & Peduzzi 1995, Middelboe & Jorgensen 2006) and are rapidly degraded, hence  
429 belonging to the pool of labile DOM (Noble & Fuhrman 1999, Middelboe & Lyck 2002). This may  
430 relieve the carbon limitation of the growth of deep-sea prokaryotes. Also, organic matter from cells  
431 shunted into the DOM pool by viral lysis is hardly accessible to higher trophic levels, thus resulting  
432 in a slower transfer of organic matter towards higher trophic levels (Fuhrman 1999). This process  
433 could sustain a high prokaryotic biomass and provide an important contribution to prokaryotic  
434 metabolism, allowing the system to cope with the severe organic resource limitation of deep-sea  
435 ecosystems as has been demonstrated for benthic and pelagic communities (Danovaro et al. 2008,  
436 Lara et al. 2017). Moreover, viral lysis may prime the biological pump and the microbial carbon  
437 pump and hence carbon sequestration in the ocean, which has significant global consequences  
438 (Suttle 2007, Brussaard et al. 2008, Jiao et al. 2011, Guidi et al. 2016).

439 *Acknowledgements.* We thank the captain and crew of the R/V ‘Pelagia’ for their support at sea, and  
440 K. Bakker, J. Hegeman, S. Gonzalez and A. Smit for help during sample processing. The comments  
441 of 2 reviewers improved a former version of the manuscript. This work was supported by a grant of  
442 the Dutch Science Foundation Earth and Life Sciences (NWO-ALW, project # 811.33.004) to G.J.H  
443 and the French Science Ministry (ANR grant AQUAPHAGE) to M.G.H.

## 444 LITERATURE CITED

- 445 <jrn>Agogue H, Lamy D, Neal PR, Sogin ML, Herndl GJ (2011) Water mass-specificity of  
446 bacterial communities in the North Atlantic revealed by massively parallel sequencing. Mol  
447 Ecol 20:258–274 PubMed doi:10.1111/j.1365-294X.2010.04932.x</jrn>
- 448 <jrn>Arístegui J, Gasol JM, Duarte CM, Herndl GJ (2009) Microbial oceanography of the dark  
449 ocean’s pelagic realm. Limnol Oceanogr 54:1501–1529 doi:10.4319/lo.2009.54.5.1501</jrn>



450 <jrn>Arrieta JM, Mayol E, Hansman RL, Herndl GJ, Dittmar T, Duarte CM (2015) Dilution limits  
 451 dissolved organic carbon utilization in the deep ocean. *Science* 348:331–333 [PubMed](#)  
 452 [doi:10.1126/science.1258955](#)</jrn>

453 <jrn>Avrani S, Schwartz DA, Lindell D (2012) Virus-host swinging party in the oceans:  
 454 incorporating biological complexity into paradigms of antagonistic coexistence. *Mob Genet*  
 455 *Elements* 2:88–95 [PubMed](#) [doi:10.4161/mge.20031](#)</jrn>

456 <jrn>Baltar F, Arístegui J, Gasol JM, Sintes E, Herndl GJ (2009) Evidence of prokaryotic  
 457 metabolism on suspended particulate organic matter in the dark waters of the subtropical North  
 458 Atlantic. *Limnol Oceanogr* 54:182–193 [doi:10.4319/lo.2009.54.1.0182](#)</jrn>

459 <jrn>Barrangou R, Fremaux C, Deveau H, Richards M and others (2007) CRISPR provides  
 460 acquired resistance against viruses in prokaryotes. *Science* 315:1709–1712 [PubMed](#)  
 461 [doi:10.1126/science.1138140](#)</jrn>

462 <jrn>Bettarel Y, Motegi C, Weinbauer MG, Mari X (2016) Colonization and release processes of  
 463 viruses and prokaryotes on artificial marine macroaggregates. *FEMS Microbiol Lett* 363:fnv216  
 464 [PubMed](#) [doi:10.1093/femsle/fnv216](#)</jrn>

465 <jrn>Binder B (1999) Reconsidering the relationship between virally induced bacterial mortality  
 466 and frequency of infected cells. *Aquat Microb Ecol* 18:207–215 [doi:10.3354/ame018207](#)</jrn>

467 <jrn>Bochdansky AB, van Aken HM, Herndl GJ (2010) Role of macroscopic particles in deep-sea  
 468 oxygen consumption. *Proc Natl Acad Sci USA* 107:8287–8291 [PubMed](#)  
 469 [doi:10.1073/pnas.0913744107](#)</jrn>

470 <jrn>Bochdansky AB, Clouse MA, Herndl GJ (2016) Dragon kings of the deep sea: Marine  
 471 particles deviate markedly from the common number-size spectrum. *Sci Rep* 6:22633 [PubMed](#)  
 472 [doi:10.1038/srep22633](#)</jrn>

473 <jrn>Boeuf D, Edwards BR, Eppley JM, Hu SK and others (2019) Biological composition and  
 474 microbial dynamics of sinking particulate organic matter at abyssal depths in the oligotrophic  
 475 open ocean. *Proc Natl Acad Sci USA* 116:11824–11832 [PubMed](#)</jrn>

476 <jrn>Bohannan BJM, Lenski RE (1997) Effect of resource enrichment on a chemostat community  
 477 of bacteria and bacteriophage. *Ecology* 78:2303–2315 [doi:10.1890/0012-](#)  
 478 [9658\(1997\)078\[2303:EOREOA\]2.0.CO;2](#)</jrn>

479 <jrn>Børsheim KY (1993) Native marine bacteriophages. *FEMS Microbiol Ecol* 102:141–159  
 480 [doi:10.1016/0378-1097\(93\)90197-A](#)</jrn>

481 <jrn>Broecker WS (1997) Thermohaline circulation, the Achilles heel of our climate system: Will  
 482 man-made CO<sub>2</sub> upset the current balance? *Science* 278:1582–1588 [PubMed](#)  
 483 [doi:10.1126/science.278.5343.1582](#)</jrn>

484 <jrn>Brussaard CPD, Wilhelm SW, Thingstad F, Weinbauer MG and others (2008) Global-scale  
 485 processes with a nanoscale drive: the role of marine viruses. *ISME J* 2:575–578 [PubMed](#)  
 486 [doi:10.1038/ismej.2008.31](#)</jrn>

487 <edb>Brussaard CPD, Payet JP, Winter C, Weinbauer MG (2010) Quantification of aquatic viruses  
 488 by flow cytometry. In: Wilhelm SW, Weinbauer MG, Suttle C (eds) *Manual of aquatic viral*  
 489 *ecology*. American Society of Limnology and Oceanography, Waco, TX, p 102–109</edb>

490 <jrn>Cottrell MT, Suttle CA (1995) Dynamics of a lytic virus infecting the photosynthetic marine  
 491 picoflagellate *Micromonas pusilla*. *Limnol Oceanogr* 40:730–739  
 492 [doi:10.4319/lo.1995.40.4.0730](#)</jrn>

493 <jrn>Danovaro R, Dell’Anno A, Corinaldesi C, Magagnini M, Noble R, Tamburin C, Weinbauer  
 494 MG (2008) Major viral impact on the functioning of benthic deep-sea ecosystems. *Nature*  
 495 454:1084–1087 [PubMed](#) [doi:10.1038/nature07268](#)</jrn>

496 <jrn>De Corte D, Sintes E, Winter C, Yokokawa T, Reinthaler T, Herndl GJ (2010) Links between  
 497 viral and prokaryotic communities throughout the water column in the (sub)tropical Atlantic  
 498 Ocean. *ISME J* 4:1431–1442 [PubMed](#) [doi:10.1038/ismej.2010.65](#)</jrn>

499 <jrn>De Corte D, Sintes E, Yokokawa T, Reinthaler T, Herndl GJ (2012) Links between viruses  
 500 and prokaryotes throughout the water column along a North Atlantic latitudinal transect. *ISME J*  
 501 6:1566–1577 [PubMed](#) [doi:10.1038/ismej.2011.214](#)</jrn>

502 <jrn>De Corte D, Sintes E, Yokokawa T, Lekunberri I, Herndl GJ (2016) Large-scale distribution  
 503 of microbial and viral populations in the South Atlantic Ocean. *Environ Microbiol Rep* 8:305–  
 504 315 [PubMed](#) [doi:10.1111/1758-2229.12381](#)</jrn>

505 <jrn>DeLong EF, Preston CM, Mincer T, Rich V and others (2006) Community genomics among  
 506 stratified microbial assemblages in the ocean’s interior. *Science* 311:496–503 [PubMed](#)  
 507 [doi:10.1126/science.1120250](#)</jrn>

508 <jrn>Fuhrman JA (1999) Marine viruses and their biogeochemical and ecological effects. *Nature*  
 509 399:541–548 [PubMed](#) [doi:10.1038/21119](#)</jrn>

510 <jrn>Fukuda R, Ogawa H, Nagata T, Koike I (1998) Direct determination of carbon and nitrogen  
 511 contents of natural bacterial assemblages in marine environments. *Appl Environ Microbiol*  
 512 64:3352–3358 [PubMed](#) [doi:10.1128/AEM.64.9.3352-3358.1998](#)</jrn>

513 <jrn>Gobler CJ, Hutchins DA, Fisher NS, Cosper EM, Sañudo-Wilhelmy SA (1997) Release and  
514 bioavailability of C, N, P, Se and Fe following viral lysis of a marine chrysophyte. *Limnol*  
515 *Oceanogr* 42:1492–1504 [doi:10.4319/lo.1997.42.7.1492](https://doi.org/10.4319/lo.1997.42.7.1492)</jrn>

516 <jrn>Gregory AC, Zayed AA, Conceicao-Neto N, Temperton B and others (2019) Marine DNA  
517 viral macro- and microdiversity from pole to pole. *Cell* 177:1109–1123 [PubMed](https://pubmed.ncbi.nlm.nih.gov/31111111/)  
518 [doi:10.1016/j.cell.2019.03.040](https://doi.org/10.1016/j.cell.2019.03.040)</jrn>

519 <jrn>Guidi L, Chaffron S, Bittner L, Eveillard D and others (2016) Plankton networks driving  
520 carbon export in the oligotrophic ocean. *Nature* 532:465–470 [PubMed](https://pubmed.ncbi.nlm.nih.gov/26511111/)  
521 [doi:10.1038/nature16942](https://doi.org/10.1038/nature16942)</jrn>

522 <jrn>Hara S, Koike I, Terauchi K, Kamiya H, Tanoue E (1996) Abundance of viruses in deep  
523 oceanic waters. *Mar Ecol Prog Ser* 145:269–277 [doi:10.3354/meps145269](https://doi.org/10.3354/meps145269)</jrn>

524 <jrn>Helton RR, Cottrell MT, Kirchman DL, Wommack KE (2005) Evaluation of incubation-based  
525 methods for estimating virioplankton production in estuaries. *Aquat Microb Ecol* 41:209–219  
526 [doi:10.3354/ame041209](https://doi.org/10.3354/ame041209)</jrn>

527 <jrn>Jiao N, Herndl GJ, Hansell DA, Benner R and others (2011) The microbial carbon pump and  
528 the oceanic recalcitrant dissolved organic matter pool. *Nat Rev Microbiol* 9:555  
529 [doi:10.1038/nrmicro2386-c5](https://doi.org/10.1038/nrmicro2386-c5)</jrn>

530 <jrn>Knowles B, Silveira CB, Bailey BA, Barott K and others (2016) Lytic to temperate switching  
531 of viral communities. *Nature* 531:466–470 [PubMed](https://pubmed.ncbi.nlm.nih.gov/26511111/) [doi:10.1038/nature17193](https://doi.org/10.1038/nature17193)</jrn>

532 <jrn>Lara E, Vaqué D, Sà EL, Boras JA and others (2017) Unveiling the role and life strategies of  
533 viruses from the surface to the dark ocean. *Sci Adv* 3:e1602565 [PubMed](https://pubmed.ncbi.nlm.nih.gov/28111111/)  
534 [doi:10.1126/sciadv.1602565](https://doi.org/10.1126/sciadv.1602565)</jrn>

535 <jrn>Li Y, Luo T, Sun J, Cai L, Liang Y, Jiao N, Zhang R (2014) Lytic viral infection of  
536 bacterioplankton in deep waters of the Western Pacific Ocean. *Biogeosciences* 11:2531–2542  
537 [doi:10.5194/bg-11-2531-2014](https://doi.org/10.5194/bg-11-2531-2014)</jrn>

538 <jrn>Liang Y, Wang L, Wang Z, Zhao J and others (2019) Metagenomic analysis of the diversity of  
539 DNA viruses in the surface and deep sea of the South China Sea. *Front Microbiol* 10:1951  
540 [PubMed](https://pubmed.ncbi.nlm.nih.gov/31111111/) [doi:10.3389/fmicb.2019.01951](https://doi.org/10.3389/fmicb.2019.01951)</jrn>

541 <jrn>Middelboe M, Jorgensen NOG (2006) Viral lysis of bacteria: an important source of dissolved  
542 amino acids and cell wall compounds. *J Mar Biol Assoc UK* 86:605–612  
543 [doi:10.1017/S0025315406013518](https://doi.org/10.1017/S0025315406013518)</jrn>

544 <jrn>Middelboe M, Lyck PG (2002) Regeneration of dissolved organic matter by viral lysis in  
545 marine microbial communities. *Aquat Microb Ecol* 27:187–194 [doi:10.3354/ame027187](https://doi.org/10.3354/ame027187)</jrn>

546 <jrn>Mizuno CM, Ghai R, Saghai A, Lopez-Garcia P, Rodriguez-Valera F (2016) Genomes of  
547 abundant and widespread viruses from the deep ocean. *MBio* 7:e00805-16 [PubMed](https://pubmed.ncbi.nlm.nih.gov/26811111/)  
548 [doi:10.1128/mBio.00805-16](https://doi.org/10.1128/mBio.00805-16)</jrn>

549 <jrn>Muck S, Griessler T, Köstner N, Klimiuk A, Winter C, Herndl GJ (2014) Fracture zones in the  
550 Mid Atlantic Ridge lead to alterations in prokaryotic and viral parameters in deep-water masses.  
551 *Front Microbiol* 5:264 [PubMed](https://pubmed.ncbi.nlm.nih.gov/25411111/) [doi:10.3389/fmicb.2014.00264](https://doi.org/10.3389/fmicb.2014.00264)</jrn>

552 <jrn>Murray AG, Jackson GA (1992) Viral dynamics: a model of the effects of size, shape, motion  
553 and abundance of single-celled planktonic organisms and other particles. *Mar Ecol Prog Ser*  
554 89:103–116 [doi:10.3354/meps089103](https://doi.org/10.3354/meps089103)</jrn>

555 <jrn>Nagata T, Fukuda H, Fukuda R, Koike I (2000) Bacterioplankton distribution and production  
556 in deep Pacific waters: large-scale geographic variations and possible coupling with sinking  
557 particle fluxes. *Limnol Oceanogr* 45:426–435 [doi:10.4319/lo.2000.45.2.0426](https://doi.org/10.4319/lo.2000.45.2.0426)</jrn>

558 <jrn>Noble RT, Fuhrman JA (1997) Virus decay and its causes in coastal waters. *Appl Environ*  
559 *Microbiol* 63:77–83 [PubMed](https://pubmed.ncbi.nlm.nih.gov/9111111/) [doi:10.1128/AEM.63.1.77-83.1997](https://doi.org/10.1128/AEM.63.1.77-83.1997)</jrn>

560 <jrn>Noble RT, Fuhrman JA (1999) Breakdown and microbial uptake of marine viruses and other  
561 lysis products. *Aquat Microb Ecol* 20:1–11 [doi:10.3354/ame020001](https://doi.org/10.3354/ame020001)</jrn>

562 <jrn>Parada V, Herndl GJ, Weinbauer MG (2006) Viral burst size of heterotrophic prokaryotes in  
563 aquatic systems. *J Mar Biol Assoc UK* 86:613–621 [doi:10.1017/S002531540601352X](https://doi.org/10.1017/S002531540601352X)</jrn>

564 <jrn>Parada V, Sintes E, van Aken HM, Weinbauer MG, Herndl GJ (2007) Viral abundance, decay  
565 and diversity in the meso- and bathypelagic waters of the North Atlantic. *Appl Environ*  
566 *Microbiol* 73:4429–4438 [PubMed](https://pubmed.ncbi.nlm.nih.gov/17111111/) [doi:10.1128/AEM.00029-07](https://doi.org/10.1128/AEM.00029-07)</jrn>

567 <jrn>Paul JH (2008) Prophages in marine bacteria: Dangerous molecular time bombs or key to the  
568 survival in the seas? *ISME J* 2:579–589 [PubMed](https://pubmed.ncbi.nlm.nih.gov/18111111/) [doi:10.1038/ismej.2008.35](https://doi.org/10.1038/ismej.2008.35)</jrn>

569 <edb>Paul JH, Weinbauer MG (2010) Detection of lysogeny in marine environments. In: Wilhelm  
570 SW, Weinbauer MG, Suttle C (eds) *Manual of aquatic viral ecology*. American Society of  
571 *Limnology and Oceanography*, Waco, TX, p 30–33</edb>

572 <jrn>Peduzzi P, Weinbauer MG (1993) Effect of concentrating the virus-rich 2-200 nm size  
573 fraction of seawater on the formation of algal flocs (marine snow). *Limnol Oceanogr* 38:1562–  
574 1565 [doi:10.4319/lo.1993.38.7.1562](https://doi.org/10.4319/lo.1993.38.7.1562)</jrn>

575 <jrn>Reinthal T, van Aken H, Veth C, Arístegui J, Robinson C, Williams PJL (2006) Prokaryotic  
576 respiration and production in the meso-and bathypelagic realm of the eastern and western North  
577 Atlantic basin. *Limnol Oceanogr* 51:1262–1273 [doi:10.4319/lo.2006.51.3.1262](https://doi.org/10.4319/lo.2006.51.3.1262)</jrn>

578 <jrn>Steward GF, Smith DC, Azam F (1996) Abundance and production of bacteria and viruses in  
579 the Bering and Chukchi Seas. *Mar Ecol Prog Ser* 131:287–300 [doi:10.3354/meps131287](https://doi.org/10.3354/meps131287)</jrn>

580 <jrn>Stewart FM, Levin BR (1984) The population biology of bacterial viruses: why be temperate.  
581 *Theor Popul Biol* 26:93–117 [PubMed](https://pubmed.ncbi.nlm.nih.gov/2441111/) [doi:10.1016/0040-5809\(84\)90026-1](https://doi.org/10.1016/0040-5809(84)90026-1)</jrn>

582 <jrn>Sullivan MB, Weitz JS, Wilhelm S (2017) Viral ecology comes of age. *Environ Microbiol*  
583 *Rep* 9:33–35 [PubMed](https://pubmed.ncbi.nlm.nih.gov/2811111/) [doi:10.1111/1758-2229.12504](https://doi.org/10.1111/1758-2229.12504)</jrn>

584 <jrn>Suttle CA (2007) Marine viruses—major players in the global ecosystem. *Nat Rev Microbiol*  
585 5:801–812 [PubMed](https://pubmed.ncbi.nlm.nih.gov/17501111/) [doi:10.1038/nrmicro1750](https://doi.org/10.1038/nrmicro1750)</jrn>

586 <jrn>Suttle CA, Chen F (1992) Mechanisms and rates of decay of marine viruses in seawater. *Appl*  
587 *Environ Microbiol* 58:3721–3729 [PubMed](https://pubmed.ncbi.nlm.nih.gov/11281111/) [doi:10.1128/AEM.58.11.3721-3729.1992](https://doi.org/10.1128/AEM.58.11.3721-3729.1992)</jrn>

588 <jrn>Swan BK, Martinez-Garcia M, Preston CM, Sczyrba A and others (2011) Potential for  
589 chemolithoautotrophy among ubiquitous bacteria lineages in the dark ocean. *Science* 333:1296–  
590 1300 [PubMed](https://pubmed.ncbi.nlm.nih.gov/21261111/) [doi:10.1126/science.1203690](https://doi.org/10.1126/science.1203690)</jrn>

591 <jrn>Teira E, Reinthal T, Pernthaler A, Pernthaler J, Herndl GJ (2004) Combining catalyzed  
592 reporter deposition-fluorescence in situ hybridization and microautoradiography to detect  
593 substrate utilization by Bacteria and Archaea in the deep ocean. *Appl Environ Microbiol*  
594 70:4411–4414 [PubMed](https://pubmed.ncbi.nlm.nih.gov/11281111/) [doi:10.1128/AEM.70.7.4411-4414.2004](https://doi.org/10.1128/AEM.70.7.4411-4414.2004)</jrn>

595 <jrn>Teira E, Lebaron P, van Aken H, Herndl GJ (2006) Distribution and activity of Bacteria and  
596 Archaea in the deep water masses of the North Atlantic. *Limnol Oceanogr* 51:2131–2144  
597 [doi:10.4319/lo.2006.51.5.2131](https://doi.org/10.4319/lo.2006.51.5.2131)</jrn>

598 <jrn>Thingstad TF, Våge S (2019) Host-virus-predator coexistence in a grey-box model with  
599 dynamic optimization of host fitness. *ISME J* 13:3102–3111 [PubMed](https://pubmed.ncbi.nlm.nih.gov/31396111/) [doi:10.1038/s41396-019-0496-7](https://doi.org/10.1038/s41396-019-0496-7)</jrn>

600

601 <jrn>Thingstad TF, Våge S, Storesund JE, Sandaa RA, Giske J (2014) A theoretical analysis of  
602 how strain-specific viruses can control microbial species diversity. *Proc Natl Acad Sci USA*  
603 111:7813–7818 [PubMed](https://pubmed.ncbi.nlm.nih.gov/24009111/) [doi:10.1073/pnas.1400909111](https://doi.org/10.1073/pnas.1400909111)</jrn>

604 Tian Y, Cai L, Xu Y, Luo T and others (2020) Stability and infectivity of  
605 allochthonous viruses in deep sea: A long-term high pressure simulation  
606 experiment.  
607 *Deep Sea Res Part I*: 103302 (in press) [doi:10.1016/j.dsr.2020.103302](https://doi.org/10.1016/j.dsr.2020.103302)



608 <jrn>Umani SF, Malisana E, Focaracci F, Magagnini M, Corinaldesi C, Danovaro R (2010)  
609 Disentangling the effect of viruses and nanoflagellates on prokaryotes in bathypelagic waters of  
610 the Mediterranean Sea. *Mar Ecol Prog Ser* 418:73–85 [doi:10.3354/meps08803](https://doi.org/10.3354/meps08803)</jrn>

611 <jrn>Våge S, Bratbak G, Egge J, Heldal M and others (2018) Simple models combining  
612 competition, defence and resource availability have broad implications in pelagic microbial food  
613 webs. *Ecol Lett* 21:1440–1452 [PubMed](https://pubmed.ncbi.nlm.nih.gov/31111111/) [doi:10.1111/ele.13122](https://doi.org/10.1111/ele.13122)</jrn>

614 <jrn>van Aken HM (2000a) The hydrography of the mid-latitude Northeast Atlantic Ocean. I: the  
615 deep water masses. *Deep Sea Res I* 47:757–788 [doi:10.1016/S0967-0637\(99\)00092-8](https://doi.org/10.1016/S0967-0637(99)00092-8)</jrn>

616 <jrn>van Aken HM (2000b) The hydrography of the mid-latitude Northeast Atlantic Ocean. II: the  
617 intermediate water masses. *Deep Sea Res I* 47:789–824 [doi:10.1016/S0967-0637\(99\)00112-](https://doi.org/10.1016/S0967-0637(99)00112-0)  
618 [0](https://doi.org/10.1016/S0967-0637(99)00112-0)</jrn>

619 <jrn>Vestergaard G, Shah SA, Bize A, Reitberger W and others (2008) *Stygiolobus* rod-shaped  
620 virus and the interplay of crenarchaeal rudiviruses with the CRISPR antiviral system. *J Bacteriol*  
621 190:6837–6845 [PubMed](https://pubmed.ncbi.nlm.nih.gov/1868376845/) [doi:10.1128/JB.00795-08](https://doi.org/10.1128/JB.00795-08)</jrn>

622 <jrn>Weinbauer MG (2004) Ecology of prokaryotic viruses. *FEMS Microbiol Rev* 28:127–181  
623 [PubMed](https://pubmed.ncbi.nlm.nih.gov/1501016/j.femsre.2003.08.001/) [doi:10.1016/j.femsre.2003.08.001](https://doi.org/10.1016/j.femsre.2003.08.001)</jrn>

624 <jrn>Weinbauer MG, Peduzzi P (1995) Effect of virus-rich high molecular weight concentrates of  
625 seawater on the dynamics of dissolved amino acids and carbohydrates. *Mar Ecol Prog Ser*  
626 127:245–253 [doi:10.3354/meps127245](https://doi.org/10.3354/meps127245)</jrn>

627 <jrn>Weinbauer MG, Suttle CA (1999) Lysogeny and prophage induction in coastal and offshore  
628 bacterial communities. *Aquat Microb Ecol* 18:217–225 [doi:10.3354/ame018217](https://doi.org/10.3354/ame018217)</jrn>

629 <jrn>Weinbauer MG, Fuks D, Peduzzi P (1993) Distribution of viruses and dissolved DNA along a  
630 coastal trophic gradient in the northern Adriatic Sea. *Appl Environ Microbiol* 59:4074–4082  
631 [PubMed](https://pubmed.ncbi.nlm.nih.gov/101128/AEM.59.12.4074-4082.1993/) [doi:10.1128/AEM.59.12.4074-4082.1993](https://doi.org/10.1128/AEM.59.12.4074-4082.1993)</jrn>

632 <jrn>Weinbauer MG, Winter C, Höfle MG (2002) Reconsidering transmission electron microscopy  
633 based estimates of viral infection of bacterioplankton using conversion factors derived from  
634 natural communities. *Aquat Microb Ecol* 27:103–110 [doi:10.3354/ame027103](https://doi.org/10.3354/ame027103)</jrn>

635 <jrn>Weinbauer MG, Brettar I, Höfle MG (2003) Lysogeny and virus-induced mortality of  
636 bacterioplankton in surface, deep, and anoxic waters. *Limnol Oceanogr* 48:1457–1465  
637 [doi:10.4319/lo.2003.48.4.1457](https://doi.org/10.4319/lo.2003.48.4.1457)</jrn>

638 <jrn>Weinbauer MG, Bettarel Y, Cattaneo R, Luef B and others (2009a) Viral ecology of organic  
639 and inorganic particles in aquatic systems: avenues for further research. *Aquat Microb Ecol*  
640 57:321–341 [PubMed](#) [doi:10.3354/ame01363](#)</jrn>

641 <jrn>Weinbauer MG, Arrieta JM, Griebler C, Herndl GJ (2009b) Enhanced viral production and  
642 infection of bacterioplankton during an iron-induced phytoplankton bloom in the Southern  
643 Ocean. *Limnol Oceanogr* 54:774–784 [doi:10.4319/lo.2009.54.3.0774](#)</jrn>

644 <edb>Weinbauer MG, Rowe JM, Wilhelm SW (2010) Determining rates of virus production in  
645 aquatic systems by the virus reduction approach. In: Wilhelm SW, Weinbauer MG, Suttle C  
646 (eds) *Manual of aquatic viral ecology*. American Society of Limnology and Oceanography,  
647 Waco, TX, p 1–8</edb>

648 <jrn>Wilhelm SW, Suttle CA (1999) Viruses and nutrient cycles in the sea. *BioScience* 49:781–788  
649 [doi:10.2307/1313569](#)</jrn>

650 <jrn>Wilhelm SW, Brigden SM, Suttle CA (2002) A dilution technique for the direct measurement  
651 of viral production: a comparison in stratified and tidally mixed coastal waters. *Microb Ecol*  
652 43:168–173 [PubMed](#) [doi:10.1007/s00248-001-1021-9](#)</jrn>

653 <edb>Williams ST (1994) Bacteriophages in soils. In: Webster RG, Granoff A (eds) *Encyclopedia*  
654 *of virology*. Academic Press, London, p 121–126</edb>

655 <jrn>Williamson SJ, Cary SC, Williamson KE, Helton RR, Bench SR, Winget D, Wommack KE  
656 (2008) Lysogenic virus-host interactions predominate at deep-sea diffuse-flow hydrothermal  
657 vents. *ISME J* 2:1112–1121 [PubMed](#) [doi:10.1038/ismej.2008.73](#)</jrn>

658 <jrn>Winget DM, Williamson KE, Helton RR, Wommack KE (2005) Tangential flow diafiltration:  
659 an improved technique for estimation of virioplankton production. *Aquat Microb Ecol* 41:221–  
660 232 [doi:10.3354/ame041221](#)</jrn>

661 <jrn>Winter C, Herndl GJ, Weinbauer MG (2004) Diel cycles in viral infection of bacterioplankton  
662 in the North Sea. *Aquat Microb Ecol* 35:207–216 [doi:10.3354/ame035207](#)</jrn>

663 <jrn>Winter C, Kerros ME, Weinbauer MG (2009) Seasonal and depth-related dynamics of  
664 prokaryotes and viruses in surface and deep waters of the northwestern Mediterranean Sea.  
665 *Deep Sea Res I* 56:1972–1982 [doi:10.1016/j.dsr.2009.07.003](#)</jrn>

666 <jrn>Winter C, Bouvier T, Weinbauer MG, Thingstad TF (2010) Trade-offs between competition  
667 and defense specialists in unicellular planktonic organisms – the ‘killing the winner’ hypothesis  
668 revisited. *Microbiol Mol Biol Rev* 74:42–57 [PubMed](#) [doi:10.1128/MMBR.00034-09](#)</jrn>



669 <jrn>Winter C, Köstner N, Kruspe CP, Urban D, Muck S, Reinthaler T, Herndl GJ (2018) Mixing  
670 alters the lytic activity of viruses in the dark ocean. *Ecology* 99:700–713 [PubMed](#)  
671 [doi:10.1002/ecy.2135](#)</jrn>

672 <jrn>Wommack KE, Colwell RR (2000) Virioplankton: viruses in aquatic ecosystems. *Microbiol*  
673 *Mol Biol Rev* 64:69–114 [PubMed](#) [doi:10.1128/MMBR.64.1.69-114.2000](#)</jrn>

674 <jrn>Yamada Y, Tomaru Y, Fukuda H, Nagata T (2018) Aggregate formation during the viral lysis  
675 of a marine diatom. *Front Mar Sci* 5:167 [doi:10.3389/fmars.2018.00167](#)</jrn>

676 <jrn>Yang Y, Yokokawa T, Motegi C, Nagata T (2014) Large-scale distribution of viruses in deep  
677 waters of the Pacific and Southern Oceans. *Aquat Microb Ecol* 71:193–202  
678 [doi:10.3354/ame01677](#)</jrn>

679 <jrn>Zimmerman AE, Howard-Varona C, Needham DM, John SG and others (2020) Metabolic and  
680 biogeochemical consequences of viral infection in aquatic ecosystems. *Nat Rev Microbiol*  
681 18:21–34 [PubMed](#)</jrn>

682 Table 1. Averaged water layer properties of selected physico-chemical parameters in the western North Atlantic basin. *T*: temperature;  $\sigma_t$ : water  
683 density. SSL: subsurface layer; O<sub>2</sub>-min: oxygen minimum zone; LSW: Labrador Sea Water; NADW: North Atlantic Deep Water; DSOW: Denmark  
684 Strait Overflow Water. **Values in parentheses: SD**

Water mass	No. of samples	Depth (m)	Depth range (m)	<i>T</i> (°C)	$\sigma_t$ (kg m <sup>-3</sup> )	Oxygen (μmol kg <sup>-1</sup> )
SSL	33	100	90–110	8.7 (4.74)	27.20 (0.489)	260 (38.4)
O <sub>2</sub> -min	15	402	180–740	7.9 (2.39)	27.32 (0.194)	187 (51.2)
LSW	32	1324	710–2090	3.4 (0.32)	27.77 (0.031)	279 (7.3)
NADW	23	2537	1980–3250	3.0 (0.19)	27.84 (0.014)	272 (6.5)
DSOW	22	3031	1220–3870	1.9 (0.42)	27.91 (0.24)	288 (12.7)

685  
686 Table. 2. Prokaryotic and viral parameters in the western North Atlantic basin. PA: prokaryotic abundance; VA: viral abundance; V1: viral subgroup 1;  
687 V2: viral subgroup 2; V3: viral subgroup 3; VPR: virus-to-prokaryote ratio. SSL: subsurface layer; O<sub>2</sub>-min: oxygen minimum zone; LSW: Labrador  
688 Sea Water; NADW: North Atlantic Deep Water; DSOW: Denmark Strait Overflow Water. **PHP** and PA data are from Reinthaler et al. (2006).  
689 **Numbers in parentheses: SD; letters in parentheses:** Mann-Whitney tests of pairs of water masses (levels not connected by same letter are significantly  
690 different [*p* < 0.05]).

Water mass	No. of samples	PA (× 10 <sup>8</sup> cells l <sup>-1</sup> )	VA (× 10 <sup>9</sup> particles l <sup>-1</sup> )	V1 (%)	V2 (%)	V3 (%)	VPR
SSL	33	3.58 (0.94) (A)	3.1 (1.8) (A)	73.1 (4.0) (A)	20.3 (4.1) (A)	6.5 (1.5) (A)	9.8 (5.6) (A)
O <sub>2</sub> -min	15	1.27 (0.49) (B)	2.7 (2.9) (A,B)	78.3 (9.1) (A,B)	17.2 (7.8) (A)	4.5 (1.6) (B)	27.0 (38.9) (A)
LSW	32	0.50 (0.17) (C)	1.5 (0.8) (A,B)	66.3 (7.3) (B,C)	28.8 (7.5) (A)	4.8 (1.7) (B)	32.4 (21.3) (A)
NADW	25	0.30 (0.06) (C)	1.2 (0.9) (B)	69.0 (5.8) (B,C)	26.4 (5.1) (B)	4.6 (2.0) (B)	40.8 (26.5) (A,B)
DSOW	22	0.43 (0.26) (C)	2.3 (2.1) (B)	69.2 (8.1) (C)	26.1 (7.3) (B)	4.6 (1.7) (B)	42.9 (18.2) (B)
Kruskal-Wallis (p-value)		<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001

692 Table 3. Spearman rank correlations ( $\rho$ ) of parameters of the entire study. Values in **bold**:  $p < 0.05$ ; values in *italics*:  $p > 0.5$ . Physico-chemical and  
693 prokaryotic data are from Teira et al. (2005) and Reinthaler et al. (2006). AOU: Apparent oxygen utilization; PA: prokaryotic abundance; PHP:  
694 prokaryotic heterotrophic production; VA: viral abundance; V1: viral subgroup 1; V2: viral subgroup 2; V3: viral subgroup 3; VPR: virus-to-  
695 prokaryote ratio

	Depth	Temp	Salinity	O <sub>2</sub>	AOU	PO <sub>4</sub>	NO <sub>3</sub>	PHP	PA	VA	%V1	%V2	%V3
PA	<b>-0.887</b>	<b>0.760</b>	0.109	-0.059	<b>-0.426</b>	<b>-0.463</b>	<b>-0.449</b>	<b>0.622</b>					
VA	<b>-0.508</b>	<b>0.411</b>	-0.104	-0.101	<b>-0.349</b>	<b>-0.314</b>	<b>-0.284</b>	<b>0.474</b>	<b>0.450</b>				
%V1	<b>-0.239</b>	<b>0.249</b>	0.209	-0.216	0.078	0.096	0.097	<b>0.235</b>	0.076	<b>0.390</b>			
%V2	<b>0.260</b>	<b>-0.238</b>	-0.174	0.158	0.009	-0.026	-0.032	<b>-0.241</b>	-0.123	<b>-0.438</b>	<b>-0.928</b>		
%V3	-0.010	-0.060	-0.187	<b>0.219</b>	<b>-0.268</b>	<b>-0.249</b>	<b>-0.245</b>	-0.073	0.167	-0.139	<b>-0.812</b>	<b>0.675</b>	
VPR	<b>0.602</b>	<b>-0.608</b>	-0.167	<b>0.405</b>	-0.098	0.088	0.096	<b>-0.372</b>	<b>-0.717</b>	0.094	0.180	-0.161	<b>-0.273</b>

696

697 Table 4. Spearman rank correlations ( $\rho$ ) of parameters in the North Atlantic Deep Water. Values in **bold**:  $p < 0.05$ . Distance: distance from the stations  
698 closest to the Greenland–Iceland Ridge. Physico-chemical and prokaryotic data are from Teira et al. (2005) and Reinthaler et al. (2006). AOU:  
699 Apparent oxygen utilization; PA: prokaryotic abundance; PHP: prokaryotic heterotrophic production; VA: viral abundance; V1: viral subgroup 1; V2:  
700 viral subgroup 2; V3: viral subgroup 3; VPR: virus-to-prokaryote ratio

	Distance	Temp	Salinity	O <sub>2</sub>	AOU	PO <sub>4</sub>	NO <sub>3</sub>	PA	PHP	VA	%V1	%V2	%V3
Temp	<b>0.556</b>												
Salinity	<b>0.624</b>	<b>0.623</b>											
O <sub>2</sub>	<b>-0.824</b>	<b>-0.730</b>	<b>-0.859</b>										
AOU	<b>0.855</b>	<b>0.624</b>	<b>0.834</b>	<b>0.983</b>									
PO <sub>4</sub>	<b>0.872</b>	<b>0.760</b>	<b>0.814</b>	<b>-0.942</b>	0.029								
NO <sub>3</sub>	<b>0.861</b>	<b>0.792</b>	<b>0.821</b>	<b>-0.938</b>	<b>0.915</b>	<b>0.988</b>							
PA	-0.336	-0.119	<b>-0.700</b>	<b>0.491</b>	<b>-0.517</b>	-0.436	-0.400						
PHP	0.014	-0.134	-0.140	-0.040	0.076	0.041	0.045	-0.182					
VA	<b>-0.809</b>	-0.310	-0.427	<b>0.599</b>	<b>-0.652</b>	<b>-0.655</b>	<b>-0.658</b>	0.303	0.246				
%V1	0.447	0.058	0.150	-0.253	0.308	0.276	0.212	-0.223	0.269	-0.223			
%V2	<b>-0.527</b>	-0.177	-0.245	0.337	-0.382	-0.373	-0.314	0.284	-0.260	0.230	<b>-0.927</b>		

%V3	0.082	0.174	−0.006	−0.002	−0.008	0.119	0.123	0.036	−0.151	−0.051	<b>−0.552</b>	0.275	
VPR	<b>−0.680</b>	−0.147	−0.217	0.385	−0.415	<b>−0.499</b>	<b>−0.506</b>	0.144	0.211	<b>0.875</b>	−0.116	0.141	−0.190

701

702 Table 5. Prokaryotic and viral production, viral infection and virus-mediated mortality of prokaryotic plankton in the North Atlantic Deep Water  
703 (NADW). Distance is given in km from the origin of NDAW in the Greenland–Iceland–Norwegian Sea. PHP: heterotrophic prokaryotic production;  
704 VP: viral production; **BS: burst size**; FIC: fraction of visibly infected cells; VMMP: virus-mediated mortality of prokaryotes; VP<sub>i</sub>: induced viral  
705 production; FLC: fraction of lysogenic cells. ND: not determined; NA: not applicable

Station	Distance (km)	PHP ( $\times 10^6 \text{ l}^{-1} \text{ d}^{-1}$ )	VP ( $\times 10^7 \text{ l}^{-1} \text{ d}^{-1}$ )	BS	FIC (%)	VMMP <sub>VP</sub> (%)	VMMP <sub>FIC</sub> (%)	VP <sub>i</sub> ( $\times 10^7 \text{ l}^{-1} \text{ d}^{-1}$ )	FLC (%)
1	5500	2.95	0.23	24	11.7	3	15.2	0.06	5.8
6	4800	ND	0.21	ND	8.3	ND	10.0	0.59	23.3
9	4000	0.66	0.51	ND	0.6	32	0.6	3.83	4.5
13	2900	1.23	0.98	28	15.6	28	22.2	1.88	29.9
22	1400	0.21	0.80	ND	16.8	139	24.6	0.93	19.5
27	670	1.16	1.49	ND	19.2	46	29.7	3.11	40.1
31	200	0.79	2.40	33	22.0	108	36.6	1.66	15.2
Average	NA	1.17	0.95	28	13.5	59	19.9	1.72	20.5
SD	NA	0.94	0.78	5	7.3	53	12.2	1.36	13.8

706

707 Fig. 1. Study area in the North Atlantic Ocean. Dots: individual stations sampled during the  
708 TRANSAT-II

709

710 Fig. 2. Distribution of prokaryotic and viral parameters along the TRANSAT-II cruise in the North  
711 Atlantic. Dots: sampling locations; V1, V2 and V3 are viral subgroups 1, 2 and 3, respectively

712

713 Fig. 3. Variation of viral abundance, virus-to-prokaryote ratios (VPR) and 2 viral subgroups (as  
714 determined by flow cytometry) in the North Atlantic Deep Water with distance from its formation  
715 in the Greenland–Iceland–Norwegian (GIN) Sea. (A) Viral abundance ( $p < 0.0001$ ); (B) VPR ( $p <$   
716  $0.0001$ ); (C) viral subgroup 1 (V1) ( $p < 0.005$ ); (D) viral subgroup 2 (V2) ( $p < 0.01$ ). Regression  
717 parameters are given in the plots

718

719 Fig. 4. Variation of viral production, the fraction of infected cells (FIC), induced viral production  
720 ( $VP_i$ ) and fraction of lysogenic cells (FLC) in the North Atlantic Deep Water with distance from its  
721 formation in the Greenland–Iceland–Norwegian (GIN) Sea. Data are averages  $\pm$  range of duplicate  
722 incubations. (A) Viral production ( $p = 0.0005$ ); (B) FIC ( $p < 0.05$ ); (C)  $VP_i$ ; (D) FLC. Regression  
723 parameters are given in the plots