



HAL
open science

EVOLUTION OF THE SHOOT DENSITY OF THE POSIDONIA SEAGRASS BED OF CALVI BAY (CORSICA)

M Soullard, I Bourge, J Fogel, D Lardinois, T Mathieu, C Veeschkens, D Bay, P Dauby, J M Bouquegneau

► **To cite this version:**

M Soullard, I Bourge, J Fogel, D Lardinois, T Mathieu, et al.. EVOLUTION OF THE SHOOT DENSITY OF THE POSIDONIA SEAGRASS BED OF CALVI BAY (CORSICA). Vie et Milieu / Life & Environment, 1994, pp.199-201. <hal-03048001>

HAL Id: hal-03048001

<https://hal.sorbonne-universite.fr/hal-03048001v1>

Submitted on 9 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

EVOLUTION OF THE SHOOT DENSITY OF THE *POSIDONIA* SEAGRASS BED OF CALVI BAY (CORSICA)

M. SOULLARD^(1,2), I. BOURGE⁽¹⁾, J. FOGEL⁽¹⁾, D. LARDINOIS⁽¹⁾,
T. MATHIEU⁽¹⁾, C. VEESCHKENS⁽¹⁾, D. BAY⁽³⁾,
P. DAUBY⁽¹⁾ and J.M. BOUQUEGNEAU⁽¹⁾

⁽¹⁾ Laboratoire d'Océanologie, Université de Liège, B6-Sart Tilman, B-4000 Liège, Belgique

⁽²⁾ Università di Corsica, Faculté des Sciences, Bât. 018, F-20250 Corte, Corse

⁽³⁾ STARESO, BP 33, F-20260 Calvi, Corse

POSIDONIA OCEANICA
DENSITÉ DE FAISCEAUX

POSIDONIA OCEANICA
SHOOT DENSITY

RÉSUMÉ – La densité de faisceaux de *Posidonia oceanica* (L.) Delile a été étudiée dans la baie de Calvi par Bay en 1975 et 1977. Nous avons effectué les mêmes mesures dans le même herbier en 1991. Aucune régression n'a été mise en évidence. Au contraire, nous avons observé une augmentation de la densité des faisceaux à 10 mètres de profondeur.

ABSTRACT – The density of seagrass shoots in the *Posidonia* bed of Calvi Bay was studied by Bay in 1975 and 1977. The present paper reports the shoot density of the same seagrass bed in 1991. No regression has been observed. On the contrary, an increase of shoot density has been observed at 10 metres depth.

INTRODUCTION

Posidonia oceanica (L.) Delile seagrass beds play an important role in building up and protecting the Mediterranean coast and they provide refuge, habitat and nourishment for many animal and plant species. Since two decades, several authors have noticed the regression of these meadows, which can correspond either to a recession of the whole community (especially in deeper areas) or to a decrease of the shoot density (Peres, 1984), in response to various human activities : urban and industrial sewage, mechanized fishing activities, yachting and dredging, rise of water turbidity, etc.

Between 1975 and 1977, one of us (D.B.) has extensively studied the structure and production of the *Posidonia* meadow in Calvi Bay (Bay, 1978, 1984). Since these works, many scuba diving were performed in that area without any notice of obvious decrease of the surface area of the bed; nevertheless, during the same period a quite complete disappearance of another marine plant, *Cystoseira balearica*, was observed by Hoffmann *et al.* (1988) who hypothesized an eutrophication effect. As no estimation of the *Posidonia* shoot density has been performed since 1977, we thought it was worth making a new estimation in order to follow the evolution of the bed during these last 15 years.

METHODS

In May 1991, we have performed countings on the same sites than those previously studied by Bay (1978), *i.e.* at 10 and 30 metres depth along a transect from the oceanographic station STARESO to the citadel of Calvi (see map on fig. 1, stations 1 and 2). Moreover, countings were also made at a distance of 50 m on both sides of station 1 on the 10 m isobath (stations 1' and ", fig. 1). The countings were performed following the same methodology than Bay's one : a metal circle (30 cm diameter) is randomly thrown down on the bed and the number of included seagrass shoots is recorded by a diver. Most of the countings were performed by three of us (I.B., D.L. & C.V.), and the reliability of their estimations has been tested against Bay's observations : the metal circle has been settled on two sites chosen for their respective low and high shoot density. The shoots have been successively counted by D.B., I.B., D.L. & C.V., after what they were uprooted and counted in the laboratory. The results are shown in Table I A, from which it appears that the difference between the observations of respectively D.B., and the other three divers does not exceed ten percent of the average value. The observations are very accurate in less dense areas, but they are underesti-




Fig. 1. - Location of the stations.

Table I. - A, Comparison between the shoot density estimations made by four different divers. B, Mean shoot number per m² at 10 m depth (May 1991). Location of the stations : see Fig. 1. n = number of counts ; m = mean density ; S.D. = standard deviation. C, Time variation of the shoot number per m² at 10 m depth. D, time variation of the shoot number per m² at 30 m depth.

Real number of shoots	Estimation by D.B	Estimation by I.B	Estimation by D.L.	Estimation by C.V.
20	18	21	20	19
46	41	42	41	40

Station	n	m	S.D.	P (t-test)	P(t-test)
1	32	500	142	0,935	0,060
1'	32	503	150		
1''	32	434	134		
1+1'+1''	96	481	143		

Year	n	m	S.D.	P(t-test)	P(t-test)	Ref.
1975	104	407	239	0,982	0,009	BAY,1984
1977	32	408	156			
1991	96	481	143			0,016

Year	n	m	S.D.	P(t-test)	P(t-test)	Ref.
1975	39	223	95	0,039	0,010	BAY, 84
1977	32	174	101			
1991	32	168	75			0,788

mated in most dense ones. However, that underestimation never exceeds 12 % of the actual value.

RESULTS AND DISCUSSION

Tables I B and C are presenting the results of the countings performed at 10 metres depth. The three sampled stations are quite similar even if station 1" is somewhat less dense than the other two ones (Table I B). Nevertheless, obtained values are 20 % higher than those recorded by Bay in 1975 and 1977 (Table I C).

On the contrary, the results obtained for 30 metres depth (Table I D) reveal a significant change in shoot density at that depth when comparing them with those of 1977. This observation suggests that these variations could be the result of a year-to-year variation of the activity of the meadow instead of the result of a long-term one.

From the statistical treatment of our results, we are tempted to conclude that the general trend of the seagrass bed of Calvi Bay, at 10 m depth, is a continuing increase at the shoot density point of view. Indeed, 15 years ago, the meadow was balancing between scattered and dense status, according to the classification of Giraud (1977 – "herbier clairsemé et dense"), but now it must be considered as a true dense one. However, it has been recently shown that the branching rate of the *Posidonia* shoots is high, implying a potential high turnover rate, with a mean two years virtual life span of the shoots (Boudouresque, personal communication). So the following questions arise, what is the actual signification of such an increase in the shoot density? Is it the consequence of a continuous increase of the density of the shoots, or is this observation to be considered as an ac-

cidental one, resulting from both an unusual high branching rate and a low mortality rate of the shoots during the two previous years? A continuing scheme of shoot density evaluations is planned for the next years to try and answer that question.

ACKNOWLEDGEMENTS – This work was supported by grants from the Belgian National Foundation for Scientific Research (FNRS n° 2 : 4508 : 88) and from the European Community (contract CEC/STEP-CT90-0063-c-CD)). We are grateful to Prof. C.F. Boudouresque and to Dr. P. Dubois & M. Warneau for critical review of the manuscript.

REFERENCES

- BAY D., 1978. Etude *in situ* de la production primaire d'un herbier de posidonies (*Posidonia oceanica* (L.) Delile) de la Baie de Calvi – Corse. Thèse Doct. Univ. Liège.
- BAY D., 1984. A field study of the growth dynamics and productivity of *Posidonia oceanica* (L.) Delile in Calvi Bay, Corsica. *Aquat. Bot.* **20** : 43-64.
- HOFFMANN L., S. CLARISSE, X. DETIENNE, A. GOFFART, R. DENARD & V. DEMOULIN, 1988. Evolution of the populations of *Cystoseira balearica* (Phaeophyceae) and epiphytic Bangiophyceae in the Bay of Calvi (Corsica) in the last eight years. *Bull. Soc. roy. Sci. Lg.* **57** : 263-273.
- GIRAUD G., 1977. Essai de classement des herbiers de *Posidonia oceanica* (Linné) Delile. *Bot. Mar.* **20** (8) : 487-491.
- PERES J.M., 1984. La régression des herbiers à *Posidonia oceanica*. International Workshop on *Posidonia oceanica* Beds, Boudouresque C.F., Jeudy De Grissac A. et Olivier J. edit., GIS Posidonie publ., Fr., **1** : 445-454.

Reçu le 15 avril 1994; received April 15, 1994
 Accepté le 10 mai 1994; accepted May 10, 1994