



HAL
open science

Identical trends of SARS-Cov-2 transmission and retail and transit mobility during non-lockdown periods

Bernard Cazelles, Catherine Comiskey, Benjamin Nguyen van Yen, Clara Champagne, Benjamin Roche

► To cite this version:

Bernard Cazelles, Catherine Comiskey, Benjamin Nguyen van Yen, Clara Champagne, Benjamin Roche. Identical trends of SARS-Cov-2 transmission and retail and transit mobility during non-lockdown periods. *International Journal of Infectious Diseases*, 2021, 10.1016/j.ijid.2021.01.067 . hal-03134842v1

HAL Id: hal-03134842

<https://hal.sorbonne-universite.fr/hal-03134842v1>

Submitted on 8 Feb 2021 (v1), last revised 9 Apr 2021 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Journal Pre-proof

Identical trends of SARS-Cov-2 transmission and retail and transit mobility during non-lockdown periods

Bernard Cazelles, Catherine Comiskey, Benjamin Nguyen Van Yen, Clara Champagne, Benjamin Roche



PII: S1201-9712(21)00079-5
DOI: <https://doi.org/10.1016/j.ijid.2021.01.067>
Reference: IJID 5076

To appear in: *International Journal of Infectious Diseases*

Received Date: 15 December 2020
Revised Date: 25 January 2021
Accepted Date: 26 January 2021

Please cite this article as: Cazelles B, Comiskey C, Nguyen Van Yen B, Champagne C, Roche B, Identical trends of SARS-Cov-2 transmission and retail and transit mobility during non-lockdown periods, *International Journal of Infectious Diseases* (2021), doi: <https://doi.org/10.1016/j.ijid.2021.01.067>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier.

Identical trends of SARS-Cov-2 transmission and retail and transit mobility during non-lockdown periods

Bernard Cazelles^{a,b}, Catherine Comiskey^c, Benjamin Nguyen Van Yen^{b,d}, Clara Champagne^{e,f} and Benjamin Roche^g

a. UMMISCO, Sorbonne Université, Paris, France

b. Eco-Evolution Mathématique, IBENS, UMR 8197, CNRS, Ecole Normale Supérieure, Paris, France

c. School of Nursing and Midwifery, Trinity College Dublin, The University of Dublin, Dublin, Ireland

d. Institut Pasteur, Unité de Génétique Fonctionnelle des Maladies Infectieuses, Paris, France

e. Swiss Tropical and Public Health Institute, Basel, Switzerland

f. University of Basel, Basel, Switzerland

g. MIVEGEC, IRD, CNRS and Université de Montpellier, Montpellier, France

Highlights

- We document a strong relationship between SARS-Cov-2 transmission and mobility during the time period between the two first waves.
- The trends of both mobility and viral transmission evolve almost in parallel during this time period.
- This suggests that the evolution of mobility patterns would help Public Health authorities to anticipate transmission increases.
- This result is important from a Public Health perspective when relaxing control measures.

Abstract

Recent literature strongly supports the idea that mobility reduction and social distancing play a crucial role in transmission of SARS-Cov-2 infections. It was shown during the first wave that mobility restrictions reduce significantly infection transmission. Here we document another relationship and show that, in the period between the first two COVID-19 waves, there exists a high positive correlation between the trends of SARS-Cov-2 transmission and mobility. These two trends oscillate simultaneously and increased mobility following lockdown relaxation has a significant positive relationship with increased transmission. From a public health perspective, these results highlight the importance of following the evolution of mobility when relaxing mitigation measures to anticipate the future evolution of the spread of the SARS-Cov-2.

Keywords: SARS-Cov-2

Transmission

Effective Reproduction Number

Mobility

Mobility of hosts and/or vectors has always had a large impact on the propagation of the transmission of diseases (Tizzoni et al., 2014). This is once again the case for the Covid-19 pandemic (Kraemer et al., 2020).

Recent literature strongly supports the idea that mobility reduction and social distancing played a crucial role in the reduction of SARS-CoV-2 infections during the first wave. Some studies concluded that mobility patterns correlate with the prevalence of COVID-19 and that travel reduction had a positive effect on reducing the transmission of SARS-CoV-2 (Badr et al., 2020; Chang et al., 2020). Some other works have used the changes in mobility patterns to estimate the effect of mitigation measures on the reproductive number. They showed that the drop in mobility, overlapping with the introduction of mitigation measures induced a sharp reduction in transmission (Lemaitre et al., 2020; Park et al., 2020). But all these interesting results are limited to the first wave, before and during the lockdown period.

Nevertheless, it remains unknown if mobility increases following lockdown relaxation is actually associated with an enhanced transmission of SARS-Cov-2. Lockdowns impact a myriad of dimensions in our society, and not only on mobility patterns. This makes it difficult to draw a causal link between mobility and viral transmission from lockdown data. Instead, to better understand this link, we propose investigating this link in the period between lockdown periods, to potentially use mobility data as an early signal for implementing public health measures.

Here for the first time, to the best of our knowledge, we document a high correlation between the trends of SARS-Cov-2 transmission and mobility in the period between the first two Covid-19 waves in some French regions and in Ireland. Our aim was to compare different geographical settings with similar population size. We quantified the change in

transmission of SARS-CoV-2 by computing the effective reproduction number $R_{eff}(t)$. The effective reproduction number is defined by the average number of secondary cases at time t arising from a primary infected case. It is an important metric for measuring time-varying transmissibility and assessing the effectiveness of different interventions. The classical statistical methods for estimating $R_{eff}(t)$ have many shortcomings due to the characteristics of the transmission of this virus: silent transmission and major time variation in the reporting of cases due to lack of timely or appropriate testing (Gostic et al., 2020). Thus we used an original mechanistic method based on a stochastic model (Fig. A1 and eqs A1-A3) with time-varying parameters (Cazelles et al., 2018) and inferred with a Bayesian method using hospital data (Cazelles et al., 2021). We used daily mobility data provided by Google, which is a proxy for real-time trends in movement patterns and human behavior (Google, 2020). These mobility data measure the percentage of change relative to the pre-pandemic baseline mobility and measure visits and the time spent in: retail and recreation, grocery and pharmacy, transit stations, workplace and residential.

Our analysis revealed that trends in transmission as estimated from well documented hospital data correlate strongly with trends in mobility patterns within and between the observed first and second epidemic waves for retail and recreation mobility, as well as transit station mobility. The $R_{eff}(t)$ increased and oscillated with mobility with a highly significant correlation between the 15th of May and the 15th of October, with Pearson correlation coefficients above 0.5 (Figure). We show that these relationships are strong, particularly for retail and recreation mobility and transit station mobility. Additionally, we computed the Cross-Correlation Functions for these datasets that revealed the correlations were maximal for a delay between 0 and 10 days, and that this delay varied depending on the region (Figs A2-A3).

Earlier findings showed that mobility reductions can dramatically reduce infections, our results now confirm the reverse relationship; increased mobility following lockdown relaxation has a significant, positive relationship with increased transmission. This suggests the importance of following the evolution of mobility data at different spatial resolutions to anticipate the future evolution of the spread of the SARS-Cov-2 and to guide public health policy.

Additionally, the strong relationship between retail and recreational mobility and increased transmissibility suggests that a return to baseline human activity poses a significant risk of increased infection. Moreover, even if a direct causal link cannot be drawn, these strong correlations indicate that an increase in retail and recreational mobility promote high-intensity contact between people and also interaction between people of different households, resulting in increased SARS-CoV-2 transmission. This strong relationship is in total agreement with those of Chang et al. (2020) who have also demonstrated that mobility network models can inform the reopening of society.

Our findings illustrate the importance of mobility at defined regional levels and perhaps more importantly, they highlight that mobility measures of transit and retail are more strongly correlated with transmission than other forms of mobility. These findings will aid the refinement of future possible lockdown policies and mitigation measures as we await the roll out of imminent vaccines.

Funding

B. C. and B. R. are partially supported by a grant ANR Flash Covid-19 from the “Agence Nationale de la Recherche” (DigEpi).

Ethical approval

Not required.

Conflict of interest

None.

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper

References

Badr HS, Du H, Marshall M, Dong E, Squire MM, Gardner LM. Association between mobility patterns and COVID-19 transmission in the USA: A mathematical modelling study. *The Lancet Infectious Diseases* 2020; 20: 1247-1254.

Cazelles B, Champagne C, Dureau J. Accounting for non-stationarity in epidemiology by embedding time-varying parameters in stochastic models. *PLoS Computational Biology* 2018; 14: e1006211.

Cazelles B, Nguyen Van Yen B, Champagne C, Comiskey C. Dynamics of the COVID-19 epidemic in Ireland under mitigation. 2021; Research Square preprint (doi:10.21203/rs.3.rs-143697/v1).

Chang S, Pierson E, Koh PW, Gerardin J, Redbird B, Grusky D, Leskovec J. Mobility network models of COVID-19 explain inequities and inform reopening. *Nature* 2021;589:82-87. (doi:10.1038/s41586-020-2923-3).

Google (2020). Google COVID-19 Community Mobility Reports.

<https://www.google.com/covid19/mobility/> Accessed: 30 Nov 2020.

Gostic KM, McGough L, Baskerville EB, Abbott S, Joshi K, Tedijanto C, Kahn R, Niehus R, Hay JA, De Salazar PM, Hellewell J, Meakin S, Munday JD, Bosse NI, Sherrat K, Thompson RN, White LF, Huisman JS, Scire J, Bonhoeffer S, Stadler T, Wallinga J, Funk S, Lipsitch M, Cobey S. Practical considerations for measuring the effective reproductive number, R_t . *PLoS Comput Biol.* 2020;16:e1008409.

Kraemer MUG, Yang CH, Gutierrez B, Wu CH, Klein B, Pigott DM; Open COVID-19 Data Working Group, du Plessis L, Faria NR, Li R, Hanage WP, Brownstein JS, Layan M, Vespignani A, Tian H, Dye C, Pybus OG, Scarpino SV. The effect of human mobility and control measures on the COVID-19 epidemic in China. *Science* 2020;368:493-497.

Lemaitre JC, Perez-Saez J, Azman A, Rinaldo A, Fellay J. Assessing the impact of non-pharmaceutical interventions on SARS-CoV-2 transmission in Switzerland. *Swiss Med Wkly* 2020; 150: w20295.

Park SW, Sun K, Viboud C, Grenfell BT, Dushoff J. Potential roles of social distancing in mitigating the spread of coronavirus disease 2019 (COVID-19) in South Korea. *Emerging Infectious Diseases* 2020; 26: 2697-2700.

Tizzoni M, Bajardi P, Decuyper A, Kon Kam King G, Schneider CM, Blondel V, Smoreda Z, González MC, Colizza V. On the Use of Human Mobility Proxies for Modeling Epidemics. PLoS Comput Biol 2014; 10: e1003716.

Figure. Time evolution of the estimated $R_{eff}(t)$ (black lines) and Google Mobility (retail and recreation mobility (continuous blue lines) and transport mobility (dashed blue lines)) in Ile-de-France region (A) and in Ireland (B). The mobility time series have been smoothed using moving average over a 7 days window. In A/ $R_{eff}(t)$ is computed for two different models with or without accounting for hospital discharges. Pearson Correlation in Ile-de-France (model with hospital discharges) with retail and recreation mobility 0.70 and with transport mobility 0.77. Pearson Correlation in Ile-de-France (model without hospital discharges) with retail and recreation mobility 0.64 and with transport mobility 0.70. Pearson Correlation in Ireland with retail and recreation mobility 0.86 and with transport mobility 0.56. Figs A2-A3 display the significance of the Cross-Correlation functions (CCF). Vertical dashed lines are main mitigation measures (lockdown and curfew) and the horizontal dashed lines are the threshold limit for $R_{eff}(t)$.

