



HAL
open science

Evidence of early circulation of SARS-CoV-2 in France: findings from the population-based “CONSTANCES” cohort

Fabrice Carrat, Julie Figoni, Joseph Henny, Jean-Claude Desenclos, Sofiane Kab, Xavier de Lamballerie, Marie Zins

► To cite this version:

Fabrice Carrat, Julie Figoni, Joseph Henny, Jean-Claude Desenclos, Sofiane Kab, et al.. Evidence of early circulation of SARS-CoV-2 in France: findings from the population-based “CONSTANCES” cohort. *European Journal of Epidemiology*, 2021, 10.1007/s10654-020-00716-2 . hal-03135488

HAL Id: hal-03135488

<https://hal.sorbonne-universite.fr/hal-03135488>

Submitted on 9 Feb 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Title

Evidence of early circulation of SARS-CoV-2 in France – findings from the population-based “CONSTANCES” cohort.

Authors

Fabrice Carrat¹✉, Julie Figoni², Joseph Henny^{3,4}, Jean-Claude Desenclos², Sofiane Kab^{3,4}, Xavier de Lamballerie⁵, Marie Zins^{3,4}.

Short title

Evidence of early circulation of SARS-CoV-2 in France

Affiliations

¹ Sorbonne Université, Inserm, Institut Pierre-Louis d’Epidémiologie et de Santé Publique, Département de Santé Publique, APHP.Sorbonne Université, Paris, France

² Santé publique France, Saint-Maurice, France

³ Paris University, Paris, France

⁴ Paris Saclay University, Inserm UMS 11 Villejuif, France

⁵ Unité des Virus Emergents, UVE: Aix Marseille Univ, IRD 190, INSERM 1207, IHU Méditerranée Infection, 13005, Marseille, France

✉ Correspondance to

Pr Fabrice Carrat, Institut Pierre Louis d’Épidémiologie et de Santé Publique, 27 rue Chaligny, 75571 PARIS CEDEX 12, France. Tel : +33 (0)1 71 97 01 10.

fabrice.carrat@iplesp.upmc.fr

ORCID

F Carrat [0000-0002-8672-7918](#)
JC Desenclos [0000-0003-0892-8252](#)
S Kab [0000-0001-6041-9602](#)
X de Lamballerie [0000-0001-7895-2720](#)
M Zins [0000-0002-4540-4282](#)

Abstract

Recent studies suggest that SARS-CoV-2 may have circulated earlier than January 2020 in Europe. To explore this hypothesis, we investigated the serological status for SARS-CoV-2 antibodies in 9,144 adults from a French general population-based cohort with serum samples routinely collected between November 4, 2019 and March 16, 2020. Samples were tested using a commercial enzyme-linked immunosorbent assay to detect antibodies directed against the S1 domain of the spike protein of the virus. Positive, indeterminate and a subset of negative specimens were also tested with a micro-neutralization assay.

We identified 353 participants with a positive anti-SARS-CoV-2 test, among whom 13 were sampled between November 2019 and January 2020 and were confirmed by neutralizing antibodies testing. Investigations in 11 of these participants revealed experience of symptoms possibly related to a SARS-CoV-2 infection in five participants or situations at risk of potential SARS-CoV-2 exposure in eight participants. Altogether our findings suggest early circulation of SARS-CoV-2 in Europe.

Key words

SARS-CoV-2, COVID-19, general population, cohort

Declarations

Funding

This study

The French National Research Agency [grant number ANR-10-COHO-06]

Santé Publique France: N°20DMIA014-0

Cohorts funding

The CONSTANCES Cohort Study is supported by the Caisse Nationale d'Assurance Maladie (CNAM), the French Ministry of Health, the Ministry of Research, the Institut national de la santé et de la recherche médicale. CONSTANCES benefits from a grant from the French National Research Agency [grant number ANR-11-INBS-0002] and is also partly funded by MSD, AstraZeneca, Lundbeck and L'Oreal.

Conflict of interest None

Ethics approval / Consent to participate

The cohort received ethical approval and all participants provided written consent to the cohort and additional consent to the current study. No consent was

Availability of data and material

In regards to data availability, data of the study are protected under the protection of health data regulation set by the French National Commission on Informatics and Liberty (Commission Nationale de l'Informatique et des Libertés, CNIL). The data can be available upon reasonable request to the corresponding author

(fabrice.carrat@iplesp.upmc.fr), after a consultation with the steering committee of the CONSTANCES cohort study. The French law forbids us to provide free access to raw data; access could however be given by the steering committee after legal

verification of the use of the data. Please, feel free to come back to us should you have any additional question.

Code availability NA

Authors' contributions

Study idea and design: Carrat, Desenclos, de Lamballerie, Zins

Data acquisition: Figoni, Henny, Kab, Zins

Data analysis and interpretation: Carrat, Desenclos, de Lamballerie, Zins

Drafting of the manuscript: Carrat

Critical revision of the manuscript for important intellectual content: All authors.

Acknowledgments

The authors warmly thank all the volunteers of the Constances cohort.

We thank Nathalie Vanhaverbeke, CEMKA, for conducting the participants interview.

We thank the staff of the Constance cohort that have worked with dedication and engagement to manage the data used for this study and to ensure continuing communication with the cohort participants.

In the virology department, we thank Dr Nadège Brisbarre and the technical staff for impeccable management of samples and serological assays.

In Santé publique France, we thank Harold Noel for his help building the investigation questionnaire and for his feedback on the study.

Introduction

The first identified cases of COVID-19 were detected on December 8, 2019, in Wuhan, China [1] and the first documented case in Europe was reported retrospectively in France in one patient with a diagnosis of pneumonia and a positive SARS-CoV-2 RT-PCR result on December 27, 2019 [2]. By April 4, 2020, local community transmission was reported in all continents and over 1 million cases of COVID-19 had been confirmed worldwide [3]. Although dedicated surveillance and contact tracing did not identify local transmission before the second half of February in Europe, there is accumulated evidence that SARS-CoV-2 circulated in early January 2020 in the East of France [4] and environmental studies suggest that the virus could have been present in December 2019 in Northern Italy [5]. A recent investigation of the presence of SARS-CoV-2 antibodies in 959 adults participating to a trial in Italy with blood samples collected between September 2019 to February 2020 identified 111 (11.6%) samples with a positive receptor-binding protein specific enzyme-linked immunosorbent assay (ELISA), among which 4 samples collected in October, 1 in November and 1 in February were also positive in a qualitative microneutralization assay [6]. This indicates that SARS-CoV-2 could have been present in Italy since the beginning of autumn 2019. However, information on antibody responses at the early stage of the SARS-CoV-2 spread in other European countries or worldwide remains scarce.

Participants and methods

We explored the serological status for SARS-CoV-2 antibodies in participants from the “CONSTANCES” cohort. Briefly, CONSTANCES is a general population-based cohort made up of a nationwide representative sample of 215,000 adults aged 18 to 69 at inclusion. Inclusion started in 2012, and serum samples are regularly collected

during the follow-up of participants for future analyses and stored in a centralized biobank. A complete description of the cohort design can be found in [3]. The cohort received ethical approval and all participants provided written consent to the cohort and additional consent to the current study.

We selected all 9144 serum samples collected between November 4, 2019 and March 16, 2020 in participants living in the 12 mainland French regions. All samples were centralized to the virology laboratory (Unité des virus Émergents, Marseille, France) for serological analysis. The serological analysis was performed using a commercial Elisa test (Euroimmun®, Lübeck, Germany) to detect anti-SARS-CoV-2 antibodies (IgG) directed against the S1 domain of the spike protein of the virus (ELISA-S). In accordance with the manufacturer's instructions a test was considered to be ELISA-S positive with an optical density ratio ≥ 1.1 , indeterminate between 0.8 and 1.1, and negative, <0.8 . All samples with an ELISA-S test ≥ 0.7 were also tested with an in-house micro-neutralization assay to detect neutralizing anti-SARS-CoV-2 antibodies (SN), as described elsewhere [7, 8]. A SN titer ≥ 40 was considered to be positive. Six replicates were performed to confirm positive SN.

Participants with both ELISA-S and SN positive tests in serum sampled before February 1, 2020 were interviewed to identify potential exposure to SARS-CoV-2 infection. A trained investigator collected standardized information on clinical details (in the participant and her/his relatives), history of possible exposure (notably history of travel in Asia), and any remarkable event in close contacts (e.g. unexplained pneumonia).

Results

Participants were aged 55 (min: 19, max: 79) years at sample collection; 4,623 (51%) were female; 1,503 (16%) were living in Ile de France, 935 (10%) in Grand-Est - these two regions being the French regions with the highest incidences of hospitalization for COVID-19 during the first semester 2020, 6,706 (73%) were living in the other 10 mainland regions.

Three-hundred and fifty-three (3.9%) participants were ELISA-S positive, 138 were undetermined and 8,653 were negative (undetermined and negative, 96.1%). The proportion of ELISA-S positive increased from 1.9% (42 of 2218) in November and 1.3% (20 of 1,534) in December to 5.0% (114 of 2,268) in January, 5.2% (114 of 2,179) in February and 6.7% (63 of 945) in the first half of March ($P < 0.001$, Trend test; Figure 1).

Neutralizing antibodies were detected in 44 (0.48%) participants (23 with a titer of 40, 12 with a titer of 80, 9 with a titer of 160), were undetermined in 15 participants, negative in 498 and not done in 8597 (Figure 2). Strikingly, 13 participants with positive ELISA-S and SN tests had been sampled between November 5, 2019 and January 30, 2020. Table 1 describes the serological results in these 13 participants, among whom 11 were interviewed. Six of those interviewed did not report any symptom during the weeks preceding the sample collection. Five participants experienced signs of viral respiratory illnesses, and 8 had close contact with persons who exhibited such signs or reported situations at risk of potential SARS-CoV-2 exposure. Of note, participant #7 who was tested positive on Nov 29, 2020 had a second serological sample collected in July 2020 with a positive ELISA-S test and negative SN test - this participant also tested positive in SARS-CoV-2 RT-PCR in September 2020 for new symptoms suggesting a possible reinfection.

Discussion

This report suggests that SARS-CoV-2 infection may have occurred as early as November 2019 in France. In several participants with both ELISA-S and SN positive results, we identified symptoms, history of possible exposures, or specific events compatible with early SARS-CoV-2 infection.

The critical issue is whether these findings might result from a lack of specificity of our serological methods. The proportion of ELISA-S positive reported may look overestimated given the time period the samples were collected. We used manufacturer-defined cutoff points for ELISA-S positivity, but the test specificity and the positive predictive values can increase by using other higher cut-off values [9]. A cut-off of 2.5 was shown to be 100% specific and would have led to select only 2 positive participants out of 13. However, all positive ELISA-S were confirmed by SN testing, and the specificity of SN was estimated at 100% over thousands of blood donors sampled in 2017-2018 when samples with a titer ≥ 40 were considered to be positive [7]. In addition, all positive SN results were confirmed in multiple replicates. Thus, also we can't exclude potential misclassifications of some participants, it is unlikely that all of them were false positive results, and detailed investigation suggested compatible history of exposure in several participants. On the opposite, it should be notice that our highly specific selection of participants might lack sensitivity and have led to excluding truly infected participants or early infected participants with waning immunity. Altogether our findings are in line with the reported Italian study [6] and suggest earlier than reported SARS-CoV-2 circulation in Europe.

References

1. Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, et al. Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus-Infected Pneumonia. *N Engl J Med*. 2020 Mar 26;382(13):1199-207. PubMed PMID: 31995857. Pubmed Central PMCID: PMC7121484. Epub 2020/01/30.
2. Deslandes A, Berti V, Tandjaoui-Lambotte Y, Alloui C, Carbonnelle E, Zahar JR, et al. SARS-CoV-2 was already spreading in France in late December 2019. *Int J Antimicrob Agents*. 2020 Jun;55(6):106006. PubMed PMID: 32371096. Pubmed Central PMCID: PMC7196402. Epub 2020/05/07.
3. Zins M, Goldberg M, team C. The French CONSTANCES population-based cohort: design, inclusion and follow-up. *Eur J Epidemiol*. 2015 Dec;30(12):1317-28. PubMed PMID: 26520638. Pubmed Central PMCID: PMC4690834. Epub 2015/11/02.
4. Gerbaud L, Guiguet-Auclair C, Breyse F, Odoul J, Ouchchane L, Peterschmitt J, et al. Hospital and Population-Based Evidence for COVID-19 Early Circulation in the East of France. *Int J Environ Res Public Health*. 2020 Sep 30;17(19). PubMed PMID: 33007976. Epub 2020/10/04.
5. La Rosa G, Mancini P, Bonanno Ferraro G, Veneri C, Iaconelli M, Bonadonna L, et al. SARS-CoV-2 has been circulating in northern Italy since December 2019: Evidence from environmental monitoring. *Sci Total Environ*. 2020 Aug 15;750:141711. PubMed PMID: 32835962. Pubmed Central PMCID: PMC7428442. Epub 2020/08/25.
6. Apolone G, Montomoli E, Manenti A, Boeri M, Sabia F, Hyseni I, et al. Unexpected detection of SARS-CoV-2 antibodies in the pre-pandemic period in Italy. *Tumori*. 2020 Nov 11:300891620974755. PubMed PMID: 33176598. Epub 2020/11/13.
7. Gallian P, Pastorino B, Morel P, Chiaroni J, Ninove L, de Lamballerie X. Lower prevalence of antibodies neutralizing SARS-CoV-2 in group O French blood donors. *Antiviral Res*. 2020 Sep;181:104880. PubMed PMID: 32679056. Pubmed Central PMCID: PMC7362788. Epub 2020/07/18.
8. Carrat F, de Lamballerie X, Rahib D, Blanché H, Lapidus N, Artaud F, et al. Seroprevalence of SARS-CoV-2 among adults in three regions of France following the lockdown and associated risk factors: a multicohort study. *MedRxiv*. 2020.
9. Meyer B, Torriani G, Yerly S, Mazza L, Calame A, Arm-Vernez I, et al. Validation of a commercially available SARS-CoV-2 serological immunoassay. *Clin Microbiol Infect*. 2020 Oct;26(10):1386-94. PubMed PMID: 32603801. Pubmed Central PMCID: PMC7320699. Epub 2020/07/01.

Table 1. Participants with positive ELISA-S and positive SN on blood samples collected between November 2019 and January 2020.

#	Age Class (yrs)	Sex	Date sample (year-month)	ELISA-S	SN	COVID-19 Symptoms	Risk factors and symptoms (travel, meetings, contact)
1	60-69	F	2019-11-	1.17	40	No	None
2	60-69	M	2019-11	2.00	40	No	None
3	30-39	F	2019-11	1.32	160	Yes	Her partner was sick with intense cough in October 2019.
4	30-39	M	2019-11	2.01	40	NA	NA
5	40-49	F	2019-11	1.16	40	No	None
6	30-39	F	2019-11	1.75	80	Yes	Travel in Spain in early November. She had daily encounters with a family member who had a respiratory illness of unknown origin between October and December. She suffered from dysgeusia, hyposmia, and cough before the sample was taken, but could not remember the date of illness
7	30-39	M	2019-11	2.50	40	Yes	The participant and his partner were sick with a severe cough in October 2019. He had a follow-up serology at the end of July, 2020. ELISA-S = 3.82; SN = 10. The participant experienced another episode of cough, fever, rhinorrhea with a SARS-CoV-2 PCR positive test in the second half of September 2020
8	40-49	M	2019-12	1.53	160	No	2-month travel in Asia between October and December, 2019.
9	30-39	F	2019-12	1.88	40	NA	NA
10	50-59	F	2019-12	1.83	80	Yes	Travel in Italy (Roma) end October - early November. Febrile illness at the end of October 2019
11	40-49	F	2020-01	1.71	40	Yes	Febrile illness during the third week of November 2019. Her husband and children were sick with febrile illness between November 10 and November 25, 2019
12	30-39	M	2020-01	2.83	40	No	Father was hospitalized for pneumonia in early December 2019
13	40-49	F	2020-01	1.23	40	No	General Practitioner in Paris

NA: not available

Figure 1. Number of blood samples tested each week for anti-SARS-CoV-2 antibodies by ELISA-S IgG test (horizontal bars) and percentages of ELISA-S positive test (red dots, with exact 95% Confidence Interval) in adult participants from the Constances cohort study, France.




Figure 2. Participants with a positive ELISA-S or a positive SN anti-SARS-CoV-2 test in the CONSTANCES cohort.

