



**HAL**  
open science

## Cognitive processes and a Centre-of-Pressure error-based moving light-touch Biofeedback

Fabien V erit e, Wael Bachta

► **To cite this version:**

Fabien V erit e, Wael Bachta. Cognitive processes and a Centre-of-Pressure error-based moving light-touch Biofeedback. *Neuroscience Letters*, 2021, pp.135743. 10.1016/j.neulet.2021.135743 . hal-03148959

**HAL Id: hal-03148959**

**<https://hal.sorbonne-universite.fr/hal-03148959>**

Submitted on 22 Feb 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin ee au d ep ot et  a la diffusion de documents scientifiques de niveau recherche, publi es ou non,  emanant des  tablissements d'enseignement et de recherche franais ou  trangers, des laboratoires publics ou priv es.

# Cognitive processes and a Centre-of-Pressure error-based moving light-touch Biofeedback

Fabien V erit e and Wael Bacht a

Sorbonne Universit e, ISIR, UMR 7222 CNRS, Agathe Group INSERM U 1150. Paris, France

---

## Abstract

Lightly touching an earth-fixed external surface with the forefinger provides somatosensory information that reduces the Center of Pressure (CoP) oscillations. If this surface were to move slowly, the Central Nervous System (CNS) would misinterpret its movement as body self-motion, and involuntary compensatory sway responses would appear, resulting in a significant coupling between finger and CoP motions. We designed a forefinger moving light-touch biofeedback based on this finding, which controls the surface velocity to drive the CoP towards a target position.

Here, we investigate this biofeedback resistance to cognitive processes. In addition to a baseline, the experimental protocol includes four main conditions. In the first, participants were utterly naive about the feedback. Then, they received additional reliable sensory information. The third condition ensured their full awareness of the external nature of the surface motion. Finally, the experimenter notified them that the external motion drives their balance and asked them to reject its influence.

Our investigation shows that despite the robustness of the proposed biofeedback, light-touch remains penetrable by cognitive processes. For participants to dramatically reduce the existing coupling between the finger and CoP motions, they should be aware of the external motion, how it impacts sway, and actively reject its influence.

The main implication of our findings is that light-touch exhibits the same cognitive flexibility as vision when artificially stimulated. This could be interpreted as a defense mechanism to re-weight these two sensory inputs in a moving environment.

*Keywords:* light touch, biofeedback, cognitive processes, voluntary control

---

## 1. Introduction

Independent artificial manipulation of sensory inputs evokes coupled postural responses. The strength of this coupling may depend on cognitive processes, including awareness, prediction of the forthcoming events, central multisensory integration, and voluntary control. When manipulating sensory inputs signalling only self-motion, the coupling remains strong regardless of the presence of cognitive processes. In contrast, cognitive processes can weaken or even preclude the coupling, during artificial manipulation of senses reporting both external and self motions.

Vestibular and kinesthetic sensory inputs signal only body self-motion. A typical way of artificially manipulating the vestibular sensory modality is to apply Galvanic Vestibular Stimulation (GVS) to the vestibular nerves [1, 2]. Once applied, the participant experiences a virtual rotation and leans in the opposite direction. GVS stimulation is immune to cognitive processes [3]. The coupling remains high regardless of the awareness of the artificial nature of the stimulus, pre-cueing of its occurrence or even its self-triggering. Applying vibrations to neuromuscular spindles at the calves muscles' level is a usual way to manipulate kinesthetic channels artificially. It induces a false sensation of falling forward [4], and an automatic backward postural response is then triggered. This artificial stimulus is also immune to cognitive processes, as reported in [5]. Prediction or self-triggering could only delay the evoked response.

Vision signals both external and self motions. People, standing in a room, whose walls are moving slowly, experience illusory self-motion. Postural reactions are then engaged in the same direction of the moving walls [6, 7]. Unlike vestibular inputs and muscle spindles, if anything alerts participants about their misinterpretation of the visual information, the evoked sway may be strongly inhibited [8].

Lightly touching a stationary surface with the forefinger is a significant sensory input to postural balance. It diminishes dramatically sway without providing any mechanical support [9]. However, if the surface moves periodically and slowly, body sway shows an automatic coupling to the stimulus trajectory [10, 11]. Like vision, a moving light touch leads, most of the time, to a perceptual ambiguity. The Central Nervous System (CNS) misinterprets the surface movement as self-motion [12]. We designed a forefinger moving light-touch biofeedback based on the finding of Jeka *et al.*, which controls the surface's velocity to drive the CoP towards a target position [13]. Our control sets the surface speed proportional to the error between the current and the target CoP positions. We tuned the control to keep the speed low with the objective of increasing the pre-mentioned sensory ambiguity.

Due to sensory re-weighting mechanisms, providing participants with a reliable additional sensory input may decrease the evoked postural responses. For example, if participants could benefit from light-touch with a stationary surface, postural

55 responses to visual stimuli [14, 15], tendon vibration [16] and 97  
 56 galvanic stimulation [17] would decrease significantly. Jeka 98  
 57 *et al.* reported that the coupling strength is also subject to 99  
 58 multisensory integration mechanisms [15] and that opening the 100  
 59 eyes can reduce coupling for moving light touch. 101

61 This paper questions our biofeedback resistance to the 103  
 62 following cognitive processes: the addition of reliable sensory 104  
 63 information, the explicit awareness of the motion’s external 105  
 64 nature, and the understanding of the potential coupling associ- 106  
 65 ated with the instruction to reject it actively. Our investigation 107  
 66 shows the robustness of the proposed feedback. For partici- 108  
 67 pants to dramatically reduce the existing coupling between the 109  
 68 finger and CoP motions, they should be aware of the external 110  
 69 motion, how it impacts sway, and actively reject its influence. 111

71 The main implication of our findings is that light-touch be- 113  
 72 haves to a large extent, like vision when taking into account 114  
 73 cognitive processes. Similarly to vision, participants could 115  
 74 drastically reduce the coupling between a moving-light touch 116  
 75 and the evoked postural responses. 117

## 76 2. Methods

### 77 2.1. Participants

78 The study, achieved at Sorbonne University, complied with 121  
 79 the Helsinki declaration relative to research involving human 122  
 80 beings and received the approval of the local ethical committee. 123

81 Forty-four healthy participants, divided into three groups, 125  
 82 were involved in the experiments. Participants did not present 126  
 83 any known neurological or postural history. Table 1 summa- 127  
 84 rizes the descriptive statistics of the three groups. 128

	GR1	GR2	GR3	Statistical
	(N=18)	(N=13)	(N=13)	Significance
Age (years old)	22 (10.5)	22 (3)	22 (2)	N.S.
BMI (kg.m <sup>-2</sup> )	22.5 (4.7)	21.1 (3.3)	21.9 (4.1)	N.S.
Gender (f/m)	7/11	5/8	5/8	N.S.

134 Table 1: Participants characteristics summary. GR1, GR2 and GR3 designate 134  
 135 three separate groups. *N* indicates the sample size of each group. BMI stands 135  
 136 for Body Mass Index. Quantitative data is presented as medians (interquartile 136  
 137 ranges). N.S. means Non Significant. 137

### 86 2.2. Experimental setup

87 Figure 1 shows a view of the experimental setup. It consists 141  
 88 of a force plate (AMTI BP400600-1000) and one Degree of 142  
 89 Freedom (DoF) translational device, which workspace is of 143  
 90 6 cm. 144  
 91 A typical trial consists of a participant standing on the top of 145  
 92 the force plate and lightly touching the translational device. 146  
 93 147

94 The force plate measures forces and torques applied by 148  
 95 standing participants, which allows the computation of the 149  
 96 CoP position. The translational device encloses a force sensor 150

which measures the applied finger’s six force components (see 1  
 top-left on figure 1). Participants hear an alarm sound each 2  
 time the applied vertical force exceeds 1N. Two Light-Emitting 3  
 Diodes (LED), are placed on the top of the translational device. 4  
 The LEDs are either off or on to indicate the direction of 5  
 movement of the translational device (see top-right on figure 6  
 1) . Participants put their finger on a double-sided tape to avoid 7  
 sliding on the translational device. 8

A DC motor drives the translational device motion, and 9  
 thus participants’ forefinger, with a linear motion resolution of 10  
 0.003 mm. Loudspeakers broadcast continuously pink noise in 11  
 the experimental room to prevent hearing the sound from motor 12  
 and associated mechanical parts. A white sheet, covering the 13  
 experimental setup, prevents participants from guessing that a 14  
 translational mechanism is in play. During the experiment, the 15  
 translational mechanism was placed in front of participants and 16  
 oriented to produce translation in the sagittal plane. 17

Custom software controls the motion of the translational 18  
 mechanism (more specifically its velocity) and collects the data 19  
 in real-time with a refresh rate of 500 Hz. 20

### 21 2.3. Moving light-touch Biofeedback design

22 In [13], we proposed moving-light touch biofeedback allow- 22  
 23 ing an automatic displacement of the CoP to a new target posi- 23  
 24 tion in the sagittal plane. 24

25 We controlled the lightly touched translational mechanism 25  
 26 velocity to be proportional to the difference between the target 26  
 27 and the current CoP positions (see bottom-left on figure 1). The 27  
 28 translational device drives *CoP* along a smooth path *CoP<sub>Ref</sub>*, 28  
 29 until reaching the final spot. 29

The control law writes:

$$V_{Finger}(t) = K(CoP_{Ref}(t) - CoP(t)) \quad (1)$$

30 Where  $V_{Finger}$  is the velocity of the finger (equal to the ve- 30  
 31 locity of the translational mechanism) at sample time  $t$ ,  $CoP_{Ref}$  31  
 32 is the reference trajectory,  $CoP(t)$  is a 0.3 Hz Butterworth 32  
 33 low-pass filtered current CoP position in the anteroposterior 33  
 34 direction. 34

35 In other words, biofeedback works as follows: if a participant 35  
 36 leaned forwards and overreached the desired value of the 36  
 37 reference trajectory (*i.e.*  $CoP > CoP_{Ref}$ ), the translational 37  
 38 mechanism would move backwards to bring back *CoP* toward 38  
 39  $CoP_{Ref}$ , and conversely. 39

40 The feedback gain  $K$  is equal to  $0.96 s^{-1}$ . We tuned it 40  
 41 empirically to maintain  $V_{Finger}$  low enough with an average 41  
 42 of about  $1 mm s^{-1}$  during our experimental session. This 42  
 43 tuning aimed at increasing the ambiguity between external and 43  
 44 self-motions. 44

45 The time-domain description of the reference trajectory 45  
 46 ( $CoP_{Ref}$ ) includes four-time intervals (in blue on Figure 1): 46




Figure 1: In the figure center, a participant is standing on the top of a force platform and lightly touching the translational device. From top to bottom and from left to right: 1/ A close view on the translational device composition, 2/ A block diagram of the closed loop, with a time domain description of  $CoP_{Ref}$  3/ The four experimental conditions, with a highlight on the illuminating LED, 4/ Two temporal representations of a closed-loop results. The first illustrates a strong coupling: the CoP in red follows the predefined path in blue, the velocity plot in purple is low. The second illustrates a weak coupling: the CoP is far from the predefined path, the moving plate reach its mechanical limits (saturation) and thus the velocity is equal to zero.

- 151 • [0 to 10]s: there is no control. The software computes the 176 average of  $CoP$ . 177
- 152 178
- 153 • [10 to 20]s:  $CoP_{Ref}$  is equal to the mean of  $CoP$  computed 179 during the previous time interval 180
- 154 181
- 155 • [20 to 30]s:  $CoP_{Ref}$  is a smooth trajectory moving 8mm 182 forward 183
- 156 184
- 157 • [30 to 60]s:  $CoP_{Ref}$  remains constant at its new value (8 185 mm away from the initial position) 186
- 158 187

#### 159 2.4. Data Collection and processing

160 For each experimental trial, we recorded  $CoP_{Ref}$  and raw 190  
 161  $CoP$  corresponding to the reference and the current  $CoP$  191  
 162 position (unfiltered) in the sagittal plane. 192

163 We introduced an evaluation criterion called  $\epsilon$  that quantifies 193  
 164 the closed-loop performance, and consequently the strength of 194  
 165 the coupling between the finger and  $CoP$  motions: 195  
 166 196

$$167 \epsilon = \frac{1}{N} \left| \sum_N (CoP_{Ref} - CoP) \right| \quad (2) \quad 197$$

168  $N$  designates the number of samples of the experiment when 200  
 169 the biofeedback was on (*i.e.* the [10 to 60]s time interval). 201  
 170 This tracking error qualifies the efficiency of the closed-loop 202  
 171 performances. The higher the tracking error is, the weaker is 203  
 172 the coupling. A high  $\epsilon$  indicates a failure in driving the  $CoP$  204  
 173 around the reference trajectory. An upper-bound of 8 was 205  
 174 assigned to the error. 206  
 175

#### 2.5. Experimental procedure

All participants were utterly naive about the goal of the experiment. For all the conditions, participants stood on the top of the force platform and touched the double-sided tape, located on the top of the translational mechanism, lightly with the index of their dominant hand. As soon as normal force exceeds 1N, an alarm sound is emitted and participants are asked to release the pressure. They held the other arm along the body. The experimenter adjusted the height of the translational device for each participant. Figure 1 illustrates the experimental protocol.

The experimenter controlled visually the participants' upper limb configuration, which they kept almost the same during the whole experiment. We also checked that the upper limb configuration was far from all joint limits.

We instructed participants to keep a neutral upright standing.

The experiment, consisting of providing participants with our moving light-touch biofeedback, included a baseline and four main conditions:

- **W/O feedback:** In each trial, we considered the [0 to 10]s time lapse where the moving-light touch feedback was off. We computed the average  $CoP$  position during the first 5 seconds, and we considered a hypothetical 8 mm forward reference for the remaining time to obtain a baseline score  $\epsilon$ . This condition is the baseline.
- **EC:** Participants kept their eyes shut. The LEDs were off.
- **EO :** Participants kept their eyes open and looked at a cross drawn on a wall located 50 cm in front of them. The LEDs were off, and the moving plate was outside their field of

view. This condition consists of *adding reliable sensory input*.

- **EOF**: Participants kept their eyes open and looked at the translational device. The experimenter told them that the translational device is moving. The LEDs were on and indicated the direction of motion of the plate (*i.e.* forward or backward). This condition consists of *adding the awareness about the external movement*.

- **AR**: Participants, aware of the external motion, are always looking at their finger, with the LEDs indicating the direction of movement of the plate. The experimenter informed them about the existing coupling between their finger motion and their postural sway. The instruction changed: in this condition, they should try to reject the coupling. This condition corresponds to *a change from a neutral standing to voluntary rejection of the coupling*.

Participants achieved each condition three times. We thus computed three tracking errors, and the average is denoted  $\epsilon$ .

Participants of **GR1** took part in the five conditions. The W/O feedback condition was always the first presented one. Then, the two second conditions (**EC** and **EO**) were presented randomly. The two remaining conditions took place in the same order: **EOF** and then **AR**. No further randomisation was possible since participants were gaining awareness progressively.

In order to check that participants of **GR1** did not benefit from learning or habituation, two other Groups were involved. In addition to the W/O feedback, participants of **GR2** and **GR3** were involved respectively in the **EOF** and **AR** conditions.

## 2.6. Statistical Analyses

Taking into account the relatively small sample sizes of the groups included in the study, we present the descriptive statistics describing the data as medians (Inter-Quantile -Range), *i.e.*  $Mdn$  ( $IQR$ ), and we use non-parametric methods for analyses.

We investigated the null hypothesis validity for gender ratio, BMI, and age between Groups using a  $\chi^2$  and two Kruskal-Wallis tests.

A Kruskal-Wallis test allowed checking the rejection of the null hypothesis for the **W/O feedback** condition between the three groups.

The investigation of the null hypothesis between the tracking error during the different conditions (**W/O feedback**, **EC**, **EO**, **EOF**, and **AR**) for **GR1** relied on a Friedmann test analysis. If the test rejected the null hypothesis, Post-hoc paired Wilcoxon tests, with a Bonferroni correction, is used.

Two Wilcoxon signed-rank tests allowed checking the existence of a significant difference between the **W/O feedback**,

**EOF** and **AR** respectively for groups **GR2** and **GR3**.

A Mann-Whitney U test allowed the comparison of the tracking error between Groups (**GR1/GR2**), and (**GR1/GR3**) for the **EOF** and **AR** conditions, respectively. A final Mann-Whitney U test allowed the comparison of the tracking error between **GR2** in the **EOF** condition and **GR3** in the **AR** condition.

The statistical level of significance has been set at  $p = 0.05$ .

## 3. Results

### 3.1. Participants

Table 1 summarises the three groups characteristics. The groups did not differ by gender,  $\chi^2(2, N = 44) = 0.001, p = 1$ . Two Kruskal Wallis tests rendered no significant difference between groups for age ( $H(2)=0.1, p = 0.95$ ) and BMI ( $H(2)=3.89, p = 0.143$ ).

### 3.2. The tracking error $\epsilon$

Figure 2 shows a Tukey outlier boxplot of the tracking error score  $\epsilon$ .

**Between groups comparisons.** A Kruskal-Wallis indicated no significant difference ( $H(2)=2.21, p=0.33$ ) for the **W/O feedback** conditions between the three groups **GR1** ( $Mdn=7.99$ ), **GR2** ( $Mdn=7.81$ ), and **GR3** ( $Mdn=7.14$ ).

A first Mann-Whitney U test indicated no significant difference in the **EOF** condition between **GR1** ( $Mdn=2.14$ ) and **GR2** ( $Mdn=0.85$ ),  $U=90, p=0.293$ .

The second Mann-Whitney U test indicated no significant difference in the **AR** condition between **GR1** ( $Mdn=4.3$ ) and **GR3** ( $Mdn=3.3$ ),  $U=87.5, p=0.242$ .

The final Mann-Whitney U test indicated a significant difference between **GR2** ( $Mdn=0.85$ ) and **GR3** ( $Mdn=3.3$ ) involved in the **EOF** and **AR** conditions respectively. The test statistic  $U$  was equal to 134.5, with  $p=0.009$ .

**Comparisons within GR2.** A Wilcoxon signed-rank test showed a significant difference between the **W/O feedback** and **EOF** conditions,  $T=91, z=-3.81$  and  $p<0.001$ .

**Comparisons within GR3.** A Wilcoxon signed-rank test showed a significant difference between the **W/O feedback** and **AR** conditions,  $T=78, z=3.3$  and  $p<0.01$ .

**Comparisons within GR1.** A Friedman’s test rendered a significant difference between the five conditions for **GR1**,  $\chi^2_F(4) = 57.9$ ,  $p < 0.001$ . Post-hoc analysis using Wilcoxon rank-sum test with a Bonferroni adjustment showed significant pairwise comparisons between the **AR** and **W/O feedback**,  $p < 0.01$ . **W/O feedback** is significantly different from the other conditions,  $p < 0.001$ . **AR** is significantly different from the other conditions,  $p < 0.01$ .




Figure 2: A Tukey outlier boxplot of the tracking error. Note that for sake of clarity, the W/O feedback of all the groups are merged. The full statistical analyses are provided in the Results section

Figure 2 summarizes the main results. Our results show that the **AR** and **W/O feedback** are significantly different, which means that the coupling is not completely rejected. The significant difference between the **AR** and the **EC**, **EO** and **EOF** conditions suggest a drastic decrease of the coupling when participants are actively rejecting it.

#### 4. Discussion

The main finding is the robustness of our proposed biofeedback to cognitive processes. Nevertheless, light-touch is still penetrable by cognitive processes. To dramatically reduce the existing coupling between the finger and CoP motions, participants should be aware of the external motion, how it impacts sway, and actively reject its influence. Results from groups **GR2** and **GR3** suggest the absence of a significant learning effect during the experimental session. We will discuss the results obtained from **GR1**.

Light-touch compares well to vision. Unlike vestibular and kinesthetic inputs, cognitive processes could reduce the coupling of evoked postural responses to their artificial manipulation. The similarity between vision and light-touch is due to their capacity to signal self and environment movements ([5, 8, 3]). These two sensory information are subject to ambiguous information, especially in a moving environment. To this regard, the re-weighting mechanism (either sensory or cognitive), could be seen as a defense mechanism allowing to

maintain an upright posture in a moving environment.

The environmental motion could either present high dynamics (high amplitude and high velocity) or low dynamics (low amplitude and low velocity). In the former, vision and light-touch can easily separate the external and self-motion. In the latter, the ambiguity increases, and the separation becomes difficult. Barela *et al.* showed, in their work [18], a group of participants not aware of the used moving-room paradigm were able to reduce the coupling after being exposed to a faster and larger moving-room motion. The increase in dynamics allows an implicit understanding of the stimulus and its influence on posture, which in turn allows a greater attenuation of the coupling than in the case of an explicit indication from the experimenter. This compares favourably to light-touch, where postural responses also depend on the stimuli dynamics. In [19], the authors displayed ten consecutive high-velocity and high-amplitude linear sagittal stimuli. In the first trial, more than half of the participants perceived by themselves the platform motion and their involuntary postural responses, which then vanished in subsequent trials. According to the authors, participants understood their overreaction and chose to actively ignore the stimulus. The cited study is in line with our results. The weaker coupling comes from the learnt awareness of the external motion and its influence on postural balance. The coupling rejection was not due to an explicit instruction from the experimenter but is instead an effect of understanding that the first evoked postural response could have threatened balance stability.

When the stimulus is periodic and presents a low velocity and low amplitude, it becomes less easy to be detected and the coupling less easy to reject. The low dynamics of the stimulus increases its ambiguity. The authors of [20] reported that participants felt that their sway was increasing, without successfully attributing it to the touched device motion. Only one participant attributed the sway increase to the external motion and thus exhibited weaker coupling. This finding compares favourably with the results of [7, 18], where the authors informed participants about visual manipulation, and this information allowed them to decrease the coupling, even without being asked to do so. One can conclude that the only awareness of an external motion may change participants’ postural control strategy and lead them to reject the coupling, but the change of strategy differs across individuals. In the study of Jeka *et al.* [10], all participants noticed by themselves that the motion of the moving touched-plate was ambiguous and failed to characterize it. Their postural sway remained strongly coupled to moving plate. One could hypothesize that the awareness of the external motion without understanding its impact on balance could be insufficient to reduce the coupling. Our proposed biofeedback highlights this hypothesis; we designed it to increase the ambiguity, decreasing the probability of guessing its effect on balance. This allows us to induce a relatively large CoP displacement without the participants’ knowledge.

In their works [21, 22], the authors studied the visual senso-

rimotor coupling under a moving-room paradigm with participants asked to "resist the room's movement". They reported two results. The first is that the active resistance condition reduces the coupling, in line with the results of this paper. Second, the reducing rate decreased when resisting the visual manipulation and performing at the same time a concurrent cognitive task, since the attentional resources should then be shared. Preliminary trials reported in [13], indicates that the proposed biofeedback performances were not significantly influenced by a concurrent cognitive task. This is a little bit contradictory with the results of [23, 24], where a concurrent cognitive task altered the assistance provided by lightly touching a stable surface. One should notice, that none of these studies required intentional resources dedicated to the coupling between posture and the stimuli. A moving-light touch paradigm with the explicit instruction to reject the coupling, associated with an additional cognitive task, needs to be addressed carefully.

Finally, our biofeedback contrasts with previous studies on moving-light touch. Unlike the results reported in [15], where the addition of a stationary visual input reduced the sensorimotor coupling significantly, our study revealed no significant difference between the EC and EO conditions. The median slightly increased when participants looked at earth grounded visual information, but to a lesser extent than expected. The design and the tuning of the biofeedback explain the difference: it is based on the current CoP position and tuned to increase the ambiguity. Any attempt of reducing the coupling, e.g. due to a piece of reliable sensory information, will result in deviation of the CoP position. This deviation would constitute an error, and the biofeedback will gently compensate for by bringing the CoP to its target position.

In conclusion, the main implication of our findings is that light-touch behaves to a large extent, like vision when taking into account cognitive processes. Similarly to vision, participants can voluntarily reduce the coupling between a moving light touch and the evoked postural responses. A plausible interpretation is that, as a defense mechanism, the CNS is able to re-weight these two sensory inputs to preserve balance in moving environments. Future research needs to focus on the attentional resources sharing when participants are asked to resist the coupling while achieving a concurrent dual-task.

## 5. Acknowledgement

This work was funded by the ANR within the i-Gait project under reference ANR-16-CE33-0012.

## References

[1] B. L. Day, a. Séverac Cauquil, L. Bartolomei, M. A. Pastor, I. N. Lyon, Human body-segment tilts induced by galvanic stimulation: a vestibularly driven balance protection mechanism., *The Journal of physiology* 500 ( Pt 3 (3) (1997) 661–672.

[2] B. L. Day, R. C. Fitzpatrick, The vestibular system., *Current biology : CB* 15 (15) (2005) R583–6.

[3] M. Guerraz, B. L. Day, Expectation and the vestibular control of balance., *Journal of cognitive neuroscience* 17 (3) (2005) 463–469.

[4] A. Kavounoudias, R. Roll, J. P. Roll, Foot sole and ankle muscle inputs contribute jointly to human erect posture regulation, *Journal of Physiology* 532 (3) (2001) 869–878.

[5] S. Caudron, F. Boy, N. Forestier, M. Guerraz, Influence of expectation on postural disturbance evoked by proprioceptive stimulation, *Experimental Brain Research* 184 (1) (2008) 53–59.

[6] D. Lee, J. Lishman, Visual proprioceptive control of stance., *Journal of Human Movement Studies* 1 (January) (1975) 87–95.

[7] P. B. Freitas Júnior, J. A. Barela, Postural control as a function of self- and object-motion perception, *Neuroscience Letters* 369 (1) (2004) 64–68.

[8] M. Guerraz, K. V. Thilo, A. M. Bronstein, M. A. Gresty, Influence of action and expectation on visual control of posture, *Cognitive Brain Research* 11 (2) (2001) 259–266.

[9] J. J. Jeka, J. R. Lackner, Fingertip contact influences human postural control., *Experimental brain research* 100 (3) (1994) 495–502.

[10] J. Jeka, K. Oie, G. Schöner, T. Dijkstra, E. Henson, Position and velocity coupling of postural sway to somatosensory drive., *Journal of Neurophysiology* 79 (4) (1998) 1661–1674.

[11] A. M. Wing, L. Johannsen, S. Endo, Light touch for balance: influence of a time-varying external driving signal., *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* 366 (1581) (2011) 3133–41.

[12] F. Vêrité, R. Reynolds, W. Bachta, Perception of haptic motion is enhanced during conditions of increased postural stability, *Gait & Posture* 76 (2020) 334 – 338.

[13] F. Vêrité, W. Bachta, G. Morel, Closed Loop Kinesthetic Feedback for Postural Control Rehabilitation, *IEEE Transactions on Haptics* 7 (2) (2014) 150–160.

[14] C. J. Hausbeck, M. J. Strong, L. S. Tamkei, W. a. Leonard, K. I. Ustinova, The effect of additional hand contact on postural stability perturbed by a moving environment., *Gait & posture* 29 (3) (2009) 509–13.

[15] J. Jeka, K. S. Oie, T. Kiemel, Multisensory information for human postural control: integrating touch and vision, *Experimental Brain Research* 134 (1) (2000) 107–125.

[16] J. R. Lackner, E. Rabin, P. DiZio, Fingertip Contact Suppresses the Destabilizing Influence of Leg Muscle Vibration, *Journal of Neurophysiology* 84 (5) (2000) 2217–2224.

[17] E. Maaswinkel, H. E. Veeger, J. H. Dieën, Interactions of touch feedback with muscle vibration and galvanic vestibular stimulation in the control of trunk posture, *Gait and Posture* 39 (2) (2014) 745–749.

[18] J. Barela, M. Weigelt, P. Polastri, D. Gofdoi, A. SA, J. Jeka, Explicit and implicit knowledge of environment states induce adaptation in postural control., *Neurosci Lett.* 566 (2014) 6–10.

[19] J. Misiaszek, J. Forero, E. Hiob, T. Urbanczyk, Automatic postural responses following rapid displacement of a light touch contact during standing, *Neuroscience* 316 (2016) 1–12.

[20] J. J. Jeka, G. Schöner, T. Dijkstra, P. Ribeiro, J. R. Lackner, Coupling of fingertip somatosensory information to head and body sway., *Experimental brain research* 113 (3) (1997) 475–83.

[21] S. A. Aguiar, K. Gramani-Say, A. G. Lopes, J. A. Barela, Dual task interferes with sensorimotor coupling in postural control, *Psychology and Neuroscience* 7 (4) (2014) 593–599.

[22] G. G. Genoves, A. M. F. Barela, C. Sanches, J. A. Barela, Attentional artifacts in sensorimotor coupling in the postural control of young adults, *Experimental Brain Research* 234 (12) (2016) 3641–3647.

[23] Y. Lee, N. Goyal, A. S. Aruin, Effect of a cognitive task and light finger touch on standing balance in healthy adults, *Experimental Brain Research* 236 (2) (2018) 399–407.

[24] D. G. dos Santos, J. M. Prado-Rico, S. R. Alouche, R. B. d. S. C. Garbus, P. B. de Freitas, S. M. S. F. de Freitas, Combined effects of the light touch and cognitive task affect the components of postural sway, *Neuroscience Letters* 703 (March) (2019) 99–103.