How fast do mobile organisms respond to stimuli? Response times from bacteria to elephants and whales

Jean-Pierre Rospars, Nicole Meyer-Vernet

► To cite this version:

Jean-Pierre Rospars, Nicole Meyer-Vernet. How fast do mobile organisms respond to stimuli? Response times from bacteria to elephants and whales. *Physical Biology*, 2021, 18 (2), 10.1088/1478-3975/abcd88 . hal-03171698

HAL Id: hal-03171698

<https://hal.sorbonne-universite.fr/hal-03171698>

Submitted on 17 Mar 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 1 HOW FAST DO MOBILE ORGANISMS RESPOND TO STIMULI?
2
3
4
5
6 2 RESPONSE TIMES FROM BACTERIA TO ELEPHANTS AND WHALES
7
8
9 3

10
11 4 Jean-Pierre Rospars^a and Nicole Meyer-Vernet^b
12
13 5

14
15 6 ^a Institute of Ecology and Environmental Sciences of Paris, INRA, route de Saint-Cyr, 78000
16
17
18 7 Versailles, France
19

20 8 ^b LESIA, Observatoire de Paris, Université PSL, CNRS, Sorbonne Université, Université de
21
22 9 Paris, 5 place Jules Janssen, 92195 Meudon, France
23
24

25 10
26
27 11 Corresponding author: Jean-Pierre Rospars
28

29 12 Emails: Jean-Pierre.Rospars@inrae.fr and Nicole.Meyer@obspm.fr
30
31

32 13
33
34 14 Manuscript length: 9800 words (title page, summary and statements not included)
35

36 15 Number of figures: 4.
37

38 16 Number of tables: 2.
39

40 17 Supplementary material: yes.
41
42

43 18
44
45 19 Keywords: response time, startle response, locomotion, biophysical modelling, biological scaling.
46
47
48 20
49
50 21

51
52
53
54 Published in Phys. Biol. 18 (2021) 026002

<https://doi.org/10.1088/1478-3975/abcd88>

SUMMARY

Quick responses to fast changes in the environment are crucial in animal behaviour and survival, for example to seize prey, escape predators, or negotiate obstacles. Here, we study the ‘simple response time’ that is the time elapsed between receptor stimulation and motor activation as typically shown in escape responses, for mobile organisms of various taxa ranging from bacteria to large vertebrates. We show that 95 % of these simple response times lie within one order of magnitude of the overall geometric mean of about 25 ms, which is similar to that of a well-studied sensory time scale, the inverse of the critical flicker fusion frequency in vision, also lying within close bounds for all the organisms studied. We find that this time scale is a few times smaller than the minimum time to move by one body length, which is known to lie also within a relatively narrow range for all moving organisms. The remarkably small 10^2 -fold range of the simple response time among so disparate life forms varying over 10^{20} -fold in body mass suggests that it is determined by basic physicochemical constraints, independently on the structure and scale of the organism. We thus propose first-principle estimates of the simple response and sensory time scales in terms of physical constants and a few basic biological properties common to mobile organisms and constraining their responses.

1
2
3 40 1. INTRODUCTION
4

5 41
6
7 42 The concept of timescale is fundamental in science. An important timescale in biology is the
8
9
10 43 minimum response time of mobile organisms to a dynamic environment. When an animal
11
12 44 suddenly encounters a prominent event such as a prey, a predator or an obstacle, it must react
13
14 45 fast enough, albeit not faster than necessary, in order not to sacrifice accuracy or waste the
15
16 46 energy or space dedicated to its sensory systems (e.g. Attwell and Laughlin 2001). For a
17
18 47 response to be fast enough, the information that produced the response should not be outdated
19
20 48 (e.g. Spence 2009), which requires in particular that the response time does not exceed the
21
22 49 movement duration and that the animal position has not changed much. For animals moving with
23
24 50 legs, the response time is often compared to the stance or step duration (e.g. More et al. 2018);
25
26 51 however, for animals not equipped with legs, a more general time scale for comparison with the
27
28 52 minimum response time is the minimum time to move by one body length, that is the ratio of
29
30 53 body length to maximum speed.
31
32
33
34

35 54 The maximum speed of terrestrial and aquatic organisms has been found to be roughly
36
37 55 proportional to their body length, from bacteria to large vertebrates (Bonner 1965, McMahon and
38
39 56 Bonner 1983), contrary to the preferred speed which is subjected to different constraints (e.g.
40
41 57 Bejan and Marden 2006). These results, later confirmed with a large data set, imply that the time
42
43 58 to move by one body length at maximum speed lies in a narrow range around one tenth of
44
45 59 second within a factor of ten, for running and swimming organisms of mass varying by 10^{20} -fold
46
47 60 in body mass. The ubiquity of this minimum locomotion time scale, holding for so different
48
49 61 organisms' structures and sizes, whereas characteristic biological timescales cover more than 12
50
51 62 orders of magnitude (e. g. Shamir et al. 2016), suggests that it is bounded by universal
52
53
54 63 constraints, and led Meyer-Vernet and Rospars (2015, 2016) to propose a tentative interpretation
55
56
57
58

based on the mass density of living organisms, the maximum specific tension exerted by molecular motors and muscles (Rospars and Meyer-Vernet 2016) and the maximum mass specific metabolic rate (Weibel and Hoppeler 2005, Glazier 2014, Makarieva et al. 2005), which constrain the maximum speed and remain within close bounds for all moving organisms.

The question therefore arises of whether a similar result could hold for the minimum time to react to stimuli. The scaling of sensorimotor delays with body mass in relation to movement duration has been studied by More and Donelan (2018) for the stretch reflex in terrestrial mammals, but there is no large-scale study of the minimum response time covering the entire mass range of mobile species. We therefore collated data from the literature for the ‘simple response time’, that is the time to detect the occurrence of a simple stimulus, as determined from behavioural and electrophysiological measurements in various taxa from free-living cells to large metazoans like sharks, turtles, and elephants, spanning 20 orders of magnitude in mass from 10^{-16} to 3900 kg.

An important constraint to fast response is set by sensory limitations that affect the ability to track fast moving objects such as prey or mates. These limitations have been studied in the case of vision, using the critical flicker fusion frequency (CFF), defined as the frequency at which a flickering light is indistinguishable from a continuous light (e.g. D’Eath 1998). The scaling of this property with body mass and metabolism has been studied by Healy et al. (2013) for vertebrates; we collated the corresponding characteristic time ($1/\text{CFF}$) and extended the data to compare it to the minimum response time.

Our aims are (i) to determine whether the minimum response time remains within close bounds across the whole of mobile life despite the diversity of structures and mechanisms, (ii) to compare it with the minimum sensory time-scale for vision (inverse of the critical flicker fusion frequency) and to the minimum time to move by one body length, and (iii) to propose an

1
2
3 88 interpretation based on basic physicochemical and biological properties. Because the response
4
5 89 times of animals are expected to be hugely different from those of microorganisms, which are
6
7
8 90 subjected to very different constraints (e.g. Martens et al. 2015), we also investigate the
9
10 91 timescales separately for single cells and for metazoans.

11
12 92 The empirical data are studied and discussed in sections 2 to 4. Section 5 proposes order-of-
13
14 93 magnitude interpretations based on fundamental physicochemical constants and basic properties
15
16
17 94 of life, with conclusions given in Section 6.

18
19 95

20
21 96 2. MATERIAL AND METHODS

22
23
24 97

25
26 98 2.1. *Data Collection*

27
28 99

29
30
31 100 We define as “simple response time” (T_s) the time to detect the occurrence of a simple and
32
33 101 sudden stimulus. These T_s are delays from the time at which the stimulus reaches the organism to
34
35 102 either the onset of movement – measured by behavioural methods (for example with high-speed
36
37 103 cinematography), or to muscle activation – measured by electrophysiological methods with direct
38
39 104 recordings from muscles (electromyogram EMG). Since our study concerns minimum response
40
41
42 105 times, it does not include the “discrimination (or identification) and choice response times” (e.g.
43
44 106 Luce 1991). In these complex tasks, the subject, whether animal or human, is presented with one
45
46
47 107 of several stimuli and has to respond to only one of them (discrimination, e.g. Blough 1978) or to
48
49 108 perform different responses depending on the stimulus presented (choice between stimuli, e.g.
50
51 109 Abraham et al. 2004). Likewise, for microorganisms, we only considered the response delay after
52
53
54 110 a stimulus (e.g. Block et al. 1982), which only involves internal time scales of the organisms and
55
56 111 is shorter than the time based on the comparison of different measurements, which depends on

1
2
3 112 ambient conditions (e.g. Mitchell 1991). Thus, our definition of simple response times T_s being
4
5 113 based on the simplicity of the task and the stimulus does not take into account the length,
6
7 114 variability or underlying sensorimotor mechanisms of the response delays (Roeder 1963, Koch
8
9 115 1999, Herberholz and Marquart 2012, Sillar et al. 2016, Roberts et al. 2019). It avoids the
10
11 116 operational difficulty to implement these multiple criteria and insures a better representativeness
12
13
14 117 of the sample.

15
16
17 118 We searched the Google Scholar database and extracted simple response times and other
18
19 119 relevant data (species names, stimuli, experimental conditions, etc.) published in refereed
20
21 120 journals. Our main objective being to investigate the interspecific variability, we kept all values
22
23 121 found for non-human species, but not for humans, for which we considered only a selection of
24
25 122 papers, either classical or illustrating the diversity of experimental paradigms. For a given species
26
27 123 and stimulus, we tabulated simple response times as the mean provided by the authors, or the
28
29 124 mean of the extremes when given as a range.

30
31
32
33 125 Our T_s sample (Table S1) includes 175 measurements on 81 species. Behavioural ($n = 134$)
34
35 126 and electromyographic (EMG, $n = 41$) measurements (column M in Table S1) are not
36
37 127 significantly different (Fig. S2A and Table S10); thus, we pooled them together in subsequent
38
39 128 analyses. The data belong to two main classes: startle and non-startle. “A startle response is an
40
41 129 abrupt response, often of relatively short latency, to a sudden stimulus that we believe to be both
42
43 130 unexpected and alarming (i.e., of high valence)” (Bullock 1984), so that threats that develop
44
45 131 gradually are excluded. Startle responses form a homogeneous class gathering the majority of
46
47 132 measurements (73 %) and of species (68 %, multicellular only). They may result in a large
48
49 133 movement translating the whole body (escape, called fast-start in fishes, 86 measurements on 50
50
51 134 species) or in small movements (called sometimes ‘eyelid, jaw, etc. reflex’ in birds and
52
53
54
55 135 mammals, depending on the muscle triggered, 42 measurements in five species). Non-startle

136 responses (27 % of measurements on 32 % of species) include taxes (in single cells only), reflex
137 control of locomotion, fast limb movements, predatory movements, and a few other fast
138 responses in multicellular organisms. Only six species were tested for more than one response
139 type.

140 For flicker fusion times T_F , we used the CFF data from Healy et al. (2013) (34 vertebrate
141 species) and Inger et al. (2014) (31 invertebrate species and 41 vertebrate species not considered
142 by Healy et al. 2013), plus 25 other CFF measurements. Our sample ($T_F = 1/\text{CFF}$, Table S2)
143 includes 130 measurements on 108 species. The values measured by behavioural ($n = 26$) and
144 electroretinographic (ERG, $n = 103$, column M in Table S2) techniques are not significantly
145 different (Fig. S2C and Table S10); so, we pooled them together.

146 The minimum times to move by one body length T_L were collected and analysed previously
147 from the measured maximum speeds $V_{\max}$ of swimming and running organisms of length L as T_L
148 $= L/V_{\max}$; this sample includes 458 measurements from 427 species (Table S3) and does not
149 include flying, whose maximum speed is not constrained by muscles (Meyer-Vernet and Rospars
150 2016).

151 For T_S and T_L , we distinguished unicellular and multicellular organisms and spermatozooids (T_L
152 only), as specified in column U of Tables S1 and S3. For the three timescales, column Cla of
153 Tables S1-S3 defines groups of phylogenetically related species; they belong to the same class in
154 multicellular organisms and to the same kingdom according to the WoRMS database (World
155 Register of Marine Species, <http://www.marinespecies.org>) in unicellular organisms (except
156 spermatozooids). Further details on Tables S1-S3 are given in Supplementary material.

157 Table S4 lists 14 mammalian species for which both minimum locomotor times T_L and
158 maximum mass specific metabolic rates (MSMR in W/kg) are known.

160 2.2. *Body Mass*

161
 162 We characterized body size by the mass M for each species. Except for vertebrates with measured
 163 CFFs from Healy et al. (2013) for which we used the mass provided by these authors, we
 164 searched the original papers for mass, length or age. When given as a range, we took the mean.
 165 We converted length in mass using either the length-mass relationship of the species when it is
 166 known or a more generic relationship, for example $M_{\text{kg}} = 11.2 L_{\text{m}}^{3.04}$ which applies to fusiform
 167 fishes (Froese et al. 2014; see other relationships in Meyer-Vernet and Rospars 2016). When no
 168 indication was provided in the papers, we searched the average mass (or length) of the species in
 169 the scientific literature (for example, Bartholomew and Heinrich 1973; Byrne et al. 1988; Niven
 170 and Scharlemann 2005, for insects; Falk-Petersen 1981, for shrimps) or in websites. We have not
 171 considered in the analyses two T_{F} data for which the mass could not be found (lines 24 and 87 in
 172 Table S2). Otherwise, all body masses and their references are given in Tables S1-S3.

174 2.3. *Temperature Effects*

175
 176 In order to check the dependence of the time scales on temperature, we used (in the Discussion
 177 section only) the Boltzmann-Arrhenius model from chemical reaction kinetics, which holds
 178 approximately for metabolic and locomotor rates (e.g. Dell et al. 2011). This model yields
 179 biological time scales inversely proportional to the Boltzmann factor, $\exp(-E/k_{\text{B}}\theta)$, where E is the
 180 activation energy of the process studied (in joules), k_{B} , the Boltzmann constant, and θ , the
 181 temperature (in kelvins). Thus, the corrected time scale T_0 , at the reference temperature θ_0 , of
 182 time T measured at temperature θ is $T_0 = T \exp(qE/k_{\text{B}})$, where $q = 1/\theta_0 - 1/\theta$. For all timescales,
 183 we chose $\theta_0 = 20^\circ\text{C}$ as reference and $E = 0.66$ eV, the mean activation energy observed in a wide

range of species and traits (Dell et al. 2001). In the special case of T_S , we also applied a finer standardization procedure distinguishing defence or movement away from a stimulus, like startle responses, and consumption or movement toward a stimulus (almost all other responses), whose mean activation energies are $E = 0.4$ and 0.7 eV, respectively (Dell et al. 2001).

2.4. Statistical Analyses

Statistics were computed on either data T or their log-transform ($\log_{10} T$). In the main text, but not in Supplementary material, all data (T and $\log_{10} T$) were averaged per species, counting separately sperm cells (in T_L data, Table S3) and late developmental stages (for *Danio rerio* and *Procambarus clarkii* in T_S data, Table S1). The data in each category (T_S , T_F and T_L) were characterized by their medians and interquartile ranges IQR. Lognormal distributions were fitted after determination of their parameters (mean μ and standard deviation σ) on log-transformed data. However, for easier readability, μ in log units was converted in seconds in text and figures (except in Supplementary material Tables S7-S10), as $\mu^* = 10^\mu$ (μ^* is the geometric mean and median of the lognormal distribution fitted to data T). Similarly, σ was expressed as a multiplicative standard deviation $\sigma^* = 10^\sigma$ (σ^* , like σ , is dimensionless; it determines the asymmetry of the distribution, and the interval $[\mu^*/\sigma^*, \mu^*\sigma^*]$ covers a probability of 68.3 %, see Limpert et al. 2001). The overall interval of variation in each timescale was expressed as the percentiles 2.5 % and 97.5 %, which are less sensitive to outliers and sampling fluctuations than the minimum and maximum; the ratio of these percentiles and its logarithm (denoted δT_{95}) estimate the multiplicative range including 95 % of values. Statistical distributions were compared with the Kolmogorov-Smirnov test. Least-square regressions of $\log_{10} T_i$ against $\log_{10} M$ and least-rectangle regressions of $\log_{10} T_L$ against $\log_{10} MSMR$ (Dagnelie 2011) were

calculated and given as scaling equations $T_i = T_0 M^\alpha$ in figures and their slope (also called scaling exponent) α as 95 % confidence intervals in the text. Tables S7-S10 provide details of ANOVA and multiple comparisons of means using Tukey-Kramer adjustment method. We used the significance level 5 % in all tests. We performed all statistics with the Matlab Statistical Toolbox (The Mathworks, Natick, USA).

3. EMPIRICAL RESULTS

3.1. Statistical Distributions of Timescales and Comparisons

Simple response times, T_s , extend from 2.5 ms in the escape behaviour of the calanoid copepod *Undinula vulgaris* to 485 ms in the acoustic response of the white whale *Delphinapterus leucas*, with median 24 ms and IQR 43 ms. The distribution is lognormal (Fig. S1A), with geometric mean $\mu^* = 26$ ms and multiplicative standard deviation $\sigma^* = 3.26$. Critical fusion times T_F range from 2.5 ms in the black fire beetle *Melanophila acuminata* to 250 ms in the crustacean isopod *Booralana tricarinata*, with median 24 ms and IQR 27 ms. The distribution is lognormal (Fig. S1C) with $\mu^* = 25$ ms and $\sigma^* = 2.14$. The times to move by one body length at maximum speed T_L extend from 5 ms for a sea urchin to 2.8 s for a large spirochetes bacterium, with median 71 ms and IQR 99 ms (Meyer-Vernet and Rospars 2016). The distribution is lognormal with $\mu^* = 78$ ms and $\sigma^* = 2.73$ (Fig. S1B). Since most of the variability in the data comes from the diversity of species, stimuli and measurement methods, such lognormal distributions are expected if these factors act in a multiplicative way.

Figure 1 compares as boxplots the (log-transformed) timescales T_F , T_s and T_L . Their relative position is indicated by the medians (central red lines). ANOVA and multiple comparisons of

1
2
3 232 means show that the sensory timescale T_F and the simple response time T_S are similar, whereas T_L
4
5 233 is significantly different and about three times longer (Table S7).
6
7
8 234

9
10 235 3.2. *Variation with Body Mass*
11
12 236

13
14 237 To study the dependence of simple response times on body mass M for the whole data set
15
16 238 encompassing life's major domains, we plotted the pairs (M, T_S) in log-log plots for 81 species
17
18 239 (Fig. 2), including microorganisms in the mass range 10^{-16} (bacterium) to 5×10^{-13} kg (green alga)
19
20 240 and multicellular organisms from 3×10^{-9} (spider) to 3860 kg (elephant). Fig. 2A distinguishes the
21
22 241 taxonomic groups whereas Fig. 2B distinguishes the types of responses. Overall, the mass M
23
24 242 varies by a factor of about 10^{20} whereas 95 % of the T_S values lie between 4.7×10^{-3} and 0.31 s
25
26 243 (Table 1, first line). Although very small, the slope of the regression line is significantly different
27
28 244 from zero (95 % confidence intervals $[-0.065, -0.010]$, Table 2), but as shown in Table 1, this
29
30 245 effect results from the larger simple response times in single cells ($\mu^* = 129$ ms) than in
31
32 246 multicellular organisms ($\mu^* = 22$ ms). Since this difference is significant (Table S8c, first line,
33
34 247 $p < 10^{-3}$), the two groups must be studied separately. The scaling exponents α of the power law
35
36 248 regressions in both groups are not significantly different from zero with 95 % confidence
37
38 249 intervals $[-0.41, 0.49]$ for single cells and $[-0.01, 0.08]$ for multicellular organisms, whereas the
39
40 250 intercepts differ by more than one order of magnitude (Table 2 and Fig 2A).
41
42
43
44
45

46
47 251 Flicker fusion times T_F could only be measured in multicellular organisms. Therefore, the
48
49 252 body mass of the 106 species shown in Figure 3 only varies from 2×10^{-6} kg (fruit fly) to 354 kg
50
51 253 (sea turtle), representing over eight orders of magnitude, whereas 95 % of the T_F values lie
52
53 254 between 2.5×10^{-3} s and 0.25 s (Table 1). No effect of body mass on T_F could be evidenced, since
54
55 255 the power law regression is $T_F = 0.026 M^\alpha$ (Fig 3), with α in the 95 % CI $[-0.01, 0.06]$ (Table 2).
56
57
58

The times to move by one body length at maximum speed T_L (Meyer-Vernet and Rospars 2016, Fig. 4) concern 426 species including microorganisms from 10^{-16} kg (bacterium) to 1.3×10^{-8} kg (ciliate eukaryotic cell) and multicellular organisms from 10^{-9} (copepod) to 1.4×10^5 kg (blue whale). So, M varies by 21 orders of magnitude whereas 95 % of the T_L values lie between 21×10^{-3} and 0.71 s. The times T_L in single cells ($\mu^* = 145$ ms) are twice longer and more variable than in multicellular organisms ($\mu^* = 69$ ms), this difference being significant (Table S8b, second line, $p < 10^{-8}$). In both groups T_L increases slightly with mass, since the 95 % CI of the slope of the regression line α is [0.020, 0.14] for single cells and [0.028, 0.060] for multicellular organisms (Table 2). However, the trend of the multicellular group results from the largest vertebrates since for species under 50 kg the slope of the regression law α is not significantly different from zero (95 % CI is [-0.030, 0.11] (Table 2).

Since the maximum metabolic rate of organisms affects their maximum speed (Meyer-Vernet and Rospars 2016), we studied also the dependence of T_L on maximum specific metabolic rates (MSMR) in 14 species of mammals for which both values were determined (Table S5). The slope of the regression lines of MSMR against body mass (Fig. S6A) and of T_L against MSMR (Fig. S6B) are not significantly different from 0 and -1 respectively, suggesting independence of MSMR on mass and inverse dependence of T_L on MSMR.

4. DISCUSSION

4.1. *Variability and Mass Dependency of Timescales*

Over the whole mass range, the variabilities of the timescales expressed by the ranges including 95 % of the data (δT_{95} , Table 1), are so small compared to the mass ranges (δM , Table 2), that

1
2
3 280 their ratios are always less than one in a million ($10^{\delta T_{95}}/10^{\delta M}$). Indeed, 95 % of the simple
4
5 281 response times T_S (Fig. 2), of the visual resolution times ($T_F = 1/\text{CFF}$, Fig. 3) and of the minimum
6
7 282 locomotor times T_L (Fig. 4) lie within less than a factor of ten, five and six respectively of their
8
9 283 geometric mean, whereas the mass varies by 20, 8, and 21 orders of magnitude for T_S , T_F and T_L
10
11 284 respectively. This is noteworthy, given the diversity of sensorimotor systems and the huge mass
12
13 285 range, and suggests that these times are strongly constrained by physics, as was previously
14
15 286 proposed for T_L . The small variation of the visual resolution times is remarkable, given the
16
17 287 variation in spatiotemporal optical quality with eye size, which is correlated to body size (e.g.
18
19 288 Currea et al. 2018), and the large diversity of visual systems (e.g. Fernald 2000) and of strategies
20
21 289 of spatiotemporal summations. Although CFF is unrivaled to quantify the ability of an organism
22
23 290 to track a moving object, whether it is representative of other sensory systems is an open question
24
25 291 for further investigation. However, as suggested by a few reports (for example in olfaction,
26
27 292 Rumbo and Kaissling 1989, Lemon and Getz 1997, Smear et al. 2011, Jacob et al. 2017),
28
29 293 temporal resolution might reflect properties common to diverse neural networks rather than
30
31 294 specificities of the receptors (Butts et al. 2007, Panzeri et al. 2010) and, hence, be similar in the
32
33 295 visual and other sensory systems.

34
35 296 Most of the overall variability results from variations at smaller mass ranges. Within uni- and
36
37 297 multi-cellular organisms considered separately, the mass-scaling exponents are not significantly
38
39 298 different from zero or result from a small-scale trend (T_L in large metazoans), so that the T_L
40
41 299 (small) mass dependency in single cells appears as an exception that will need further data to be
42
43 300 interpreted. For T_S and T_L , differences between single cells and metazoans contribute to the
44
45 301 overall variability. The twice longer geometric mean of T_L in single cells (Table 1) is well
46
47 302 documented ($n = 70$ species) but the six times longer one of T_S ($n = 7$) should be confirmed on a
48
49 303 larger sample.

Although outside the scope of this paper, we examined narrower mass ranges, i.e. groups of related species of the same class (or kingdom in microorganisms) including more than three species (Fig. S3, S4, S5 and Table S6). For most groups, the mass scaling exponents are positive with intercepts decreasing with mass, so that all the data still lie in a relatively narrow range, as found previously for the mass specific metabolism (Makarieva et al. 2008, Hatton 2019) and its maximum value (Makarieva et al. 2005). For simple response times T_s , the only exponent that reaches statistical significance is that of mammals (Fig. S3) where the trend explains 42% of the variance and agrees with previous studies (section 4.3). Although not significant, the trend in bacteria stands out owing to the proportion of variance it explains (81 %) and its steep slope ($\alpha = 0.66$) which agrees with first-principle derivations (section 5). For the flicker fusion time, the mass exponent of the small-scale regressions are not significantly different from zero. Finally, for the minimum time to move by one body length, the positive mass scaling exponent of mammals stems from large masses (the inflection point near 50 kg is apparent in Fig. 4), as noted and interpreted by Meyer-Vernet and Rospars (2016); we shall return to this point in section 5.

4.2. Possible Errors Resulting from Biological, Methodological and Experimental Factors

Are our data and analyses adequate for supporting our conclusion that the simple response time T_s lies within less than a factor of ten from the mean for organisms varying by more than 20 orders of magnitude in mass, as previously shown for the minimum time to move by one body length T_L (Meyer-Vernet and Rospars 2016)? That a similar finding holds for T_F over the smaller 10^8 -fold mass range for which this time could be measured? And that these time scales display

1
2
3 326 small or no systematic variation with body mass in uni- and in multicellular organisms? Several
4
5 327 criticisms could be raised.
6

7
8 328 First, one could object that the diversity of organisms, of sensory and motor mechanisms, and
9
10 329 of experimental procedures hides any trend in the data. However, it is unlikely that the restriction
11
12 330 to more homogeneous data would reveal trends presently hidden because the smaller size of
13
14 331 samples would decrease the statistical significance. On the contrary, the diversity of species,
15
16 332 measurements and systems is indispensable to estimate reliably the variability and mean of the
17
18 333 timescales, independently of the specializations and limitations of taxa and systems. Our aim is to
19
20 334 transcend mass scalings holding within groups by considering large ranges of mass and
21
22 335 taxonomic groups.
23

24
25
26 336 Second, the evolution of different traits is correlated throughout a phylogenetic lineage so that
27
28 337 species values do not represent statistically independent data (Felsenstein 1985). This leads to
29
30 338 overestimation of degrees of freedom, which artificially narrows confidence intervals. Our major
31
32 339 finding that timescales lie within close bounds and that for unicellular (for T_s) and multicellular
33
34 340 organisms (for all timescales) they do not significantly depend on mass would not be adversely
35
36 341 affected by any widening of confidence intervals (on the contrary for T_L in single cells) and is
37
38 342 therefore impervious to this criticism.
39

40
41
42 343 A third possible objection is that several experimental factors affecting the measurements have
43
44 344 not been considered explicitly. The electrical or behavioural measurement methods play a minor
45
46 345 role and correcting them would have practically no effect (Fig. S2 and Table S10 show that their
47
48 346 difference is not significant). However, the effect of temperature, which is an important
49
50 347 determinant of metabolism and behaviour, should be checked. We thus studied how correcting
51
52 348 the time scales of ectotherms to a standard temperature $\theta_0 = 20^\circ\text{C}$ would change our results (see
53
54 349 Methods). First, the temperature being unknown in several measurements, we considered the
55

350 worst-case scenario assuming that short time scales ($T < \mu^*$) were measured at 10 °C, and would
 351 thus be still shorter if they had been measured at θ_0 , and that long time scales ($T \geq \mu^*$) were
 352 measured at 30 °C, these two temperatures being close to the extremes in our data (Fig. S7A). For
 353 this preliminary test, we used the mean activation energy $E = 0.66$ eV observed in a wide range
 354 of species and traits (Dell et al. 2001), and studied how the percentage of timescales lying outside
 355 the range $[\mu^*/10, \mu^* \times 10]$ would be affected by temperature in this worst-case scenario. For all
 356 timescales, this percentage is less than 3 % without standardization (Table S11a) and less than
 357 10 % with standardization (Table S11b), except for T_s where it is 5 % and 22 % respectively.
 358 Next, noting the greater sensitivity of T_s , we applied a finer standardization procedure to this
 359 timescale, for which the temperature is known in 72 % of our data for ectothermic species, based
 360 on the different mean activation energies of movements away from and toward a stimulus (see
 361 Methods). This procedure leads to no significant change neither in the distribution of T_s values
 362 (Fig. S7B) nor in the number of species beyond the limits (it remains the same, i. e. four, Fig.
 363 S7). This indicates that our results in mean, variability and overall trend are robust with respect to
 364 temperature.

4.3. *Comparison with Previous Work*

368 Several studies across a wide range of mass and life forms have found organisms' properties to
 369 lie in a relatively narrow range, without mass scaling across groups, despite the size scaling
 370 observed within groups. This is the case for the mass-specific metabolic rate (Makarieva et al.
 371 2008, Hatton et al. 2019), its maximum value per unit of active mass (Makarieva et al. 2005), the
 372 cross-section-specific forces exerted by muscles and molecular motors (Marden and Allen 2002,

1
2
3 373 Marden 2005, Rospars and Meyer-Vernet 2016), and the minimum time to move by one body
4
5 374 length $T_L = L/V_{\max}$ (McMahon and Bonner 1983, Meyer-Vernet and Rospars 2015, 2016).
6
7
8 375 However, for the simple response time T_s and the critical flicker fusion time T_F , previous
9
10 376 studies concern only relatively small mass ranges and we have included these data. The mass
11
12 377 scaling of sensorimotor delays has been studied by More and Donelan (2018) in the particular
13
14 378 case of the stretch reflex – a monosynaptic reflex that governs the fastest neural response to
15
16 379 peripheral stimuli in terrestrial mammals, with a determination of the different components and a
17
18 380 comparison with stance and stride durations. In our T_s sample, we have used the sum of these
19
20 381 components except the force generation delay (time between force onset and peak), in coherence
21
22 382 with the rest of our data, for which the response is measured by the onset of force production or
23
24 383 the onset of movement (which can begin before the production of peak force). These data points
25
26 384 lie within the range of our general data set T_s for multicellular organisms, where the nerve
27
28 385 conduction delay is expected to be mainly responsible for the increase of T_s shown in figure 2 at
29
30 386 large mass. Indeed, the increase in nerve conduction speed due to increase in fibre diameter and
31
32 387 to myelination is too moderate to compensate for the increase in distance of conduction, because
33
34 388 of the trade-off between responsiveness and compactness (Castelfranco and Hartline 2016),
35
36 389 resolution (More et al. 2013), and energy cost (Perge et al. 2012). Let us compare the nerve
37
38 390 conduction delay estimated by More and Donelan (2018) $T_{\text{cond}} = 5.3 M^{0.3}$ ms to the geometric
39
40 391 mean ($T_s = 22$ ms) that we find for multicellular species (Table 1). This yields
41
42 392 $T_{\text{cond}}/T_s = 0.24 M^{0.3}$, which shows that the conduction delay becomes important only for large M ,
43
44 393 although even for the largest mammal studied (the elephant), T_s is only four times larger than its
45
46 394 average value.
47
48
49
50
51
52
53
54 395

396 5. QUANTITATIVE INTERPRETATIONS FROM FIRST PRINCIPLES

397
398 The variation of the simple response time, as well as the other time scales studied here, by less
399 than a factor of ten around their mean for organisms so diverse in structure and size suggests that
400 it may be determined by basic constraints set by the universal properties of living matter. We will
401 thus derive first-principle estimates, trying to capture the essential processes at play, whereas
402 neglecting specific details that should be considered in scaling studies over narrow ranges of
403 mass and taxa. Such simple analytic calculations are expected to yield only order-of-magnitude
404 results, that is to within a 10-fold or so accuracy, similar to the variability of the time scales in
405 our data, in the line of the so-called “Fermi problems” or of Weisskopf’s physics courses
406 (Weisskopf 1975, 1989).

408 5.1. *Simple Response Time*

409
410 Consider first the simple response time T_s of microorganisms. It includes the transmission delay
411 from the sensor(s) receiving external stimuli to the motor apparatus producing the response by
412 regulating the swimming behaviour. A basic process is the transport of a signalling molecule
413 through the cytoplasm (e.g. Bitbol and Wingreen 2015) via diffusion (e.g., Purcell 1977,
414 Dusenberry 2009). In water, of viscosity $\eta \simeq 10^{-3} \text{ kg m}^{-1} \text{ s}^{-1}$, a sphere of diameter d at
415 temperature θ has a diffusion coefficient $D = k_B \theta / (3\pi \eta d)$, where k_B is Boltzmann constant.
416 Signalling molecules are small proteins of typical size $d \simeq 3 \text{ nm}$, like the key signalling protein
417 CheY (Bren and Eisenbach 2000), yielding a diffusion coefficient in water $D \simeq 150 \mu\text{m}^2/\text{s}$.
418 Assuming that the crowding of the cytoplasm decreases D by one order of magnitude (e.g. Dill et

1
2
3 419 al. 2011, Mika and Poolman 2011) and that the sensor-to-motor distance equals the organism's
4
5 420 length L , we find the three-dimensional diffusion time $L^2/6D$
6
7
8 421 $\tau_{\text{dif}} \simeq 10 L_{\mu\text{m}}^2 \text{ ms.}$ (1)
9
10 422 This timescale is the minimum response time T_s of a microorganism of size L if information is
11
12 423 transmitted through the cytoplasm by diffusion of signalling proteins, which is the most basic
13
14 424 process. With $L \propto M^{1/3}$, this would yield a mass variation $T_s \propto M^{2/3}$, as observed for bacteria (see
15
16
17 425 the leftmost regression line in Figure S3 with slope $\alpha = 0.66$, as given in Table S5). The median
18
19 426 mass of a bacterium of length L being $M \simeq (3.3L)^3$ (Meyer-Vernet and Rospars 2016), the length
20
21 427 of a bacterium of mass 10^{-15} kg (μ_M^* , Table S6) is $L \simeq 3 \mu\text{m}$. We deduce from Eq. (1) $T_s = 90$
22
23 428 ms, which agrees to better than a factor of two with the empirical value for bacteria ($\mu^* = 144 \text{ ms}$
24
25 429 in Table S5) and is close to the response time given by their regression line ($T_s = 100 \text{ ms}$ for
26
27 430 $M = 10^{-15} \text{ kg}$, Table S6).

31 431 Consider now multicellular organisms. For most of them, the responses are mediated by
32
33 432 conduction of information via electric pulses propagating through neurons, whose membrane
34
35 433 regulates the permeation of ions via an insulating lipid bilayer and proteins. The proteins act as
36
37 434 active ion channels and pumps and the lipid bilayer acts as a capacitance that enables charges to
38
39 435 accumulate and produce a potential across it. Biological membranes are also involved in sense
40
41 436 organs, via the concentration gradients they enable which produce active transport, whereas ion
42
43 437 channels play an essential role in signal transduction (Martinac and Cox 2016). The role of ions
44
45 438 to alter charge and thus protein conformation is essential in signal transduction in both uni- and
46
47 439 multicellular organisms (Clapham 2007). For example, paramecia use Ca^{2+} ions and show
48
49 440 dynamic changes in the electrical properties of their membrane in response to stimuli, as do
50
51
52
53
54
55
56
57
58

neurons, and the structures of their receptors are similar to those of vertebrates (e.g. Maegawa 2017).

A tentative first principle estimate of T_S can be obtained from the inverse of the number of action potentials sent along an axon per second, since this is the shortest possible time to send a bit of information along an axon. Let us first evaluate the cost of generating an action potential U per surface S of membrane, of width a and dielectric constant ϵ_m , so that the capacitance is

$$C \simeq \epsilon_0 \epsilon_m S/a, \quad (2)$$

where ϵ_0 is vacuum permittivity. With the electric charge across the membrane CU , of energy CU^2 , the cost of an action potential is obtained from (2) as $\epsilon_0 \epsilon_m S U^2/a$, which agrees with values in the literature (Aiello 2000). Using two biological properties common to living matter: the mass density $\rho \simeq 10^3 \text{ kg/m}^3$ and the maximum metabolic rate per unit mass of active tissue $b_M \simeq 2 \times 10^3 \text{ W/kg}$ (Makarieva et al. 2005), the maximum power available to this surface S of membrane of mass $\rho a S$ is

$$b_M \rho a S \quad (3)$$

Dividing this maximum power by the cost of an action potential estimated above and using the energy corresponding to one monovalent ion crossing the membrane $eU \simeq W_0$, where $e = 1.6 \times 10^{-19} \text{ C}$ is the electron charge and W_0 is the energy released by one ATP molecule, we deduce the maximum number of action potentials per unit time, whose inverse yields

$$T_S \simeq \epsilon_0 \epsilon_m W_0^2 / (\rho b_M e^2 a^2) \quad (4)$$

Substituting the typical dielectric constant of lipids $\epsilon_m \simeq 2$, a membrane width $a \simeq 6 \text{ nm}$ – similar to the typical protein size (e.g. Erickson 2009), the above value of b_M , and the first-principles relation (Meyer-Vernet and Rospars 2016)

$$W_0 \simeq e^2 / (4\pi \epsilon_0 a) \quad (5)$$

1
2
3 464 we obtain $T_S \simeq 14$ ms, close to the empirical mean simple response time T_S of metazoans (22
4
5 465 ms).

6 466
7
8
9
10 467 *5.2. Flicker Fusion Time*
11

12 468
13
14
15 469 Let us now estimate the timescale T_F , assuming that it is limited by the membrane time constant
16
17 470 τ_m . For a surface S of membrane of width a and resistivity r_m , the electrical resistance is
18
19 471 $R = r_m a/S$ and the capacitance is given by (2), so that the time constant $\tau_m = RC$ is given by
20
21
22 472
$$\tau_m = r_m \epsilon_0 \epsilon_m. \tag{6}$$

23

24 473 To estimate a basic value for the resistivity r_m , independent on the details of the system, we
25
26 474 consider again energy constraints. In an electrical circuit of resistance R , submitted to the
27
28
29 475 potential difference U , the power is U^2/R . Using the maximum power available (3), we deduce
30
31 476 the resistance $R \simeq U^2/(b_M \rho a S)$, whence the resistivity $r_m \simeq U^2/(\rho b_M a^2)$. Using again the order
32
33 477 of magnitude $U \simeq W_0/e$, we deduce τ_m , which yields the flicker fusion time $T_F \simeq \tau_m$ (reflecting the
34
35 478 ability of the membrane to resolve a time-varying signal)

36
37
38 479
$$T_F \simeq \epsilon_0 \epsilon_m W_0^2 / (\rho b_M e^2 a^2) \simeq T_S \simeq 14 \text{ ms} \tag{7}$$

39

40
41 480 close to the empirical value of T_F (25 ms), and in agreement with our empirical result $T_F \simeq T_S$.
42
43 481 The absence of significant increase of T_F for the largest masses (Fig. 3), contrary to what is
44
45 482 observed for T_S (Fig. 2) suggests that the signal conduction time and hence the organism length L
46
47 483 plays a minor role in T_F , contrary to T_S .
48
49

50 484
51
52 485 *5.3. Minimum Locomotion Time*
53
54 486

487 An order of magnitude estimate of the minimum time to move by one body length $T_L = L/V_{\max}$
 488 was derived by Meyer-Vernet and Rospars (2016) from the invariance over the whole domain of
 489 life of the force per cross-sectional area exerted by molecular motors and muscles ($f \simeq W_0/a^3 \simeq$
 490 $2 \times 10^5 \text{ N/m}^2$, Rospars and Meyer-Vernet 2016), and the two basic quantities considered above,
 491 mass density ρ and mass-specific (per unit mass of active tissue) metabolic rate at maximum
 492 activity b_M . The maximum speed was estimated as $V_{\max} \simeq L \rho b_M / f$ for swimming and running
 493 organisms (including microorganisms) of length L satisfying $L \lesssim (f/\rho)^{3/2}/b_M \simeq 1.4 \text{ m}$ –
 494 corresponding roughly to $M \simeq 50 \text{ kg}$ (Meyer-Vernet and Rospars 2016). This yields $T_L \simeq L/V_{\max}$,
 495 the minimum time to move by L

$$T_L \simeq f/(\rho b_M) \simeq W_0/(\rho b_M a^3) \simeq 100 \text{ ms} \quad (8)$$

497 for $M \lesssim 50 \text{ kg}$, which agrees to better than a factor of two with the empirical value (59 ms,
 498 geometric mean of T_L for all organisms of mass $M < 50 \text{ kg}$, Table 1). The increase in T_L observed
 499 for larger organisms (Fig. 4) has been interpreted by dynamic constraints, yielding

$$T_L \simeq L (\rho/f)^{1/2} \quad (9)$$

501 (Meyer-Vernet and Rospars 2016), in agreement with the known maximum speed of about 15
 502 m/s (e.g. McMahon and Bonner 1983, Garland, 1983, Iriarte-Diaz 2002).

503 It is interesting to note that our estimates of the minimum times for response (4), for vision
 504 (7), and for moving by one body length (8) all yield timescales proportional to the inverse of the
 505 maximum mass-specific metabolic rate (per unit of active mass). This suggests that, since the
 506 metabolic rate varies with mass within groups of related species, these time scales may depend
 507 not only on body mass within groups of related species but also on metabolic rates. This is shown
 508 for T_L in mammals (Fig. S6B), which is inversely proportional to the maximum mass-specific

1
2
3 509 metabolic rate, as expected from (8) if the proportion of active tissue in the body mass does not
4
5 510 vary in the range considered in the regression.
6

7
8 511 Finally, it is noteworthy that Eqs. (4), (5) and (8) yield $T_L/T_S \simeq 4\pi/\varepsilon_m \simeq 6$ in order of
9
10 512 magnitude, to be compared to the empirical ratio $T_L/T_S = 78/26 = 3$, suggesting that fast reacting
11
12 513 species are also fast moving.
13

14 514
15
16
17 515 6. CONCLUSIONS
18

19 516
20
21 517 We have shown that across the whole of mobile life, from bacteria to large vertebrates, the simple
22
23 518 response time T_S lies in a relatively narrow range, with 95 % of species reacting in a time that
24
25 519 differs by less than one order of magnitude from the mean, in striking contrast to the 20 orders of
26
27 520 magnitude difference in body mass. The simple response time does not display significant scaling
28
29 521 with body mass across groups in unicellular organisms nor in multicellular ones, although it is
30
31 522 almost six times larger in the former life form. The absence of large-scale trend does not preclude
32
33 523 – and is indeed compatible with various scalings valid in narrower ranges of size and taxa, as is
34
35 524 known for the specific metabolic rate. However, within narrow ranges, the detailed characteristics
36
37 525 of the organisms must be considered and it is only on larger scales that these variations can be
38
39 526 transcended.
40
41
42
43

44 527 This simple response time T_S is close to a well-studied sensory timescale (the inverse of the
45
46 528 critical flicker fusion frequency for vision) T_F , and is a few times smaller than the minimum time
47
48 529 to move by one body length T_L , with all time scales lying in a relatively narrow range. Since this
49
50 530 narrow range suggests that these time scales may be strongly constrained by physics and basic
51
52 531 properties of life, independently of the structure or mass of the organism, we have performed
53
54
55
56
57
58

tentative simple estimates of T_S and T_F based on fundamental physicochemical constants and basic properties common to motile organisms.

It is fair to note that the agreement between these simple estimates of minimum time scales (for sensing, for reacting and for moving) and the measurements is only indicative of the dominant constraints in play (e.g. Phillips and Milo 2009), since several mechanisms may be operating in parallel. For example, one tenth of second is in the middle of the range of protein folding time scales, which however spans six orders of magnitude (e.g. Lane and Pande 2013) and close to the maximal turn-over rate of the most abundant enzyme in the biosphere (e.g. Flamholz et al. 2019). The relation $T_S \simeq T_F < T_L$ found from the data (Fig. 1) and tentatively interpreted from first-principles, is reminiscent of symmorphosis (Weibel et al. 1991) and expected to be favoured by evolution since it is indicative of an optimum state. Indeed, if $T_S < T_F$ the sensory resolution would be in excess over motor control whereas if $T_S > T_F$ the response would be limited by sensory performance, so that in both cases, at least one component of the nervous system would be out of tune and wasting energy or space (Laughlin 2001). Likewise, the evolution is expected to favour organisms for which the sensory and response organs act fast enough with respect to their moving performances. It is noteworthy that since the first-principle estimate (8) of T_L holds for both uni- and multi-cellular organisms (Meyer-Vernet and Rospars 2016), the estimate (1) of the minimal response time T_S yields a maximal length L of a few micrometres in order to ensure $T_S \leq T_L$ when the response is mediated by diffusion, as observed for bacteria

These preliminary results encourage further research on the response times of mobile organisms and their fundamental bases. They might also possibly be used to infer properties of extinct species since the reconstructed speed (e.g. Hutchinson and Garcia 2002) and size (e.g.

1
2
3 555 Hutchinson et al. 2011) of the giant dinosaur *Tyrannosaurus rex* yield a time to move by one
4
5 556 body length $T_L \simeq 1$ s for a mass $M \simeq 10^4$ kg, which is close to the value expected from (9) and
6
7
8 557 would put *Tyrannosaurus rex* in the middle of the data of extant animals of this size in figure 4.
9
10 558

559 *Ethical Statement*

560 The present article is based on a meta-analysis. No live animals were used.

561

562 *Data Accessibility*

563 The datasets supporting the article are included in the Supplementary Material.

564

565 *Competing Interests*

566 We have no competing interests.

567

568 *Author's Contributions*

569 J.-P.R. and N.M.-V. each made significant and substantial contributions to this study in terms of

570 the conception, design, data collection and interpretation of results, as well as preparing the

571 manuscript. J.-P.R. contributed primarily to the statistical analyses and N.M.-V. to the physical

572 analyses.

573

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

REFERENCES

Abraham, N. M., Spors, H., Carleton, A., Margrie, T. W., Kuner, T. and Schaefer, A. T. (2004).
Maintaining accuracy at the expense of speed: stimulus similarity defines odor
discrimination time in mice. *Neuron*, 44 (5), 865-876

Agutter, P. S. and Wheatley, D. N. (2004) Metabolic scaling: consensus or controversy? *Theor.*
Biol. Med. Model. 1 (1), 13

Aiello, G. L. and Bach-Y-Rita, P. (2000) The cost of an action potential. *J. Neuroscience*
Methods 103, 145-149

Attwell, D. and Laughlin, S. B. (2001) An energy budget for signaling in the grey matter of the
brain. *J. Cerebral Blood Flow Metabol.* 21 (10), 1133-1145

Bartholomew, G. A. and Heinrich, B. (1973) A field study of flight temperatures in moths in
relation to body weight and wing loading. *J. Exp. Biol.* 58, 123-135

Bejan, A. and Marden, J. H. (2006) Unifying constructal theory for scale effects in running,
swimming and flying. *J. Exp. Biol.* 209 (2), 238-248

Bitbol, A. F. and Wingreen, N. S. (2015) Fundamental constraints on the abundances of
chemotaxis proteins. *Biophys. J.* 108 (5), 1293-1305

Block, S. M., Segall, J. E. and Berg, H. C. (1982) Impulse responses in bacterial chemotaxis.
Cell, 31 (1), 215-226

Blough, D. S. (1978) Reaction times of pigeons on a wavelength discrimination task. *J. Exp.*
Anal. Behav., 30 (2), 163-167

Bonner, J T (1965) *Size and Cycle: An Essay on the Structure of Biology* (Princeton N J:
Princeton University Press)

- 597 Bren, A. and Eisenbach, M. (2000) How signals are heard during bacterial chemotaxis: protein-
 598 protein interactions in sensory signal propagation. *J. Bacteriol.* 182 (24), 6865-6873
- 599 Bullock, T. H. (1984) Comparative neuroethology of startle, rapid escape, and giant fiber-
 600 mediated responses. In *Neural Mechanisms of Startle Behavior* (ed. R.C. Eaton), pp. 1-13.
 601 Boston: Springer Science and Business Media
- 602 Butts, D. A., Weng, C., Jin, J., Yeh, C. I., Lesica, N. A., Alonso, J. M. and Stanley, G. B. (2007)
 603 Temporal precision in the neural code and the timescales of natural vision. *Nature* 449, 92–
 604 95
- 605 Byrne, D.N., Buchmann, S.L. and Spangler, H.G. (1988) Relationship between wing loading,
 606 wingbeat frequency and body mass in homopterous insects. *J. Exp. Biol.* 135, 9-23.
- 607 Castelfranco, A. M. and Hartline, D. K. (2016) Evolution of rapid nerve conduction. *Brain Res.*
 608 1641, 11-33.
- 609 Clapham, D.E. (2007) Calcium signaling. *Cell*, 131 (6), 1047-1058
- 610 Currea, J. P., Smith, J. L. and Theobald, J. C. (2018) Small fruit flies sacrifice temporal acuity to
 611 maintain contrast sensitivity. *Vision Res.* 149, 1-8
- 612 Dagnelie, P. (2011) *Statistique théorique et appliquée. Tome 2. Inférence statistique à une et à*
 613 *deux dimensions.* Bruxelles, De Boeck
- 614 D'Eath, R. B. (1998) Can video images imitate real stimuli in animal behaviour experiments?.
 615 *Biol. Rev.* 73 (3), 267-292.
- 616 Dell, A. I., Pawar, S., and Savage, V. M. (2001) Systematic variation in the temperature
 617 dependence of physiological and ecological traits. *Proc. Natl. Acad. Sci.* 108 (26), 10591-
 618 10596.
- 619 Dill, K. A., Ghosh, K. and Schmidt, D. (2011) Physical limits of cells and proteomes *Proc. Nat.*
 620 *Acad. Sci. USA* 108 17876-17882

1
2
3 621 Dusenberry, D. B. (2009) *Living at Micro Scale. The Unexpected Physics of Being Small*.
4
5 622 Cambridge, MA: Harvard University Press
6
7 623 Erickson, H. P. (2009) Size and shape of protein molecules at the nanometer level determined by
8
9 624 sedimentation, gel filtration, and electron microscopy, in *Biological procedures Online* ed
10
11 625 Shulin Li vol 11, 1
12
13
14 626 Falk-Petersen, S. (1981) Ecological investigations on the zooplankton community of Balsfjorden,
15
16 627 Northern Norway: seasonal changes in body weight and the main biochemical composition
17
18 628 of *Thysanoessa inermis* (Krøyer), *T. raschii* (M. Sars), and *Meganyctiphanes norvegica* (M.
19
20 629 Sars) in relation to environmental factors. *J. Exp. Mar. Biol. Ecol.* 49, 103-120
21
22
23 630 Felsenstein, J. (1985) Phylogenies and the comparative method. *The American Naturalist*, 125(1),
24
25 631 1-15
26
27
28 632 Fernald, R. D. (2000) Evolution of eyes. *Curr. Opin. Neurobiol.* 10 (4), 444-450
29
30
31 633 Flamholz, A. I., Prywes, N., Moran, U., Davidi, D., Bar-On, Y. M., Oltrogge, L. M., Alves, R.,
32
33 634 Savage, D. and Milo, R. (2019) Revisiting trade-offs between Rubisco kinetic parameters
34
35 635 *Biochemistry* 58, 3365-76
36
37
38 636 Froese, R., Thorson, J. T. and Reyes, R. B. (2014) A Bayesian approach for estimating length-
39
40 637 weight relationships in fishes. *J. Applied Ichthyol.* 30, 78-85
41
42 638 Garland, T. Jr. (1983) The relation between maximal running speed and body mass in terrestrial
43
44 639 mammals. *J. Zool.*, 199, 157-70
45
46
47 640 Glazier, D. (2014) Metabolic scaling in complex living systems. *Systems*, 2 (4), 451-540
48
49 641 Hatton, I. A., Dobson, A. P., Storch, D., Galbraith, E. D. and Loreau, M. (2019) Linking scaling
50
51 642 laws across eukaryotes. *Proc. Natl. Acad. Sci.* DOI: 10.1073/pnas.19004921161900492116
52
53
54 643 Healy, K., McNally, L., Ruxton, G. D., Cooper, N. and Jackson, A. L. (2013) Metabolic rate and
55
56 644 body size are linked with perception of temporal information. *Animal Behav.* 86 (4), 685-696
57
58

- Herberholz, J. and Marquart, G. D. (2012) Decision making and behavioral choice during predator avoidance. *Front. Neurosci.* 6, 125
- Hutchinson, J. R. and Garcia, M. (2002) *Tyrannosaurus* was not a fast runner. *Nature* 415, 1018-1022
- Hutchinson, J. R., Bates, K. T., Molnar, J., Allen, V. and Makovicky, P. J. (2011) A computational analysis of limb and body dimensions in *Tyrannosaurus rex* with implications for locomotion, ontogeny, and growth. *PLoS One*, 6, e26037
- Inger, R., Bennie, J., Davies, T. and Gaston, K. (2014) Potential biological and ecological effects of flickering artificial light. *PLoS One* 9, e98631
- Iriarte-Diaz, J. (2002) Differential scaling of locomotor performance in small and large terrestrial mammals. *J. Exp. Biol.* 205, 2897-2908
- Jacob, V., Monsempès, C., Rospars, J.-P., Masson, J.-B. and Lucas, P. (2017) Olfactory coding in the turbulent realm. *PLoS Comput. Biol.* 13 (12): e1005870
- Koch, M. (1999) The neurobiology of startle. *Progress in neurobiology*, 59 (2), 107-128
- Lane, T. J. and Pande V. S. (2013) Inferring the rate-length law of protein folding. *PLoS One* e78606
- Laughlin, S. B. (2001) Energy as a constraint on the coding and processing of sensory information. *Curr. Opinion in Neurobiol.*, 11, 475e480
- Lemon, W. and Getz, W. (1997) Temporal resolution of general odor pulses by olfactory sensory neurons in American cockroaches. *J. Exp. Biol.*, 200 (12), 1809-1819
- Limpert, E., Stahel, W. A. and Abbt, M. (2001) Log-normal distributions across the sciences: keys and clues. *BioSci.* 51 (5), 341-352
- Luce, R. D. (1991) *Response Times*. New York: Oxford University Press.

1
2
3 668 Makarieva, A. M., Gorshkov, V. G. and Li, B. L. (2005) Energetics of the smallest: do bacteria
4
5 669 breathe at the same rate as whales?. *Proc. R. Soc. B: Biol. Sci.* 272 (1577), 2219-2224
6
7 670 Makarieva, A. M., Gorshkov, V. G., Li, B. L., Chown, S. L., Reich, P. B. and Gavrilov, V. M.
8
9 671 (2008) Mean mass-specific metabolic rates are strikingly similar across life's major domains:
10
11 672 Evidence for life's metabolic optimum. *Proc. Natl. Acad. Sci.* 105, 16994-16999
12
13
14 673 Maegawa, S. (2017) Molecular characteristics of neuron-like functions in single-cell organisms,
15
16 674 In Shigeno et al. eds *Brain Evolution by Design: from Neutral Origin to Cognitive*
17
18 675 *Architecture*, Springer
19
20
21 676 Marden, J. H. and Allen, L R. (2002) Molecules, muscles, and machines: universal performance
22
23 677 characteristics of motors. *Proc. Natl. Acad. Sci. USA* 99, 4161-4166
24
25
26 678 Marden, J. H. (2005) Review: Scaling of maximum net force output by motors used for
27
28 679 locomotion. *J. Exp. Biol.* 208, 1653-64
29
30
31 680 McMahon, T. A. and Bonner, J. T. (1983) *On Size and Life* (New York: Scientific American
32
33 681 Library)
34
35 682 Mika, J. T. and Poolman, B. (2011) Macromolecule diffusion and confinement in prokaryotic
36
37 683 cells. *Current Opinion in Biotechnology* 22, 117-126
38
39
40 684 Martens, E.A., Wadhwa, N., Jacobsen, N. S., Lindemann, C., Andersen, K. H. and Visser, A.
41
42 685 (2015) Size structures sensory hierarchy in ocean life. *Proc. R. Soc. B*, 282, 20151346
43
44 686 Martinac, B. and Cox, C. D. (2016) Mechanosensory transduction: Focus on ion channels.
45
46 687 *Comprehensive Biophysics*, Elsevier
47
48
49 688 Medina, J. M., Wong, W., Diaz, J. A. and Colonius, H. (2015) Advances in modern mental
50
51 689 chronometry. *Front. Human Neurosci.* 9, 256
52
53
54 690 Meyer-Vernet, N. and Rospars, J. P. (2015) How fast do living organisms move: Maximum
55
56 691 speeds from bacteria to elephants and whales. *Am. J. Phys.* 83 (8), 719-722
57
58

- 692 Meyer-Vernet, N. and Rospars, J. P. (2016) Maximum relative speeds of living organisms: Why
 693 do bacteria perform as fast as ostriches? *Phys. Biol.* 13 (6), 066006
- 694 Mitchell, J. G. (1991). The influence of cell size on marine bacterial motility and energetics.
 695 *Microbial Ecology*, 22 (1), 227-238
- 696 More, H. L. and Donelan, J. M. (2018) Scaling of sensorimotor delays in terrestrial mammals.
 697 *Proc. R. Soc. B: Biol. Sci.* 285 (1885), 20180613
- 698 More, H. L., Hutchinson, J. R., Collins, D. F., Weber, D. J., Aung, S. K. H. and Donelan, J. M.
 699 (2010) Scaling of sensorimotor control in terrestrial mammals. *Proc. R. Soc. B* 277, 3563-
 700 3568
- 701 Niven, J. E. and Scharlemann, J. P. W. (2005) Does metabolic rate at rest and during flight scale
 702 with body mass in insects? *Biol. Lett.* 1, 346-349
- 703 Panzeri, S., Brunel, N., Logothetis, N. K. and Kayser, C. (2010) Sensory neural codes using
 704 multiplexed temporal scales. *Trends Neurosci.* 33 (3), 111-120
- 705 Perge, J. A., Niven, J. E., Mugnaini, E., Balasubramanian, V. and Sterling, P. (2012) Why do
 706 axons differ in caliber? *J. Neurosci.* 32 (2), 626-638
- 707 Phillips, R. and Milo, R. (2009) A feeling for the numbers in biology *Proc. Natl. Acad. Sci.* 106,
 708 21465-21471
- 709 Purcell, E. M. (1977) Life at low Reynolds number. *Am. J. Phys.* 45 (1), 3-11
- 710 Roberts, A., Borisyuk, R., Buhl, E., Ferrario, A., Koutsikou, S., Li, W. C. and Soffe, S. R. (2019)
 711 The decision to move: response times, neuronal circuits and sensory memory in a simple
 712 vertebrate. *Proc. R. Soc. B*, 286 (1899), 20190297
- 713 Roeder, K. D. (1963) *Nerve Cells and Insect Behavior*. Cambridge, USA: Harvard University
 714 Press, 188 pp.

1
2
3 715 Rospars, J. P. and Meyer-Vernet, N. (2016) Force per cross-sectional area from molecules to
4
5 716 muscles: a general property of biological motors. *R. Soc. Open Sci.* 3 (7), 160313
6
7 717 Rumbo, E. R. and Kaissling, K. E. (1989) Temporal resolution of odour pulses by three types of
8
9 718 pheromone receptor cells in *Antheraea polyphemus*. *J. Comp. Physiol. A*, 165 (3), 281-291
10
11 719 Sillar, K. T., Picton, L. D. and Heitler, W. J. (2016) *The Neuroethology of Predation and Escape*.
12
13 John Wiley & Sons
14
15 720 Shamir, M., Bar-On, Y., Phillips, R. and Milo, R. (2016) Snapshot: timescales in cell biology.
16
17 721 *Cell* 164, 1302-1303
18
19 722 Smear, M., Shusterman, R., O’connor, R., Bozza, T. and Rinberg, D. (2011) Perception of sniff
20
21 723 phase in mouse olfaction. *Nature*, 479 (7373), 397-400
22
23 724 Spence, A. J. (2009) Scaling in biology. *Current Biology* 19, R57-R61
24
25 725 Weibel, E. R., Taylor, C. R. and Hoppeler, H. (1991) The concept of symmorphosis: A testable
26
27 726 hypothesis of structure-function relationship *Proc. Natl. Acad. Sci. USA* 88, 10357-10361
28
29 727 Weibel, E. R. and Hoppeler, H. (2005) Exercise-induced maximal metabolic rate scales with
30
31 728 muscle aerobic capacity. *J. Exp. Biol.* 208 (9), 1635-1644
32
33 729 Weisskopf, V. F. (1975) Of atoms, mountains, and stars: a study in qualitative physics. *Science*,
34
35 730 187 (4177), 605-612
36
37 731 Weisskopf, V. F. (1989) *The Privilege of Being a Physicist* (W. H. Freeman, New York, 1989)
38
39 732
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

TABLE 1. Summary statistics of timescales (in ms) for unicellular, multicellular and all species

Timescale	n	min	max	$Q_{2.5}$	$Q_{97.5}$	r_{95}	δT_{95}	med	IQR	μ^*	s^*	out ₉₅	out ₁₀
T_S all	81	2.5	485	4.7	311	66	1.82	24	43	26	3.26	4	4
T_S unicell	7	18.0	350	18.0	350	19	1.29	200	195	129	3.17	0	0
T_S multicell	74	2.5	485	4.7	238	51	1.71	23	35	22	2.93	4	3
T_S multic<50kg	68	2.5	240	4.6	148	32	1.51	18	28	19	2.64	4	1
T_F all multic	106	2.5	250	4.2	100	24	1.37	24	27	25	2.14	5	1
T_L all	427	5.3	2778	20.9	706	34	1.53	71	99	78	2.73	22	11
T_L unicell	70	5.3	2778	11.9	2778	234	2.37	123	213	145	3.86	2	9
T_L multicell	357	12.5	2525	22.3	567	25	1.41	61	82	69	2.40	18	3
T_L multic<50kg	300	12.5	657	21.5	408	19	1.28	50	61	59	2.22	14	3

Rows: T_S , simple response time; T_F , flicker fusion time; T_L , minimum time to move by one body length.

Columns: n , number of species; minimum; maximum; $Q_{2.5}$, percentile 2.5 %; $Q_{97.5}$, percentile 97.5 %; r_{95} , ratio $Q_{97.5}/Q_{2.5}$ including 95 % values; $\delta T_{95} = \log_{10}(r_{25})$; med, median; IQR, interquartile range; μ^* , geometric mean (ms); s^* , multiplicative standard deviation; out₉₅, number of values outside $[Q_{2.5}, Q_{97.5}]$; out₁₀, number of values outside $[\mu^*/10, \mu^* \times 10]$.

TABLE 2. Dependence of timescales on body mass for unicellular, multicellular and all species

		<i>n</i>	$\log_{10}T_0$	α	IC ₁	IC ₂	μ_M^*	δM	r_α	δT_α	r^2	<i>P</i>	Sig
<i>T_S</i> all		81	-1.69	-0.033	-0.06	-0.01	-3.10	19.59	0.2	-0.64	7	0.02	1
<i>T_S</i> unicell		7	-0.35	0.038	-0.41	0.49	-14.17	3.70	1.4	0.14	1	0.83	0
<i>T_S</i> multicell		74	-1.58	0.036	-0.01	0.08	-2.05	12.11	2.7	0.44	3	0.12	0
<i>T_S</i> multic<50kg		68	-1.77	-0.024	-0.08	0.03	-2.45	9.89	0.6	-0.24	1	0.36	0
<i>T_F</i> all multic		106	-1.57	0.023	-0.01	0.06	-1.93	8.25	1.6	0.19	2	0.19	0
<i>T_L</i> all		427	-1.13	-0.010	-0.02	-0.00	-2.86	21.05	0.6	-0.20	1	0.03	1
<i>T_L</i> unicell		70	0.10	0.078	0.02	0.14	-12.01	8.01	4.2	0.63	10	0.01	1
<i>T_L</i> multice		357	-1.11	0.044	0.03	0.06	-1.06	14.15	4.2	0.62	7	0.00	1
<i>T_L</i> multic<50kg		300	-1.25	-0.010	-0.03	0.01	-1.72	10.70	0.8	-0.10	0	0.34	0

Rows: same as in Table 1.

Columns: *n*, number of species; $\log_{10}T_0$, intercept of least square regression line, $\log_{10} T = \log_{10}T_0 + \alpha \log_{10} M$; α , allometric coefficient; [IC₁, IC₂], 95% confidence intervals of slope α ; μ_M^* , mean $\log_{10}M$; δM , mass range of the category, $\delta M = \log_{10}(M_{\max}/M_{\min})$, with $M_{\max}$ and $M_{\min}$ masses of heaviest and lightest species in the category; r_α , fitted ratio $T_{\max}/T_{\min}$ with $T_{\max} = T_0(M_{\max})^\alpha$ and $T_{\min} = T_0(M_{\min})^\alpha$; $\delta T_\alpha = \log_{10} r_\alpha$; r^2 , coefficient of determination (percent); *P*, p-value of test of slope $\alpha = 0$; Sig, slope α of regression line significantly different from zero (1) or not (0).

Figure 1. Boxplots of log-transformed flicker fusion times T_F , simple response times T_S , and times to move by one body length at maximum speed T_L . The boxes extend from the lower quartile to the upper quartile values with the medians (red line) in between. The whiskers extend to the most extreme data values within $1.5 \times \text{IQR}$. Outliers (red crosses) are values beyond the end of the upper whiskers. ANOVA and multiple comparisons of means (Supplementary material, Table S7): $T_F = T_S \neq T_L$.

Figure 2. Simple response times T_s versus cell or body mass M ($n = 81$). For clarity, the scale on the y-axis is 1.5 times larger than on the x-axis .

A. Taxonomic groups. Groups with three species or less shown as empty symbols, other groups as filled symbols. Bacteria (Bac, $n = 4$), Echinodermata (Ech, 1, sperm), Plantae (Pla, 2), Arachnida (Ara, 1), Insecta (Ins, 16), Hexanauplia (Hex, 1; copepods), Cnidaria (Cni, 3), Malacostraca (Mal, 2; crustaceans), Polychaeta (Pol, 1; bristle annelids), Oligochaeta (Oli, 1; earthworms), Actinopterygii (Act, 28; cartilaginous fishes), Amphibia (Amp, 1), Aves (Ave, 1; birds), Mammalia (Mam, 18). The horizontal dashed line is the geometric mean μ^* (26 ms) with values larger and smaller by one order of magnitude dotted. Vertical dashed line separates unicellular from multicellular organisms. Intermediate-scale regression laws for, from left to right, single cells (solid black line), multicellular organisms above 50 kg excluded (dashed black line) and included (solid black line). Slopes not significantly different from zero.

B. Types of response. Species tested for a single type of response shown in filled symbols: cell chemotaxis and phototaxis (Tax, $n = 7$), startle with escape (Esc, 46), sensory control of locomotion (Loc, 11), predatory movement (Pre, 6), small movement in startle of birds and mammals (Sma, 2), other behaviour (Oth, 3; see Table S1). Species tested for two or more types of response (empty symbol, Mul, 6). Dashed and dotted horizontal lines as in A.

Figure 3. Flicker fusion times T_F versus body mass M ($n = 106$) according to taxonomic groups, shown as colour of empty and filled symbols as in Fig. 2. Arachnida (Ara, $n = 2$), Insecta (Ins, 21), Malacostraca (Mal, 29; crustaceans), Cho (Chondrichthyes, 5; cartilaginous fishes), Actinopterygii (Act, 11; ray-finned fishes), Amphibia (Amp, 3), Reptilia (Rep, 15), Aves (Ave, 7; birds), Mammalia (Mam, 13). The horizontal dashed line is the geometric mean μ^* (25 ms) with values larger and smaller by one order of magnitude dotted. Scaling regression law in inset. Slope not significantly different from zero.

Figure 4. Times to move by one body length at maximum speed T_L versus cell or body mass ($n = 426$) according to taxonomic groups. For clarity, the scale on the y-axis is 1.5 times larger than on the x-axis. Groups with three species or less shown as empty symbols, other groups as filled symbols. Bacteria (Bac, $n = 17$), Protozoa (Pro, 7; flagellates), Bivalvia (Biv, 1, sperm), Planta (Pla, 3), Echinodermata (Ech, 3, sperm), Polychaeta (Pol, 1, sperm; bristle annelids), Ascidiacea (Asc, 1, sperm; sea squirts), Chromista (Chr, 32; ciliates), Hexanauplia (Hex, 6; copepods), Arachnida (Ara, 10), Insecta (Ins, 20; 2 sperm), Malacostraca (Mal, 3; crustaceans), Cephalopoda (Cep, 1), Actinopterygii (Act, 55; cartilaginous fishes), Reptilia (Rep, 96), Aves (Ave, 9; birds), Mammalia (Mam, 161; 3 sperm). The horizontal dashed line is the geometric mean μ^* (78 ms) with values larger and smaller by one order of magnitude dotted. Vertical dashed line approximately separates unicellular and multicellular organisms. Intermediate-scale regression laws in inset as in Fig. 2. Slopes significantly different from zero, except for multicellular organisms under 50 kg.