



HAL
open science

Longitudinal Cytokine Profiling in Severe COVID-19 Patients on ECMO and Haemoadsorption

Guillaume Lebreton, Karim Dorgham, Paul Quentric, Alain Combes, Guy Gorochov, Matthieu Schmidt

► **To cite this version:**

Guillaume Lebreton, Karim Dorgham, Paul Quentric, Alain Combes, Guy Gorochov, et al.. Longitudinal Cytokine Profiling in Severe COVID-19 Patients on ECMO and Haemoadsorption. American Journal of Respiratory and Critical Care Medicine, 2021, 10.1164/rccm.202011-4140le . hal-03172057

HAL Id: hal-03172057

<https://hal.sorbonne-universite.fr/hal-03172057>

Submitted on 17 Mar 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Longitudinal Cytokine Profiling in Severe COVID-19 Patients on ECMO and Haemoadsorption

Guillaume Lebreton, MD, PhD¹⁻², Karim Dorgham, PhD³, Paul Quentric, MD, MSc³,

Alain Combes, MD, PhD¹⁻⁴⁻⁵, Guy Gorochov, MD, PhD³⁻⁶, Matthieu Schmidt, MD, PhD¹⁻⁴⁻⁵

¹ Sorbonne Université INSERM Unité Mixte de Recherche (UMRS) 1166, Institute of
Cardiometabolism and Nutrition, Paris, France

² Service de chirurgie cardiaque, Institut de Cardiologie, APHP Sorbonne Université Hôpital
Pitié–Salpêtrière, F-75013 PARIS, France.

³ Sorbonne Université, Inserm, Centre d’Immunologie et des Maladies Infectieuses, CIMI-
Paris, F-75013, Paris, France.

⁴ Service de Médecine Intensive-Réanimation, Institut de Cardiologie, Assistance Publique-
Hôpitaux de Paris (APHP) Hôpital Pitié-Salpêtrière, Paris, France

⁵ Sorbonne Université, GRC 30, RESPIRE, Assistance Publique-Hôpitaux de Paris (APHP)
Hôpital Pitié-Salpêtrière, Paris, France

⁶ Assistance Publique-Hôpitaux de Paris (AP-HP), Hôpital Pitié-Salpêtrière, Département
d’Immunologie, F-75013, Paris, France.

Source of funding: The study was partially supported by a clinical research program (PHRC
N 2020 "Digital Covid") funded by AP-HP, Paris.

Contributors:

GL, AC, and MS designed the study and collected the data; KD, PQ, and GG performed the

cytokines analyses; KD and PQ performed the statistical analysis; MS wrote the first draft of the manuscript; GL, AC, and GG provided critical revisions of the manuscript. All authors gave critical comments on the manuscript.

Corresponding author:

Matthieu Schmidt, MD, PhD

Service de Médecine Intensive–Réanimation

Hôpital Pitié–Salpêtrière, Assistance Publique–Hôpitaux de Paris

Sorbonne Université, INSERM, UMRS_1166-ICAN, Institute of Cardiometabolism and Nutrition

47, boulevard de l’Hôpital, F-75013 Paris, France

Phone: +33 142 162 937 ; e-mail: matthieu.schmidt@aphp.fr

This article is open access and distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives License 4.0 (<http://creativecommons.org/licenses/by-nc-nd/4.0/>). For commercial usage and reprints please contact Diane Gern (dgern@thoracic.org).

Dear Editor,

Early in the COVID-19 pandemic, the description of a cytokine storm associated with the most severe forms of the disease elicited consideration of anti-cytokine therapies. However, more recent data showed that inflammatory cytokine concentrations in patients with critical COVID-19 are markedly less than those reported in sepsis or acute respiratory distress syndrome (ARDS) unrelated to COVID-19 (1). In the most severe forms of COVID-19 which required extracorporeal membrane oxygenation (ECMO), concerns were raised about the potential harm of ECMO itself (2, 3) that may increase serum and bronchoalveolar lavage fluid interleukine-6 (IL-6) and tumor necrosis factor- α (TNF- α) concentrations (4) while significantly higher mortality had been associated with higher serum IL-6 in COVID-19 patients (5). Besides, TNF- α and IL-8 concentrations rose rapidly during the first 2 hours of ECMO in a neonatal porcine ECMO model (6). Although estimated 60-day survival of ECMO-rescued patients with COVID-19 was similar to that of studies published in the past 2 years on ECMO for severe ARDS (7), the contribution of ECMO itself to the cytokine release syndrome observed in some COVID patients remains a matter of debate.

CytoSorb® (CytoSorbents Europe, Berlin, Germany) is a haemoadsorption cartridge containing polystyrene divinylbenzene beads coated with polyvinylpyrrolidone designed to remove cytokines from the blood. Small case series have suggested that cytokine removal with this device that can be simply connected to the ECMO circuit may improve the outcomes of severe ARDS or cardiac surgery patients (8).

In this context, we studied 22 COVID-19 ECMO patients. The first consecutive 11 patients were included prospectively with a CytoSorb® adsorber being combined to ECMO. During that period, 4 ECMO patients were not included because a pre-ECMO sample was not possible. Then, 11 non-contemporaneous ECMO patients without CytoSorb® (i.e control

group) were subsequently included. All of these patients were included without any selection based on clinical criteria, elevated levels of IL-6, or other biomarkers of inflammation. Whole blood was collected before ECMO, 4 hours after ECMO start without CytoSorb® adsorber, and after 12 and 48 hours of CytoSorb® and ECMO or after 12 and 48 hours on ECMO alone in the control group, respectively. We used highly sensitive, classical or digital, multiplex enzyme-linked immunosorbent assay technologies, to directly analyze combined cytokine production profiles (IL-1 β , IL-6, IL-8, IL-22, IL-10, IL-17A, IL-18, GM-CSF, IFN- α , IFN- γ , TNF- α , and IFN- β) in serum of patients at these different time points. Serum samples were also obtained at the same time points in 11 control patients (i.e COVID-19 patients on ECMO who did not receive CytoSorb®). We compared cytokine profiles in patients with ECMO and CytoSorb® (CytoSorb group) or ECMO alone (control group), between 1) before and after four hours on ECMO (i.e direct impact of the ECMO) and 2) four and 48 hours on ECMO. Comparisons were performed by using a Wilcoxon test for non-parametric variables (SPSS version 21.0, IBM Corp, 2012). The study was performed at the Pitie Salpetrière hospital and approved by the local ethical committee, Comité d’Ethique de la Recherche of Sorbonne University (#CER-SU-2020-21 and -31).

Twenty-two COVID-19 ECMO patients (11 with ECMO and CytoSorb®) were included: 16 males, median (range) age 49 (33-65), SAPS II 46 (17-92), and time between intubation and ECMO start of 3 (1-11) days. The 48-hour study period was completed for 8/11 patients with Cytosorb® and all control group patients, respectively. Pre-ECMO, only IL-1 β was significantly lower in the Cytosorb group compared to the control group ($p=0.022$). Levels of IL-6, IL-8, and IL-18 were very high in all patients, and higher than serum concentrations of other cytokines tested. Importantly, serum cytokine levels, and specifically IL-6, did not increase after 4 hours on ECMO (Figure 1A and 1B). Furthermore, IL-10 and IFN- γ concentrations decreased after 48 hours of CytoSorb® treatment ($p=0.008$

and $p=0.02$, respectively). IL-6 levels also decreased from baseline levels, although the difference did not reach statistical significance ($p=0.08$) (Figure 1B). Other cytokine-levels were not altered by haemoabsorption. However, IL-6, IL-8, and IL-10 concentrations also significantly decreased in the 48 hours following ECMO initiation in the 11 patients who did not receive CytoSorb® (Figure 1A). The eight 60-day survivors in the Cytosorb® group were on ECMO for 25 (6-56) days whereas the 7 survivors of the control group spent 20 (1-42) days on ECMO.

Our findings suggest that ECMO itself does not exacerbate cytokine release in COVID-19 patients, contrary to what was previously suggested (2). Our results also question the actual impact of the CytoSorb® treatment to decrease serum concentrations of IL-10, IFN- γ , and IL-6 in this context. Indeed, a prompt switch to “ultra-protective” mechanical ventilation aiming to markedly reduce tidal volume and the driving pressure (9, 10) or the spontaneous evolution of the disease may explain the significant decrease of IL-6, IL-8, and IL-10 also observed in controls in the 48 hours post-ECMO initiation (Figure 1A). However, cytokine adsorption was associated with a more pronounced decrease of serum IL-6 in a recent series of four COVID-19 patients on ECMO (11). While these preliminary results need confirmation in larger cohorts, it should be mentioned that the non-selective reduction of cytokines with the CytoSorb® absorber could lead to paradoxical effects. Indeed, decreasing cytokine levels of IL-10, which is considered to dampen inflammation (12), may exacerbate COVID-19 associated organ damage. On the other hand, the reduction of IFN- γ , the main driver of macrophage activation, could contribute to control haemophagocytic lymphohistocytosis-like features associated with organ damage in some severe cases of COVID-19 pneumonia.

Our study has some limitations. First, the two groups were not randomly selected and the control group was non contemporaneous. Second, timing of initiation of Cytosorb® was heterogeneous. We cannot rule out that the cytokine profile/response with the Cytosorb® could have been different if we had selected patients with higher level of inflammation before ECMO or if we had protocolized the timing of the Cytosorb® initiation. Third, the number of patients included is small, which limits the interpretation and generalization of our results. The ongoing randomized, multicenter controlled trial evaluating cytokine adsorption in COVID-19 patients on ECMO (NCT04385771) may help to clarify whether this strategy improves the outcomes of these severe patients.

In conclusion, ECMO does not exacerbate cytokine release in COVID-19 patients whereas IL-6, IL-8, and IL-10 decrease after 48 hours on ECMO with ultra-protective mechanical ventilation. To what extent combining a Cytosorb® adsorber with ECMO could enhance the decrease of these cytokines and improve outcomes, warrants further investigations.

Disclosures:

Dr. Combes reports grants and personal fees from Getinge, personal fees from Baxter, personal fees from Xenios, outside the submitted work. Dr Schmidt reports lectures fees from Getinge, Drager, and Xenios, outside the submitted work. Other authors have no conflicts of interest to disclose.

Acknowledgments:

The authors wish to thank the doctors and nurses from the ICU department of Pitié-Salpêtrière hospital who made this study possible, all members from the Cimi-COVID-19 consortium, and particularly Christophe Parizot that took care of sample processing, clinical

data mining and biobanking for this part of the Cimi COVID-19 effort focused on ECMO patients.

References

1. Leisman DE, Ronner L, Pinotti R, Taylor MD, Sinha P, Calfee CS, Hirayama AV, Mastroiani F, Turtle CJ, Harhay MO, Legrand M, Deutschman CS. Cytokine elevation in severe and critical COVID-19: a rapid systematic review, meta-analysis, and comparison with other inflammatory syndromes. *Lancet Respir Med* 2020;8:1233–1244.
2. Henry BM. COVID-19, ECMO, and lymphopenia: a word of caution. *Lancet Respir Med* 2020;8:e24.
3. Chen Q, Yu W, Shi J, Shen J, Hu Y, Gao T, Zhang J, Xi F, Gong J, Li J, Li N. The effect of venovenous extra-corporeal membrane oxygenation (ECMO) therapy on immune inflammatory response of cerebral tissues in porcine model. *J Cardiothorac Surg* 2013;8:186.
4. Shi J, Chen Q, Yu W, Shen J, Gong J, He C, Hu Y, Zhang J, Gao T, Xi F, Li J. Continuous renal replacement therapy reduces the systemic and pulmonary inflammation induced by venovenous extracorporeal membrane oxygenation in a porcine model. *Artif Organs* 2014;38:215–223.
5. Risnes I, Wagner K, Ueland T, Mollnes T, Aukrust P, Svennevig J. Interleukin-6 may predict survival in extracorporeal membrane oxygenation treatment. *Perfusion* 2008;23:173–178.
6. McILwain RB, Timpa JG, Kurundkar AR, Holt DW, Kelly DR, Hartman YE, Neel ML, Karnatak RK, Schelonka RL, Anantharamaiah GM, Killingsworth CR, Maheshwari A. Plasma concentrations of inflammatory cytokines rise rapidly during ECMO-related SIRS due to the release of preformed stores in the intestine. *Lab Invest* 2010;90:128–139.

7. Schmidt M, Hajage D, Lebreton G, Monsel A, Voiriot G, Levy D, Baron E, Beurton A, Chommeloux J, Meng P, Nemlaghi S, Bay P, Leprince P, Demoule A, Guidet B, Constantin JM, Fartoukh M, Dres M, Combes A, Groupe de Recherche Clinique en REanimation et Soins intensifs du Patient en Insuffisance Respiratoire aiguE (GRC-RESPIRE) Sorbonne Université, Paris-Sorbonne ECMO-COVID investigators. Extracorporeal membrane oxygenation for severe acute respiratory distress syndrome associated with COVID-19: a retrospective cohort study. *Lancet Respir Med* 2020.
8. Kogelmann K, Scheller M, Drüner M, Jarczak D. Use of hemoadsorption in sepsis-associated ECMO-dependent severe ARDS: A case series. *J Intensive Care Soc* 2020;21:183–190.
9. Schmidt M, Pham T, Arcadipane A, Agerstrand C, Ohshimo S, Pellegrino V, Vuylsteke A, Guervilly C, McGuinness S, Pierard S, Breeding J, Stewart C, Ching SSW, Camuso JM, Stephens RS, King B, Herr D, Schultz MJ, Neuville M, Zogheib E, Mira J-P, Rozé H, Pierrot M, Tobin A, Hodgson C, Chevret S, Brodie D, Combes A. Mechanical Ventilation Management during Extracorporeal Membrane Oxygenation for Acute Respiratory Distress Syndrome. An International Multicenter Prospective Cohort. *Am J Respir Crit Care Med* 2019;200:1002–1012.
10. Rozenchwajg S, Guihot A, Franchineau G, Lescroat M, Bréchet N, Hékimian G, Lebreton G, Autran B, Luyt C-E, Combes A, Schmidt M. Ultra-Protective Ventilation Reduces Biotrauma in Patients on Venovenous Extracorporeal Membrane Oxygenation for Severe Acute Respiratory Distress Syndrome. *Crit Care Med* 2019;47:1505–1512.
11. Rieder M, Wengenmayer T, Staudacher D, Duerschmied D, Supady A. Cytokine adsorption in patients with severe COVID-19 pneumonia requiring extracorporeal membrane oxygenation. *Crit Care* 2020;24:435.

12. Jankovic D, Kullberg MC, Feng CG, Goldszmid RS, Collazo CM, Wilson M, Wynn TA, Kamanaka M, Flavell RA, Sher A. Conventional T-bet(+)Foxp3(-) Th1 cells are the major source of host-protective regulatory IL-10 during intracellular protozoan infection. *J Exp Med* 2007;204:273–283.

Figure 1: Cytokine production profiles in COVID-19 patients A) before, after 4, 12, and 48 hours on ECMO alone (n=11), and B) before, after 4 hours on ECMO, 12, and 48 hours with the CytoSorb® adsorber combined with ECMO (n=11).

Symbols represent individual subjects; bars show the median. Statistical analyses were conducted using the Wilcoxon signed-rank test.

Comparisons were performed between before ECMO and after 4 hours on ECMO as well as between after ECMO and either 48 hours on ECMO (control group) or 48 hours on Cytosorb® (Cytosorb group)




Figure 1: Cytokine production profiles in COVID-19 patients A) before, after 4, 12, and 48 hours on ECMO alone (n=11), and B) before, after 4 hours on ECMO, 12, and 48 hours with the CytoSorb® adsorber combined with ECMO (n=11).

190x254mm (96 x 96 DPI)