



**HAL**  
open science

# PREDATION BY COD (GADUS MORHUA) ON AMPHIPOD CRUSTACEANS IN THE NORTHWESTERN ATLANTIC

D H Steele, G R Lilly

► **To cite this version:**

D H Steele, G R Lilly. PREDATION BY COD (GADUS MORHUA) ON AMPHIPOD CRUSTACEANS IN THE NORTHWESTERN ATLANTIC. *Vie et Milieu / Life & Environment*, 1999, pp.309-316. hal-03180876

**HAL Id: hal-03180876**

**<https://hal.sorbonne-universite.fr/hal-03180876>**

Submitted on 25 Mar 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## PREDATION BY COD (*GADUS MORHUA*) ON AMPHIPOD CRUSTACEANS IN THE NORTHWESTERN ATLANTIC

D.H. STEELE\*, G.R. LILLY\*\*

\* Memorial University of Newfoundland, St. John's, Newfoundland, Canada A1B 3X9

\*\* Department of Fisheries and Oceans, P.O. Box 5667, St. John's, Newfoundland, Canada, A1C 5X1

COD  
PREDATION  
AMPHIPODS  
NORTHWESTERN ATLANTIC

**ABSTRACT.** – Cod of all sizes caught offshore in the northwest Atlantic fed on amphipods. Eighty two species were identified as prey. Many were collected only at restricted depths and locations. Cod on the Southeast Shoal of the Grand Bank preyed on shallow water species that have a relict distribution. The sizes of amphipods consumed increased rapidly up to 36 cm long cod, but there was little further increase to the size of amphipods consumed by larger cod. The preponderance of mature males amphipods and the occurrence of sympagic (under ice) species in the cod indicates that they forage extensively off the bottom and may at times rise to near the surface under the pack ice.

MORUE  
AMPHIPODES  
PREDATION  
ATLANTIQUE NORD-OUEST

**RÉSUMÉ.** – Les Morues de toutes tailles recueillies au large, dans l'Atlantique du nord-ouest, se nourrissent d'Amphipodes. Quarante vingt deux espèces ont été identifiées comme proies. De nombreuses espèces ne peuvent être récoltées qu'à des profondeurs et des emplacements limités. Les Morues du haut-fond sud-est du Grand Banc capturent les espèces d'eau peu profonde qui ont une distribution relictuelle. La taille des Amphipodes consommés augmente rapidement jusqu'à ce que les Morues atteignent 36 cm, puis elle n'augmente que modérément pour les Morues de taille plus élevées. La prépondérance des mâles d'Amphipodes mûrs et l'occurrence d'espèces sympagiques (sous la glace) parmi les Morues indique qu'elles capturent bien leurs proies au fond et qu'il peut leur arriver de chasser aussi presque jusqu'à la surface.

### INTRODUCTION

The cod stock complex known as 'Northern Cod' found in Northwest Atlantic Fisheries Organization (NAFO) Subareas 2J and 3KL and the Grand Bank cod, Subareas 3N and 3O were the largest stocks in the northwestern Atlantic. A collapse of these stocks resulted in a fishing moratorium being declared in 1992, but since then cod have not recovered to any significant degree in the offshore waters. The present study deals with fish caught offshore (Fig. 1) before the moratorium was invoked.

Adult offshore cod spawn primarily but not exclusively along the margin of the continental slope from Labrador to the northern margin of the Grand Bank, from February to May, depending upon the latitude. The eggs and larvae drift south with the Labrador Current. As juveniles, cod associate with pelagic cnidarians (Sars 1879, J.M. Green pers obs) and by August, they appear inshore (< 25 m) where they are attracted to algal beds and other types of cover (Keats *et al* 1986). However, even at this time they continue to feed

primarily on pelagic copepods (Keats *et al* 1986). There is a gradual shift to benthic prey such as harpacticoid copepods, amphipods, annelids and molluscs (Keats & Steele 1992). As they age (one and two years old) and grow in size, they occur in deeper water (Gregory *et al* 1996), but migrate inshore at night (Keats 1990). With increasing size they become increasingly piscivorous. For 'Northern Cod' of 40-70 cm length, capelin and sand lance become the dominant foods (Lilly & Fleming 1981) because of their abundance and relatively small size. Thus 'Northern Cod' tend to become predominantly piscivorous at a younger age and smaller size than do cod in more southern areas where herring, a larger fish, is the dominant prey (Powles 1958).

Although predominantly piscivorous, even large cod (> 70 cm) continue to feed on crustaceans (Powles 1958, Lilly 1991) if they are abundant or when fish are scarce (Templeman 1965; see however Fahrig *et al.* (1993) for evidence of lack of switching). Cod are opportunistic predators and can always find some kind of food, but their behaviour is complex and variable.




Fig. 1. – Gammaridean (left) and Hyperidean (right) sample locations.

Following offshore spawning, spent 'Northern Cod' migrate inshore (Templeman 1965, Rose 1993) and in early summer feed intensely on the spawning shoals of capelin. At the time of their inshore migration they were the basis of the historic Newfoundland inshore fishery.

The inshore-offshore migrations of 'Northern Cod' became the subject of considerable speculation after the offshore spawning aggregations were discovered. They formed the basis of a massive offshore fishery in the 1960s and inshore catches declined. However, it was uncertain whether there was a causal relationship, or whether the cod were simply failing to come inshore. After spawning in the relatively warm water off the continental slope, the fish have to cross an intermediate layer of cold water (CIL) before they reach the warm coastal water near shore where the spawning capelin shoals are found in early summer. The CIL could therefore form a potential thermal barrier to the inshore migrations.

The interactions between 'Northern Cod' and capelin have been considered in detail because of the presumed influence on the concentrations and movements of 'Northern Cod' (Lilly 1984). Little or no attention has been given to other types of prey. The present study considers the predatory activities of cod on amphipod crustaceans and what this can indicate about cod behaviour. It also provides some information on the occurrences of

amphipods from an area for which there is relatively little published information.

## MATERIALS AND METHODS

297 of the cod caught on research cruises made between 1979 and 1992 contained amphipods. Most were obtained in the spring (March-May) (Table I). They were caught at depths between 40 and 450 m, between southern Labrador and the tail of the Grand Bank (Fig. 1), with peaks at 60-100, 180-220 and 300-400 m (Fig. 2 top). Cod with amphipods ranged in length from 20 to 126 cm, but most were 30 to 70 cm in length (Fig. 2 middle).

Since the size of prey consumed is constrained primarily by the mouth size of the fish, fish length has been converted to mouth size, estimated as the product of the length of the mandible x the distance between the right and left mandibles. The relationship between mouth size and body length is allometric (Fig. 2 bottom) so that large fish have proportionately larger mouths than do small.

Table I. – Seasonal distribution of samples.

Month	Dec.-Feb.	Mar.-May	June-Aug.	Sept.-Nov.
Number	16	100	23	46




Fig. 2. – Top, depth distribution of cod samples. Middle, length frequency distribution of containing amphipods. Bottom, length-mouth area relationship of cod.

RESULTS

82 species of amphipods (76 gammaridean, 2 caprellideans, 4 hyperiideans) were identified in the stomachs. A few species were found in cod at all depths but most occurred in depth-defined patterns (Table II). Representative distributions are shown in Fig. 3, 4, 5.

The Southeast Shoal of the Grand Bank has shallow depths and variable habitats. Thus it

Table II. – Distributional patterns of Gammaridean and Caprellidean amphipods in cod stomachs.

Shallow (Southeast Shoal)	
Fine Sediments	Rocks and Algae
<i>Anonyx sarsi</i>	<i>Pontogeneia inermis</i>
<i>Psammonyx terranova</i>	<i>Metopa alderi</i>
<i>Hippomedon serratus</i>	<i>Ischyrocerus anguipes</i>
<i>Monoculodes edwardsi</i>	<i>Corophium bonelli</i>
<i>Monoculodes tuberculatus</i>	<i>Paramphithoe hystrix</i>
<i>Monoculodes intermedius</i>	<i>Pardaliscia cuspidatus</i>
<i>Unciola irrorata</i>	<i>Tiron acanthurus</i>
<i>Amphiporeia lawrenciana</i>	<i>Caprella septentrionalis</i>
<i>Acanthoastorius spinosus</i>	<i>Aeginina longicornis</i>
Sympagic (Ice associated)	
<i>Pseudalibrotus nanseni</i>	<i>Pseudalibrotus glacialis</i>
<i>Gammarus wilkitzkii</i>	<i>Gammaracanthus loricatus</i>
Grand Bank (except SE Shoal)	
<i>Anonyx nugax</i>	<i>Onisimus edwardsi</i>
<i>Anonyx laticoxae</i>	<i>Haploops tubicola</i>
<i>Anonyx lilljeborgi</i>	<i>Arrhis phylloxy</i>
<i>Eusirus cuspidatus</i>	<i>Paroediceros lynceus</i>
<i>Rozinante fragilis</i>	<i>Melita dentata</i>
<i>Unciola leucopis</i>	<i>Ischyrocerus commensalis</i>
Cold Intermediate Layer	
<i>Anonyx makarovi</i>	
Continental Slope	
<i>Anonyx nugax</i>	<i>Orchomene serratus</i>
<i>Anonyx ochoticus</i>	<i>Uristes umbonatus</i>
<i>Anonyx compactus</i>	<i>Aristias tumidus</i>
<i>Tmetonyx 'cicada'</i>	<i>Stegocephalus inflatus</i>
<i>Haploops setosa</i>	<i>Haliragoides inermis</i>
<i>Stenopleustes latipes</i>	<i>Eurysteus melanops</i>
<i>Eusirus holmi</i>	<i>Lilljeborgia fissicornis</i>
<i>Maera loveni</i>	<i>Protomeдея stephensi</i>
<i>Goesia depressa</i>	<i>Neohela monstrosa</i>
<i>Ischyrocerus megacheir</i>	<i>Ischyrocerus latipes</i>
All Depths (except SE Shoal)	
<i>Tmetonyx albidus</i>	<i>Ampelisca macrocephala</i>
<i>Ampelisca eschrichti</i>	<i>Byblis gaimardi</i>
<i>Syrrhoe crenulatus</i>	<i>Rhachotropis aculeata</i>
<i>Erichthonius hunteri</i>	

supports a diverse assemblage of shallow water species that either burrow into sandy sediments or that live on rocks and algae. Most of these shallow water species such as *Anonyx sarsi* (Fig. 3 bottom) also occur at shallow depths around Newfoundland (Frost 1936, Fenwick & Steele 1983, Kennedy 1985) but *Acanthoastorius spinosus*, and *Hippomedon serratus* which are endemic to the northwestern Atlantic were not known previously east of coastal Nova Scotia. Their presence is probably due to the fact that the Shoal is shallow enough (< 50 m) that warm surface water reaches the bottom in the summer. Similar offshore shallow habitats with warm water in the summer occur on Banquereau Bank and at Sable Island.




Fig. 3. - Top right, distribution of *Ampelisca* spp. and *Byblis gaimardi* top left distribution of *Haploops setosa*. Bottom right, distribution of *Anonyx* spp. Bottom left, distribution of *Tmetonyx* spp.

At the other extreme, a number of species (e.g. *Tmetonyx* spp, Fig. 3 bottom) were obtained only in relatively deep water off the edges of the banks (Table II), where the water is also relatively warm (slope water). *Anonyx nugax* was found here and

on the top of the Grand Bank, but in the CIL on the northeast side of the bank it was replaced by *Anonyx makarovi*. This depth defined *Anonyx* distribution is similar to what is found in the Gulf of St. Lawrence (pers obs).




Fig. 4. – Left, distribution of *Eusirus cuspidatus*. Right, distribution of *Rhachotropis aculeata*.

The ampeliscid *Haploops setosa* (Fig. 3 top) was also found only in cod caught off the northeast slope of the bank. This is probably related to the substratum in this region since it is not restrained in its depth distribution. *Ampelisca* spp and *Byblis gaimardi* were found at all depths (Fig. 3 top).

The benthopelagic predaceous *Rhachotropis aculeata* and *Eusirus cuspidatus* were also found to be widely distributed, except on the Southeast Shoal (Fig. 4).

Pelagic hyperiidean amphipods occurred in many stomachs and in large numbers (Fig. 5). Hyperiideans occurred almost exclusively in cod caught on the slopes of the banks and consisted mainly of large adults.

Four species of pelagic gammaridean amphipods (*Gammaracanthus loricatus*, *Gammarus wilkitzkii*, *Pseudalibrotus glacialis*, *Pseudalibrotus nansenii*) were also found in cod stomachs (Fig. 5 left). These are Arctic sympagic species, normally associated with the undersurface of multiyear sea ice.

#### Prey Selection

##### Size of Amphipods

There is a rapid increase in the diameter of the amphipods consumed up to a mouth size of 1,000 mm<sup>2</sup> (Fig. 6, 7). The maximum diameter of the

amphipod prey increases rapidly with an increase in mouth size, but the minimum size increases much more slowly. Large cod (up to 150 cm) continue to feed on amphipods although there is little further increase in the sizes of the amphipod prey since larger species do not exist in the cod habitat. The dominant prey of the larger cod are decapod crustaceans (shrimp and crabs) and fish.

A mouth size of 1,000 mm<sup>2</sup> corresponds to a cod body length of 36 cm and is approximately the size at which they cod begin to feed intensively on capelin (100 mm) and sand lance (150 mm) (Lilly & Fleming 1981). These fish prey have corresponding diameters of 13-33 and 13-22 mm respectively (Fig. 6 top). The sizes of the fish prey are not correlated with the sizes of the cod (Lilly & Fleming *ibid*).

#### Sex and maturity of the amphipods

The males of some benthic species become pelagic at maturity and search for females in the water column, at times as far up as the surface (pers obs). When they achieve maturity they develop a distinctive morphology (Steele 1995). Such sexually mature males with their distinctive morphology were more abundant in cod stomachs than the immatures which tend to remain close to the bottom. While this bias could be due to size selection for amphipods of larger size, the fact that the mature specimens were mostly males

Table III. - Reproductive stage of Gammaridean amphipods.

Species	Total	Imm.+ Adult	%	Imm. Female	%	Mat. Female	%	Imm. Male	%	Mat. Male	%
Benthic Searching Males											
<i>Anonyx nugax, sarsi,</i> <i>makarovi, lilljeborgi</i>	200	152	58.0	17	11.2	1	0.7	14	9.2	120	79.0
<i>Ampelisca eschrichti,</i> <i>macrocephala</i> <i>Byblis gaimardi</i> <i>Haploops setosa</i>	243	217	89.0	79	36.4	43	19.8	15	6.9	80	36.9
Benthopelagic Species											
<i>Rhachotropis</i> <i>aculeata</i> <i>Eusirus cuspidatus</i>	55	40	82.0	13	28.9	15	33.3	6	13.3	11	24.4
Holobenthic Species											
<i>Melita dentata</i> <i>Maera loveni</i> <i>Unciola irrorata,</i> <i>leucopis</i> <i>Erichthonius</i> <i>hunteri</i>	45	40	89.0	5	12.5	10	22.2	15	37.5	10	22.2

(Table III) indicates that the latter were more vulnerable to cod predation. Similar numbers of mature males and females of the benthopelagic and holobenthic species were found in the stomachs (Table II).

DISCUSSION

Although traditionally classed as a groundfish and caught commercially on the bottom when in excess of 45 mm in length, it is apparent from




Fig. 5. - From left to right, distribution of sympagic species, of *Parathemisto libellula*, and of *Hyperia galba* and *Parathemisto* spp.




Fig. 6. - Top, diameter/body length relationships of capelin and sand lance. Asterisks indicate the body lengths at which cod begin to feed intensively on these fish. Below, size frequencies (diameter) of amphipod prey in cod of varying mouth size categories.

their predation on capelin and sand lance (Lilly & Fleming 1981) as well as on the pelagic stages of bottom dwelling gammaridean and pelagic hyperiidean amphipods, that cod forage extensively above the bottom and probably undergo vertical feeding migrations above the bottom, as discussed by Brunel (1965). Their predation on sympagic species indicates the CIL is not a barrier to their movements. This behaviour has also been extensively documented by De Blois & Rose (1995) when cod were feeding intensively on benthopelagic shrimp.

The present study demonstrates that information on cod behaviour can be obtained from analysis of the sizes and reproductive states of the amphipod prey, as well as the species composition. The presence of adult hyperiid and the pelagic stages of bottom dwelling gammarideans provides further evidence in support of pelagic foraging.

Although planktonic hyperiidean amphipods are common in the region (Frost 1936, Bousfield 1951, Strong 1981, Buchanan & Browne 1981) and are an important food source of sea birds and marine mammals, their distribution has not been described in any detail, and the species and their




Fig. 7. - Mean diameter of amphipod prey related to cod mouth size.

reproductive stages have not usually been differentiated. The cod in the present study fed preferentially on the larger adults and those cod with many hyperiideans were obtained off the edges of the banks (Fig. 5). This is consistent with the observations of Strong (1981) who found adult hyperiideans only beyond the 200 m depth contour.

Many of the samples of 'Northern Cod' were obtained in deep water and when the surface of the ocean was covered by pack ice (Table I, Fig. 1, 2). It is therefore of interest that some cod contained sympagic gammaridean species that live on the under surfaces of multiyear pack ice. They include *Gammarus wilkitzkii* which Gulliksen & Lonne (1989) consider to be autochthonous sympagic and not normally benthic or pelagic. Moreover Gulliksen & Lonne (1989) also consider the hyperiidean *Parathemisto libellula* to be associated with sea ice at times and consider it to be allochthonous sympagic. The occurrence of these ice associated gammarideans and hyperiideans suggests therefore that some cod forage, at times, almost up to the surface under the pack ice. To do so, they must tolerate cold water. Thus the

Cold Intermediate Layer cannot be a significant impediment to their later inshore migration.

The occurrence of a number of southern species of amphipods in cod foraging on the shallow Southeast Shoal of the Grand Bank, some of which are not known in the shallow coastal waters of Newfoundland suggests that these species have a relict distribution and have persisted on the Grand Bank since these shallow water sites were emergent (see also Steele 1983, Carscadden *et al.* 1989, Hutcheson & Stewart 1994). The Southeast Shoal of the Grand Bank has a number of unique features and is important for the Grand Bank ecosystem (see Whitehead & Glass (1985) for its significance to cetaceans). On the deeper, colder portions of the Bank, the amphipods are typical of northern habitats, as is the benthic fauna generally (Hutcheson *et al.* 1981, Steele 1983).

#### LITERATURE CITED

- Bousfield EL 1951. Pelagic amphipods of the Belle Isle Strait region. *J Fish Res Bd Canada* 8(3) : 134-162.
- Brunel P 1965. Food as a factor or indicator of vertical migrations of cod in the western Gulf of St. Lawrence. International Commission of the Northwest Atlantic. Spec Pub 6 : 439-448.
- Buchanan RA, Browne SM 1981. Zooplankton of the Labrador coast and shelf during summer, 1979. LGL Limited Report for Petro-Canada Explorations Ltd., Calgary, Alberta : 1-78.
- Carscadden JE, Frank KT, Miller DS 1989. Capelin (*Mallotus villosus*) spawning on the Southeast Shoal : influence of physical factors past and present. *Can J Fish Aquat Sci* 46 : 1743-1754.
- DeBlois EM, Rose GA 1995. Effect of foraging activity on the shoal structure of cod (*Gadus morhua*). *Can J Fish Aquat Sci* 52 : 2377-2387.
- Fahrig L, Lilly GR, Miller DS 1993. Predator stomachs as sampling-tools for prey distribution : Atlantic cod (*Gadus morhua*) and capelin (*Mallotus villosus*). *Can J Fish Aquat Sci* 50 : 1541-1547.
- Fenwick GD, Steele DH 1983. Amphipods of Placentia Bay, Newfoundland. Memorial University of Newfoundland, Occas. Papers in Biol. No. 7 : 1-22.
- Frost N 1936. Amphipoda from Newfoundland waters, with a description of a new species. Nfld. Dept. Nat. Res., Div Fish Res Rept : Faunistic Series 1 : 4-9.
- Gregory RS, Andersen JT, Dalley EL 1996. Use of habitat information in conducting assessments of juvenile cod abundance. Northwest Atlant. Fish. Org., Sci Coun Res Doc 96/23 : 1-14.
- Gulliksen B, Lonne OJ 1989. Distribution, abundance and ecological importance of marine sympagic fauna in the Arctic. *Rapp P.-v. Réun Cons int Explor Mer* 188 : 133-138.
- Hutcheson MS, Stewart PL, Spry JM 1981. The biology of benthic communities of the Grand Banks of Newfoundland (including the Hibernia area). Grand Banks Oceanographic Studies. Final Report by McLaren Plansearch for Mobil Oil Canada, Ltd. 3 : 1-99.
- Hutcheson MS, Stewart PL 1994. A possible relict population of *Mesodesma deauratum* (Turton : Bivalvia (Mesodesmatidae) from the southeast Shoal, Grand Bank of Newfoundland. *Can J Fish Aquat Sci* 51 : 1162-1168.
- Keats DW 1990. A nocturnal inshore movement of juvenile cod *Gadus morhua* L. in eastern Newfoundland. *J Exp Mar Biol Ecol* 139 : 167-173.
- Keats DW, Steele DH, South GR 1986. The role of fleshy macroalgae in the ecology of juvenile cod (*Gadus morhua* L.), in inshore waters off eastern Newfoundland. *Can J Zool* 65 : 49-53.
- Keats DW, Steele DH 1992. Diurnal feeding of juvenile cod (*Gadus morhua*) which migrate into shallow water at night in eastern Newfoundland. *J Northw Atl Fish Sci* 13 : 7-14.
- Kennedy VS 1985. A summer benthic survey in Conception Bay, Newfoundland, emphasizing zoogeography of annelids and amphipods. *Can J Zool* 63 : 1863-1869.
- Lilly GR 1984. Influence of the Labrador Current on predation by Cod on Capelin and Sand Lance off eastern Newfoundland. Northw. Atl. Fish. Org. Sci. Coun.Stud. 3 : 77-82.
- Lilly GR 1991. Interannual variability in predation by cod (*Gadus morhua*) on capelin (*Mallotus villosus*) and other prey off southern Labrador and north-eastern Newfoundland. Cons. int. Explor. Mer Sci. Symp. 193 : 133-146.
- Lilly GR, Fleming AM 1981. Size relationships in predation by Atlantic Cod, *Gadus morhua*, on Capelin, *Mallotus villosus*, and Sand Lance, *Ammodytes dubius*, in the Newfoundland area. Northw. Atl. Fish. Org. Sci. Coun. Stud. 1 : 41-45.
- Powles PM 1958. Studies of reproduction and feeding of Atlantic cod (*Gadus callarias* L. in the southwestern Gulf of St. Lawrence. *J Fish Res Bd Canada* 15(6) : 1383-1402.
- Rose GA 1903. Cod spawning on a migration highway in the northwest Atlantic. *Nature* 366 : 458-461.
- Sars GO 1879. Report of practical and scientific investigations of the cod fisheries near the Loffoden Islands made during the years 1864-1869. U.S. Comm. Fish and Fisheries Part 5, Rept. Commiss. for 1877 : 565-611. 612-681.
- Steele DH 1983. Marine ecology and zoogeography. In GR South ed. Biogeography and ecology of the island of Newfoundland. *Mon Biol* 48 : 421-465.
- Steele DH 1995. Sexual dimorphism in mature gammaridean amphipods. *Pol Arch Hydrobiol* 42(4) : 303-317.
- Strong KW 1981. Seasonal occurrence and distribution of zooplankton in waters over the Grand Banks of Newfoundland. Grand Banks Oceanogr. Stud. Final Report by McLaren Plansearch for Mobil Oil Canada Ltd. 1 : 1-132.
- Templeman W 1965. Some instances of cod and haddock behaviour and concentrations in the Newfoundland and Labrador area in relation to food. Inter. Comm. Northw. Atl. Fish. Spec Pub 6 : 49-62.
- Whitehead H, Glass C 1985. The significance of the Southeast Shoal of the Grand Bank to humpback whales and other cetacean species. *Can J Zool* 63 : 2617-2625.

Reçu le 12 mai 1999; received May 12, 1999

Accepté le 5 octobre 1999; accepted October 5, 1999