



**HAL**  
open science

# THE MEIOBENTHOS OF THE MOLLOY DEEP (5 600 M), FRAM STRAIT, ARCTIC OCEAN

T Soltwedel, M Miljutina, V Mokievsky, D Thistle, K Vopel

► **To cite this version:**

T Soltwedel, M Miljutina, V Mokievsky, D Thistle, K Vopel. THE MEIOBENTHOS OF THE MOLLOY DEEP (5 600 M), FRAM STRAIT, ARCTIC OCEAN. *Vie et Milieu / Life & Environment*, 2003, pp.1-13. hal-03205078

**HAL Id: hal-03205078**

<https://hal.sorbonne-universite.fr/hal-03205078v1>

Submitted on 22 Apr 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## THE MEIOBENTHOS OF THE MOLLOY DEEP (5 600 M), FRAM STRAIT, ARCTIC OCEAN

T. SOLTWEDEL<sup>1\*</sup>, M. MILJUTINA<sup>2</sup>, V. MOKIEVSKY<sup>3</sup>, D. THISTLE<sup>4</sup>, K. VOPEL<sup>4</sup>

<sup>1</sup>Alfred-Wegener-Institute for Polar and Marine Research,  
Columbusstraße, 27568 Bremerhaven, Germany

<sup>2</sup>Moscow State University, Faculty of Biology, Vorobjovy Gory, Moscow 119899, Russia

<sup>3</sup>P.P. Shirshov Institute of Oceanology, Russian Academy of Sciences,  
36, Nakhimov Prospect, Moscow 117218, Russia

<sup>4</sup>Florida State University, Department of Oceanography, West Call Street, Tallahassee,  
FL 32306-4320, USA

\* Corresponding author: [tsoltwedel@awi-bremerhaven.de](mailto:tsoltwedel@awi-bremerhaven.de).

MEIOBENTHOS  
NEMATODES  
DEEP-SEA  
MOLLOY DEEP  
ABYSSAL PLAINS  
TRENCHES

**ABSTRACT.** – The meiobenthos (including foraminiferans) of the Molloy Deep (Fram Strait, Arctic Ocean) was studied along a 15 km transect crossing the deep in NW-SE direction. Four stations between 5 416 and 5 569 m water depth were sampled during summer months between 1997 and 2001. In comparison with other abyssal and hadal regions of the World Ocean, meiofauna abundances were extremely high, ranging from 2 153 to 2 968 ind./10 cm<sup>2</sup> (values for the uppermost 5 cm of the sediments). The analysis of biogenic sediment compounds (e.g. chloroplastic pigments, particulate proteins) confirmed comparably high amounts of organic matter in the sediments, presumably favouring increased faunal densities and biomasses. Subsurface peaks in meiobenthic abundances at 1-2 cm sediment depth may be due to substantial disturbance and/or predation by dense herds of small holothurians (*Elpidia glacialis*), obviously inhabiting the entire Molloy Deep in very high numbers. Faunal composition of the meiobenthic community of the Molloy Deep was similar to other deep-sea regions. Foraminiferans were the dominant taxa of the total meiobenthos (48.5-59.9%), whereas nematodes dominated the metazoan meiofauna (91.7-95.8%). The total meiofauna of the Molloy Deep consisted of relatively small organisms compared to other/shallower oceanic regions, which could not be explained by reduced food availability to the benthos.

MÉIOBENTHOS  
NÉMATODES  
MER PROFONDE  
DÉPRESSION DE MOLLEY  
PLAINES ABYSSALES  
FOSSÉS

**RÉSUMÉ.** – Le méiobenthos (incluant les Foraminifères) de la dépression de Molloy (Déroit de Fram, Océan Arctique) a été étudié le long d'un transect de 15 km qui recoupe cette dépression selon la direction NO-SE. Quatre stations situées entre 5 416 et 5 569 m de profondeur ont été échantillonnées pendant les mois d'été entre 1997 et 2001. En comparaison avec d'autres zones abyssales et hadales de l'Océan mondial, les abondances de la méiofaune, qui s'échelonnent de 2 153 à 2 968 ind./10 cm<sup>2</sup> (dans les 5 cm supérieurs des sédiments), sont extrêmement élevées. L'analyse des composants biogènes des sédiments (pigments des chloroplastes, protéines particulières) confirment les taux relativement élevés de matière organique des sédiments, qui pourraient favoriser une augmentation des densités et des biomasses élevées. Les pics d'abondance de la méiofaune proche de la surface, à 1-2 cm de profondeur dans le sédiment, pourraient être dus à une perturbation notable et/ou à la prédation de denses concentrations de petites Holothuries (*Elpidia glacialis*), qui habitent toute la dépression de Molloy en très grand nombre. La composition de la communauté méiofaunique de la dépression de Molloy est similaire de celle des autres régions de mer profonde. Les Foraminifères constituent le taxon dominant du méiobenthos total (48.5-59.9 %), tandis que les Nématodes dominent la fraction des Métazoaires (91.7-95.8 %). La méiofaune totale de Molloy Deep comprend des organismes de taille relativement petite par rapport aux autres zones océaniques moins profondes, ce qui ne peut s'expliquer par des ressources nutritives réduites disponibles pour le benthos.

## INTRODUCTION

The presence and persistence of life at the ocean floor can be seen as a response to organic matter input from the upper water levels. Abundance of benthic organisms generally tends to decrease with ocean depth from continental margins to the hadal zone (Rowe 1983, Thiel 1983). Decreasing food supply with increasing water depth and distance from land masses appear to be one of the primary causes for reduced faunal abundance in the deep oceanic basins and on the abyssal plains. Some deep-sea areas, however, support unusually high densities of benthic organisms. These high densities have been related to high surface productivity (e.g. upwelling areas) with subsequently enhanced input of organic matter to the seafloor (Thiel 1982, Soltwedel 1997, 2000), or to the accumulation of organic material in low-energy bottoms (Thiel 1971, 1983).

Hadal regions like trenches are generally located in the vicinity of the continental margins. Such settings adjacent to land masses enhance a natural dumping of terrestrial and coastal, inorganic and organic material. Trenches as well as fracture zones are also characterised by steep walls, which may promote a rapid down slope transport of any kind of material. In this way, enhanced availability of organic matter might promote rich communities of benthic organisms, even at hadal depths.

In the past decades, knowledge of the standing stocks of deep-sea meiobenthos has increased considerably. Due to logistical reasons and limitations in ship time, information on the meiobenthos from abyssal depths deeper than 5 000 m, and especially from hadal regions of the World Ocean is still rather limited. To date about 30 ultra-deep sites have been visited to investigate meiobenthic communities. Most of them are situated in subtropical and tropical regions; except for the study by Vinogradova *et al.* (1993) there is, to our knowledge, no information on ultra-deep meiobenthic communities from higher latitudes.

This paper describes the meiobenthic communities (with special attention on the free-living Nematoda) in the Molloy Deep, the deepest region of the Arctic Ocean (max. depth 5 607 m). On the basis of the Molloy Deep data set, and in comparison with published meiofauna data from other ultra-deep parts of the World Ocean, we address the following questions: Does meiofauna abundance generally correlate with indicators of organic-matter input? Does variation in organic matter availability correlate with changes in the taxonomic composition of the meiofauna? Does the size structure correlate with organic matter input?

## MATERIALS AND METHODS

*Area of investigation:* The seafloor of Fram Strait is part of a complex transform zone between the Knipovich Ridge of the Greenland Sea and the Gakkel Ridge of the Arctic Ocean. The Molloy Deep is a nodal basin formed at the intersection of the Molloy Fracture Zone and the Molloy Ridge (Fig. 1). Especially to the east and to the south, the deep has steep flanks with up to 30° inclination between 3 000 and 5 000 m water depth. The floor of the Molloy Deep as defined by the 5 200 m isobath, measures 12 km (SW-NE) by 22 km (NW-SE). According to Thiede *et al.* (1990), the deep has a maximum water depth of 5 607 m at 79°08.5'N and 2°47.0'E.

In Fram Strait, Atlantic waters flow northward in the West Spitzbergen Current. To the west, surface waters of polar origin flow southwards along the western Greenland margin. The bathymetry of Fram Strait plays a guiding role in water mass modifications (Quadfasel *et al.* 1987). The "Molloy Deep Eddy", a cyclonic gyre of 80-100 km in diameter, is located persistently just north of the Molloy Deep at 79°30'N and 3°00'E. It is induced either by the deep itself (Wadhams & Squire 1983, Smith *et al.* 1984) or by the Molloy Ridge just north of the deep (Quadfasel *et al.* 1987). Based on a modelling study, Schlichtholz & Houssais (1999) confirmed a cyclonic circulation of Eurasian Basin Deep Water in the Molloy Deep region. These specific hydrographic circumstances may favour the sedimentation of particulate matter in the centre of the gyre (see Discussion).

The overall circulation pattern in Fram Strait results in variable sea-ice cover with permanent ice-covered areas in the west, permanently ice-free areas in the south-east, and seasonally varying ice conditions in the central and eastern parts of the area (Vinje 1977, 1985). For most of the year, the Molloy Deep is located in the marginal ice zone, an area characterised by enhanced primary productivity (Sakshaug & Skjoldal 1989).

Photo and video observations (in 1997 with a towed camera system, "Ocean Floor Observation System", in 1999 with the help of the French deep-diving Remotely Operated Vehicle VICTOR 6000, and in 2001 with a video-guided Multiple Corer) gave an impression of the epibenthos, inhabiting the deepest parts of the Molloy Deep (Ritter & Soltwedel, unpubl. data). The epibenthic community is characterised by three species: a holothurian (*Elpidia glacialis*), a scavenging amphipod (most probably *Uristes* sp.), and the sea anemone *Bathypheilia margaritacea* (Riemann-Zürneck, pers. comm). The holothurian *E. glacialis* is by far the dominating epibenthic organism. Abundances of the 1-2 cm tall holothurian reached 20-30 ind/m<sup>2</sup>. Similar numbers at Sites 1997, 1999 and 2001 let us infer that comparable densities occur all over the deepest part of the Molloy Deep. The abundance of the scavenging amphipod was 2-3 ind/m<sup>2</sup>. *B. margaritacea* was found, only where hard substrate (clay aggregations?, wooden pieces, drop stones, human debris) occurred on the visually homogeneous silty sediments.

*Sampling and sample processing:* Bottom samples were taken between 1997 and 2001 at 4 sites along a 15 km transect that crossed the Molloy Deep in NW-SE direction (Fig. 1). Water depths ranged from 5 416 to 5 569 m (Table 1). Figure 2 shows the depth profile of the Molloy Deep floor along the transect, positions of




Fig. 1. – Sampling sites between 1997 and 2001 along a NW-SE transect crossing the Molloy Deep, Fram Strait.

the sampling sites, and distances between these sites. Inclinations between outer sampling sites and the bottom of the deep are about 3 degree.

Sediments were sampled with a Multiple Corer (MUC) (Barnett *et al.* 1984) in 1997, 2000 and 2001, and with the French Remotely Operated Vehicle (ROV) VICTOR 6000 in 1999 (Table I). Due to limitations in ship time only one MUC haul could be performed per sampling site. Subsamples for meiobenthic investigations were taken from the core samples by inserting small plastic disposal syringes of 2-cm inner diameter and 5-cm length with cut-off anterior ends. To minimise effects of small-scale variability in meiofauna dispersion at the different sampling sites, we analysed three “pseudo” replicates (Hurlbert 1984) from different MUC cores, and 4 “real” replicates from different push-cores sampled with the ROV, respectively.

Subsamples for meiofauna investigations were split into five 1-cm-thick layers and preserved separately in 4% buffered Formalin. In the laboratory, each sediment sample was washed through a set of sieves with different mesh sizes (500, 250, 125, 63 and 32  $\mu\text{m}$ ), stained with Rose Bengal, and sorted under a low-power stereo microscope. Organisms were identified to major taxa, i.e. Foraminifera, Nematoda, Polychaeta, Harpacticoida/

Nauplii. Taxa that occurred in minor quantities (e.g. Kinorhyncha, Isopoda, Gastrotricha, Bivalvia, Gastropoda) were pooled in *one* category, classified as the ‘Others’.

Special attention was paid to the nematodes because they generally dominate the benthic metazoan meiofauna. To estimate nematode biomasses, 676 to 988 specimens covering all sieve size fractions and sediment layers were chosen from each of the four sampling sites and transferred to slides for further investigations. Nematode lengths (excluding the filiform tail) and widths were measured by a semi-automated image-analysis system (analySIS<sup>®</sup> 2.1). Nematode volumes and biomasses (wet weights) were calculated with equations given by Andrassy (1956). Nematoda wet weights were converted to carbon biomass by assuming that 100% wet weight corresponds to 12.4% carbon weight (Jensen 1984).

Subsamples (three “pseudo” replicates from each MUC; Hurlbert 1984) for abiotic and biotic sediment parameters (water content, sediment-bound pigments, as well as two biomass parameters, i.e. phospholipids and particulate proteins) were also taken with modified plastic syringes (1.2 or 2 cm in diameter, and 5 cm in length), and split in five 1-cm-thick layers to investigate vertical gradients in the sediment. Due to limitations in

ship time it was not possible to investigate environmental parameters for the 1999 samples.

Sediment water contents were estimated by weight loss of wet sediment samples dried at 60°C. Concentrations of chloroplastic pigments (chlorophyll *a* and its degradation products) were studied to estimate the amount of plant organic matter reaching the seafloor. Chloroplastic pigments were extracted in 90% acetone and measured with a TURNER fluorometer according to Yentsch & Menzel (1963) and Holm-Hansen *et al.* (1965). This method does not discriminate among individual breakdown products, but distinguishes intact chlorophyll *a* from the bulk of pheopigments. Because the pigment composition in sediments is not completely understood, the term chloroplastic pigment equivalents (CPE) for the bulk of pigments registered with this method is used. Differences in the “total microbial biomass” (TMB), comprising all organisms from bacteria to meiofauna (i.e. fungi, yeasts, flagellates, ciliates, protozoans and small metazoans) within the sediment samples, were assessed with biochemical procedures. Phospholipids (PL), a measure of the amount of cell-wall material, were determined by the method of Findlay *et al.* (1989) with slight modifications as described in Boetius & Lochte (1994). PL concentrations were used

to calculate the total microbial biomass (TMB) in terms of organic carbon by applying a conversion factor of  $100 \mu\text{mol P (g C)}^{-1}$  (Findlay & Dobbs 1993). Particulate proteins (PP) were determined to estimate the bulk of “living” plus “dead biomass”, i.e. organisms and the proportion of detrital organic matter in the sediments. PP, operationally defined as  $\gamma$ -globulin-equivalents, were measured following instructions given by Greiser & Faubel (1988).

*Statistical analyses:* To test variances between stations we performed the Mann-Whitney U-test. SPEARMAN rank-correlation analyses were used to identify relationships between single parameters. All statistical analyses were performed using the computer program STATISTICA®.

## RESULTS

### *Environmental parameters*

Data on abiotic and biotic parameters for the uppermost centimetre of the sediment, and cumulative values for the upper 5 cm are summarised in Table I. Water contents, indicating the porosity of sediments, ranged between 54.6 and 63.1% (mean values for 0-5 cm). The highest values were found in the central part of the Molloy Deep (Site 2000). No significant differences among sampling sites were found for cumulative values of the bulk of sediment-bound pigments (CPE) for the upper 5 cm of the sediments ( $18.4\text{--}21.4 \mu\text{g}/5 \text{ cm}^3$ ), but, in the south-easterly part of the deep (Site 1997), CPE concentrations in the uppermost centimetre of the sediments were significantly higher ( $p = 0.024$ ) than those at the other sampling sites. The relative proportion of comparably “fresh” phytodetrital matter at Site 1997 was an order of magnitude greater (15.2% Chl. *a*) than at all other sampling sites, indicating a significant input of organic mat-




Fig. 2. – Bathymetry of the Molloy Deep along the NW-SE transect.

Table I. – Station data and environmental information. Values for abiotic and biotic parameters are for the uppermost centimetre with cumulative values for 0-5 cm in parentheses. % H<sub>2</sub>O = water content of the sediments (\*means for 0-5 cm. Chl.a = chlorophyll *a*. Pheo. = pheopigments. CPE = chloroplastic pigment equivalents (Chl.a + Pheo.). % Chl.a = relative proportion of chlorophyll *a*. FDA = bacterial exoenzymatic activity. PL = phospholipids. PP = particulate proteins.

| Station Data |              |              |               |           |      | Porosity        | Organic Matter Availability |                                     |                                     |                                   | Activity     | Biomass        |                              |
|--------------|--------------|--------------|---------------|-----------|------|-----------------|-----------------------------|-------------------------------------|-------------------------------------|-----------------------------------|--------------|----------------|------------------------------|
| Site [year]  | Date [dd/mm] | Latitude [N] | Longitude [E] | Depth [m] | Gear |                 | H <sub>2</sub> O [%]        | Chl.a [ $\mu\text{g}/\text{cm}^3$ ] | Pheo. [ $\mu\text{g}/\text{cm}^3$ ] | CPE [ $\mu\text{g}/\text{cm}^3$ ] |              | % Chl.a [%]    | FDA [nmol/cm <sup>3</sup> h] |
| 1997         | 04.08.       | 79°06.1'     | 3°02.6'       | 5445      | MUC  | 66.7<br>(58.0)* | 1.4<br>(2.5)                | 7.6<br>(17.1)                       | 9.0<br>(19.6)                       | 15.2<br>(12.8)                    | 1.4<br>(1.8) | 10.6<br>(25.7) | 0.6<br>(2.9)                 |
| 1999         | 03.07.       | 79°08.6'     | 2°49.5'       | 5552      | ROV  | ---             | ---                         | ---                                 | ---                                 | ---                               | ---          | ---            | ---                          |
| 2000         | 16.08.       | 79°08.2'     | 2°53.6'       | 5569      | MUC  | 73.4<br>(63.1)* | 0.1<br>(0.3)                | 6.5<br>(21.1)                       | 6.6<br>(21.4)                       | 1.6<br>(1.3)                      | 1.3<br>(2.1) | 18.1<br>(62.9) | 0.2<br>(1.6)                 |
| 2001         | 13.07.       | 79°12.0'     | 2°34.5'       | 5416      | MUC  | 53.3<br>(54.6)* | 0.1<br>(0.2)                | 6.0<br>(18.2)                       | 6.1<br>(18.4)                       | 1.2<br>(0.9)                      | 1.2<br>(3.0) | 14.1<br>(36.8) | 0.3<br>(1.7)                 |

ter from phytoplankton production shortly prior to sampling.

Bacterial exoenzymatic activities (FDA) in the uppermost centimetre of the sediments slightly (non-significantly) decreased in SE-NW, whereas FDA values for 0-5 cm showed an opposite trend. As for the water contents (see above), phospholipid concentrations were greatest in central parts and lower in peripheral areas of the deep. Concentrations of sediment-bound particulate proteins (PP) mirrored the distribution patterns of intact Chl. *a* in the sediments with significantly higher values ( $p = 0.024$ ) in the SE part of the Molloy Deep, suggesting that enhanced PP concentrations also indicated comparably "fresh" phytodetrital matter at Site 1997.

### *Meiofaunal abundances and biomasses*

Total meiofauna densities (including foraminiferans) at the different sampling sites ranged from  $2\,153 \pm 483$  ind./10 cm<sup>2</sup> to  $2\,968 \pm 613$  ind./10 cm<sup>2</sup> (Table II); the highest numbers were found at Site 1999, and the lowest values appeared at Site 2000. There was no trend in meiofauna densities along the transect: both the highest and lowest values were found at central sampling sites of the Molloy Deep (distance between stations only about 1.5 km). Peripheral sites in the NW and SE had intermediate abundances ( $2\,436 \pm 322$  ind./10 cm<sup>2</sup> and  $2\,636 \pm 420$  ind./10 cm<sup>2</sup>, respectively).

Except for Site 1999, metazoan abundances within the sediment columns had distinct sub-surface peaks at 1-2 cm sediment depth inhabiting almost one third of the total metazoan assemblages (number of individuals in 0-5 cm = 100%), whereas foraminiferans generally declined with increasing sediment depth (Fig. 3). That rather high meiofauna densities found in 4-5 cm sediment depth ( $49-94$  ind./10 cm<sup>2</sup>) suggest that significant numbers of

meiofauna live deeper than 5 cm in the sediment, but, in this study, no attempts were made to quantify these densities.

Free-living nematodes were by far the dominant taxon within the metazoans (91.7-95.8%) and therefore were of special interest. Nematode lengths and widths were measured to calculate nematode biomasses at the different sampling sites and for comparison with estimates of total microbial biomass (TMB) calculated from biochemical analyses (phospholipid measurements, see above). Mean nematode biomass varied from 0.026 to 0.037 µgC (Table III) and showed no significant differences between the four sampling sites. Total nematode biomass per sampling site generally increased from SE to NW sites (Table III). The relative proportion of nematode biomass from the TMB per site ranged between 0.51 and 1.17% (Table III), with lowest values at a central Molloy Deep site.

### *Faunal composition and size spectra*

Meiobenthic assemblages of the Molloy Deep were dominated by foraminiferans (48.5-59.9%) and nematodes (36.7-49.4%). The relative proportion of all other taxa ranged between 2.1 and 3.4% (mean values for polychaetes: 0.1%, harpacticoids: 1.0%, nauplii: 0.9%, others: 1.0%). A comparison of the faunal composition at the different sampling sites showed some clear trends. Relative proportions of nematodes along the transect declined from NW to SE, whereas that of foraminiferans increased in NW-SE direction (Fig. 4). An U-test confirmed significant differences ( $p = 0.049$ ) in the relative abundance of nematodes and foraminiferans between the outer sites of the transect (Sites 1997 and 2001, respectively). The relative proportion of other meiofaunal taxa showed no significant differences between the sampling sites.

Table II. – Meiofauna densities (ind./10 cm<sup>2</sup>) for the uppermost centimetre and cumulative values for 0-5 cm (in parentheses).

| Year/<br>Site | Foraminifera  | Nematoda      | Polychaeta | Nauplii    | Harpacticoida | Others     | Total<br>Metazoans | Total<br>Meiofauna |
|---------------|---------------|---------------|------------|------------|---------------|------------|--------------------|--------------------|
| 1997          | 763<br>(1582) | 238<br>(966)  | 2<br>(2)   | 6<br>(16)  | 6<br>(29)     | 4<br>(41)  | 256<br>(1054)      | 1019<br>(2636)     |
| 1999          | 963<br>(1673) | 427<br>(1187) | 1<br>(1)   | 13<br>(28) | 10<br>(31)    | 11<br>(48) | 462<br>(1295)      | 1421<br>(2968)     |
| 2000          | 424<br>(1218) | 168<br>(875)  | 0<br>(6)   | 2<br>(23)  | 4<br>(23)     | 1<br>(5)   | 175<br>(935)       | 599<br>(2153)      |
| 2001          | 466<br>(1183) | 308<br>(1201) | 0<br>(0)   | 4<br>(26)  | 4<br>(15)     | 4<br>(11)  | 322<br>(1253)      | 788<br>(2436)      |




Fig. 3. – Vertical profiles of foraminifera and nematodes in the uppermost 5 cm of the sediments (mean values from 3-4 subsamples).

Table III. – Relative proportion of nematodes of the total meiobenthos (i.e. including foraminiferans), mean biomass per nematode, total nematode biomass per site, and the proportion of the total microbial biomass estimated from biochemical analyses that is nematode biomass.

| Year | rel. proportion of nematodes from the total meiobenthos [%] | mean biomass of nematodes [ $\mu\text{g C}$ ] | total nematode biomass per site [ $\mu\text{g C} / 10 \text{ cm}^2$ ] | total microbial biomass from biochemical analyses [ $\text{mg C} / 10 \text{ cm}^2$ ] | rel. proportion of nematodes from TMB [%] |
|------|-------------------------------------------------------------|-----------------------------------------------|-----------------------------------------------------------------------|---------------------------------------------------------------------------------------|-------------------------------------------|
| 1997 | 36.65                                                       | 0.029                                         | 29.95                                                                 | 2.57                                                                                  | 1.17                                      |
| 1999 | 39.99                                                       | 0.026                                         | 30.86                                                                 | ---                                                                                   | ---                                       |
| 2000 | 40.64                                                       | 0.037                                         | 32.38                                                                 | 6.29                                                                                  | 0.51                                      |
| 2001 | 49.30                                                       | 0.032                                         | 39.02                                                                 | 3.67                                                                                  | 1.06                                      |

Size spectra for the different taxa from the various sampling site were rather similar. For foraminiferans, 95.3% of individuals appeared in size fractions 32-63  $\mu\text{m}$  and 63-125  $\mu\text{m}$  (42.8% and 52.5%, respectively). Nematodes showed a peak (55.7%) in size fraction 63-125  $\mu\text{m}$ , with rather similar proportions in size fractions 32-63  $\mu\text{m}$  and 125-250  $\mu\text{m}$  (24.5% and 17.5%, respectively); only about 2.5% of nematodes were found in sieve sizes > 250  $\mu\text{m}$ . Approximately 90% of all other meiofaunal taxa appeared in size fractions < 250  $\mu\text{m}$  (i.e. 32-63  $\mu\text{m}$ : 22.7%; 63-125  $\mu\text{m}$ : 31.8%; 125-250  $\mu\text{m}$ : 36.4%).

## DISCUSSION

### Quality of the data set

Due to limitations in ship time it was impossible to sample the different sites during a single expedition. Results presented in this study derive from samples obtained in four different years (1997-2001) with sampling dates from the beginning of




Fig. 4. – Relative proportion of foraminifera, nematoda and other meiofauna taxa along the NW-SE transect in relation to concentrations of sediment-bound chloroplast pigments (? = no pigment data for Site 1999).

July to the middle of August. Therefore, a comparison of data from the different sampling sites could be affected by seasonal and interannual variability,

for example in phytoplankton production or organic-matter sedimentation. The time lag between the most recent input of organic matter to the seafloor and the sampling date might be of crucial importance. Gooday (1988), Pfannkuche (1993) and Gooday *et al.* (1996) described a rapid and pronounced response of deep-sea foraminiferans to the pulsed input of phytodetrital matter in some areas of the NE Atlantic. Lamshead *et al.* (1995) and Galéron *et al.* (2001) reported that meiobenthic organisms move vertically in the sediment in response to the impoverishment of surface layers and the downward mixing of organic matter by the bioturbation of larger benthic organisms. De Bovée *et al.* (1990) even detected changes in the taxonomic composition of the metazoan meiofauna at deep sampling sites (> 2000 m) following the productive phase in the north-western Mediterranean Sea. Soltwedel *et al.* (1996) found indications that deep-sea nematode assemblages might show growth responses to an episodic food supply. A measurable response in metazoan abundances following the input of particulate organic matter, however, has yet to be observed in oceanic regions (Pfannkuche 1993, Gooday *et al.* 1996, Galéron *et al.* 2001).

Sediment sampling during 1997, 2000, and 2001 was done with a multiple corer (MUC). In 1999, however, we were able to use a Remotely Operated Vehicle (ROV) and push-corers to sample. As deduced from the comparison of box corers and multiple corers, different sampling equipment can influence quantitative estimates of the deep-sea meiobenthos (Bett *et al.* 1994). Compared to the use of any kind of coring gear lowered to the seafloor at a certain speed (thereby producing a bow wave), push-coring with the help of an ROV is probably the best way of getting undisturbed sediment samples from the deep sea. In 1999, the ROV allowed targeted sampling of areas of the seabed that appeared visually to be perfectly undisturbed, i.e. also without any kind of (larger) biogenic structuring. In contrast, MUC hauls delivered a set of ("pseudo") replicates generally including cores with plain sediments and sediment surfaces modified by larger organisms. Therefore, push-coring with a ROV, and the use of a MUC for sediment sampling might lead to different results.

Looking at vertical gradients of meiobenthic organisms we, in fact, found striking differences between ROV and MUC samples. Whereas at Site 1999, metazoan densities gradually decreased with increasing sediment depth, all other (MUC) sampling sites had subsurface peaks for metazoans abundance at 1-2 cm depth. A bow-wave effect of the MUC might have blown away the uppermost millimetres of the sediments (including associated organisms) and therefore could be responsible for reduced densities in the uppermost sediment layer at Sites 1997, 2000, and 2001. Subsurface peaks

were, however, only found for metazoans, whereas foraminiferans generally decreased with increasing sediment depth. Because all abiotic and biotic parameters investigated in this study also did not show consistent subsurface peaks at any of the sampling sites, we suggest that the observed differences in meiofauna gradients are not caused by different sampling methods but come out for other reasons, as discussed in one of the following sections.

### *Meiobenthic assemblages in the Molloy Deep*

Our investigations discovered extremely high meiofaunal densities for all sites visited along the transect across the Molloy Deep (mean: ~ 2 500 ind./10 cm<sup>2</sup>). Meiobenthic abundances showed no correlation with station depth, and no general trend along the transect. Looking at relative proportions of foraminiferans and metazoans at the different sites, however, we found significantly increasing values for foraminiferans in south-easterly direction, and an opposite trend for nematodes (Fig. 4). Differences among sites in organic-matter supply (food source) to the benthos might explain these trends. Although the pigment inventory of the sediments (CPE values for the upper 0-5 cm of the sediments) showed no significant differences between sampling sites, pigment concentrations in the uppermost centimetre of the sediments (and especially the relative proportion of Chl. *a*) increased in south-easterly direction. These differences in food-supply markers could be explained by differences among sites in ice-cover because of generally enhanced primary production in ice-free areas. According to long-term observations, the marginal ice zone is, for most of the year, located in the Molloy Deep area. The probability of ice-free surface water increases towards the south-east.

Higher relative proportions of protozoans in those areas could be explained by their ability to react comparably fast to any kind of perturbation (including an episodic input of potential food source; Gooday & Lamshead 1989); protozoans (as well as bacteria) can multiply in hours and can also activate dormant cells and resting spores. This ability might be an advantage in areas with pulsed organic matter input (e.g. the SE part of the Molloy Deep), compared to almost permanently ice-covered regions with a more regular (and comparably low) food supply to the benthos, as expected for the north-western part of the Molloy Deep.

Except for Site 1999, vertical gradients of nematodes (and also other metazoans) in the sediment generally revealed distinct subsurface peaks in 1-2 cm depth, whereas densities of foraminiferans, in general, continuously decreased with increasing sediment depth. The occurrence of subsurface peaks



can probably be explained by a high predation pressure by dense herds of holothurians (*Elpidia glacialis*), intensively reworking the upper few millimetres of the sediments. Extremely high abundances of *E. glacialis* (20-30 ind./m<sup>2</sup>) were found for Sites 1997 and 1999 (results from photo and video observations; Ritter & Soltwedel, unpubl data), but were also assumed for the two other sites. Highly motile metazoan taxa (e.g. nematodes) are able to escape predation by vertical migration to deeper sediment horizons, whereas most foraminiferans are rather immobile and only passively moved, e.g. by the bioturbation of larger benthic organisms. This fact might result in different vertical distribution patterns within the sediments. No explanation, however, could be given for the lack of a subsurface peak for metazoans at Site 1999.

Looking at size spectra for meiofauna communities of the Molloy Deep, we found no significant differences between the four sites, but, the proportion of individuals in the smallest sieve size fraction (< 63 µm) was relatively high (30-38%). Using the same sizes of sieves (and also including foraminiferans), Schewe & Soltwedel (1999) and Schewe (2001) found similar high proportions (24-40%) in the size class < 63 µm for a number of sampling sites from the central Arctic Ocean (depth range: 864 to 4 187 m). Relative high proportions of small meiofauna individuals in both studies were explained by a tendency towards dwarfism in an area of reduced food availability to the benthos, due to limited primary production in a permanently ice-covered region. As indicated by concentrations of sediment-bound pigments and particulate proteins, food availability in the Molloy Deep should be up to 20-times higher than those at sampling sites in the central Arctic Ocean. Thus, in contrast to suggestions originally given by Thiel (1975), reduced food availability is obviously not the only factor explaining dwarfism in deep-sea meiofauna communities. When comparing meiofauna size spectra with biochemical data indicating food availability (quantity and quality) in Atacama Trench off Chile (7 800 m), Danovaro *et al.* (2002) came to the same conclusion and presented different suggestions to explain the disproportion between meiofauna sizes and food availability. To our mind, pressure-related effects on metabolic processes, and selection of small-sized barophilic or barotolerant organisms are the most probable explanations, but, further investigations are needed to confirm these hypotheses.

Direct biomass measurement were carried out for nematodes exclusively. As expected from sieve size spectra, mean biomasses per nematode were comparably low (Soltwedel *et al.* 2000) and showed no significant differences between sampling sites, however, as the relative proportion of nematodes from the total biomass clearly increased

in north-westerly direction (see above), total nematode biomass per site followed the same trend. The relative proportion of nematode biomass from the total microbial biomass (TMB) estimated from biochemical analyses (concentrations of phospholipids, indicating the total biomass of the bulk of sediment-inhabiting organisms from bacteria to meiofauna) revealed lowest values at a central Molloy Deep site (Table III). Highest TMB values together with intermediate densities of foraminiferans at this site presume a relatively high proportion of nanofauna organisms, i.e. fungi, yeasts, flagellates and ciliates, in the central Molloy Deep.

### ***Meiobenthos in ultra-deep regions of the World Ocean***

For logistical reasons (e.g. insufficient cable lengths, lack of availability of deep-diving instruments, limitations in ship time), information on the meiobenthos from abyssal depths deeper than 5 000 m, and especially from hadal regions of the world's oceans is still rather limited. Some 30 ultra-deep sites have been visited to investigate meiobenthic communities, most of them situated in subtropical and tropical regions (Table IV). Except for investigations by Vinogradova *et al.* (1993) in the Orkney Trench (Antarctica) our study is, to our knowledge, the only work on ultra-deep meiobenthic communities from higher latitudes.

Table IV summarises data on densities and taxonomic composition for ultra-deep meiobenthic assemblages from deep-sea basins and abyssal plains (20 sites) and deep-sea trenches and fracture zones (13 sites). Information on methods (i.e. lower mesh size limits, total sediment depth investigated) were included for better assessment of data comparability. Most investigations focussed on the metazoan meiofauna; only 7 studies included meiobenthic foraminiferans.

Due to lack in standardisation of sampling and sorting techniques, a comparison of meiobenthic standing stocks in the literature is fraught with problems. The use of different sampling devices, subsampling strategies and extraction methods makes a direct comparison of meiofauna data given by different authors quite a challenge. Due to known trends in dwarfism of deep-sea fauna, lower sieve size limits play an especially crucial role for meiobenthos abundance estimates. When comparing results from this study with data from the literature (see below), we tried to take into account these limitations.

A comparison with results from other regions revealed surprisingly high meiofauna densities for the Molloy Deep; mean numbers were about 10-times higher than in other ultra-deep regions of the world ocean. Similarly high numbers were only found by Shirayama & Kojima (1994) for the Japan Trench. These authors investigated the metazoan

Table IV. – Comparison of meiofauna densities from different deep-sea sites > 5 000 m water depth, separated into investigations of abyssal plains and basins, and investigations of trenches and fracture zones. Lower Size Limit refers to the opening of the smallest sieve used. FORAM = foraminifera. NEMA = nematodes. META = metazoan meiofauna. MEIO = total meiofauna.

| Authors                          | Area                 | Region                           | Water Depth [m] | Sediment Depth [cm] | Lower Size Limit [µm] | FORAM Abundances [Ind./10cm <sup>2</sup> ] | NEMA Abundances [Ind./10cm <sup>2</sup> ] | META Abundances [Ind./10cm <sup>2</sup> ] | MEIO Abundances [Ind./10cm <sup>2</sup> ] |
|----------------------------------|----------------------|----------------------------------|-----------------|---------------------|-----------------------|--------------------------------------------|-------------------------------------------|-------------------------------------------|-------------------------------------------|
| <b>Abyssal Plains, Basins:</b>   |                      |                                  |                 |                     |                       |                                            |                                           |                                           |                                           |
| Dinet (1973)                     | SE Atlantic          | Angola Basin                     | 5170            | 5                   | 40                    | -                                          | 294                                       | 313                                       | -                                         |
| Ingole et al. (2000)             | Central Indian Ocean | Central Indian Basin             | 5300 - 5330     | 10                  | 45                    | -                                          | 20 - 24                                   | 35 - 45                                   | -                                         |
| Namenskaya (1991)                | Southern Atlantic    | Namibia Abyssal Plain            | 5230            | 3                   | 90                    | -                                          | 4                                         | 6                                         | -                                         |
| Kaneko et al. (1997)             | Central Pacific      | Central Pacific Basin            | 5300            | 5                   | 30                    | -                                          | 95                                        | 101                                       | -                                         |
| Parulekar et al. (1982)          | NW Indian Ocean      | Arabian Basin                    | 5100            | 10                  | 44                    | -                                          | 9                                         | 12                                        | -                                         |
| Parulekar et al. (1982)          | NW Indian Ocean      | Somali Basin                     | 5400            | 10                  | 44                    | -                                          | 5                                         | 7                                         | -                                         |
| Parulekar et al. (1992)          | Central Indian Ocean | Central Indian Basin             | 5250 - 5750     | ?                   | 63                    | -                                          | 30 - 34                                   | 50 - 55                                   | -                                         |
| Rachor (1975)                    | NE Atlantic          | Tejo Abyssal Plain               | 5111            | 5                   | 50                    | -                                          | 23                                        | 28                                        | -                                         |
| Rachor (1975)                    | NE Atlantic          | Canary Basin                     | 5510            | 6                   | 50                    | -                                          | 76                                        | 103                                       | -                                         |
| Shirayama (1984)                 | Western Pacific      | Eastern Marianan Basin           | 5820            | 10                  | 37                    | 87                                         | 78                                        | 89                                        | 176                                       |
| Shirayama (1984)                 | NW Pacific           | Northwestern Pacific Basin       | 5580 - 5730     | 10                  | 37                    | 39 - 214                                   | 31 - 257                                  | 38 - 317                                  | 77 - 317                                  |
| Sibuet et al. (1989)             | NE Atlantic          | Cape Verde Basin                 | 5190            | ?                   | 40                    | -                                          | -                                         | 87                                        | -                                         |
| Sibuet et al. (1989)             | NE Atlantic          | Central Pacific Basin            | 5800            | 25 - 55             | 42                    | 100                                        | 92                                        | 102                                       | 202                                       |
| Soltwedel & Thiel (1995)         | Eastern Atlantic     | Guinea Basin                     | 5213            | 5                   | 42                    | 112                                        | 193                                       | 219                                       | 331                                       |
| Soltwedel (1993)                 | SE Atlantic          | Angola Basin                     | 5027            | 5                   | 42                    | 156                                        | 93                                        | 115                                       | 271                                       |
| Thiel (1975)                     | NE Atlantic          | Iberian Basin                    | 5500            | 4                   | 40                    | -                                          | -                                         | 210                                       | -                                         |
| Thiel (1975)                     | Western Indian Ocean | Somali Basin                     | 5000            | 4                   | 40                    | -                                          | -                                         | 50                                        | -                                         |
| Thiel (1975)                     | Central Pacific      | Central Pacific Basin            | 5000            | 4                   | 40                    | -                                          | -                                         | 100                                       | -                                         |
| Tieljen et al. (1989)            | NW Atlantic          | Haiteras Abyssal Plain           | 5411            | 15                  | 44                    | -                                          | 85                                        | 114                                       | -                                         |
| Woods & Tieljen (1985)           | Caribbean Sea        | Venezuela Basin                  | 5050            | 6                   | 44                    | -                                          | 44                                        | 49                                        | -                                         |
| <b>Trenches, Fracture Zones:</b> |                      |                                  |                 |                     |                       |                                            |                                           |                                           |                                           |
| Danovaro et al. (2002)           | SE Pacific           | Atacama Trench                   | 7800            | 6                   | 20                    | -                                          | 5072                                      | 6378                                      | -                                         |
| George & Higgins (1979)          | NW Atlantic          | Puerto Rico Trench               | 8570            | 7                   | 62                    | -                                          | 12                                        | 17                                        | -                                         |
| Olu et al. (1997)                | W Atlantic           | Barbados Trench                  | 5000            | ?                   | ?                     | -                                          | 790 - 8336                                | 845 - 8438                                | -                                         |
| Renaud-Mornant & Goubault (1990) | Eastern Pacific      | Clarion-Clipperton Fracture Zone | 5015            | 2.5                 | 40                    | -                                          | 78                                        | 82                                        | -                                         |
| Saidova (1970)                   | NW Pacific           | Kurile-Kamtscharka Trench        | 5000 - 9000     | 1                   | ?                     | 19 - 94                                    | -                                         | -                                         | -                                         |
| Shirayama (1984)                 | NW Pacific           | Ogasawara Trench                 | 8260            | 10                  | 37                    | 132                                        | 389                                       | 430                                       | 562                                       |
| Shirayama & Kojima (1994)        | NW Pacific           | Japan Trench                     | 5370 - 7460     | 3                   | 63                    | -                                          | 357 - 470                                 | 373 - 488                                 | -                                         |
| Sibuet et al. (1989)             | Central Atlantic     | Vema Fracture Zone               | 5100            | ?                   | 40                    | -                                          | -                                         | 130                                       | -                                         |
| Soltwedel (unpubl. data)         | Central Atlantic     | Romanche Fracture Zone           | 7628            | 5                   | 42                    | 43                                         | 152                                       | 155                                       | -                                         |
| Richardson et al. (1995)         | NW Atlantic          | Puerto Rico Trench               | 8380            | 20                  | 44                    | -                                          | 66                                        | 69                                        | -                                         |
| Thiel (1979)                     | Western Pacific      | Philippine Trench                | 9807            | ?                   | 40                    | 125                                        | ?                                         | 30                                        | 155                                       |
| Tieljen et al. (1989)            | NW Atlantic          | Puerto Rico Trench               | 7460 - 8189     | 15                  | 44                    | -                                          | 41 - 93                                   | 44 - 96                                   | -                                         |
| Vinogradova et al. (1993)        | Antarctica           | Orkney Trench                    | 5150 - 6160     | 3 - 4               | 40                    | -                                          | 5 - 6                                     | 7 - 11                                    | -                                         |
| <b>This Study</b>                | Arctic Ocean         | Molloy Deep                      | 5445 - 5569     | 5                   | 32                    | 1183 - 1673                                | 875 - 1201                                | 935 - 1295                                | 2153 - 2968                               |

meiofauna with a lower size limit of 63  $\mu\text{m}$  and found 373-488 ind./10  $\text{cm}^2$ . When adapting our results from the Molloy Deep to the same size range (simply by subtracting the number of meiofauna individuals found in mesh size class 32-63  $\mu\text{m}$ ), we found comparably high densities (205-311 ind./10  $\text{cm}^2$ ). Such an adaptation of data, of course, assumes identical meiofauna size spectra for the different sampling site. Because we do not have information on the entire size spectra from the Japan Trench, these calculations are somewhat speculative.

To our knowledge only one paper (Danovaro *et al.* 2002) reports higher meiofaunal densities than those given in this study. These authors found metazoan meiofauna densities of  $6\,378 \pm 3\,061$  ind./10  $\text{cm}^2$  for samples from the Atacama Trench (7 800 m) in the upwelling sector off Chile, SE Pacific (Table IV). The lower sieve-size limit used in their study was 20  $\mu\text{m}$ , but even when adapting their results to the lower size limit we used (assuming that more meiofauna individuals would be found when further reducing the lower size limit), metazoan meiofauna abundances in the Atacama Trench were still 6-times higher than those of the Molloy Deep and thus by far the highest abundances ever measured for deep-sea

sites. Although Olu *et al.* (1997) presented comparably high metazoan meiofauna densities from a cold-seep area seaward of the Barbados accretionary prism at about 5 000 m water depth (Table IV), these data should not be considered for direct comparison with results from “normal” deep-sea regions.

When plotting meiofauna abundances from the papers listed in Table IV against water depth, we found major differences between stations from basins or abyssal plains, and trenches or fracture zones (Fig. 5). Whereas there is no obvious relationship between metazoan densities and water depth on the plains (Fig. 5A), data from trenches show an exponential decrease of individuals with increasing depth (Fig. 5B). The weak correlation between metazoan meiofauna densities and water depth on abyssal plains could be explained by the varying productivity of the overlying water (and a subsequently varying food supply to the benthos) in the different regions considered in this comparison. The same should hold for trench regions, however, because of the specific topographic settings of trenches (it should make a difference, whether the samples came from the central parts or from the flanks of a trench) variability should probably be even greater. The significant correlation ( $R^2 =$




Fig. 5. – Graphs of metazoan meiofaunal densities against water depth and primary productivity data (World Ocean Atlas 1998) for abyssal plains (a, c) and trenches (b, d) on the basis of faunal data listed in Table IV (data of Olu *et al.* 1997 and Danovaro *et al.* 2002, excluded). The open circles indicate the data from this study.

0.709) between meiofauna numbers and water depth in the data set for trenches (Table IV, neglecting the extremely high values for Atacama trench and the Barbados cold seep area; Fig. 5B) was probably caused by the immense range in water depth (~ 5 000-9 800 m) covered by the different studies.

Benthic standing stocks are known to generally decrease with increasing water depth, and a number of studies suggest that food supply is one of the major controlling factors (e.g. Thiel 1983, Sibuet *et al.* 1989, Vincx *et al.* 1994). Bathymetric settings of the Molloy Deep and the special hydrographic conditions in the area of investigation together with enhanced primary productivity associated with processes in the marginal ice zone favour a strong input of particulate organic matter to the seafloor. The whole system might act as a huge sediment trap accumulating organic matter at the bottom of the Molloy Deep, consequently allowing a rich benthic community with surprisingly high meiofauna numbers but also allowing high standing stocks of megafaunal organisms (e.g. dense herds of small holothurians; see above). Comparably high remineralisation rates (3.74 mg C/m<sup>2</sup> d), obtained during the deployment of a free-falling grab respirometer in summer 2000 in the central Molloy Deep (Soltwedel unpubl data), and steep oxygen gradients in the uppermost sediment layers obtained with a micro-profiling unit in summer 2002 (Sauter pers comm) confirm the picture of an unusually eutrophic region at hadal depths.

Only a very few investigations on meiofauna assemblages from ultra-deep regions included parameters assessing environmental settings, e.g. food availability, grain sizes, oxygen contents of bottom waters. To study relationships between meiofauna densities and food availability at ultra-deep sites, we compared primary-production estimates based on satellite-derived data (World Ocean Atlas 1998) from the different regions with the meiofauna data given in the papers listed in Table IV. Assuming that local primary productivity supports benthic assemblages in the same region (which for current-driven lateral advection might not always be the case; Soltwedel 1997), we found good correlations ( $R^2 = 0.384$  and  $R^2 = 0.897$ , respectively, when neglecting data of Olu *et al.* 1997 and Danovaro *et al.* 2002) between chlorophyll concentrations in the upper water layers and meiofauna numbers at abyssal depths (Fig. 5C), and trenches (Fig. 5D), further demonstrating that food supply plays a major role in determining meiobenthic standing stocks.

## CONCLUSIONS

In comparison with other abyssal and hadal regions of the World Ocean, meiofauna abundances

in the Molloy Deep were extremely high. Analyses of biogenic sediment compounds confirmed comparably high amounts of organic matter in the sediments, presumably favouring increased faunal densities and biomasses. Variations in organic matter availability correlate with changes in the taxonomic composition of the meiofauna. Subsurface peaks in meiobenthic abundances are most probably due to substantial disturbance and/or predation by dense herds of small holothurians (*Elpidia glacialis*), obviously inhabiting the entire Molloy Deep in very high numbers. The total meiofauna of the Molloy Deep consisted of relatively small organisms compared to other/shallower oceanic regions, which could not be explained by reduced food availability to the benthos. Pressure-related effects on metabolic processes, and the selection of small-sized barophilic or barotolerant organisms might explain the occurrence of comparably small meiofauna individuals in the Molloy Deep.

ACKNOWLEDGEMENTS. – We wish to thank the crew of R/V POLARSTERN for their helpful support during the expeditions in summers 1997 and 1999-2001. We gratefully acknowledge the help of E Siebert, S Brückner, F Wolfarth and M Weber for help with the measurements of various biogenic sediment compounds and the meiofauna sortings. We also wish to thank two anonymous reviewers for their valuable comments on the manuscript. This is publication “awi-n10414” of the Alfred-Wegener-Institute for Polar and Marine Research.

## REFERENCES

- Andrassy I 1956. The determination of volume and weight of nematodes. *Acta Zool (Hungarian Acad Sci)* 2 (1-3): 1-15.
- Barnett PRO, Watson J, Connelly D 1984. A multiple corer for taking virtually undisturbed samples from shelf, bathyal and abyssal sediments. *Oceanol Acta* 7: 399-408.
- Bett BJ, Vanreusel A, Vincx M, Soltwedel T, Pfannkuche O, Lamshead PJD, Gooday AJ, Ferrero T, Dinét A 1994. Sampler bias in the quantitative study of deep-sea meiobenthos. *Mar Ecol Prog Ser* 104: 197-203.
- Boetius A, Lochte K 1994. Regulation of microbial enzymatic degradation of OM in deep-sea sediments. *Mar Ecol Prog Ser* 104: 299-307.
- Bovée F de, Guidi LD, Soyer J 1990. Quantitative distribution of deep-sea meiobenthos in the Northwestern Mediterranean (Gulf of Lions). *Cont Shelf Res* 10 (9-11): 1123-1145.
- Danovaro R, Gambi C, Della Groce N 2002. Meiofauna hotspot in the Atacama Trench, eastern South Pacific Ocean. *Deep-Sea Res* 49: 843-857.
- Dinét A 1973. Distribution quantitative du méiobenthos profond dans la région de la dorsale Walvis (Sud-ouest africain). *Mar Biol* 20: 20-26.
- Findlay RH, King GM, Watling L 1989. Efficiency of phospholipid analysis in determining microbial bio-

- mass in sediments. *Appl Environ Microbiol* 55: 2888-2893.
- Findlay RH, Dobbs FC 1993. Quantitative description of microbial communities using lipid analysis. In Handbook of methods in aquatic microbial ecology. Edited by P F Kemp, Lewis Publishers, Boca Raton FL: 271-284.
- Galéron J, Sibuet M, Vanreusel A, Mackenzie K, Gooday AJ, Dinet A, Wolff GA 2001. Temporal patterns among meiofauna and macrofauna taxa related to changes in sediment geochemistry at an abyssal NE Atlantic site. *Prog Oceanogr* 50: 303-324.
- George RY, Higgins RP 1979. Eutrophic hadal benthic community in the Puerto Rico Trench. *Ambio Spec Rep* 6: 51-58.
- Gooday AJ 1988. A response by benthic foraminifera to the deposition of phytodetritus in the deep sea. *Nature* (London) 332: 70-73.
- Gooday AJ, Lamshead PJD 1989. Influence of seasonally deposited phytodetritus on benthic foraminiferal populations in the bathyal northeast Atlantic: the species response. *Mar Ecol Prog Ser* 5: 53-67.
- Gooday AJ, Pfannkuche O, Lamshead PJD 1996. An apparent lack of response by metazoan meiofauna to phytodetritus deposition in the bathyal North-eastern Atlantic. *J Mar Biol Ass UK* 76: 297-310.
- Greiser N, Faubel A 1988. Biotic factors. In Introduction to the study of meiofauna. Edited by RP Higgins & H Thiel, Smithsonian Institution Press, Washington D.C., London: 79-114.
- Holm-Hansen O, Lorenzen CJ, Holmes RW, Strickland JDH 1965. Fluorometric determination of chlorophyll. *J Cons Perm-Int Explor Mer* 30 (1): 3-15.
- Hurlbert SH 1984. Pseudoreplication and the design of ecological field experiments. *Ecol Monogr* 54 (2): 187-211.
- Ingole BS, Ansari ZA, Rathod V, Rodrigues N 2000. Response of meiofauna to immediate benthic disturbance in the Central Indian Ocean Basin. *Mar Geores Geotech* 18: 263-272.
- Jensen P 1984. Measuring carbon content in nematodes. *Helgol Meeresunters* 38: 83-86.
- Kamenskaya OE 1991. Study of abyssal multicell meiobenthos of the South Atlantic (in Russian). *Ecol Moria* 39: 82-85.
- Kaneko T, Maejima Y 1997. The abundance and vertical distribution of abyssal benthic fauna in the Japan deep-sea impact experiment. *Proc Int Offshore Polar Engineering Conf*, Honolulu, USA: 475-480.
- Lamshead PJD, Ferrero TJ, Wolff GA 1995. Comparison of the vertical distribution of nematodes from two contrasting abyssal sites in the northeast Atlantic subject to different seasonal inputs of phytodetritus. *Int Rev ges Hydrobiol* 80: 327-331.
- Olu K, Lance S, Sibuet M, Henry P, Fiala-Médioni A, Dinet A 1997. Cold seep communities as indicators of fluid expulsion patterns through mud volcanoes seaward of the Barbados accretionary prism. *Deep-Sea Res* 44 (5): 811-841.
- Parulekar AH, Harkantra SN, Ansari ZA, Matondkar SGP 1982. Abyssal benthos of the central Indian Ocean. *Deep-Sea Res* 29 (12A): 1531-1537.
- Parulekar AH, Ingole BS, Harkantra SN, Ansari ZA 1992. Deep sea benthos of the Western and Central Indian Ocean. In Oceanography of the Indian Ocean. Edited by BT Densai, Balkema, Rotterdam: 261-267.
- Pfannkuche O 1993. Benthic response to the sedimentation of particulate organic matter at the BIOTRANS station, 47°N, 20°W. *Deep-Sea Res* 40: 135-149.
- Quadfasel D, Gascard JC, Koltermann KP 1987. Large scale oceanography in Fram Strait during MIZEX '84. *J Geophys Res* 92 (C7): 6719-6728.
- Rachor E 1975. Quantitative Untersuchungen über das Meiobenthos der nordostatlantischen Tiefsee. *Meteor Forsch Ergeb, Reihe D*, 21: 1-10.
- Renaud-Mornant J, Gourgault N 1990. Evaluation of abyssal meiobenthos in the eastern central Pacific (Clarion-Clipperton fracture zone). *Prog Oceanogr* 24: 317-329.
- Richardson MD, Briggs KB, Bowles FA, Tietjen JH 1995. A depauperate benthic assemblage from the nutrient-poor sediments of the Puerto Rico Trench. *Deep-Sea Res* 42 (3): 351-364.
- Rowe GT 1983. Biomass and production of the deep-sea macrobenthos. In Deep-sea biology. Edited by GT Rowe, John Wiley & Sons, New York: 97-121.
- Saidova KM 1970. Benthic foraminifers of the Kurile Kamtchatka Trench area. *Trudy akad Nauk SSSR* 86: 144-176.
- Sakshaug E, Skjoldal HR 1989. Life at the ice edge. *Ambio* 18 (1): 60-67.
- Schewe I 2001. Small-sized benthic organisms of the Alpha Ridge, central Arctic Ocean. *Internat Rev Hydrobiol* 86 (3): 317-335.
- Schewe I, Soltwedel T 1999. Deep-sea meiobenthos of the central Arctic Ocean: distribution patterns and size-structure under extreme oligotrophic conditions. *Vie Milieu* 49 (2/3): 79-92.
- Schlichtholz P, Houssais M-N 1999. An inverse modeling study in Fram Strait. Part II: water mass distribution and transports. *Deep-Sea Res II* 46: 1137-1168.
- Shirayama Y 1984. The abundance of deep sea meiobenthos in the Western Pacific in relation to environmental factors. *Oceanol Acta* 7: 113-121.
- Shirayama Y, Kojima S 1994. Abundance of deep-sea meiobenthos off Sanriku, Northeastern Japan. *Japan J Oceanogr* 50: 109-117.
- Sibuet M, Lambert CE, Chesselet R, Laubier L 1989. Density of the major size groups of benthic fauna and trophic input in deep basins of the Atlantic Ocean. *J Mar Res* 47: 851-867.
- Smith DC, Morison J, Johannessen JA, Untersteiner N 1984. Topographic generation of an eddy at the edge of the East Greenland Current. *J Geophys Res* 89: 8205-8208.
- Snider LJ, Burnett BR, Hessler RR 1984. The composition and distribution of meiofauna and nanobiota in a central North Pacific deep-sea area. *Deep-Sea Res* 31: 1225-1249.
- Soltwedel T 1993. Meiobenthos und biogene Sedimentkomponenten im tropischen Ost-Atlantik. Ber Zentrum für Meeres- und Klimaforsch, Reihe E, 6, 192 p.
- Soltwedel T 1997. Meiobenthos distribution pattern in the tropical East Atlantic: indication for a fractionated sedimentation of organic matter to the sea floor? *Mar Biol* 129: 747-756.
- Soltwedel T 2000. Metazoan meiobenthos along continental margins: a review. *Prog Oceanogr* 46: 59-84.

- Soltwedel T, Mokievsky V, Schewe I 2000. Benthic activity and biomass on the Yermak Plateau and in adjacent deep-sea regions northwest of Svålbard. *Deep-Sea Res* 47 (9): 1761-1785.
- Soltwedel T, Thiel H 1995. Biogenic sediment compounds in relation to marine meiofaunal abundances. *Int Revue ges Hydrobiol* 80 (2): 297-311.
- Soltwedel T, Pfannkuche O, Thiel H 1996. The size structure of deep-sea meiobenthos in the North-eastern Atlantic: nematode size spectra in relation to environmental variables. *J Mar Biol Ass UK* 76: 327-344.
- Thiede J, Pfirman S, Schenke HW, Reil W 1990. Bathymetry of Molloy Deep: Fram strait between Svalbard and Greenland. *Mar Geophys Res* 12: 197-214.
- Thiel H 1971. Häufigkeit und Verteilung der Meiofauna im Bereich des Island-Faröer-Rückens. *Ber Dtsch Wiss Komm Meeresforsch* 22: 99-128.
- Thiel H 1975. The size structure of the deep-sea benthos. *Int Revue ges Hydrobiol* 60: 575-606.
- Thiel H 1979. First quantitative data on the deep Red Sea benthos. *Mar Ecol Prog Ser* 1: 347-350.
- Thiel H 1982. Zoobenthos of the CINECA area and other upwelling regions. *Rapp P-V Réun Cons Int Explor Mer* 180: 323-334.
- Thiel H 1983. Meiobenthos and nanobenthos of the deep sea. *In The Sea*. Edited by GT Rowe, Vol. 8, John Wiley & Sons, New York: 167-230.
- Tietjen JH, Deming JW, Rowe GT, Macko S 1989. Meiobenthos of the Hatteras Abyssal Plain and Puerto Rico Trench: abundance, biomass and associations with bacteria and particulate fluxes. *Deep-Sea Res* 36 (10): 1567-1577.
- Vincx M, Bett BJ, Dinet A, Ferrero T, Gooday AJ, Lamshead PJD, Pfannkuche O, Soltwedel T, Vanreusel A 1994. Meiobenthos of the deep Northeast Atlantic. *In Advances in Marine Biology*. Volume 30. Edited by JHS Blaxter AJ Southward, Academic Press: 2-88.
- Vinje TE 1977. Sea ice conditions in the European sector of the marginal seas of the Arctic. 1966-75. *Norsk Polarinst Årbok* 1975: 163-174.
- Vinje TE 1985. Sea ice distribution 1971-80. *Norsk Polarinst Skrifter* 179D.
- Vinogradova NG, Belyaev GM, Gebruk AV, Zhivago AV, Kamenskaya OE, Levitan MA, Romanov VN 1993. The deep-sea bottom fauna in the southern part of the Atlantic Ocean. *In Investigations of the Orkney Trench during the 43rd cruise of R/V "Dmitri Mendeleev": Geomorphology, bottom sediments and bottom fauna (in Russian)*. *Trans PP Shirshov Inst Oceanol* 127: 9-33.
- Wadhams P, Squire VA 1983. An ice-water vortex at the edge of the East Greenland Current. *J Geophys Res* 88: 2770.2780.
- World Ocean Atlas 1998. [Http://www.nodc.noaa.gov/OC5/WOA98F/woaf\\_cd/search.html](http://www.nodc.noaa.gov/OC5/WOA98F/woaf_cd/search.html)
- Woods DR, Tietjen JH 1985. Horizontal and vertical distribution of meiofauna in the Venezuela Basin. *Mar Geol* 68: 233-241.
- Yentsch CS, Menzel DW 1963. A method for the determination of phytoplankton chlorophyll and phaeophytin by fluorescence. *Deep-Sea Res* 10: 221-231.

Reçu le 5 septembre 2002; received September 5, 2002  
Accepté le 8 janvier 2003; accepted January 8, 2003

