



Language barrier as a risk factor for obstetric anal sphincter injury – A case-control study

Stephanie Schrot-Sanyan, Kamila Kolanska, Yousra Haimeur, Valentin Varlas,
Laure Parisot-Liance, Emile Daraï, Marie Bornes

► To cite this version:

Stephanie Schrot-Sanyan, Kamila Kolanska, Yousra Haimeur, Valentin Varlas, Laure Parisot-Liance, et al.. Language barrier as a risk factor for obstetric anal sphincter injury – A case-control study. Journal of Gynecology Obstetrics and Human Reproduction, 2021, 50 (8), pp.102138. 10.1016/j.jogoh.2021.102138 . hal-03212493

HAL Id: hal-03212493

<https://hal.sorbonne-universite.fr/hal-03212493v1>

Submitted on 29 Apr 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Language barrier as a risk factor for obstetric anal sphincter injury – a Case-Control Study

Short title: Obstetric anal sphincter injury

Stephanie Schrot-Sanyan^a, Kamila Kolanska^a, Yousra Haimeur^a, Valentin Varlas^a, Laure Parisot-Liance^a, Emile Daraï^a, Marie Bornes^a

^a Service de Gynécologie Obstétrique et Médecine de la Reproduction, AP-HP Sorbonne

Université site Tenon, 4 rue de la Chine, 75020 Paris, France

Corresponding author :

Kamila Kolanska

Service de Gynécologie Obstétrique et Médecine de la Reproduction

Hôpital Tenon

4 rue de la Chine

75020 Paris

0033 (0) 1 56 01 73 18

email : kamila.kolanska@aphp.fr

ABSTRACT:

Introduction

The incidence of grade 3-4 perineal tears, also known as obstetric anal sphincter injury (OASI), is reported to be between 0.5 and 2.5%. Beyond the medico-economic burden, the consequences of OASI on a woman's emotional, psychological, sexual, and physical wellbeing are considerable. Among the various risk factors of OASI, few data are available about the impact of a language barrier on its incidence.

Material and Methods

We conducted a case-control study to evaluate the effect of language barriers on the risk of OASI comparing 171 women with OASI and 163 matched controls. The matched criteria included ethnicity, age, previous vaginal delivery, delivery mode, prophylactic episiotomy and birthweight. Patients' characteristics were compared and crude ORs and 95% CIs estimated using unadjusted logistic models. Multivariate analysis was performed with recognized potential confounders.

Results

All of the cases had grade 3 tears. Language barrier was a determinant factor of OASI with an OR of 3.32 [1.36-8.90], $p=0.01$. Other risk factors were occipito-posterior delivery, African origin and prolonged labor duration (OR 6.33, 95% CI: 2.04-27.78, $p=0.004$, OR 1.85, 95% CI: 1.08-3.19, $p=0.03$ and OR 1.03, 95% CI: 1.01-1.05, $p=0.004$, respectively).

Conclusion

Our data suggest that language barrier is an independent risk factor of OASI. Physicians and midwives should attempt to identify patients with a language barrier during prenatal visits. Education about simple terms used during delivery could decrease the incidence of this complication.

KEY WORDS: obstetric anal sphincter injury; perineal tears; language barrier; risk factors

1 INTRODUCTION

2 Spontaneous perineal tears during vaginal birth occur in up to 85% of women during
3 exiting of the baby's head [1,2]. Higher rates of perineal tears are reported after first vaginal
4 deliveries especially if the delivery is assisted [3]. Both the anterior and posterior perineum
5 can be involved and the tears are classified according to the depth of the injury: from grade 1
6 (least serious) to grade 4 (most serious) [2,4]. The incidence of grades 3-4, also known as
7 obstetric anal sphincter injury (OASI), is reported to be between 0.5 and 2.5% after a
8 spontaneous delivery and may be higher for instrumental births [5,6]. In France, the incidence
9 of OASI is estimated at 0.8% [7]. The differences in incidence of perineal tears from one
10 country to another are partly due to various definitions and practices [5,6].

11 The most common consequence of perineal tears in the immediate postpartum period
12 is severe perineal pain [8], followed by wound infection and suture breakdown [9]. Pain is
13 still reported by one third of women 2 weeks after delivery and, while it usually resolves
14 within 2 months, up to 7% still experience pain at 3 months [1,9]. A relation between pain and
15 tear grade has been demonstrated [10]. Other long term consequences of perineal injury are
16 dyspareunia [11] and anal incontinence [12] which can affect women in their daily tasks and
17 their sexuality, and has a major psychological impact [13].

18 After OASI, 8% of women experience fecal and 45% flatus incontinence [14]. These
19 complications depend on the experience of the operator and the type of suture [15,16], but not
20 on the extent of the anal sphincter injury [17]. Half of obstetric recto-vaginal fistulae are
21 related to OASI [18]. Finally, in addition to the medico-economic burden, the consequences
22 of perineal tear on women's emotional, psychological, sexual and physical wellbeing,
23 especially when the anal sphincter is injured, must be considered [13,19].

Some risk factors for OASI, such as assisted deliveries, primiparity, birth weight and ethnicity, are well identified [20–22]. Additional risk factors, such as episiotomy, gestational or maternal age, are more debatable [23,24]. In contrast, some protective factors, including perineal massage before delivery and other perineum management techniques during delivery, have been suggested but their impact on perineal tears is debatable [6,25]. Language barriers could negatively impact perineal injuries, since the collaboration between the obstetrician, midwife and the parturient herself is important during the delivery [26] and lack of communication could be a risk factor for grade 3 tears [27]. In contrast to anatomical conditions and obstetrical procedures, few data are available about the impact of a language barrier on the occurrence of perineal tears.

Therefore, the aim of the present retrospective study was to evaluate the impact of language barrier on the risk of OASI in a secondary university maternity unit.

MATERIALS AND METHODS

Data collection and variable definition

This retrospective study was conducted in Tenon University Hospital in Paris. Among 35 912 women aged 18 years or more who underwent vaginal delivery between January 2001 and March 2016, women with grade 3-4 tears were identified using procedure codes from our prospective database. The control group was established on the basis of the antichronological list of deliveries between January 2001 and March 2016. The controls were matched on available known risk factors identified in the literature: ethnicity (Asian/others), age (+/- 3 years), previous vaginal delivery (yes/no), delivery mode (assisted or not), prophylactic episiotomy (yes/no) and birth weight (+/-150 g) in order to include 140 women, that is 1:1 ratio of case and controls. Women with the history of grade 3-4 tears were excluded from the analysis. Further data were extracted from hospital records for each patient: language difficulties, labor duration, ethnicity (defined as maternal birthplace), occipito-posterior delivery, and newborn outcomes.

Perineal tears were classified according to the depth of the injury [4] (grade 1 – vaginal mucosa injury, grade 2 – perineal muscle injury, grade 3 – external anal sphincter rupture (3a: <50% of fibers torn or involved, 3b: ≥50% of fibers torn, 3c: injury of internal sphincter), grade 4 – rectal mucosa injury) [2,4]. As no consensus exists on the definition of language difficulties and barriers, we defined a language barrier as being present if it was impossible to dialogue with the patient without a translator during prenatal visits, and if the woman could not understand important words such as "push" and "breath" or simple anatomical terms or instructions for body positions. This information is systematically reported in the patient's file during the first prenatal visit and is defined as the necessity of translator during prenatal visits and the labor. The incidence of women with a language

barrier in our department is estimated at 5% due to the multiple ethnic origins of our population. Language difficulties are clearly identified during prenatal visits by the midwife, obstetrician and/or anesthesiologist and reported in the hospital records.

According to the birth policy in our department, uncomplicated vaginal deliveries are usually performed by midwives and assisted deliveries by obstetricians. During the study period, epidural anesthesia was performed in 81% of the patients. Mediolateral episiotomy is not systematic for either assisted or non-assisted deliveries. OASI is systematically managed by an obstetrician. The procedures used in this retrospective study were in accordance with the guidelines of the Helsinki Declaration on Human Experimentation and the Good Clinical Practice (CGP) and approved by the IRB (CEROG 020-GYN-1102).

Statistical analysis

The primary objective of the study was to analyze the incidence of OASI in women presenting a language barrier. A sample size of 140 pairs was calculated for a language difficulty prevalence expected to be 5% for controls and 15% for cases with a power (β) of 0.80 and α 0.05 with one control for one case. To provide for potential missing values, we decided to include 200 pairs. In this study, 66 patients (37 controls and 29 cases) lacked primary outcome or matching data and were excluded from the analysis (Figure 1).

First, a descriptive analysis of all the patients was performed. Patients' characteristics were compared using the Chi square or Fisher's exact tests for categorical values, and Wilcoxon's tests for numerical data. Crude ORs and 95% CIs were estimated using unadjusted logistic models. Multivariate analysis was performed with all known potential confounders (match variables and others including ethnicity, labor duration, and occipito-posterior delivery). All non-contributive variables ($p < 0.05$) were stepwise excluded. Bilateral tests

85 were computed and the significance level was set at 0.05; OR and 95%IC were calculated. We
86 used R 3.3.0 to perform statistical analysis.

87

RESULTS

Epidemiological characteristics of the population

The matching criteria of the 334 women included in the study (171 women with OASI and 163 controls without OASI) are summarized in Table 1. All the women in the OASI group had grade 3 perineal laceration. The most commonly reported maternal ethnic groups were African (47.6%), Caucasian (27.5%) and Asian (20.4%), with similar rates between the two groups ($p=0.15$). Previous vaginal delivery was reported for 35.6% of the women without difference between the groups. There was no statistical difference in any of the matching variables (Table 1).

Univariate and multivariate analysis

Univariate analysis (Table 2) identified a significant difference between the groups in the proportion of women with language barrier (12.3% in the OASI group versus 4.9% in the control group, $p=0.03$). Labor induction and gestational age at delivery were not different between the groups ($p=0.22$ and $p=0.96$, respectively). Labor duration was longer and occipito-posterior delivery was more frequent in the women with OASI compared to the controls (17.0 (11.0-26.0) versus 22.0 (13.0-31.0), $p=0.01$ and 1.8% versus 11.1%, $p<0.001$, respectively). These variables affected the rate of OASI in the final multivariate analysis model (OR 1.03, 95%CI: 1.01-1.05, $p=0.004$ and OR 6.33, 95%CI: 2.04-27.78, $p=0.004$, respectively). Concerning major indicators of the newborn status, an Apgar score under 7 at 5 minutes and cord arterial pH were comparable between the groups ($p=0.17$ and $p=0.83$, respectively).

Age, previous vaginal delivery, delivery mode, prophylactic episiotomy and birth weight had no impact on the model construction and were stepwise excluded from the multivariate analysis. Even if the ethnic origin was not different between both groups, it was

retained in the model. Labor duration and occipito-posterior delivery had a major impact on the risk of OASI and were included in the analysis.

The results of the multivariate analysis are reported in Table 3. Occipito-posterior delivery was the most important risk factor of OASI (OR 6.33, 95%CI: 2.04-27.78, $p=0.004$). African origin and prolonged labor duration remained risk factors of OASI (OR 1.85, 95%CI: 1.08-3.19, $p=0.03$ and OR 1.03, 95%CI: 1.01-1.05, $p=0.004$, respectively). Language difficulties also remained significant with an OR of 3.32 [1.36 - 8.90], $p=0.01$.

DISCUSSION

The present study, using both uni- and multivariable analyses, showed that a language barrier was a significant factor of OASI with an OR of 3.32. Our results are in agreement with those of Esscher et al demonstrating that foreign-born women in Sweden were at a higher risk of maternal, fetal and newborn morbidity and mortality during the various steps of pregnancy and delivery than Swedish-born women [28]. This was essentially related to language barriers which increased the time it took to report to the right medical department, to explain their health concerns, and to be correctly understood [28]. During delivery, language barrier negatively impacted perineal injuries, since communication between the midwife and the parturient woman is essential as the fetal head stretches the pelvic floor [26]. In 2007, Dahlen et al. reported that "a lack of communication" mentioned by midwives was a risk factor for grade 3 tears [27]. However, no objective data were available to support this conclusion. In the presence of communication difficulties, the contribution of an interpreter could be precious [29,30]. However, the availability of an interpreter is limited, especially when the delivery occurs at night or during the week-end [29]. Similarly, the contribution of relatives is not always possible, especially if their knowledge of French is insufficient which limits their contribution in a stressful situation. No study has evaluated the impact of motivating patients during pregnancy to learn simple instructions such as "push" and "stop pushing". Beyond the linguistic issues, the increased maternal morbidity and mortality in women with OASI might be associated with a less favorable economic and social status which hampers optimal prenatal and obstetrical management [31]. Moreover, language barrier might reflect other communication barriers connected to health literacy and the vision of the body anatomy, which could represent an interesting research perspective. The differentiated care could play an important role in avoiding this severe obstetrical complication if the language barrier could be identified during the first stages of the pregnancy follow-up.

Multivariate analysis found that an occipito-posterior fetal presentation was a major risk factor of OASI in our study (adjusted OR 6.33 [2.04-27.78], $p=0.004$), even if it was observed in only 6.6% of the births. This fetal head malposition has already been reported as a major risk factor for OASI with an OR between 5.64 and 13.7 [32,33], especially for primiparous women [34].

In our study, African ethnicity was a risk factor of OASI compared to Caucasian women (OR 1.85 [1.08-3.19], $p=0.03$). This observation has already been reported in previous studies and can partly be explained by the perineal status of immigrant women from eastern African countries who have undergone female genital mutilation [21,35]. Women of Asian origin, as well as other ethnical origins, were not at higher risk in our study. Asian origin has been largely studied showing an OR of between 1.8 and 8.9 for OASI [27,36,37], but literature data concerning other origins are scarce [21,38]. Other risk factors of OASI, such as labor characteristics, have also been identified: labor duration emerged as a minor risk factor in our multivariate analysis (OR 1.03, 95%CI: 1.01-1.05, $p=0.004$) although its impact on OASI is debatable [34,39,40].

The matching variables were chosen based on data from the literature. Among major risk factors of OASI, a first vaginal delivery is well recognized with an OR between 2.4 and 7.55 compared to multiparous women [21,36,41]. Assisted delivery has also been well described in large studies and remains a significant risk factor with an OR between 1.9 and 10.2 [21,32,37,42]. The particular risk of each instrument is more debatable: use of a ventouse seems to have a lower OR than forceps use (1.9-2.7 vs 3.9-10.2) but remains a significant risk factor compared to non-assisted delivery [20,23,25]. The risk of sphincter injury related to spatulas, which are often used in France, does not seem to be higher than a forceps delivery and not less than a ventouse delivery [43–45]. Fetal birth weight is a risk factor for grade 3 perineal tear [46,47], especially if the fetal birth weight exceeds 4000g (OR between 1.86 and

3.01) [33,34,38]. Correlated variables such as a head circumference greater than 35 cm [21], shoulder dystocia [37,38,47], gestational age at delivery [21,48] or diabetes [21,34] are also reported as risk factors.

As the advantage of episiotomy in the prevention of OASI is debatable, we decided to match this variable. In the literature prophylactic episiotomy was considered a risk factor in one study [20], a protective factor in two studies [34,49] and without significant effect in others [24,39,47,50]. These contradictory results could be explained by the existence of associated confounding factors [51], such as the length and the angle of the episiotomy [52], with a maximum risk for a vertical cut (median episiotomy) [41,42,53]. A recent meta-analysis performed by Verghese et al showed a low overall protective effect and no effect in the nulliparous group [23]. The authors claim that performing 65 episiotomies could spare one sphincter injury. Some other factors are more debatable, and seem to be minor risk factors, such as maternal age [21,38,39,47], some specific maternal positions during delivery, or giving birth at night (from 3 to 6 a.m.) [41]. Some authors have studied other factors which do not seem to impact the risk of OASI, such as epidural anesthesia [34,39,41], use of oxytocin [39], or pushing duration [32]. On the contrary, antenatal perineal massage or the use of warm compresses during delivery could reduce by half the risk of sphincter laceration [6].

Some limits of the present study need to be underlined. First, the incidence of OASI varies from hospital to hospital. However, during the study period, the incidence of sphincter injury in our department was 0.5% which is in accordance with a study reporting an estimated incidence of around 0.8% in France and 0.1-4% in Europe [7]. Second, despite the case-control study design, the retrospective nature of the current analysis cannot exclude all possible biases. Third, the relatively long study period could be a potential cause of bias: the guidelines on management of low-risk delivery patients changed during this period to reduce the indications of systematic prophylactic episiotomy. However, as previously mentioned, the

protective effect of episiotomy on perineal tears is debatable [24,28,38,39,47,49,50]. Fourth, the absence of patients with grade 4 tears in our population means that we are unable to draw a definitive conclusion as to whether a language barrier is a risk factor of this severe complication which exposes patients to the risk of rectovaginal fistula. However, the low incidence of OASI, estimated at 0.8% in the French population, would require a multicenter study. Fifth, about half of the population of the present study was composed of African women but without clear information on excision and infibulation that could impact on the incidence OASI representing a limit in the interpretation of our results. However, as previously mentioned, no difference in ethnicity was noted between the groups. Finally, it was not possible to exclude additional confounding factors of perineal tears, such as malnutrition, diabetes and obesity that are often associated with low economic conditions and that are well recognized as a source of inequality for access to health care.

CONCLUSION

Despite some limits of the present study, our data support the fact that, in addition to well-recognized factors of high-grade perineal tears, a language barrier is an independent risk factor. Physicians and midwives should try to identify patients who have language difficulties during the prenatal visits. The education of simple terms that are used during delivery and the implication of education care providers could decrease the incidence of this complication.

214 **DECLARATION OF INTEREST:** None

215 **FUNDING:**

216 This research did not receive any specific grant from funding agencies in the public,
217 commercial, or not-for-profit sectors.

218 **ACKNOWLEDGEMENTS:**

219 We thank Felicity Neilson for editing the English of this paper.

220 **BIBLIOGRAPHY**

- 221 [1] McCandlish R, Bowler U, van Asten H, Berridge G, Winter C, Sames L, et al.
222 A randomised controlled trial of care of the perineum during second stage of normal
223 labour. *Br J Obstet Gynaecol* 1998;105:1262–72. [https://doi.org/10.1111/j.1471-](https://doi.org/10.1111/j.1471-0528.1998.tb10004.x)
224 0528.1998.tb10004.x.
- 225 [2] Goh R, Goh D, Ellepola H. Perineal tears - A review. *Aust J Gen Pract*
226 2018;47:35–8. <https://doi.org/10.31128/AFP-09-17-4333>.
- 227 [3] Christianson LM, Bovbjerg VE, McDavitt EC, Hullfish KL. Risk factors for
228 perineal injury during delivery. *Am J Obstet Gynecol* 2003;189:255–60.
229 <https://doi.org/10.1067/mob.2003.547>.
- 230 [4] Frohlich J, Kettle C. Perineal care. *BMJ Clin Evid* 2015;2015.
- 231 [5] Byrd LM, Hobbiss J, Tasker M. Is it possible to predict or prevent third degree
232 tears? *Colorectal Dis Off J Assoc Coloproctology G B Irel* 2005;7:311–8.
233 <https://doi.org/10.1111/j.1463-1318.2005.00801.x>.
- 234 [6] Aasheim V, Nilsen ABV, Reinart LM, Lukasse M. Perineal techniques during
235 the second stage of labour for reducing perineal trauma. *Cochrane Database Syst*
236 *Rev* 2017;6:CD006672. <https://doi.org/10.1002/14651858.CD006672.pub3>.
- 237 [7] Blondel B, Alexander S, Bjarnadóttir RI, Gissler M, Langhoff-Roos J, Novak-
238 Antolič Ž, et al. Variations in rates of severe perineal tears and episiotomies in 20
239 European countries: a study based on routine national data in Euro-Peristat Project.
240 *Acta Obstet Gynecol Scand* 2016;95:746–54. <https://doi.org/10.1111/aogs.12894>.
- 241 [8] Macarthur AJ, Macarthur C. Incidence, severity, and determinants of perineal
242 pain after vaginal delivery: a prospective cohort study. *Am J Obstet Gynecol*
243 2004;191:1199–204. <https://doi.org/10.1016/j.ajog.2004.02.064>.
- 244 [9] Lewicky-Gaupp C, Leader-Cramer A, Johnson LL, Kenton K, Gossett DR.

245 Wound complications after obstetric anal sphincter injuries. *Obstet Gynecol*
246 2015;125:1088–93. <https://doi.org/10.1097/AOG.0000000000000833>.

247 [10] Andrews V, Thakar R, Sultan AH, Jones PW. Evaluation of postpartum
248 perineal pain and dyspareunia--a prospective study. *Eur J Obstet Gynecol Reprod*
249 *Biol* 2008;137:152–6. <https://doi.org/10.1016/j.ejogrb.2007.06.005>.

250 [11] Barrett G, Pendry E, Peacock J, Victor C, Thakar R, Manyonda I. Women's
251 sexual health after childbirth. *BJOG Int J Obstet Gynaecol* 2000;107:186–95.
252 <https://doi.org/10.1111/j.1471-0528.2000.tb11689.x>.

253 [12] Sultan AH, Thakar R. Lower genital tract and anal sphincter trauma. *Best*
254 *Pract Res Clin Obstet Gynaecol* 2002;16:99–115.
255 <https://doi.org/10.1053/beog.2002.0258>.

256 [13] Tucker J, Clifton V, Wilson A. Teetering near the edge; women's experiences
257 of anal incontinence following obstetric anal sphincter injury: an interpretive
258 phenomenological research study. *Aust N Z J Obstet Gynaecol* 2014;54:377–81.
259 <https://doi.org/10.1111/ajo.12230>.

260 [14] Norderval S, Nsubuga D, Bjelke C, Frasurek J, Myklebust I, Vonen B. Anal
261 incontinence after obstetric sphincter tears: incidence in a Norwegian county. *Acta*
262 *Obstet Gynecol Scand* 2004;83:989–94. [https://doi.org/10.1111/j.0001-](https://doi.org/10.1111/j.0001-6349.2004.00647.x)
263 [6349.2004.00647.x](https://doi.org/10.1111/j.0001-6349.2004.00647.x).

264 [15] Fernando RJ, Sultan AH, Kettle C, Thakar R. Methods of repair for obstetric
265 anal sphincter injury. *Cochrane Database Syst Rev* 2013:CD002866.
266 <https://doi.org/10.1002/14651858.CD002866.pub3>.

267 [16] Kettle C, Hills RK, Jones P, Darby L, Gray R, Johanson R. Continuous versus
268 interrupted perineal repair with standard or rapidly absorbed sutures after
269 spontaneous vaginal birth: a randomised controlled trial. *Lancet Lond Engl*

270 2002;359:2217–23. [https://doi.org/10.1016/S0140-6736\(02\)09312-1](https://doi.org/10.1016/S0140-6736(02)09312-1).

271 [17] Luciano L, Bouvier M, Baumstarck K, Vitton V. Is the extent of obstetric anal
 272 sphincter injury correlated with the severity of fecal incontinence in the long term?
 273 Tech Coloproctology 2020;24:49–55. <https://doi.org/10.1007/s10151-019-02128-1>.

274 [18] Trovik J, Thornhill HF, Kiserud T. Incidence of obstetric fistula in Norway: a
 275 population-based prospective cohort study. Acta Obstet Gynecol Scand
 276 2016;95:405–10. <https://doi.org/10.1111/aogs.12845>.

277 [19] Darmody E, Bradshaw C, Atkinson S. Women's experience of obstetric anal
 278 sphincter injury following childbirth: An integrated review. Midwifery 2020;91:102820.
 279 <https://doi.org/10.1016/j.midw.2020.102820>.

280 [20] Dandolu V, Chatwani A, Harmanli O, Floro C, Gaughan JP, Hernandez E. Risk
 281 factors for obstetrical anal sphincter lacerations. Int Urogynecol J Pelvic Floor
 282 Dysfunct 2005;16:304–7. <https://doi.org/10.1007/s00192-005-1297-2>.

283 [21] Baghestan E, Irgens LM, Børdaahl PE, Rasmussen S. Trends in risk factors for
 284 obstetric anal sphincter injuries in Norway. Obstet Gynecol 2010;116:25–34.
 285 <https://doi.org/10.1097/AOG.0b013e3181e2f50b>.

286 [22] Sideris M, McCaughey T, Hanrahan JG, Arroyo-Manzano D, Zamora J, Jha S,
 287 et al. Risk of obstetric anal sphincter injuries (OASIS) and anal incontinence: A meta-
 288 analysis. Eur J Obstet Gynecol Reprod Biol 2020;252:303–12.
 289 <https://doi.org/10.1016/j.ejogrb.2020.06.048>.

290 [23] Verghese TS, Champaneria R, Kapoor DS, Latthe PM. Obstetric anal
 291 sphincter injuries after episiotomy: systematic review and meta-analysis. Int
 292 Urogynecology J 2016;27:1459–67. <https://doi.org/10.1007/s00192-016-2956-1>.

293 [24] Carroli G, Mignini L. Episiotomy for vaginal birth. Cochrane Database Syst
 294 Rev 2009:CD000081. <https://doi.org/10.1002/14651858.CD000081.pub2>.

295 [25] Eason E, Labrecque M, Wells G, Feldman P. Preventing perineal trauma
 296 during childbirth: a systematic review. *Obstet Gynecol* 2000;95:464–71.
 297 [https://doi.org/10.1016/s0029-7844\(99\)00560-8](https://doi.org/10.1016/s0029-7844(99)00560-8).

298 [26] Kopas ML. A review of evidence-based practices for management of the
 299 second stage of labor. *J Midwifery Womens Health* 2014;59:264–76.
 300 <https://doi.org/10.1111/jmwh.12199>.

301 [27] Dahlen HG, Ryan M, Homer CSE, Cooke M. An Australian prospective cohort
 302 study of risk factors for severe perineal trauma during childbirth. *Midwifery*
 303 2007;23:196–203. <https://doi.org/10.1016/j.midw.2006.04.004>.

304 [28] Esscher A, Binder-Finnema P, Bødker B, Högberg U, Mulic-Lutvica A, Essén
 305 B. Suboptimal care and maternal mortality among foreign-born women in Sweden:
 306 maternal death audit with application of the “migration three delays” model. *BMC*
 307 *Pregnancy Childbirth* 2014;14:141. <https://doi.org/10.1186/1471-2393-14-141>.

308 [29] Jones D, Gill P. Breaking down language barriers. The NHS needs to provide
 309 accessible interpreting services for all. *BMJ* 1998;316:1476.
 310 <https://doi.org/10.1136/bmj.316.7143.1476>.

311 [30] Phelan M, Parkman S. How to work with an interpreter. *BMJ* 1995;311:555–7.
 312 <https://doi.org/10.1136/bmj.311.7004.555>.

313 [31] Azria E, Stewart Z, Gonthier C, Estellat C, Deneux-Tharaux C. [Social
 314 inequalities in maternal health]. *Gynecol Obstet Fertil* 2015;43:676–82.
 315 <https://doi.org/10.1016/j.gyobfe.2015.09.004>.

316 [32] Burrell M, Dilgir S, Patton V, Parkin K, Karantanis E. Risk factors for obstetric
 317 anal sphincter injuries and postpartum anal and urinary incontinence: a case-control
 318 trial. *Int Urogynecology J* 2015;26:383–9. <https://doi.org/10.1007/s00192-014-2478-7>.

319 [33] Aukee P, Sundström H, Kairaluoma MV. The role of mediolateral episiotomy

320 during labour: analysis of risk factors for obstetric anal sphincter tears. *Acta Obstet*
321 *Gynecol Scand* 2006;85:856–60. <https://doi.org/10.1080/00016340500408283>.

322 [34] Hauck YL, Lewis L, Nathan EA, White C, Doherty DA. Risk factors for severe
323 perineal trauma during vaginal childbirth: a Western Australian retrospective cohort
324 study. *Women Birth J Aust Coll Midwives* 2015;28:16–20.
325 <https://doi.org/10.1016/j.wombi.2014.10.007>.

326 [35] Belihu FB, Small R, Davey M-A. Episiotomy and severe perineal trauma
327 among Eastern African immigrant women giving birth in public maternity care: A
328 population based study in Victoria, Australia. *Women Birth J Aust Coll Midwives*
329 2017;30:282–90. <https://doi.org/10.1016/j.wombi.2016.11.008>.

330 [36] Groutz A, Hasson J, Wengier A, Gold R, Skornick-Rapaport A, Lessing JB, et
331 al. Third- and fourth-degree perineal tears: prevalence and risk factors in the third
332 millennium. *Am J Obstet Gynecol* 2011;204:347.e1-4.
333 <https://doi.org/10.1016/j.ajog.2010.11.019>.

334 [37] Baghurst PA, Antoniou G. Risk models for benchmarking severe perineal tears
335 during vaginal childbirth: a cross-sectional study of public hospitals in South
336 Australia, 2002-08. *Paediatr Perinat Epidemiol* 2012;26:430–7.
337 <https://doi.org/10.1111/j.1365-3016.2012.01300.x>.

338 [38] Gurol-Urganci I, Cromwell DA, Edozien LC, Mahmood TA, Adams EJ,
339 Richmond DH, et al. Third- and fourth-degree perineal tears among primiparous
340 women in England between 2000 and 2012: time trends and risk factors. *BJOG Int J*
341 *Obstet Gynaecol* 2013;120:1516–25. <https://doi.org/10.1111/1471-0528.12363>.

342 [39] Altman D, Ragnar I, Ekström A, Tydén T, Olsson S-E. Anal sphincter
343 lacerations and upright delivery postures--a risk analysis from a randomized
344 controlled trial. *Int Urogynecol J Pelvic Floor Dysfunct* 2007;18:141–6.

345 <https://doi.org/10.1007/s00192-006-0123-9>.

346 [40] Hsieh W-C, Liang C-C, Wu D, Chang S-D, Chueh H-Y, Chao A-S. Prevalence
347 and contributing factors of severe perineal damage following episiotomy-assisted
348 vaginal delivery. *Taiwan J Obstet Gynecol* 2014;53:481–5.
349 <https://doi.org/10.1016/j.tjog.2013.07.002>.

350 [41] Jandér C, Lyrenäs S. Third and fourth degree perineal tears. Predictor factors
351 in a referral hospital. *Acta Obstet Gynecol Scand* 2001;80:229–34.

352 [42] Combs CA, Robertson PA, Laros RK. Risk factors for third-degree and fourth-
353 degree perineal lacerations in forceps and vacuum deliveries. *Am J Obstet Gynecol*
354 1990;163:100–4. [https://doi.org/10.1016/s0002-9378\(11\)90678-4](https://doi.org/10.1016/s0002-9378(11)90678-4).

355 [43] Boucoiran I, Valerio L, Bafghi A, Delotte J, Bongain A. Spatula-assisted
356 deliveries: a large cohort of 1065 cases. *Eur J Obstet Gynecol Reprod Biol*
357 2010;151:46–51. <https://doi.org/10.1016/j.ejogrb.2010.03.024>.

358 [44] Grisot C, Mancini J, de Troyer J, Rua S, Boubli L, d’Ercole C, et al. [Perineal
359 morbidity of operative vaginal delivery using spatulas and vacuum: what’s the truth?].
360 *J Gynecol Obstet Biol Reprod (Paris)* 2011;40:348–58.
361 <https://doi.org/10.1016/j.jgyn.2011.03.007>.

362 [45] Menard J-P, Provansal M, Heckenroth H, Gamberre M, Bretelle F, Mazouni C.
363 [Maternal morbidity after Thierry’s spatulas and vacuum deliveries]. *Gynecol Obstet*
364 *Fertil* 2008;36:623–7. <https://doi.org/10.1016/j.gyobfe.2008.03.013>.

365 [46] Revicky V, Nirmal D, Mukhopadhyay S, Morris EP, Nieto JJ. Could a
366 mediolateral episiotomy prevent obstetric anal sphincter injury? *Eur J Obstet Gynecol*
367 *Reprod Biol* 2010;150:142–6. <https://doi.org/10.1016/j.ejogrb.2010.03.002>.

368 [47] Hudelist G, Gelle’n J, Singer C, Ruecklinger E, Czerwenka K, Kandolf O, et al.
369 Factors predicting severe perineal trauma during childbirth: role of forceps delivery

370 routinely combined with mediolateral episiotomy. *Am J Obstet Gynecol*
371 2005;192:875–81. <https://doi.org/10.1016/j.ajog.2004.09.035>.

372 [48] Andrews V, Sultan AH, Thakar R, Jones PW. Risk factors for obstetric anal
373 sphincter injury: a prospective study. *Birth Berkeley Calif* 2006;33:117–22.
374 <https://doi.org/10.1111/j.0730-7659.2006.00088.x>.

375 [49] Räisänen SH, Vehviläinen-Julkunen K, Gissler M, Heinonen S. Lateral
376 episiotomy protects primiparous but not multiparous women from obstetric anal
377 sphincter rupture. *Acta Obstet Gynecol Scand* 2009;88:1365–72.
378 <https://doi.org/10.3109/00016340903295626>.

379 [50] Buekens P, Bernard N, Blondel B, Grandjean H, Huisjes H, Kaminski M, et al.
380 [Episiotomy and prevention of complete and complicated tears. A study in 3
381 European countries]. *J Gynecol Obstet Biol Reprod (Paris)* 1987;16:513–7.

382 [51] Macleod M, Strachan B, Bahl R, Howarth L, Goyder K, Van de Venne M, et al.
383 A prospective cohort study of maternal and neonatal morbidity in relation to use of
384 episiotomy at operative vaginal delivery. *BJOG Int J Obstet Gynaecol*
385 2008;115:1688–94. <https://doi.org/10.1111/j.1471-0528.2008.01961.x>.

386 [52] Stedenfeldt M, Pirhonen J, Blix E, Wilsgaard T, Vonen B, Øian P. Episiotomy
387 characteristics and risks for obstetric anal sphincter injuries: a case-control study.
388 *BJOG Int J Obstet Gynaecol* 2012;119:724–30. <https://doi.org/10.1111/j.1471-0528.2012.03293.x>.

389 [53] Eogan M, Daly L, O’Connell PR, O’Herlihy C. Does the angle of episiotomy
390 affect the incidence of anal sphincter injury? *BJOG Int J Obstet Gynaecol*
391 2006;113:190–4. <https://doi.org/10.1111/j.1471-0528.2005.00835.x>.

392
393

Figure 1. Flow chart.

Table 1. Matching criteria of the two study groups.

	Women without OASI (n=163)	Women with OASI (n=171)	p-value
Age (years), median (IQR)	29.0 (26.0-34.0)	30.0 (25.5-34.0)	0.85
History of previous vaginal delivery, n (%)	63 (38.7)	56 (32.7)	0.28
Ethnicity			
Caucasian, n (%)	53 (32.5)	39 (22.8)	0.15
African, n (%)	68 (41.7)	91(53.2)	
Asian, n (%)	35 (21.5)	33 (19.3)	
Other, n (%)	7 (4.3)	7 (4.1)	
Delivery mode			
Spontaneous, n (%)	63 (38.7)	69 (40.4)	0.75
Ventouse, n (%)	20 (12.3)	12 (7.0)	
Spatulas, n (%)	59 (36.2)	57 (33.3)	
Forceps, n (%)	18 (11.0)	26 (15.2)	
Prophylactic episiotomy, n (%)	73 (44.8)	79 (46.2)	0.80
Birth weight (g), median (IQR)	3400 (3090-3780)	3380 (3100-3780)	0.82

IQR – interquartile range

OASI – obstetric anal sphincter injury

Table 2. Univariate analysis

	Women without OASI (n=163)	Women with OASI (n=171)	p-value
Language barrier, n (%)	8 (4.9)	20 (11.7)	0.03
Labor induction, n (%)	115 (70.6)	130 (76.0)	0.22
Labor duration (h), median (IQR)	17.0 (11.0-26.0)	22.0 (13.0-31.0)	0.01
Gestational age at delivery, median (IQR)	40.0 (39.4-40.7)	40.1 (39.1-40.7)	0.96
Occipito-posterior delivery, n (%)	3 (1.8)	19 (11.1)	< 0.001
Apgar <7 at 5 min, n (%)	4 (2.5)	1 (0.6)	0.17
Cord arterial pH, median (IQR)	7.23 (7.19-7.28)	7.24 (7.19-7.28)	0.83

IQR – interquartile range

OASI – obstetric anal sphincter injury

Table 3. Multivariate analysis

	Univariate analysis			Multivariate analysis		
	OR	95% CI	p-value	OR	95% CI	p-value
Language barrier	2.57	1.14-6.36	0.03	3.32	1.36-8.90	0.01
Labor duration	1.03	1.01-1.05	0.001	1.03	1.01-1.05	0.004
Ethnicity						
Caucasian	ref.	-	-	ref.	-	-
African	1.82	1.08-3.07	0.02	1.85	1.08-3.19	0.03
Asian	1.28	0.68-2.41	0.44	0.95	0.47-1.92	0.89
Others	1.36	0.43-4.28	0.59	1.28	0.37-4.29	0.69
Occipito-posterior delivery	6.62	2.20-28.60	0.003	6.33	2.04-27.78	0.004

CI – confidence interval

OR – odd ratio