



**HAL**  
open science

## The Potential of FGF-2 in Craniofacial Bone Tissue Engineering: A Review

Anita Novais, Eirini Chatzopoulou, Catherine Chaussain, Caroline Gorin

► **To cite this version:**

Anita Novais, Eirini Chatzopoulou, Catherine Chaussain, Caroline Gorin. The Potential of FGF-2 in Craniofacial Bone Tissue Engineering: A Review. *Cells*, 2021, 10 (4), pp.932. 10.3390/cells10040932 . hal-03215638

**HAL Id: hal-03215638**

**<https://hal.sorbonne-universite.fr/hal-03215638v1>**

Submitted on 3 May 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Review

# The Potential of FGF-2 in Craniofacial Bone Tissue Engineering: A Review

Anita Novais<sup>1,2</sup>, Eirini Chatzopoulou<sup>1,2,3</sup>, Catherine Chaussain<sup>1,2</sup>  and Caroline Gorin<sup>1,2,\*</sup> 

- <sup>1</sup> Pathologies, Imagerie et Biothérapies Orofaciales, Université de Paris, URP2496, 1 rue Maurice Arnoux, 92120 Montrouge, France; anitarsnovais@gmail.com (A.N.); eirini.chatzopoulou@gmail.com (E.C.); catherine.chaussain@u-paris.fr (C.C.)
- <sup>2</sup> AP-HP Département d'Odontologie, Services d'odontologie, GH Pitié Salpêtrière, Henri Mondor, Paris Nord, Hôpital Rothschild, Paris, France
- <sup>3</sup> Département de Parodontologie, Université de Paris, UFR Odontologie-Garancière, 75006 Paris, France
- \* Correspondence: caroline.gorin@u-paris.fr; Tel./Fax: +33-(0)1-5807-6724

**Abstract:** Bone is a hard-vascularized tissue, which renews itself continuously to adapt to the mechanical and metabolic demands of the body. The craniofacial area is prone to trauma and pathologies that often result in large bone damage, these leading to both aesthetic and functional complications for patients. The “gold standard” for treating these large defects is autologous bone grafting, which has some drawbacks including the requirement for a second surgical site with quantity of bone limitations, pain and other surgical complications. Indeed, tissue engineering combining a biomaterial with the appropriate cells and molecules of interest would allow a new therapeutic approach to treat large bone defects while avoiding complications associated with a second surgical site. This review first outlines the current knowledge of bone remodeling and the different signaling pathways involved seeking to improve our understanding of the roles of each to be able to stimulate or inhibit them. Secondly, it highlights the interesting characteristics of one growth factor in particular, FGF-2, and its role in bone homeostasis, before then analyzing its potential usefulness in craniofacial bone tissue engineering because of its proliferative, pro-angiogenic and pro-osteogenic effects depending on its spatial-temporal use, dose and mode of administration.

**Keywords:** FGF-2; bone tissue engineering; angiogenesis; mineralization; signaling pathways



**Citation:** Novais, A.; Chatzopoulou, E.; Chaussain, C.; Gorin, C. The Potential of FGF-2 in Craniofacial Bone Tissue Engineering: A Review. *Cells* **2021**, *10*, 932. <https://doi.org/10.3390/cells10040932>

Academic Editor: Barbara Kaltschmid

Received: 5 March 2021

Accepted: 15 April 2021

Published: 17 April 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

The skeletal system is dynamic, metabolically active and functionally diverse. As well as a structural function, it has a metabolic role. It supports and protects the vital internal organs and is the site of synthesis of the hematopoietic marrow and provides sites of muscle attachment for locomotion. Bone is involved in both mineral metabolism, via calcium and phosphate homeostasis, and acid–base balance, via the buffering of hydrogen ions [1]. Moreover, it has been suggested that bone has other important endocrine functions in fertility, glucose metabolism, appetite regulation and muscle function [2,3]. The craniofacial area is prone to trauma and pathologies that often result in large bone damage, these leading to both aesthetic and functional complications for patients. Throughout life, the craniofacial area is at risk of complex injuries that require bone grafting to restore function. The etiology of such injuries may be accidental (e.g., acute trauma), congenital (e.g., birth defects or deformities), pathological (e.g., maxillofacial tumors, such as ameloblastoma, or infection) or surgical. Whenever the lesions are extensive, they cause large bone defects that cannot self-repair because they exceed the body's natural regenerative capacity. The “gold standard” for treating these large defects is autologous bone grafting. Since the defects are extensive, harvesting of the necessary bone at the donor site can cause major morbidity, including bone deformity, as well as pain and occasionally continuous progressive resorption. Tissue engineering has made it possible to approach these issues from another angle.

Before talking about tissue engineering; however, it is necessary to describe the tissue of interest. This review is, therefore, organized into three sections, the first describing general bone physiology (e.g., its composition, functioning and signaling pathways), the second the implication of FGF-2 in bone physiology, and the third highlighting the interesting characteristics of FGF-2 for craniofacial bone engineering and its various current potential uses for promoting bone repair.

## 2. Background on Bone Physiology


In the skeleton, there are two types of bone differentiated by their structure—cortical and trabecular. Although both types have an identical chemical composition, they differ both macroscopically and microscopically [4] (Appendix A Figure A1).

### 2.1. Bone Composition

Bone is a mineralized connective tissue composed of bone cells, vessels, and an extracellular matrix (ECM), which is produced by the bone cells. The proportion of these components varies with bone type (long or flat bone) and anatomical site [5]. The mineralized component of bone tissue gives rigidity and hardness to the material, while the organic components of the ECM provide flexibility [6].

#### 2.1.1. Bone Cells

The osteoblast, a specialized bone-forming cell, mostly differentiated from mesenchymal stem cells (MSCs) [7,8], produces and secretes the major bone matrix protein essential for matrix mineralization, namely, type I collagen [9,10]. These cells work in clusters on the bone surface [5], becoming committed to one of various possible fates, thanks to some key proteins and pathway signaling, such as runt-related transcription factor 2 (Runx-2) and Osterix (Figure 1).



**Figure 1.** Bone remodeling regulation can be paracrine or endocrine. Several factors participate in paracrine regulation including cytokines (IL-1, IL-6, TNF-alpha, IL-4 and interferon-gamma), PGE2, VEGF, and hypoxia, as well as bone cells. There are three main cell types involved: osteoblasts, osteocytes and osteoclasts. Osteoblasts, which differentiate from mesenchymal progenitors thanks to certain proteins (Runx2, Osx and Wnt) and FGF signaling pathways, are responsible for bone formation. They can also become osteocytes able to regulate osteoblastogenesis through production of inhibitors (DKK-1 and SOST), that inhibit Wnt signaling. Lastly, osteoclasts, involved in bone resorption are activated through RANK-RANKL-OPG signaling pathway cross-talk. Whenever there

is a need for bone resorption, osteoblasts and osteocytes express RANKL on their surface, and this then binds to RANK in osteoclast precursors, activating their differentiation. OPG is then secreted to stop bone resorption binding to RANKL blocking the possibility of RANK-RANKL binding and preventing bone resorption. Once activated, mature osteoclasts bind to the bone matrix, becoming polarized. Their cytoskeleton organizes into actin rings forming the sealing zone, which provides an isolated acidic microenvironment, to dissolve minerals and digest the selected bone matrix thanks to the ruffle border (RB). After resorption, the osteoclasts endocytose the degraded collagen fragments, and the calcium and phosphate released are then transported through the cell and liberated at the functional secretory domain before being released into the bloodstream. Bone formation and resorption are also influenced by endocrine regulation. Various factors may be involved, for example, PTH, 1,25(OH) Vitamin D, calcitonin and thyroid hormone.

Osteocytes, the most abundant cell type, are found in mature bone and are long-lived [11]. They are distributed throughout the bone matrix and can interact with other osteocytes or osteoblasts on the bone surface, vasculature and bone cells on the surface of bones, in a complex intercellular network [6,12,13]. Osteocyte-transduced signals seem to orchestrate bone response by regulating the synchronized action of osteoblasts and osteoclasts [14–16] (Figure 1).

The osteoclasts, multinucleated cells formed by the fusion of precursors derived from the hematopoietic cells of the mononuclear lineage [17,18], are the only cells known to be capable of resorbing bone (Figure 1).

Other types of cells, such as chondrocytes are found within the bone. These are derived from pluripotent MSCs, and secrete type II collagen, participating in the endochondral ossification process [19].

### 2.1.2. Bone Extracellular Matrix

Bone ECM contains both a mineral and an organic phase, the latter representing approximately 90% of the organic content of bone tissue. The mineral portion is largely calcium phosphate in the form of hydroxyapatite crystals deposited in an osteoid matrix and is responsible for bone's mechanical rigidity and load-bearing strength. The organic portion, which contains water, non-collagenous proteins, lipids and specialized bone cells, gives flexibility and elasticity to bone tissue [1,6].

Non-collagenous proteins constitute 10% to 15% of total bone protein content. Almost 25% of non-collagenous proteins are adsorbed from serum and due to their acidic properties are able to bind to the matrix [20], which allows strengthening of the collagen structure and regulating its mineralization [4]. Osteocalcin is the major non-collagenous protein, and is involved in calcium binding, stabilization of hydroxyapatite in the matrix and the regulation of bone formation, acting as a negative regulator, inhibiting premature or inappropriate mineralization [21,22].

## 2.2. Bone Formation

Bones form through two different processes: bone modeling and bone remodeling (Figure 2). Bone modeling occurs primarily during growth and development in childhood, although it can also appear after the skeleton has matured [23,24].


In contrast, bone remodeling occurs after the skeleton has reached maturity, during adulthood, involving resorption of old or damaged bone, and its replacement by newly formed bone. Here, we focus on bone remodeling.

### 2.2.1. Bone Remodeling Process

Bone remodeling is essential for structural integrity, biomechanical stability, bone volume and calcium/phosphate homeostasis [1,4,25]. In normal adult bone, there is a homeostatic balance in which bone resorption is followed by bone formation for maintaining bone strength and mineral homeostasis, keeping the overall bone volume and structure unchanged [26,27]. By this process, about 10% of the skeleton may be renewed every

year [28]. That is, bone remodeling only takes place when it is required, either because the specific area is damaged and/or old.

The bone remodeling cycle has several phases: cell activation, bone resorption, reversal, bone formation and termination (Figure 2).



**Figure 2.** Bone metabolism: Modeling vs. Remodeling—While bone modeling implies a change in bone shape or size since resorption and formation occur independently at distinct sites: osteoblasts secrete osteoid matrix in the opposite site where osteoclasts resorb bone. Bone remodeling involves the resorption and formation of bone, one after the other, at the same site to replace old and/or damaged bone by newly formed bone. An initiating remodeling signal, such as hormonal or mechanical signal, is detected by the bone, inducing the release of paracrine factors that lead to retraction of the bone lining cells which exposes the bone surface, allowing recruitment of osteoclast precursors from the capillaries directly into the basic multicellular unit. MSC-F and RANKL, secreted by osteocytes, induce recruitment of precursor cells of hematopoietic lineage, initiating their differentiation to multinucleated osteoclasts. The differentiated attached osteoclasts rearrange their cytoskeleton to adhere to the bone surface, decreasing the pH to as low as 4.5, this dissolving the bone mineral. Once resorption is finished, the osteoclasts go through apoptosis. After resorption, mononuclear cells are recruited to remove collagen fragments from the surface, and then new osteoblasts begin collagen deposition, forming what is known as osteoid matrix, until the cavities are filled. Osteoblasts produce new bone, and some of them become buried within the newly formed bone matrix turning into osteocytes with their extensive canalicular network connecting them to the bone surface lining cells, osteoblasts and other osteocytes. The osteoid mineralizes, and the bone enters into a quiescent phase.

### 2.2.2. Regulation of Bone Remodeling

The remodeling cycle is tightly regulated to achieve a balance between bone formation and resorption. Considering that remodeling can occur in several locations simultaneously, local regulation is critical to achieving this balance.

- Major signaling pathways
  1. RANKL/RANK/OPG

One of the major signaling pathways that regulates bone remodeling involves three proteins: RANKL, receptor activator of nuclear factor kappa-B (RANK) and osteoprotegerin (OPG). The interaction between these proteins determines whether, at a specific location, bone resorption or bone formation occur. RANKL is a cytokine expressed on the surface of osteoblasts, osteocytes and chondrocytes. It activates nuclear factor kappa-B (NF- $\kappa$ B) and other signaling pathways through the interaction with its receptor, RANK, located on osteoclast precursors. RANKL/RANK activation has an important role in osteoclast

differentiation, allowing osteoclast formation, activation and survival [13,29–32]. OPG, a soluble decoy receptor for RANKL expressed by osteoblasts and osteocytes, binds to RANKL with high affinity, preventing it from binding to its receptor RANK. Thus, OPG is a natural inhibitor of RANKL. The RANKL/OPG expression rate regulates the extent of osteoclast formation and activity [33–35] (Figure 1).

## 2. Wnt signaling

Wingless-related integration site (Wnt) molecules are cysteine-rich glycoproteins involved in controlling cell proliferation, cell-fate specification, gene expression and cell survival. Wnt signaling pathways are involved in bone formation, having an anabolic effect, and increasing bone density and strength, by regulation of osteoblast differentiation and function [36,37] (Figure 1).

Wnt pathways also play a major role in osteoclast differentiation. Specifically, Wnt canonical signaling up-regulates OPG and downregulates RANKL, which inhibits osteoclast formation and therefore bone resorption [36,38]. In contrast, activation of the noncanonical pathway in osteoclast precursors enhances RANKL-induced osteoclastic differentiation [36].

Wnt signaling is inhibited by secreted proteins such as sclerostin and Dickkopf-related protein 1–4 (DKK-3/4) synthesized by osteocytes [39,40] (Figure 1). Nevertheless, during bone remodeling, osteocytes decrease the expression of sclerostin and Dickkopf-related protein 1 and 2 (DKK-1/2), allowing osteoblast bone formation to occur after bone resorption. After the completion of remodeling, newly formed osteocytes become entombed within the bone matrix and start expressing Wnt inhibitors, stopping further bone formation [12].

- Endocrine regulation

Bone turnover by osteoblasts and osteoclasts is essential for the maintenance of bone strength and morphology. Due to its importance, this process must be thoroughly regulated to prevent malfunction of the remodeling process at any stage of the cycle. Several hormonal agents are involved in this regulation, including PTH [10], 1,25 (OH) Vitamin D [4], calcitonin [41], thyroid hormone [42], growth hormone [43], glucocorticoids [44] and sex hormones [45].

- Paracrine regulation

Some cytokines may have stimulatory and inhibitory effects on bone metabolism. Cytokines like interleukin 1 (IL-1), IL-6, and tumor necrosis factor (TNF $\alpha$ ) can increase osteoclastic resorption, whereas others, such as interleukin 4 (IL-4) and gamma interferon, decrease osteoclast proliferation and differentiation [46,47].

Prostaglandins may also influence bone formation, although their exact role remains unclear. Prostaglandin E<sub>2</sub> (PGE<sub>2</sub>) is a major inducer of bone resorption and is thought to increase the RANKL/OPG ratio to improve osteoclastogenesis (Figure 1). At the same time, it has been hypothesized to stimulate osteoblast proliferation and differentiation thereby enhancing bone formation [48]. Inside the bone matrix, there are growth factors that affect bone metabolism. The main families involved are the transforming growth factor  $\beta$  (TGF $\beta$ ) family (TGF $\beta$  and BMPs) and the fibroblast growth factor (FGF) family.

### 2.3. Bone Vascularization

There is an intimate link between osteogenesis and angiogenesis, both processes needing to be tightly coupled for optimal physiological bone function [49]. In the event of a critical bone defect, early vascularization is necessary for osteogenic reconstruction, to allow the nutritional support for the bone grafts [50–52]. The close relationship between blood vessels and bone cells is also well illustrated by abnormalities resulting from inappropriate vascularization, these leading to the appearance of skeletal malformation, such as craniofacial dysmorphology [53].

In the close connection between these two processes, several factors have been described as being both angiogenic and osteogenic. Notably, it has been demonstrated that

hypoxia (oxygen tension) and the vascular endothelial growth factor (VEGF) family affect endochondral angiogenesis as well as cells from the bone lineage [54–56].

Indeed, several pro-angiogenic factors are involved in bone repair. Some of these factors have direct effects, both having angiogenic properties and regulating osteogenic molecules, like BMPs, angiopoietin (Ang), platelet-derived growth factor (PDGF) and insulin-like growth factor (IGF) family members (Appendix A Table A2).

Others are well known to indirectly enhance bone repair using their pro-angiogenic properties. VEGF, an endothelial cell (EC)-specific mitogen, is secreted by cells involved in skeletal development and repair, such as hypertrophic chondrocytes or differentiating mesenchymal cells, osteoblasts, and ECs [57,58]. It can be a chemoattractant molecule, engaging ECs into bone tissue and tightly controlling the differentiation and functions of osteoblasts and osteoclasts [55,59–62]. It is also involved both in endochondral ossification promoting vessel invasion and cell recruitment [59,63], and in intramembranous ossification, by affecting bone cell activity [52,53,64]. It has been reported that VEGF upregulates the RANK receptor in ECs and strongly stimulates angiogenesis [65]. In turn, RANKL may have an important role in enabling EC survival via the phosphatidylinositol 3-kinase/protein kinase B (PI3K/Akt) signaling pathway [66,67], one of the major signaling pathways triggered by activated VEGF receptor [68] (Figure 1).

The hypoxic signaling pathway has been reported to directly enhance VEGF expression, being described as a major regulator of VEGF expression [69,70]. Indeed, hypoxia-inducible transcription factors (HIFs) are expressed in osteogenic cells, notably osteoblasts, and hence, hypoxia upregulates VEGF expression, thereby promoting angiogenesis and osteogenesis. Thus, hypoxia and VEGF signaling are involved in the coupling of angiogenesis and osteogenesis [54,55,71].

This review focuses on the FGF family and specifically fibroblast growth factor 2 (FGF-2) and its potential usefulness in bone tissue engineering.

### 3. FGF-2 in Bone Homeostasis

#### 3.1. FGF/FGFR Signaling in Bone


Various growth factors act within the bone matrix and influence bone metabolism. The main families involved are the TGF $\beta$  (TGF $\beta$  and BMPs) and FGF families. The latter is also involved in angiogenesis, which makes it interesting to study more in detail. In this review, we focus on the great potential of FGF-2 in bone metabolism and discuss the interest in its use in bone tissue engineering.

Members of the FGF family are single-chain polypeptide growth factors of approximately 20–35 kDa. At least 23 members of this family have been described in mammals [72], from FGF-1 to FGF-23. The secreted FGFs are differentially expressed in almost all tissues of the developing embryo, functioning as essential regulators of the earliest stages of embryonic development. They are also expressed in postnatal and adult tissues, fulfilling essential roles in tissue homeostasis, repair, regeneration, angiogenesis and bone metabolism [61,73–78]. There are four FGF receptors: FGFR-1 to FGFR-4 [79,80].

Given the ubiquitous roles of FGF signals in development, homeostasis, and disease, tight regulation of the pathways is essential through cofactors that participate in the affinity and specificity of FGFR-binding, and also in specifying signaling activities [81,82]. The endocrine regulation of FGFs, including FGF-19, FGF-21 and FGF-23, is mediated by their binding with the Klotho protein family ( $\alpha$  or  $\beta$ Klotho or KLPH) [83–85], which allows them to circulate through the matrix without being trapped and stored [82], thereby acting in an endocrine-like fashion.

Canonical FGFs exert their pleiotropic effects by binding and activating the FGFR subfamily of receptor tyrosine kinases that are coded by four genes (FGFR-1, FGFR-2, FGFR-3, and FGFR-4) in mammals. [86]. The four FGFRs have distinct ligand specificity and are expressed in a tissue-specific manner [87]. Cofactors such as heparan sulfate and klotho are low-affinity receptors that do not induce a biological signal but rather are used as auxiliary proteins to regulate FGF binding and subsequent phosphorylation of adaptor proteins for

four major intracellular pathways [88–91]: rat sarcoma-mitogen-activated protein kinase-extracellular signal-regulated kinase 1 and 2 (RAS-MAPK-ERK1/2), PI3K-AKT-glycogen synthase kinase 3 (GSK3), phospholipase C $\gamma$ -protein kinase C (PLC $\gamma$ -PKC), and signal transducer and activator of transcription proteins-Janus kinase (STAT-Jak) [81,86,92,93]. The main regulation pathway remains the paracrine one, involving FGF-2, which binds FGFRs through a heparan sulphate glycosaminoglycan binding site, limiting their diffusion through the ECM [88,90,94]. Upon binding of FGF to its receptor, receptor dimerization and transautophosphorylation of the kinase domain take place [95] (Figure 3).



**Figure 3.** FGF/FGFR signaling—FGFs can bind to FGFRs with the help of heparan sulfate, a co-factor, and thereby induce their biological effects through activation of four major signaling pathways: RAS-MAPK-ERK1/2, PI3K-AKT-GSK3, PLC $\gamma$ -PKC, and STAT-Jak.

FGFs can activate several MAPKs such as C-Jun N-terminal kinase (JNK), ERK, and p38MAPK [96], that share many structural similarities, while inducing different responses [97,98]. The ERK1/2 branch, one of the major routes for FGF signaling [99], promotes a mitogenic response and is observed in all cell types, while p38 and JNK kinase are usually associated with inflammatory and stress responses [98,100] and are mainly involved in the cell cycle, cytoskeleton and cell migration [101]. For instance, FGF-2 regulates mesenchymal stem cell migration [102] via the PI3K-AKT pathway, neural progenitor cell proliferation via PI3K/GSK3 signaling [103], and fibroblast migration via PI3 kinase, Ras-related C3 botulinum toxin substrate 1 (Rac1) and JNK [104], as well as promoting bone marrow MSC proliferation through the ERK1/2 pathway [105]. Activation of the MAPK pathway, in response to FGF-2 signaling, is key in determining the activity of RUNX-2, a master transcription factor of bone formation [106]. FGF-2 promotes osteoblast proliferation and differentiation [107], through the activation of the ERK1/2 pathway [108–110] and a crucial role of this pathway has been identified in the differentiation of osteoblasts and chondrocytes [107,111–113]. Specifically, MAPK phosphorylates Runx-2 Ser/Thr residues, a critical step for Runx2 acetylation and stabilization against degradation [114]. Sprouty (SPRY) and Sprouty related (SPRED) are two antagonists of this pathway [115,116] that act as intracellular antagonists of FGFR signaling, the first by suppressing ERK1/2 activation [117] and the second by suppressing rapidly accelerated fibrosarcoma (Raf) activation [118] (Figure 3).

The PI3K/AKT pathway is implicated in cell polarization, migration, cell fate determination and apoptosis. For instance, FGF-2 enhances the migratory activity of periodontal ligament cells (PDLSCs) through this pathway. In addition, FGF-2 induced Akt phosphory-



lation promotes proliferation in neural progenitor cells [103] and MSCs [105]. Sprouty2 has also been associated with the PI3K/Akt pathway, suppressing Akt phosphorylation [101].

The FGFR tyrosine kinase domain can also directly phosphorylate PLC $\gamma$ , which leads to the hydrolysis of phosphatidylinositol 4,5-bisphosphate to produce inositol triphosphate (IP3) and diacylglycerol. Subsequently, IP3 increases intracellular calcium ion levels, while diacylglycerol activates PKC. Furthermore, it has been shown that FGFs can induce expression of receptor of activated protein C kinase 1 (RACK-1), a protein that further stabilizes activated PKC [119]. It has been shown that FGFR-1 and FGFR-2 can directly bind to activate PLC $\gamma$ 1 [120]. In fact, FGF-2-induced activation of the PLC $\gamma$  pathway is involved both in increased expression and transcriptional activity of RUNX2 [121]. In dental tissue-derived mesenchymal stem cells, it has been shown that neurogenic differentiation in human dental pulp stem cells (DPSCs) is induced by FGF-2 via the PLC $\gamma$  signaling pathway [122], which is already known for its role in the differentiation of neuronal cells [123].

Other pathways such as STAT [96] and ribosomal S6 kinase 2 (RSK2) pathways [124] have been shown to be involved in FGF-2 signaling. The activated FGFR also phosphorylates and activates STAT1, STAT3, and STAT5, to regulate STAT pathway target gene expression [79]. These pathways control the steps of osteoblastogenesis resulting in the modulation of bone cell proliferation, differentiation, and apoptosis, depending on the stage of cell differentiation [125,126]. Therefore, normal FGFR activity is critical for the development of numerous types of tissue, including the craniofacial skeleton, as observed in several genetic diseases causing abnormalities in bone and cartilage formation due to mutations in the genes encoding FGFs or their receptors [127,128].

### 3.2. FGF-2: An Essential Regulator in Skeletal Tissue


FGF-2, also known as basic fibroblast growth factor, is a canonical FGF that belongs to the FGF-1 subfamily. It is encoded by *FGF-2*, on chromosome 4 [129], and is a wide-spectrum angiogenic, mitogenic and neurotrophic factor expressed by many types of cells in both adult and developmental stages [130]. It is a regulator of proliferation [131], migration [132], differentiation [122,129,133], cell survival [134], and stemness in human stem cells [135,136]. Apart from its key role in angiogenesis, FGF-2 is a major player in skeletal development, bone formation, and fracture repair [137–139]. During embryogenesis, it is a strong mesodermal inducer, and its receptors are strongly expressed in developing bones [140,141]. Throughout life, it is constantly expressed in osteoblasts and stored in the ECM [142]. Taken together, these properties make FGF-2 an attractive molecule for clinical and pharmaceutical applications in bone regeneration.

FGF-2 is highly expressed in bone tissues, with several isoforms due to alternative start codons for FGF-2 mRNA translation initiation [143]. The high molecular weight FGF-2 (HMW FGF-2) isoforms (24, 23, and 22 kD) are localized in the nucleus, whereas low molecular weight FGF-2 (LMW FGF-2) (18 kD) is cytoplasmic, and membrane associated. This differential intracellular trafficking also reflects a difference in function. Specifically, the LMW FGF-2 promotes osteoblast differentiation and mineralization via the activation of the Wnt pathway [144] and via synergistic actions with bone morphogenetic protein 2 (BMP-2). Indeed, endogenous FGF/FGFR signaling is a positive upstream regulator of the BMP-2 gene in calvarial osteoblasts [145]. In contrast, HMW FGF-2 acts in the nucleus as a transcriptional factor that upregulates the expression of genes associated with impaired mineralization, such as *SOST* and *FGF-23* [146].

Experimental in vitro evidence regarding FGFR-2 mutations associated with craniosynostosis syndromes highlights the major contribution of FGF-2 to bone cell fate. The genetic inactivation of *FGFR-2* causes reduced osteoblast proliferation and increased osteopenia, while *FGFR-1* has been associated with stage-dependent regulation of osteoblast proliferation and differentiation [147–149].

Exogenous FGF-2 rescued reduced bone nodule formation by upregulating the Wnt/ $\beta$ -catenin pathway in FGF-2 $^{-/-}$  osteoblast cultures. [150,151]. Furthermore, when two FGFR-2

mutants from the Apert and Crouzon syndromes were expressed in immature osteoblasts, both inhibited osteoblastic differentiation but also increased apoptosis [152] downregulating many Wnt targets and inducing SRY-box 2 (SOX-2), a transcription factor that maintains the undifferentiated state of cells [153] (Figure 4). Furthermore, in a line of murine mesenchymal progenitor cells, induced overexpression of FGFR-2 led to increased cell proliferation and osteogenic differentiation through ERK1/2 by FGFR-2 signaling [154]. Inversely, primary calvarial osteoblasts from *Fgf2*<sup>-/-</sup> mice showed reduced BMP-2 induced periosteal bone formation [155]. In more mature cells, ERK1/2 activation by FGF-2 enhances acetylation and stabilization of RUNX-2, a key transcription factor involved in osteoblastogenesis and bone formation [114,156,157]. In addition, FGFR-2 signaling is crucial for the induction of apoptosis when osteoblasts are well differentiated [158,159], an important step for bone homeostasis. These results underline the dual temporal role of FGF-2 signaling in bone development, this protein promoting proliferation in immature osteoblasts and differentiation in mature ones. Furthermore, it should be underlined that since FGF-2 can bind equally to FGFR-1 and FGFR-2, differential activation of one of them may be responsible for signal transduction towards the proliferation of progenitors or towards osteogenic differentiation in post-proliferating cells [154].



**Figure 4.** FGF/FGFR signaling in bone—FGF-2 is highly expressed in bone tissues. There is a high molecular weight (HMW) form, located in the nucleus, that acts as a transcriptional factor, and upregulates the expression of *SOST* and *FGF-23* responsible for inducing mineralization. The low molecular weight (LMW) form is cytoplasmic or membrane associated. The latter can promote osteoblast differentiation and mineralization through the Wnt pathway, BMP-2 signaling, or synergistic action with BMP-2. By activation of FGFR signaling, LMW FGF-2 also activates MAPK-ERK/2, which acts as a transcriptional factor that upregulates mineralization genes such as *RUNX2*.

The important dual role of FGF-2 in bone formation is saliently demonstrated by transgenic mouse models. Conditional deletion of *fgf-2* in mice yields a skeletal dwarfism phenotype and reduced bone formation [147]. Furthermore, *fgf-2* haploinsufficient mice are characterized by generalized osteopenia [151] and *fgf-2* knockout mice display greatly reduced trabecular plate-like structures and loss of connecting rods [160]. This reduced bone formation is due to defective osteoblast differentiation and alteration of progenitor cell lineage commitment, FGF-2 deficiency resulting in increased bone marrow adipogenesis and reduced osteogenesis [161]. Nonetheless, overexpression of *fgf-2* in mice also gives rise to skeletal defects, including a shortening and flattening of long bones, with a decrease in osteoblast differentiation, impaired bone formation, and moderate macrocephaly [162,163].

#### 4. FGF-2 in Craniofacial Bone Tissue Engineering

In recent decades, tissue engineering has emerged as a new biomedical field with advanced approaches for tissue regeneration and healing [164]. It was initially defined as “the application of the principles of biology and engineering to the development of functional substitutes for damaged tissue” [165]. Bone tissue engineering needs to restore distinct functions: structural (e.g., bone, cartilage), barrier- and transport-related (e.g., blood vessels), and/or biochemical and secretory (e.g., hematopoiesis, calcium metabolism).

Tissue engineering is based on the combination of three basic tools: cells, biomaterials, and suitable biochemical and physical factors, seeking to mimic the physical and functional properties of the natural tissue creating a tissue-like structure [166,167]. The last factor to consider is the presence of exogenous chemical and mechanical stimuli, such as soluble growth and differentiation factors (e.g., BMP, FGF-2, VEGF and TGF- $\beta$ ), and mechanical forces. These factors can be incorporated into a construct during scaffold fabrication itself or included in the culture medium to facilitate the survival, proliferation, and differentiation of the implanted cells and their integration into the host. This section focuses on FGF-2 as a good candidate for craniofacial bone tissue engineering by acting on both angiogenic and osteogenic processes.

##### 4.1. FGF-2: An Exogenous Factor *In Vitro*

Exogenous FGF-2 administration *in vitro* has multiple effects on bone cell fate. Firstly, it has been shown that FGF-2 maintains the osteoblast precursor proliferative state [142,168]. Its anabolic effect is evident through the stimulation of bone marrow stem cells, sustaining their osteogenic potential by maintaining the cells in an immature state with a fibroblast-like morphology expressing less alkaline phosphatase (ALP) [169]. DPSCs, stem cells from human exfoliated deciduous teeth (SHEDs) and stem cells from apical papilla (SCAPs) treated with FGF-2 express stemness-related markers including STRO-1 and CD-146 [170–172]. Recently, it has been reported that DPSCs/SHEDs displayed increased and prolonged proliferation upon FGF-2 treatment *in vitro*, together with delayed type 1 collagen expression [173]. On the other hand, human calvaria bone cells grown in mineralizing medium for several weeks with FGF-2 showed no stimulation of proliferation, suggesting that mature bone cells do not respond to FGFs' mitogenic signal [174]. In fact, the decrease in intrinsic proliferation potential of human mesenchyme-derived progenitor cells with age was partially attributed to a reduction in FGF-2 expression, as observed in elderly humans [175]. In line with this, there is evidence that the efficacy of FGF-2 in inducing bone formation might be maximized if targeting younger cells, such as juvenile osteoblasts [176].

Concomitant with prolonged stemness, the abolition of mineralization by FGF-2 has been shown in various stem cell lines. Shimabukuro et al. [177] demonstrated that treatment of human DPSCs with FGF-2 increased their migration and proliferation ability but also impaired mineralization. On the contrary, if hDPSCs were only FGF-2 pretreated for a short period of time but left to differentiate under normal osteogenic conditions, ALP activity and nodule formation increased. Likewise, other studies confirm that pretreatment with FGF-2 during the proliferation phase leads to increased ALP activity, formation of mineralized nodules and expression of dentin sialoprotein and dentin matrix protein 1 (DMP-1) in hDPSCs [178]; dentin sialophosphoprotein (DSPP) and bone sialoprotein (BSP) expression in immature adult rat incisor dental pulp cells [179]; BSP, osteocalcin (OCN), osteopontin (OPN) and matrix Gla protein (MGP) in cementoblasts [180]; and hyaluronan in hDPSCs [181]. Nauman et al. [182] showed that *in vitro* administration of FGF-2 to rat osteoprogenitor cells accelerated the mineralization process through alkaline phosphatase and OCN expression and lowered the phosphate threshold needed for successful bone nodule formation. These observations suggest that FGF-2 enhances cell growth at early stages, a step that is crucial for accelerated differentiation at later time points. On the other hand, the spatiotemporal patterns of FGF signaling *in vivo* may differ from those found in culture conditions. Another possibility is that cell responses to FGF signaling *in vivo*

are determined by, or are coordinated with, signaling from other cytokines in situ, such as BMPs and WNT [75,183].

#### 4.2. FGF-2: An Exogenous Factor In Vivo

FGF-2 alone or in conjunction with other molecules and in combination with several types of scaffold has been assessed in various preclinical models for craniofacial bone regeneration. The rationale of FGF-2 administration is, first, its direct impact on bone cell proliferation and differentiation and, second, its pro-angiogenic action at the site of bone regeneration.

FGF-2 seems equally beneficial when tested in bone regeneration models of intramembranous or endochondral ossification. FGF-2 can be expressed by differentiating osteoblasts at sites of intramembranous ossification or by growth plate chondrocytes [128]. Its administration in vivo promotes regeneration of cranial [184,185], and periodontal [186–188] bone defects, as well as being associated with a shorter timeline of craniofacial bone repair [173]. Indeed, it has also been shown that FGF-2 was able to partially restore the lost cancellous bone mass in the ovariectomized rat [138]. The addition of FGF-2 significantly increased central defect bone filling in aged mice, leading to qualitatively superior bone formation. This suggests that FGF-2 is a good candidate for boosting bone regeneration in areas with impaired angiogenic potential or small numbers of native osteoprogenitor cells [189]. Similar benefits in angiogenesis and bone formation have been found in critical size defects in rat calvaria [190,191]. Furthermore, controlled delivery of FGF-2 in combination with a low dose of BMP-2 improved aged murine calvaria bone defect healing as compared to treatment with BMP-2 alone and the use of bone substitute impregnated with BMP-2 and FGF-2 promoted periodontal regeneration in non-human primates [189,192].

Vascularization plays a crucial role in bone tissue engineering seeking to replace large tissue losses due to trauma, surgery, or other clinical scenarios where spontaneous bone repair is not feasible. Indeed, using a radiotracer, Collignon et al. observed a correlation between early angiogenesis assessed by positron emission tomography and bone formation determined by micro-computed tomography within mouse calvarial bone critical size defects [193]. Recently, Novais et al. found that FGF-2 priming of DPSCs/SHEDs boosted intramembranous bone formation in critical size calvaria defects in immunodeficient mice [173]. Indeed, priming these cells with FGF-2 greatly enhanced stem cell early proliferation leading to increased bone regeneration concomitant with the expression of mineralization markers such as OPN, DMP1, or ALP.

These results suggest the importance of vascularization in bone regeneration. In fact, upon implantation in vivo, a major challenge is the maintenance of cell viability in the bone graft core, which critically depends on rapid invasion by host blood vessels. A functionally perfused vascular network will ensure oxygen and nutrient transport and waste removal. In this regard, endothelial cells play a key role in tissue regeneration and remodeling, since they can facilitate the recruitment of osteoprogenitors and immune cells, through the secretion of osteogenic factors, such as BMPs [194]. A recent study, with involvement of our research group, has shown implantation of a pre-vascularized scaffold network engineered in vitro to be a promising strategy for promoting blood supply deep into the graft, relying on inosculation with the host vasculature [195]. In particular, it has demonstrated the importance of grafting a mature microvascular network, displaying perivascular recruitment through the PDGF-BB pathway and basement membrane remodeling, taking advantage of the angiogenic properties of DPSCs/SHEDs and allowing self-assembly of endothelial cells into capillaries. Interestingly, we previously reported that the subcutaneous implantation of tissue-engineered constructs seeded with DPSCs/SHEDs primed with FGF-2, greatly enhanced vascularization within constructs thanks to the capacity of DPSCs/SHEDs to constitutively secrete hepatocyte growth factor (HGF) [170]. We have also demonstrated that FGF-2 is instrumental in promoting both VEGF and HGF secretion by DPSCs/SHEDs. Indeed, recruited stem cells participated in the deposition of vascular basement membrane

and vessel maturation in athymic nude mice demonstrated the importance of in vitro production of mature microvasculature for improving cell-based therapies [195].

#### 4.3. Human Clinical Applications of FGF-2 in the Craniofacial Area

The observations described above have been replicated in human clinical studies more recently.

Indeed, rhFGF-2 is already used in the clinical treatment of orofacial tissues. Kitamura et al. [196] performed a double-blind randomized controlled trial with 253 patients receiving rhFGF-2 0.2%, 0.3% or 0.4% or placebo during surgical management of periodontal intrabony defects. At 36 weeks, all FGF-2-treated groups demonstrated significantly higher radiographic bone fill than the placebo group, 0.3% being the best concentration. In addition, secondary analysis in a subgroup of patients showed very low levels of FGF-2 in serum and no adverse effects were reported. A randomized controlled trial of 30 patients showed an improvement in pocket depth reduction and more clinical attachment gain compared to control sites [197]. Application of FGF-2 for the treatment of intrabony defects has been studied in another randomized controlled trial, where various concentrations of FGF-2 were used mounted in  $\beta$ -tricalcium phosphate scaffolds. At 6 months, patients treated with 0.3% or 0.4% rhFGF-2 showed 71% success for the combined outcome of attachment gain of 1.5 mm and bone fill of 2.5 mm compared to 45% success in the 0.1% FGF-2 and control groups [198]. A meta-analysis of studies using recombinant human FGF-2 for the treatment of deep intrabony periodontal defects demonstrated a clinical benefit of FGF-2 in terms of bone fill [199]. A more recent meta-analysis of six randomized controlled trials shows that administration of 0.4% rhFGF-2 yielded 22% higher bone fill of periodontal defects than control treatment, though this result was not statistically significant. It also indicated that the impact of the treatment was dose dependent, with higher FGF-2 concentrations producing better bone regeneration outcomes [200]. To date, however, there is still no consensus on the optimal dose or delivery scaffolding method for the use of FGF-2 in the field of bone regeneration.

#### 4.4. FGF-2 as an Exogenous Factor: A Synthesis of Current Knowledge

It is evident from animal models and human clinical application studies that exogenous administration of FGF-2 is a promising method for accelerating craniofacial bone regeneration. Given the spatiotemporal effect of FGF-2 on mineralization, a key challenge is to determine the optimal application strategy. Parameters such as dose, length of exposure, administration mode, and type of scaffolding, as well as the origin and differentiation state of the stem cells employed for craniofacial bone tissue engineering appear to be crucial. These parameters are discussed in the following sections and summarized in Table 1.

**Table 1.** Overview of FGF-2 used in craniofacial studies arranged by research type: dose, timing, administration mode and scaffolding.

Study	Application	Dosage and Timing	Administration and/or Scaffolding	Results
in vitro studies				
Gromolak et al. 2020	Ovine bone marrow MSCs	FGF-2 (20 ng/mL) alone or in combination with BMP-2 (100 ng/mL)	In culture medium	Osteogenic differentiation induced by BMP-2 is amplified with FGF-2 supplementation. FGF-2 alone boosted proliferation of smaller cells, but without osteoblast-like structures in culture and decreased expression of osteogenic genes.

Table 1. Cont.

Study	Application	Dosage and Timing	Administration and/or Scaffolding	Results
Li et al. 2014	Murine calvarial osteoblasts	FGF-2 (1, 10, 20, 60 ng/mL)	In culture medium	Doses $\leq 10$ ng/mL yielded higher cell proliferation Doses $> 10$ ng/mL decreased proliferation Increased mineralization at all doses
Sukarawan et al. 2014	Stem cells from human exfoliated deciduous teeth (SHEDs)	FGF-2 (10 ng/mL)	In culture medium	FGF-2 maintains cell stemness
Ou et al. 2010	Murine calvarial and femur osteoprogenitor cells Human cancellous bone osteoprogenitor cells from young and old patients	rhFGF-2 (0.0016, 0.016, 0.16, or 1.6 ng/mL) 4, 24, 48, and 72 h	In culture medium	Accelerated proliferation at all doses FGF-2 induced proliferation diminished with age
Varkey et al. 2006	Rat bone marrow cells	FGF-2 (2, 10, 50 ng/mL) and BMP-2 (50, 150, 500 ng/mL) over 3 weeks	In culture medium	Accelerated mineralization at 10 ng/mL but reduced at 50 ng/mL of FGF-2 Synergistic role of FGF-2 and BMP-2 in old rat cells
Animal models				
Novais et al. 2019	Critical calvarial bone defects in nude mice	FGF (10 ng/mL) over 72 h	SHEDs in dense collagen matrices in osteogenic culture medium	Enhanced bone formation in calvarial critical size defect
Wang et al. 2019	Mandibular defects in non-human primates	FGF-2 (0.25 $\mu\text{g}/\mu\text{L}$ )	Calcium phosphate cement for BMP-2 carrier PGA gel for FGF-2 carrier	Promotion of periodontal regeneration
Anzai et al. 2016	2-wall periodontal defects in Beagle dogs	FGF-2 (3 mg/mL) vs. vehicle	Cellulose solution Direct injection in the defect	FGF-2 promoted regeneration in alveolar bone, cementum and periodontal ligament.
Charles et al. 2015	Calvarial bone defects in old mice	FGF-2 (5 ng) and BMP-2 (2 $\mu\text{g}$ )	Collagen hydroxyapatite discs	Enhanced bone filling in the central bone defect area when BMP-2 was supplemented with FGF-2
Akita et al. 2004	Calvarial defects in nude mice	FGF-2 (2.5 ng/mL) and BMP-2 (50 ng/mL) vs. FGF-2 alone, BMP-2 alone or vehicle	Transfected human MSCs in gelatin sponge carrier	Combination of FGF-2 and BMP-2 showed the most advanced bone formation within the defects
Clinical studies				
Cochran et al. 2016	Patients with periodontal intrabony defects	rhFGF-2 (0.1%, 0.3%, 0.4%) or no application	$\beta$ -TCP	Increased clinical attachment gain and bone fill at concentrations of 0.4% and 0.3%
De Santana et al. 2015	Patients with periodontal intrabony defects	rhFGF-2 (4 mg/mL) vs. no application	Sodium hyaluronate gel	Enhanced clinical parameters of wound healing compared to negative control
Kitamura et al. 2001	Patients with periodontal intrabony defects	rhFGF-2 (0.2%, 0.3%, 0.4%) or placebo	3% hydroxypropyl cellulose gel	At 36 weeks, all defects showed bone fill except placebo 0.3% dose had the best radiographic outcomes

#### 4.4.1. Dosage (In Vitro/In Vivo)

In vitro, FGF action is complex, and the biological effect of FGF-2 may depend on the dose, length and mode of exposure. Mouse bone chip outgrowth cells that were primed with FGF-2 (0, 0.0016, 0.016 or 0.16 ng/mL) demonstrated dose-dependent expression of mesenchymal markers, suggesting dose-dependent anabolic action of FGF-2 on proliferation [201]. Human mesenchyme-derived progenitor cells from cancellous bone were harvested from young and old patients and cultured under various FGF-2 concentrations (0.0016, 0.016, 0.16 and 1.6 ng/mL) for 4, 24, 28 and 72 h. Responsiveness regarding proliferation was dose and age dependent, with the proliferation rate diminishing with age [202]. Despite the inhibitory effect on differentiation and mineralization, the addition of FGF-2 to culture medium maintains stemness in SHEDs and embryonic stem cells [203], and it has been shown to be necessary to maintain the cells in a pluripotent state [204]. Indeed, continuous treatment of cultured osteoprogenitors with 10 ng/mL of rhFGF-2 over 2 days significantly reduced expression of alkaline phosphatase, osteopontin and collagen I expression, though RUNX-2 mRNA levels were not altered, indicating that cells treated with FGF-2 retain their osteogenic commitment [145]. Threshold doses vary between studies (e.g., Varkey et al. demonstrated that concentrations higher than 2 ng/mL inhibit proliferation and differentiation of bone marrow mesenchymal cells) [205].

In vivo, several studies have been conducted to assess the performance of FGF-2 in various administration modes in vivo with varying dosages depending on the animal model and bone defect configuration. A single local injection of FGF-2 directly at the site of interest has shown to increase periosteal bone formation in murine calvaria [206] and facilitate the healing of bone fracture and segmental bone defect in rats [137], rabbits [207,208], dogs [209], and non-human primates [210,211]. Kamo et al. compared the effect of a single local injection with that of cyclical injections of FGF-2 on a cancellous bone defect in the femoral condyle of rabbits. Only the high-dose single injection (1.2 µg/µL), and not the low-dose single injection (0.4 µg/µL) or cyclical injections (0.4 µg/µL, 3 times), significantly increased cancellous bone volume as measured by bone histomorphometry [212]. These results indicate that the effect of local injections of FGF-2 on cancellous bone regeneration is greater at the very early stage of bone healing. Thus, the positive effect on proliferation is useful, but only if it is temporary and regulated. A similar conclusion can be drawn from the results of Novais et al. with short FGF-2 priming of SHEDs/DPSCs before their implantation in the calvaria defect [173].

On the other hand, some animal studies have shown the efficacy of FGF-2 with a prolonged administration mode. The anabolic effect of FGF-2 was observed with daily systemic administration for 2 weeks in a rat bone defect model [213–215]. Lane et al. found that FGF-2 treatment for 14 days (1 mg/kg/rat) was associated with the development of new trabecular elements, but with the withdrawal of FGF-2 injections, the new trabeculae were rapidly lost thorough accelerated resorption [216]. Furthermore, it has been reported that continuous local infusion of bFGF using an osmotic pump is able to shorten the consolidation phase of limb lengthening in rabbits [217].

#### 4.4.2. Scaffolding and Stabilization by Administration Mode

The dose is not the only parameter to consider. The administration mode influences the stability of FGF-2. Indeed, the inherent instability of this protein in aqueous solutions necessitates its delivery via biomimetic scaffolds that can stabilize and maximize its biological activity for a defined period of time [218]. FGF-2 exhibits a short half-life of 12 h in vivo due to degradation by proteolytic enzymes but also because of its instability as soon as it is thawed [219,220]. A crucial factor for the sustained release of FGF-2 is the resorption rate of the scaffold in which the growth factor is impregnated. A well-documented method is the use of gelatin hydrogels for the fabrication of FGF-2-loaded scaffolds for tissue regeneration applications since they can mimic the manner in which FGF-2 is stored in the ECM. Indeed, when FGF-2 in solution was directly injected ectopically in mice, the vascularization process remained unchanged, whereas the incorporation of FGF-2 into

a gelatin hydrogel greatly enhanced neovascularization. Moreover, hydrogels with less water content (77.5% vs. 95.9%) were more efficacious in sustaining FGF-2 release due to their slower resorption rate [221]. Biodegradable gelatin hydrogel incorporating rhFGF-2 has been developed successfully in Japan and shown to restore bone [219,222,223].

FGF-2-mediated tissue regeneration has also been tested with chitosan/collagen scaffolding. FGF-2 controlled release by a chitosan/fucoidan complex hydrogel is presumed to immobilize it, prolong its biological half-life time and protect it from inactivation by heat or proteolysis. After injection of this hydrogel into an ectopic murine model, significant neovascularization was observed, attributed to both slow diffusion of FGF-2 and controlled biodegradation of the hydrogel [183]. Similar findings were reported with the use of heparin/protamine water-insoluble microparticles for FGF-2 delivery [224]. Various other heparin-mimicking molecules have been tested for FGF-2 administration, such as heparin mimetic peptide nanofibers [225], sulfated peptides [226], sulfonated dextrans [227] and polysulfonated polymers [228,229]. Combining poly (lactic-co-glycolic acid) (PLGA) and poly (vinyl alcohol) (PVA) to produce FGF-2-loaded microspheres has also shown to sustain the release and stability of FGF-2 for up to 4 days in a culture of human embryonic stem cells. Furthermore, FGF-2-loaded polycaprolactone (PCL) microspheres have been reported to enhance angiogenesis in vivo [230], and the use of microspheres constructed from combined alginate/collagen hydrogels yields a scaffold that provides controlled release of FGF-2 and enhances angiogenesis [231].

Radomsky et al. showed that a single local injection of FGF-2 in a hyaluronan gel, an ECM component, significantly promoted fracture healing of the fibulae in baboons, as evidenced by increased callus formation and mechanical strength [211]. Tabata et al. reported that FGF-2 incorporated into gelatin hydrogel induced bone formation at the site of a skull defect in non-human primates [184]. More recently, Murahashi et al. [232] have developed multi-layered FGF-2-loaded poly L-lactic acid nanosheets. Their subcutaneous application allowed the sustained release of loaded rhFGF-2 for about 2 weeks and enhanced bone regeneration upon implantation at critical size femoral murine defects. Controlling rhFGF-2 stability and delivery will make it possible to adapt the dose and length of exposure necessary to the bone defect to be repaired.

#### 4.4.3. BMP-2: An Interesting Cytokine for Combined Treatments

FGF action is complex, and the biological effect of FGF-2 may depend on its interaction with other cytokines. The combination of FGF-2 and BMP-2 has already been tested at various concentrations and kinetics actions. A recent in vitro study suggests that sheep BMSCs supplemented with 20 ng/mL of FGF-2 and 100 ng/mL of BMP-2 may be a feasible cellular therapy for bone regeneration [233]. In vivo, the association of 10 µg of FGF-2 and 10 µg of BMP-2 in transfected human MSCs yielded a significantly increased bone regeneration of critical size calvarial defects of nude mice [234]. Indeed, implanted calcium phosphate ceramic tubes loaded with rat marrow MSCs preconditioned with both FGF-2 and BMP-2 yielded better bone formation than FGF-2 or BMP-2 treatment alone [235]. In addition, a novel biomimetic coating scaffold (calcium phosphate/polyelectrolyte multi-layer (bCaP-PEM)) capable of sequential delivery indicated that FGF-2 delivery followed by BMP-2 increased bone regeneration in adult mouse calvarial bone defects more than delivery of BMP2 alone [201]. The combined action of FGF-2 and BMP-2 is also dose dependent. Notably, scaffolds loaded with 2 µg of BMP-2 on collagen disks enhanced osteoinduction when FGF-2 was in the range of 16–400 ng but inhibited with 10 or 50 µg of FGF-2 [236]. Similarly, using implants in rats, Takita et al. [237] found that 100 ng of FGF-2 was able to enhance BMP-2 (0.8 µg), inducing ectopic bone formation, whereas a higher dose of FGF-2 (10 µg) exerted an inhibitory effect. Higher doses may keep cells in an undifferentiated state for a longer time, this negatively impacting mineralization [173]. A possible mode of action is that high doses of FGF-2 do not increase the expression of BMP receptors, whereas low doses of FGF-2 strengthen bone formation via BMP-2 signaling as



shown by increased Smad 1 expression, a major downstream effector in the BMP signaling pathway [238].

## 5. Discussion, Conclusions and Perspectives

Bone is a constantly evolving mineralized and vascularized connective tissue that can be repaired by simple immobilization in the case of non-displaced fractures. During major trauma, infection or cancer; however, bone loses its capacity to self-repair.

Tissue engineering is considered a therapy of the future, making it possible to clinically overcome the many limitations of current autologous graft therapies (notably, the limited quantity of tissue available and risk associated with several surgical sites on a single patient). The combination of biomaterials, cells and molecules of interest may allow great advances at the clinical level by stimulating the integration of grafts through vascularization and mineralization. The close relationship between blood vessels and bone cells has been demonstrated in studies on skeletal malformation, such as craniofacial dysmorphism.

FGF-2, by virtue of its proliferative, pro-angiogenic and pro-osteogenic properties, is one of the molecules studied in this line of research. This growth factor is a ubiquitous molecule present from the embryonic stage and throughout life and is involved in the formation of the ECM. It plays an important role in the homeostasis, repair and metabolism of bone tissue by regulating the proliferation and differentiation of osteoblasts, accelerating the healing of fractures and the repair of skeletal defects.

The effects of FGF-2 differ depending on the type and stage of cell differentiation. Some results appear to be contradictory but can be explained by differences in dose, mode of administration and lengths of exposure, as well as by the models used in *in vitro* or *in vivo* studies, each of which has certain biases.

It appears that high dose and/or long-term treatment inhibit bone regeneration. This could be explained by a positive effect on proliferation, which remains essential at the beginning, but must be temporary, and therefore regulated. Thus, low dose and/or short-term treatment may provide the best conditions for bone regeneration. There is a need to explore further the positive or negative effects on bone regeneration of other parameters, such as the type of culture medium used and any supplements added to boost the osteogenic effect (ascorbic acid, dexamethasone,  $\beta$ -GP, etc.). The way in which FGF-2 is delivered should also be considered, since it may affect bone repair, potentially leading to unwanted side effects. Moreover, the delivery mode seems to have a non-negligible impact on the stability of this cytokine, and therefore, must be carefully planned and tested before clinical use. Identifying the ideal dose and how to deliver it over a given time within a specific repair time frame remain the key challenges in this type of tissue-engineered therapy.

**Author Contributions:** A.N.: investigation, writing, and visualization; E.C.: Investigation, and writing; C.C.: supervision, and writing—review; and C.G.: investigation, and writing—original draft and review. All authors have read and agreed to the published version of the manuscript.

**Funding:** From Université de Paris, ANR PulpCell and Fondation Gueules Cassées for URP2496, and Fondation pour la Recherche Médicale (PhD grant to EC).


**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Conflicts of Interest:** The authors declare no conflict of interest.

Appendix A



**Figure A1.** Morphology of cortical and trabecular bone—The cortical bone is dense and compact with penetrating vascular canals, that has slow turnover rate and high resistance to torsional and bending forces. It constitutes 80% of the skeleton, and it makes up the outer part of skeletal structures. This bone has an outer periosteal surface containing blood vessels, nerve endings, osteoblasts, and osteoclasts and an inner endosteal surface adjacent to the marrow. This tissue is arranged in osteons (A), which are concentric layers, composed of collagen fibers. These are made up of three helical chains and combine to form fibrils, which are interwoven and bound by crosslinks, providing bone elasticity, flexibility, and tensile strength. Cortical bone provides mechanical strength and protection, and it may also participate in metabolic responses, particularly when there is a long-lasting mineral deficit. In contrast, trabecular bone represents just 20% of the skeletal mass, but 80% of the bone surface. This type of bone, which is less dense and more elastic, has a higher turnover rate than cortical bone and has high resistance to compression. It provides mechanical support, helping to maintain skeletal strength and integrity with its rods and plates aligned in a pattern that provides maximal strength. It also exhibits greater metabolic activity than cortical bone, having a larger surface area for mineral exchange. These properties explain it being found inside the long bones, throughout the bodies of the vertebrae, and in the inner portions of the pelvis and other flat bones.

**Table A1.** Comparison of endochondral and intramembranous ossification

	Endochondral Ossification	Intramembranous Ossification
Process	Replacement of cartilage with bone	Direct conversion of mesenchyme to bone
Site	Mainly long bones	Mainly flat bones
Cellular embryonic origin	Neural crest-derived mesenchymal cells	Mesoderm-derived mesenchymal cells
Functional cells	Chondrocytes that secrete ECM to form cartilage	Osteoblasts that secrete osteoid matrix

There are two different processes of bone modeling: endochondral ossification, involving cartilage as an intermediate stage, and intramembranous ossification, the chondrocyte-to-osteoblast trans-differentiation, which involves direct differentiation of MSCs into osteoblasts. The former is typical for long bones, while the latter is the main mechanism for the development of flat bones, such as those of the craniofacial skeleton [239,240].

**Table A2.** Molecules with angiogenic and osteogenic effects.

	Angiogenic Effect	Osteogenic Effect	Function
BMP	Yes (indirectly) [241]	Yes [242]	Differentiation of osteoblast-like cells chemoattractant for neighboring endothelial cells (ECs) [241,243,244]; proliferation and differentiation of mesenchymal osteoprogenitors [245]; BMP-2 and -7 enhance bone formation and repair, through induction of vascular endothelial growth factor (VEGF) expression and angiogenesis stimulation [246,247]
TGF- $\beta$	Yes, but role not well understood [248]	Yes [247]	Chemoattractant for mesenchymal stem cells, differentiation of osteoblasts [249–251], chondroblast, and osteoprogenitor cells and matrix production stimulation in healing process [252,253]
PDGF	Yes	Yes	Chemoattractant for and mitogenic stimulation of osteoblasts [251,254]
MMP	Yes	Yes	Matrix metalloproteinase 9 (MMP9) induces vascularization in bone formation by the release of VEGF from ECM [255]; while matrix metalloproteinase 13 (MMP13) induces osteogenesis [256]
Notch signaling	Yes	Yes	Notch pathway modulates the angiogenic effect of VEGF in ECs. Activation of Notch signaling in bone ECs promotes local angiogenesis and osteogenesis [257].

## References

- Clarke, B. Normal Bone Anatomy and Physiology. *Clin. J. Am. Soc. Nephrol.* **2008**, *3*, S131–S139. [[CrossRef](#)] [[PubMed](#)]
- DiGirolamo, D.J.; Clemens, T.L.; Kousteni, S. The skeleton as an endocrine organ. *Nat. Rev. Rheumatol.* **2012**, *8*, 674–683. [[CrossRef](#)] [[PubMed](#)]
- Oldknow, K.J.; Macrae, V.E.; Farquharson, C. Endocrine role of bone: Recent and emerging perspectives beyond osteocalcin. *J. Endocrinol.* **2015**, *225*, R1–R19. [[CrossRef](#)] [[PubMed](#)]
- Kenkre, J.S.; Bassett, J.H. The bone remodelling cycle. *Ann. Clin. Biochem. Int. J. Lab. Med.* **2018**, *55*, 308–327. [[CrossRef](#)] [[PubMed](#)]
- Hadjidakis, D.J.; Androulakis, I.I. Bone remodeling. *Ann. N. Y. Acad. Sci.* **2006**, *1092*, 385–396. [[CrossRef](#)] [[PubMed](#)]
- Alford, A.I.; Kozloff, K.M.; Hankenson, K.D. Extracellular matrix networks in bone remodeling. *Int. J. Biochem. Cell Biol.* **2015**, *65*, 20–31. [[CrossRef](#)]
- Yang, G.; Zhu, L.; Hou, N.; Lan, Y.; Wu, X.-M.; Zhou, B.; Teng, Y.; Yang, X. Osteogenic fate of hypertrophic chondrocytes. *Cell Res.* **2014**, *24*, 1266–1269. [[CrossRef](#)] [[PubMed](#)]
- Matic, I.; Matthews, B.G.; Wang, X.; Dymont, N.A.; Worthley, D.L.; Rowe, D.W.; Grcevic, D.; Kalajzic, I. Quiescent Bone Lining Cells Are a Major Source of Osteoblasts During Adulthood. *Stem Cells* **2016**, *34*, 2930–2942. [[CrossRef](#)]
- Murshed, M.; Harmey, D.; Millán, J.L.; McKee, M.D.; Karsenty, G. Unique coexpression in osteoblasts of broadly expressed genes accounts for the spatial restriction of ECM mineralization to bone. *Genes Dev.* **2005**, *19*, 1093–1104. [[CrossRef](#)]
- Katsimbri, P. The biology of normal bone remodelling. *Eur. J. Cancer Care* **2017**, *26*, e12740. [[CrossRef](#)]
- Franz-Odenaal, T.A.; Hall, B.K.; Witten, P.E. Buried alive: How osteoblasts become osteocytes. *Dev. Dyn.* **2005**, *235*, 176–190. [[CrossRef](#)]
- Bonewald, L.F. The amazing osteocyte. *J. Bone Miner. Res.* **2011**, *26*, 229–238. [[CrossRef](#)]
- Xiao, W.; Wang, Y.; Pacios, S.; Li, S.; Graves, D.T. Cellular and Molecular Aspects of Bone Remodeling. *Craniofac. Sutures* **2015**, *18*, 9–16. [[CrossRef](#)]
- Bonewald, L.F.; Johnson, M.L. Osteocytes, mechanosensing and Wnt signaling. *Bone* **2008**, *42*, 606–615. [[CrossRef](#)]
- Dallas, S.L.; Prideaux, M.; Bonewald, L.F. The Osteocyte: An Endocrine Cell . . . and More. *Endocr. Rev.* **2013**, *34*, 658–690. [[CrossRef](#)] [[PubMed](#)]
- Lanyon, L.E. Osteocytes, strain detection, bone modeling and remodeling. *Calcif. Tissue Int.* **1993**, *53*, S102–S107. [[CrossRef](#)]
- Sims, A.N.; Martin, T.J. Coupling the activities of bone formation and resorption: A multitude of signals within the basic multicellular unit. *Bonekey Rep.* **2014**, *3*, 481. [[CrossRef](#)] [[PubMed](#)]
- Teitelbaum, S.L. Bone Resorption by Osteoclasts. *Science* **2000**, *289*, 1504–1508. [[CrossRef](#)] [[PubMed](#)]
- Mackie, E.J.; Tatarczuch, L.; Mirams, M. The skeleton: A multi-functional complex organ. The growth plate chondrocyte and endochondral ossification. *J. Endocrinol.* **2011**, *211*, 109–121. [[CrossRef](#)] [[PubMed](#)]
- Brodsky, B.; Persikov, A.V. Molecular Structure of the Collagen Triple Helix. *Adv. Protein Chem.* **2005**, *70*, 301–339. [[CrossRef](#)] [[PubMed](#)]
- Ducy, P.; Desbois, C.; Boyce, B.; Pinero, G.; Story, B.; Dunstan, C.; Smith, E.; Bonadio, J.; Goldstein, S.; Gundberg, C.; et al. Increased bone formation in osteocalcin-deficient mice. *Nat. Cell Biol.* **1996**, *382*, 448–452. [[CrossRef](#)]
- Luo, G.; Ducy, P.; McKee, M.D.; Pinero, G.J.; Loyer, E.; Behringer, R.R.; Karsenty, G. Spontaneous calcification of arteries and cartilage in mice lacking matrix GLA protein. *Nat. Cell Biol.* **1997**, *386*, 78–81. [[CrossRef](#)] [[PubMed](#)]

23. Burr, D.; Schaffler, M.; Yang, K.; Wu, D.; Lukoschek, M.; Kandzari, D.; Sivaneri, N.; Blaha, J.; Radin, E. The effects of altered strain environments on bone tissue kinetics. *Bone* **1989**, *10*, 215–221. [[CrossRef](#)]
24. Krahl, H.; Michaelis, U.; Pieper, H.G.; Quack, G.; Montag, M. Stimulation of bone growth through sports. A radiologic investigation of the upper extremities in professional tennis players. *Am. J. Sports Med.* **1994**, *22*, 751–757. [[CrossRef](#)] [[PubMed](#)]
25. Feng, X.; McDonald, J.M. Disorders of Bone Remodeling. *Annu. Rev. Pathol. Mech. Dis.* **2011**, *6*, 121–145. [[CrossRef](#)]
26. Kobayashi, S.; Takahashi, H.; Ito, A.; Saito, N.; Nawata, M.; Horiuchi, H.; Ohta, H.; Iorio, R.; Yamamoto, N.; Takaoka, K. Trabecular minimodeling in human iliac bone. *Bone* **2003**, *32*, 163–169. [[CrossRef](#)]
27. Frost, H.M. Skeletal structural adaptations to mechanical usage (SATMU): Mechanical influences on intact fibrous tissues. *Anat. Rec. Adv. Integr. Anat. Evol. Biol.* **1990**, *226*, 433–439. [[CrossRef](#)]
28. Manolagas, S.C. Birth and death of bone cells: Basic regulatory mechanisms and implications for the pathogenesis and treatment of osteoporosis. *Endocr. Rev.* **2000**, *21*, 115–137. [[PubMed](#)]
29. Takayanagi, H. Osteoimmunology: Shared mechanisms and crosstalk between the immune and bone systems. *Nat. Rev. Immunol.* **2007**, *7*, 292–304. [[CrossRef](#)] [[PubMed](#)]
30. Boyle, W.J.; Simonet, W.S.; Lacey, D.L. Osteoclast differentiation and activation. *Nat. Cell Biol.* **2003**, *423*, 337–342. [[CrossRef](#)] [[PubMed](#)]
31. Hsu, H.; Lacey, D.L.; Dunstan, C.R.; Solovyev, I.; Colombero, A.; Timms, E.; Tan, H.-L.; Elliott, G.; Kelley, M.J.; Sarosi, I.; et al. Tumor necrosis factor receptor family member RANK mediates osteoclast differentiation and activation induced by osteoprotegerin ligand. *Proc. Natl. Acad. Sci. USA* **1999**, *96*, 3540–3545. [[CrossRef](#)]
32. Kong, Y.-Y.; Yoshida, H.; Sarosi, I.; Tan, H.-L.; Timms, E.; Capparelli, C.; Morony, S.; Oliveira-Dos-Santos, A.J.; Van, G.; Itie, A.; et al. OPG is a key regulator of osteoclastogenesis, lymphocyte development and lymph-node organogenesis. *Nat. Cell Biol.* **1999**, *397*, 315–323. [[CrossRef](#)] [[PubMed](#)]
33. Belibasakis, G.N.; Bostanci, N. The RANKL-OPG system in clinical periodontology. *J. Clin. Periodontol.* **2012**, *39*, 239–248. [[CrossRef](#)]
34. Simonet, W.; Lacey, D.; Dunstan, C.; Kelley, M.; Chang, M.-S.; Lüthy, R.; Nguyen, H.; Wooden, S.; Bennett, L.; Boone, T.; et al. Osteoprotegerin: A Novel Secreted Protein Involved in the Regulation of Bone Density. *Cell* **1997**, *89*, 309–319. [[CrossRef](#)]
35. Hofbauer, L.C.; Khosla, S.; Dunstan, C.R.; Lacey, D.L.; Boyle, W.J.; Riggs, B.L. The Roles of Osteoprotegerin and Osteoprotegerin Ligand in the Paracrine Regulation of Bone Resorption. *J. Bone Miner. Res.* **2000**, *15*, 2–12. [[CrossRef](#)] [[PubMed](#)]
36. Takahashi, N.; Maeda, K.; Ishihara, A.; Uehara, S.; Kobayashi, Y. Regulatory mechanism of osteoclastogenesis by RANKL and Wnt signals. *Front. Biosci.* **2011**, *16*, 21–30. [[CrossRef](#)] [[PubMed](#)]
37. Day, T.F.; Guo, X.; Garrett-Beal, L.; Yang, Y. Wnt/beta-catenin signaling in mesenchymal progenitors controls osteoblast and chondrocyte differentiation during vertebrate skeletogenesis. *Dev. Cell* **2005**, *8*, 739–750. [[CrossRef](#)] [[PubMed](#)]
38. Glass, D.A., II.; Bialek, P.; Ahn, J.D.; Starbuck, M.; Patel, M.S.; Clevers, H.; Taketo, M.M.; Long, F.; McMahon, A.P.; Lang, R.A.; et al. Canonical Wnt signaling in differentiated osteoblasts controls osteoclast differentiation. *Dev. Cell* **2005**, *8*, 751–764. [[CrossRef](#)] [[PubMed](#)]
39. Daoussis, D.; Andonopoulos, A.P. The Emerging Role of Dickkopf-1 in Bone Biology: Is It the Main Switch Controlling Bone and Joint Remodeling? *Semin. Arthritis Rheum.* **2011**, *41*, 170–177. [[CrossRef](#)]
40. Caetano-Lopes, J.; Canhão, H.; Fonseca, J.E. Osteoblasts and bone formation. *Acta Reumatol. Port.* **2007**, *32*, 103–110.
41. Carter, P.H.; Schipani, E. The roles of parathyroid hormone and calcitonin in bone remodeling: Prospects for novel therapeutics. *Endocr. Metab. Immune Disord. Drug Targets* **2006**, *6*, 59–76. [[CrossRef](#)]
42. Bassett, J.H.D.; Williams, G.R. Role of Thyroid Hormones in Skeletal Development and Bone Maintenance. *Endocr. Rev.* **2016**, *37*, 135–187. [[CrossRef](#)] [[PubMed](#)]
43. Iglesias, L.; Yeh, J.K.; Castro-Magana, M.; Aloia, J.F. Effects of growth hormone on bone modeling and remodeling in hypophysectomized young female rats: A bone histomorphometric study. *J. Bone Miner. Metab.* **2010**, *29*, 159–167. [[CrossRef](#)] [[PubMed](#)]
44. Weinstein, R.S.; Jilka, R.L.; Parfitt, A.M.; Manolagas, S.C. Inhibition of osteoblastogenesis and promotion of apoptosis of osteoblasts and osteocytes by glucocorticoids. Potential mechanisms of their deleterious effects on bone. *J. Clin. Investig.* **1998**, *102*, 274–282. [[CrossRef](#)]
45. Nakamura, T.; Imai, Y.; Matsumoto, T.; Sato, S.; Takeuchi, K.; Igarashi, K.; Harada, Y.; Azuma, Y.; Krust, A.; Yamamoto, Y.; et al. Estrogen Prevents Bone Loss via Estrogen Receptor  $\alpha$  and Induction of Fas Ligand in Osteoclasts. *Cell* **2007**, *130*, 811–823. [[CrossRef](#)]
46. Roodman, G.D. Role of cytokines in the regulation of bone resorption. *Calcif. Tissue Int.* **1993**, *53*, S94–S98. [[CrossRef](#)] [[PubMed](#)]
47. Graves, D.T.; Oates, T.; Garlet, G.P. Review of osteoimmunology and the host response in endodontic and periodontal lesions. *J. Oral Microbiol.* **2011**, *3*, 3. [[CrossRef](#)] [[PubMed](#)]
48. Raisz, L. Prostaglandins and bone: Physiology and pathophysiology. *Osteoarthr. Cartil.* **1999**, *7*, 419–421. [[CrossRef](#)] [[PubMed](#)]
49. Brandi, M.L.; Collin-Osdoby, P. Vascular Biology and the Skeleton. *J. Bone Miner. Res.* **2006**, *21*, 183–192. [[CrossRef](#)]
50. Fang, T.D.; Salim, A.; Xia, W.; Nacamuli, R.P.; Guccione, S.; Song, H.M.; Carano, R.A.; Filvaroff, E.H.; Bednarski, M.D.; Giaccia, A.J.; et al. Angiogenesis Is Required for Successful Bone Induction During Distraction Osteogenesis. *J. Bone Miner. Res.* **2005**, *20*, 1114–1124. [[CrossRef](#)] [[PubMed](#)]

51. Seebach, C.; Henrich, D.; Wilhelm, K.; Barker, J.H.; Marzi, I. Endothelial Progenitor Cells Improve Directly and Indirectly Early Vascularization of Mesenchymal Stem Cell-Driven Bone Regeneration in a Critical Bone Defect in Rats. *Cell Transplant.* **2012**, *21*, 1667–1677. [[CrossRef](#)]
52. Carvalho, R.; Einhorn, T.; Lehmann, W.; Edgar, C.; Al-Yamani, A.; Apazidis, A.; Pacicca, D.; Clemens, T.; Gerstenfeld, L. The role of angiogenesis in a murine tibial model of distraction osteogenesis. *Bone* **2004**, *34*, 849–861. [[CrossRef](#)]
53. Percival, C.J.; Richtsmeier, J.T. Angiogenesis and intramembranous osteogenesis. *Dev. Dyn.* **2013**, *242*, 909–922. [[CrossRef](#)] [[PubMed](#)]
54. Maes, C.; Clemens, T.L. Angiogenic–osteogenic coupling: The endothelial perspective. *Bonekey Rep.* **2014**, *3*, 578. [[CrossRef](#)]
55. Schipani, E.; Maes, C.; Carmeliet, G.; Semenza, G.L. Regulation of Osteogenesis–Angiogenesis Coupling by HIFs and VEGF. *J. Bone Miner. Res.* **2009**, *24*, 1347–1353. [[CrossRef](#)] [[PubMed](#)]
56. Sivaraj, K.K.; Adams, R.H. Blood vessel formation and function in bone. *Development* **2016**, *143*, 2706–2715. [[CrossRef](#)] [[PubMed](#)]
57. Joško, J.; Gwóźdź, B.; Jedrzejowska-Szypułka, H.; Hendryk, S. Vascular endothelial growth factor (VEGF) and its effect on angiogenesis. *Med. Sci. Monit.* **2001**, *6*, 1047–1052.
58. Bluteau, G.; Julien, M.; Magne, D.; Mallein-Gerin, F.; Weiss, P.; Daculsi, G.; Guicheux, J. VEGF and VEGF receptors are differentially expressed in chondrocytes. *Bone* **2007**, *40*, 568–576. [[CrossRef](#)] [[PubMed](#)]
59. Hu, K.; Olsen, B.R. Osteoblast-derived VEGF regulates osteoblast differentiation and bone formation during bone repair. *J. Clin. Investig.* **2016**, *126*, 509–526. [[CrossRef](#)] [[PubMed](#)]
60. Bouletreau, P.J.; Warren, S.M.; Spector, J.A.; Peled, Z.M.; Gerrets, R.P.; Greenwald, J.A.; Longaker, M.T. Hypoxia and VEGF Up-Regulate BMP-2 mRNA and Protein Expression in Microvascular Endothelial Cells: Implications for Fracture Healing. *Plast. Reconstr. Surg.* **2002**, *109*, 2384–2397. [[CrossRef](#)] [[PubMed](#)]
61. Mayr-Wohlfart, U.; Waltenberger, J.; Hausser, H.; Kessler, S.; Günther, K.-P.; Dehio, C.; Puhl, W.; Brenner, R. Vascular endothelial growth factor stimulates chemotactic migration of primary human osteoblasts. *Bone* **2002**, *30*, 472–477. [[CrossRef](#)]
62. Henriksen, K.; Karsdal, M.; Delaissé, J.M.; Engsig, M.T. RANKL and vascular endothelial growth factor (VEGF) induce osteoclast chemotaxis through an ERK1/2-dependent mechanism. *J. Biol. Chem.* **2003**, *278*, 48745–48753. [[CrossRef](#)]
63. Carlevaro, M.F.; Cermelli, S.; Cancedda, R.; Cancedda, F.D. Vascular endothelial growth factor (VEGF) in cartilage neovascularization and chondrocyte differentiation: Auto-paracrine role during endochondral bone formation. *J. Cell Sci.* **2000**, *113*, 59–69.
64. Street, J.; Bao, M.; DeGuzman, L.; Bunting, S.; Peale, F.V., Jr.; Ferrara, N.; Steinmetz, H.; Hoeffel, J.; Cleland, J.L.; Daugherty, A.; et al. Vascular endothelial growth factor stimulates bone repair by promoting angiogenesis and bone turnover. *Proc. Natl. Acad. Sci. USA* **2002**, *99*, 9656–9661. [[CrossRef](#)]
65. Kim, Y.-M.; Lee, Y.M.; Kim, H.-S.; Kim, J.D.; Choi, Y.; Kim, K.-W.; Lee, S.-Y.; Kwon, Y.-G. TNF-related Activation-induced Cytokine (TRANCE) Induces Angiogenesis through the Activation of Src and Phospholipase C (PLC) in Human Endothelial Cells. *J. Biol. Chem.* **2002**, *277*, 6799–6805. [[CrossRef](#)] [[PubMed](#)]
66. Min, J.K.; Kim, Y.M.; Kim, Y.M.; Kim, E.C.; Gho, Y.S.; Kang, I.J.; Lee, S.Y.; Kong, Y.Y.; Kwon, Y.G. Vascular endothelial growth factor up-regulates expression of receptor activator of NF-kappa B (RANK) in endothelial cells. Concomitant increase of angiogenic responses to RANK ligand. *J. Biol. Chem.* **2003**, *278*, 39548–39557. [[CrossRef](#)]
67. Kim, H.-H.; Shin, H.S.; Kwak, H.J.; Ahn, K.Y.; Kim, J.-H.; Lee, H.J.; Lee, M.-S.; Lee, Z.H.; Koh, G.Y. RANKL regulates endothelial cell survival through the phosphatidylinositol 3'-kinase/Akt signal transduction pathway. *FASEB J.* **2003**, *17*, 1–17. [[CrossRef](#)]
68. Ulici, V.; Hoenselaar, K.D.; Agoston, H.; McErlain, D.D.; Umoh, J.; Chakrabarti, S.; Holdsworth, D.W.; Beier, F. The role of Akt1 in terminal stages of endochondral bone formation: Angiogenesis and ossification. *Bone* **2009**, *45*, 1133–1145. [[CrossRef](#)] [[PubMed](#)]
69. Aragónés, J.; Fraisl, P.; Baes, M.; Carmeliet, P. Oxygen Sensors at the Crossroad of Metabolism. *Cell Metab.* **2009**, *9*, 11–22. [[CrossRef](#)]
70. Wang, Y.; Wan, C.; Deng, L.; Liu, X.; Cao, X.; Gilbert, S.R.; Bouxsein, M.L.; Faugere, M.-C.; Guldberg, R.E.; Gerstenfeld, L.C.; et al. The hypoxia-inducible factor  $\alpha$  pathway couples angiogenesis to osteogenesis during skeletal development. *J. Clin. Investig.* **2007**, *117*, 1616–1626. [[CrossRef](#)]
71. Riddle, R.C.; Khatri, R.; Schipani, E.; Clemens, T.L. Role of hypoxia-inducible factor-1 $\alpha$  in angiogenic–osteogenic coupling. *J. Mol. Med.* **2009**, *87*, 583–590. [[CrossRef](#)] [[PubMed](#)]
72. Ornitz, D.M. FGFs, heparan sulfate and FGFRs: Complex interactions essential for development. *BioEssays* **2000**, *22*, 108–112. [[CrossRef](#)]
73. Collin-Osdoby, P.; Rothe, L.; Bekker, S.; Anderson, F.; Huang, Y.; Osdoby, P. Basic fibroblast growth factor stimulates osteoclast recruitment, development, and bone pit resorption in association with angiogenesis in vivo on the chick chorioallantoic membrane and activates isolated avian osteoclast resorption in vitro. *J. Bone Miner. Res.* **2002**, *17*, 1859–1871. [[CrossRef](#)]
74. De Moerloose, L.; Spencer-Dene, B.; Revest, J.M.; Hajihosseini, M.; Rosewell, I.; Dickson, C. An important role for the IIIb isoform of fibroblast growth factor receptor 2 (FGFR2) in mesenchymal-epithelial signalling during mouse organogenesis. *Dev.* **2000**, *127*, 483–492.
75. Miraoui, H.; Marie, P.J. Fibroblast Growth Factor Receptor Signaling Crosstalk in Skeletogenesis. *Sci. Signal.* **2010**, *3*, re9. [[CrossRef](#)]
76. Marie, P.J.; Miraoui, H.; Severe, N. FGF/FGFR signaling in bone formation: Progress and perspectives. *Growth Factors* **2012**, *30*, 117–123. [[CrossRef](#)] [[PubMed](#)]

77. Kim, J.; Park, J.-C.; Kim, S.-H.; Im, G.-I.; Kim, B.-S.; Lee, J.-B.; Choi, E.-Y.; Song, J.-S.; Cho, K.-S.; Kim, C.-S. Treatment of FGF-2 on stem cells from inflamed dental pulp tissue from human deciduous teeth. *Oral Dis.* **2014**, *20*, 191–204. [[CrossRef](#)]
78. Hatch, N.E. FGF Signaling in Craniofacial Biological Control and Pathological Craniofacial Development. *Crit. Rev. Eukaryot. Gene Expr.* **2010**, *20*, 295–311. [[CrossRef](#)]
79. Ornitz, D.M.; Itoh, N. The Fibroblast Growth Factor signaling pathway. *Wiley Interdiscip. Rev. Dev. Biol.* **2015**, *4*, 215–266. [[CrossRef](#)] [[PubMed](#)]
80. Turner, N.; Grose, R. Fibroblast growth factor signalling: From development to cancer. *Nat. Rev. Cancer* **2010**, *10*, 116–129. [[CrossRef](#)]
81. Eswarakumar, V.; Lax, I.; Schlessinger, J. Cellular signaling by fibroblast growth factor receptors. *Cytokine Growth Factor Rev.* **2005**, *16*, 139–149. [[CrossRef](#)]
82. Li, X.; Wang, C.; Xiao, J.; McKeehan, W.L.; Wang, F. Fibroblast growth factors, old kids on the new block. *Semin. Cell Dev. Biol.* **2016**, *53*, 155–167. [[CrossRef](#)]
83. Potthoff, J.M.; Kliewer, S.A.; Mangelsdorf, D.J. Endocrine fibroblast growth factors 15/19 and 21: From feast to famine. *Genes Dev.* **2012**, *26*, 312–324. [[CrossRef](#)] [[PubMed](#)]
84. Tomlinson, E.; Fu, L.; John, L.; Hultgren, B.; Huang, X.; Renz, M.; Stephan, J.P.; Tsai, S.P.; Powell-Braxton, L.; French, D.; et al. Transgenic mice expressing human fibroblast growth factor-19 display increased metabolic rate and decreased adiposity. *Endocrinology* **2002**, *143*, 1741–1747. [[CrossRef](#)] [[PubMed](#)]
85. Smith, E.R.; McMahan, L.P.; Holt, S.G. Fibroblast growth factor. *Ann. Clin. Biochem. Int. J. Lab. Med.* **2014**, *51*, 203–227. [[CrossRef](#)]
86. Belov, A.A.; Mohammadi, M. Molecular Mechanisms of Fibroblast Growth Factor Signaling in Physiology and Pathology. *Cold Spring Harb. Perspect. Biol.* **2013**, *5*, a015958. [[CrossRef](#)]
87. Ornitz, D.M.; Marie, P.J. Fibroblast growth factor signaling in skeletal development and disease. *Genes Dev.* **2015**, *29*, 1463–1486. [[CrossRef](#)] [[PubMed](#)]
88. Yayon, A.; Klagsbrun, M.; Esko, J.D.; Leder, P.; Ornitz, D.M. Cell surface, heparin-like molecules are required for binding of basic fibroblast growth factor to its high affinity receptor. *Cell* **1991**, *64*, 841–848. [[CrossRef](#)]
89. Rapraeger, C.A.; Krufka, A.; Olwin, B.B. Requirement of heparan sulfate for bFGF-mediated fibroblast growth and myoblast differentiation. *Science* **1991**, *252*, 1705–1708. [[CrossRef](#)]
90. Ornitz, D.M.; Yayon, A.; Flanagan, J.G.; Svahn, C.M.; Levi, E.; Leder, P. Heparin is required for cell-free binding of basic fibroblast growth factor to a soluble receptor and for mitogenesis in whole cells. *Mol. Cell. Biol.* **1992**, *12*, 240–247. [[CrossRef](#)]
91. Spivak-Kroizman, T.; Lemmon, M.; Dikic, I.; Ladbury, J.; Pinchasi, D.; Huang, J.; Jaye, M.; Crumley, G.; Schlessinger, J.; Lax, I. Heparin-induced oligomerization of FGF molecules is responsible for FGF receptor dimerization, activation, and cell proliferation. *Cell* **1994**, *79*, 1015–1024. [[CrossRef](#)]
92. Beenken, A.; Mohammadi, M. The FGF family: Biology, pathophysiology and therapy. *Nat. Rev. Drug Discov.* **2009**, *8*, 235–253. [[CrossRef](#)]
93. Matsuo, I.; Kimura-Yoshida, C. Extracellular modulation of Fibroblast Growth Factor signaling through heparan sulfate proteoglycans in mammalian development. *Curr. Opin. Genet. Dev.* **2013**, *23*, 399–407. [[CrossRef](#)] [[PubMed](#)]
94. Xu, R.; Ori, A.; Rudd, T.R.; Uniewicz, K.A.; Ahmed, Y.A.; Guimond, S.E.; Skidmore, M.A.; Siligardi, G.; Yates, E.A.; Fernig, D.G. Diversification of the structural determinants of fibroblast growth factor-heparin interactions: Implications for binding specificity. *J. Biol. Chem.* **2012**, *287*, 40061–40073. [[CrossRef](#)]
95. Goetz, R.; Mohammadi, M. Exploring mechanisms of FGF signalling through the lens of structural biology. *Nat. Rev. Mol. Cell Biol.* **2013**, *14*, 166–180. [[CrossRef](#)]
96. Hart, K.C.; Robertson, S.C.; Kanemitsu, M.Y.; Meyer, A.N.; Tynan, J.A.; Donoghue, D.J. Transformation and Stat activation by derivatives of FGFR1, FGFR3, and FGFR4. *Oncogene* **2000**, *19*, 3309–3320. [[CrossRef](#)]
97. Shao, D.; Lazar, M.A. Modulating nuclear receptor function: May the phos be with you. *J. Clin. Investig.* **1999**, *103*, 1617–1618. [[CrossRef](#)]
98. Yun, Y.-R.; Won, J.E.; Jeon, E.; Lee, S.; Kang, W.; Jo, H.; Jang, J.H.; Shin, U.S.; Kim, H.W. Fibroblast Growth Factors: Biology, Function, and Application for Tissue Regeneration. *J. Tissue Eng.* **2010**, *2010*, 218142. [[CrossRef](#)] [[PubMed](#)]
99. Nugent, M.A.; Iozzo, R.V. Fibroblast growth factor-2. *Int. J. Biochem. Cell Biol.* **2000**, *32*, 115–120. [[CrossRef](#)]
100. Johnson, L.G.; Lapadat, R. Mitogen-activated protein kinase pathways mediated by ERK, JNK, and p38 protein kinases. *Science* **2002**, *298*, 1911–1912. [[CrossRef](#)] [[PubMed](#)]
101. Taketomi, T.; Onimura, T.; Yoshiga, D.; Muratsu, D.; Sanui, T.; Fukuda, T.; Kusukawa, J.; Nakamura, S. Sprouty2 is involved in the control of osteoblast proliferation and differentiation through the FGF and BMP signaling pathways. *Cell Biol. Int.* **2017**, *42*, 1106–1114. [[CrossRef](#)]
102. Schmidt, A.; Ladage, D.; Schinköthe, T.; Klausmann, U.; Ulrichs, C.; Klinz, F.-J.; Brixius, K.; Arnhold, S.; Desai, B.; Mehlhorn, U.; et al. Basic Fibroblast Growth Factor Controls Migration in Human Mesenchymal Stem Cells. *Stem Cells* **2006**, *24*, 1750–1758. [[CrossRef](#)]
103. Jin, L.; Hu, X.; Feng, L. NT3 inhibits FGF2-induced neural progenitor cell proliferation via the PI3K/GSK3 pathway. *J. Neurochem.* **2005**, *93*, 1251–1261. [[CrossRef](#)] [[PubMed](#)]

104. Kanazawa, S.; Fujiwara, T.; Matsuzaki, S.; Shingaki, K.; Taniguchi, M.; Miyata, S.; Tohyama, M.; Sakai, Y.; Yano, K.; Hosokawa, K.; et al. bFGF Regulates PI3-Kinase-Rac1-JNK Pathway and Promotes Fibroblast Migration in Wound Healing. *PLoS ONE* **2010**, *5*, e12228. [[CrossRef](#)]
105. Choi, S.C.; Kim, S.J.; Choi, J.H.; Park, C.Y.; Shim, W.J.; Lim, D.S. Fibroblast growth factor-2 and -4 promote the proliferation of bone marrow mesenchymal stem cells by the activation of the PI3K-Akt and ERK1/2 signaling pathways. *Stem Cells Dev.* **2008**, *17*, 725–736. [[CrossRef](#)] [[PubMed](#)]
106. Yoon, H.; Kim, H.-L.; Chun, Y.-S.; Shin, D.H.; Lee, K.-H.; Shin, C.S.; Lee, D.Y.; Kim, H.-H.; Lee, Z.H.; Ryoo, H.-M.; et al. NAA10 controls osteoblast differentiation and bone formation as a feedback regulator of Runx2. *Nat. Commun.* **2014**, *5*, 5176. [[CrossRef](#)] [[PubMed](#)]
107. Lai, C.-F.; Chaudhary, L.; Fausto, A.; Halstead, L.R.; Ory, D.S.; Avioli, L.V.; Cheng, S.-L. Erk Is Essential for Growth, Differentiation, Integrin Expression, and Cell Function in Human Osteoblastic Cells. *J. Biol. Chem.* **2001**, *276*, 14443–14450. [[CrossRef](#)] [[PubMed](#)]
108. Marie, P. Fibroblast growth factor signaling controlling osteoblast differentiation. *Gene* **2003**, *316*, 23–32. [[CrossRef](#)]
109. Chaudhary, L.R.; Avioli, L.V. Extracellular-signal regulated kinase signaling pathway mediates downregulation of type I procollagen gene expression by FGF-2, PDGF-BB, and okadaic acid in osteoblastic cells. *J. Cell. Biochem.* **2000**, *76*, 354–359. [[CrossRef](#)]
110. Ge, C.; Xiao, G.; Jiang, D.; Franceschi, R.T. Critical role of the extracellular signal-regulated kinase-MAPK pathway in osteoblast differentiation and skeletal development. *J. Cell Biol.* **2007**, *176*, 709–718. [[CrossRef](#)]
111. Marshak, D.R.; Jaiswal, R.K.; Jaiswal, N.; Bruder, S.P.; Mbalaviele, G.; Pittenger, M.F. Adult Human Mesenchymal Stem Cell Differentiation to the Osteogenic or Adipogenic Lineage Is Regulated by Mitogen-activated Protein Kinase. *J. Biol. Chem.* **2000**, *275*, 9645–9652. [[CrossRef](#)] [[PubMed](#)]
112. Kratchmarova, I.; Blagoev, B.; Haack-Sorensen, M.; Kassem, M.; Mann, M. Mechanism of Divergent Growth Factor Effects in Mesenchymal Stem Cell Differentiation. *Science* **2005**, *308*, 1472–1477. [[CrossRef](#)]
113. Matsushita, T.; Chan, Y.Y.; Kawanami, A.; Balmes, G.; Landreth, G.E.; Murakami, S. Extracellular Signal-Regulated Kinase 1 (ERK1) and ERK2 Play Essential Roles in Osteoblast Differentiation and in Supporting Osteoclastogenesis. *Mol. Cell. Biol.* **2009**, *29*, 5843–5857. [[CrossRef](#)] [[PubMed](#)]
114. Park, E.-S.; Lind, A.-K.; Dahm-Kähler, P.; Brännström, M.; Carletti, M.Z.; Christenson, L.K.; Curry, T.E.; Jo, M. RUNX2 Transcription Factor Regulates Gene Expression in Luteinizing Granulosa Cells of Rat Ovaries. *Mol. Endocrinol.* **2010**, *24*, 846–858. [[CrossRef](#)] [[PubMed](#)]
115. Bundschu, K.; Knobloch, K.-P.; Ullrich, M.; Schinke, T.; Amling, M.; Engelhardt, C.M.; Renné, T.; Walter, U.; Schuh, K. Gene Disruption of Spred-2 Causes Dwarfism. *J. Biol. Chem.* **2005**, *280*, 28572–28580. [[CrossRef](#)]
116. Taniguchi, K.; Ayada, T.; Ichiyama, K.; Kohno, R.-I.; Yonemitsu, Y.; Minami, Y.; Kikuchi, A.; Maehara, Y.; Yoshimura, A. Sprouty2 and Sprouty4 are essential for embryonic morphogenesis and regulation of FGF signaling. *Biochem. Biophys. Res. Commun.* **2007**, *352*, 896–902. [[CrossRef](#)]
117. Ozaki, K.-I.; Miyazaki, S.; Tanimura, S.; Kohno, M. Efficient suppression of FGF-2-induced ERK activation by the cooperative interaction among mammalian Sprouty isoforms. *J. Cell Sci.* **2005**, *118*, 5861–5871. [[CrossRef](#)]
118. Yoshimura, A. Regulation of Cytokine Signaling by the SOCS and Spred Family Proteins. *Keio J. Med.* **2009**, *58*, 73–83. [[CrossRef](#)] [[PubMed](#)]
119. Lu, H.C.; Swindell, E.C.; Sierralta, W.D.; Eichele, G.; Thaller, C. Evidence for a role of protein kinase C in FGF signal transduction in the developing chick limb bud. *Development* **2001**, *128*, 2451–2460.
120. Mohammadi, M.; Schlessinger, J.; Hubbard, S.R. Structure of the FGF Receptor Tyrosine Kinase Domain Reveals a Novel Autoinhibitory Mechanism. *Cell* **1996**, *86*, 577–587. [[CrossRef](#)]
121. Kim, H.-J.; Kim, J.-H.; Bae, S.-C.; Choi, J.-Y.; Ryoo, H.-M. The Protein Kinase C Pathway Plays a Central Role in the Fibroblast Growth Factor-stimulated Expression and Transactivation Activity of Runx2. *J. Biol. Chem.* **2003**, *278*, 319–326. [[CrossRef](#)]
122. Osathanon, T.; Nowwarote, N.; Pavasant, P. Basic fibroblast growth factor inhibits mineralization but induces neuronal differentiation by human dental pulp stem cells through a FGFR and PLC $\gamma$  signaling pathway. *J. Cell. Biochem.* **2011**, *112*, 1807–1816. [[CrossRef](#)] [[PubMed](#)]
123. Spivak-Kroizman, T.; Mohammadi, M.; Hu, P.; Jaye, M.; Schlessinger, J.; Lax, I. Point mutation in the fibroblast growth factor receptor eliminates phosphatidylinositol hydrolysis without affecting neuronal differentiation of PC12 cells. *J. Biol. Chem.* **1994**, *269*, 14419–14423. [[CrossRef](#)]
124. Kang, S.; Elf, S.; Dong, S.; Hitosugi, T.; Lythgoe, K.; Guo, A.; Ruan, H.; Lonial, S.; Khoury, H.J.; Williams, I.R.; et al. Fibroblast Growth Factor Receptor 3 Associates with and Tyrosine Phosphorylates p90 RSK2, Leading to RSK2 Activation That Mediates Hematopoietic Transformation. *Mol. Cell. Biol.* **2009**, *29*, 2105–2117. [[CrossRef](#)] [[PubMed](#)]
125. Dailey, L.; Ambrosetti, D.; Mansukhani, A.; Basilico, C. Mechanisms underlying differential responses to FGF signaling. *Cytokine Growth Factor Rev.* **2005**, *16*, 233–247. [[CrossRef](#)]
126. Marie, P.; Coffin, J.; Hurley, M. FGF and FGFR signaling in chondrodysplasias and craniosynostosis. *J. Cell. Biochem.* **2005**, *96*, 888–896. [[CrossRef](#)] [[PubMed](#)]
127. Muenke, M.; Schell, U. Fibroblast-growth-factor receptor mutations in human skeletal disorders. *Trends Genet.* **1995**, *11*, 308–313. [[CrossRef](#)]

128. Ornitz, M.D.; Marie, P.J. FGF signaling pathways in endochondral and intramembranous bone development and human genetic disease. *Genes Dev.* **2002**, *16*, 1446–1465. [[CrossRef](#)] [[PubMed](#)]
129. Ornitz, D.M.; Itoh, N. Fibroblast growth factors. *Genome Biol.* **2001**, *2*, 3005. [[CrossRef](#)] [[PubMed](#)]
130. Coffin, J.D.; Homer-Bouthiette, C.; Hurley, M.M. Fibroblast Growth Factor 2 and Its Receptors in Bone Biology and Disease. *J. Endocr. Soc.* **2018**, *2*, 657–671. [[CrossRef](#)] [[PubMed](#)]
131. Du, E.; Xiao, L.; Hurley, M.M. FGF23 Neutralizing Antibody Ameliorates Hypophosphatemia and Impaired FGF Receptor Signaling in Kidneys of HMWFGF2 Transgenic Mice. *J. Cell. Physiol.* **2016**, *232*, 610–616. [[CrossRef](#)]
132. Yang, J.; Meyer, M.; Müller, A.-K.; Böhm, F.; Grose, R.; Dauwalder, T.; Verrey, F.; Kopf, M.; Partanen, J.; Bloch, W.; et al. Fibroblast growth factor receptors 1 and 2 in keratinocytes control the epidermal barrier and cutaneous homeostasis. *J. Cell Biol.* **2010**, *188*, 935–952. [[CrossRef](#)]
133. Sufen, G.; Xianghong, Y.; Yongxia, C.; Qian, P. bFGF and PDGF-BB have a synergistic effect on the proliferation, migration and VEGF release of endothelial progenitor cells. *Cell Biol. Int.* **2011**, *35*, 545–551. [[CrossRef](#)]
134. Hill, P.A.; Tumber, A.; Meikle, M.C. Multiple Extracellular Signals Promote Osteoblast Survival and Apoptosis. *Endocrinology* **1997**, *138*, 3849–3858. [[CrossRef](#)]
135. Nowwarote, N.; Sukarawan, W.; Pavasant, P.; Osathanon, T. Basic Fibroblast Growth Factor Regulates REX1 Expression Via IL-6 In Stem Cells Isolated From Human Exfoliated Deciduous Teeth. *J. Cell. Biochem.* **2017**, *118*, 1480–1488. [[CrossRef](#)]
136. Wang, G.; Zhang, H.; Zhao, Y.; Li, J.; Cai, J.; Wang, P.; Meng, S.; Feng, J.; Miao, C.; Ding, M.; et al. Noggin and bFGF cooperate to maintain the pluripotency of human embryonic stem cells in the absence of feeder layers. *Biochem. Biophys. Res. Commun.* **2005**, *330*, 934–942. [[CrossRef](#)]
137. Kawaguchi, H.; Kurokawa, T.; Hanada, K.; Hiyama, Y.; Tamura, M.; Ogata, E.; Matsumoto, T. Stimulation of fracture repair by recombinant human basic fibroblast growth factor in normal and streptozotocin-diabetic rats. *Endocrinology* **1994**, *135*, 774–781. [[CrossRef](#)]
138. Liang, H.; Pun, S.; Wronski, T.J. Bone anabolic effects of basic fibroblast growth factor in ovariectomized rats. *Endocrinology* **1999**, *140*, 5780–5788. [[CrossRef](#)]
139. Yamaguchi, T.P.; Rossant, J. Fibroblast growth factors in mammalian development. *Curr. Opin. Genet. Dev.* **1995**, *5*, 485–491. [[CrossRef](#)]
140. Kimelman, D.; Kirschner, M. Synergistic induction of mesoderm by FGF and TGF-beta and the identification of an mRNA coding for FGF in the early *Xenopus* embryo. *Cell* **1987**, *51*, 869–877. [[CrossRef](#)]
141. Peters, K.; Ornitz, D.; Werner, S.; Williams, L. Unique Expression Pattern of the FGF Receptor 3 Gene during Mouse Organogenesis. *Dev. Biol.* **1993**, *155*, 423–430. [[CrossRef](#)]
142. Hurley, M.M.; Abreu, C.; Gronowicz, G.; Kawaguchi, H.; Lorenzo, J. Expression and regulation of basic fibroblast growth factor mRNA levels in mouse osteoblastic MC3T3-E1 cells. *J. Biol. Chem.* **1994**, *269*, 9392–9396. [[CrossRef](#)]
143. Vagner, S.; Gensac, M.C.; Maret, A.; Bayard, F.; Amalric, F.; Prats, H.; Prats, A.C. Alternative translation of human fibroblast growth factor 2 mRNA occurs by internal entry of ribosomes. *Mol. Cell. Biol.* **1995**, *15*, 35–44. [[CrossRef](#)] [[PubMed](#)]
144. Ding, V.M.; Ling, L.; Natarajan, S.; Yap, M.G.; Cool, S.M.; Choo, A.B. FGF-2 modulates Wnt signaling in undifferentiated hESC and iPS cells through activated PI3-K/GSK3beta signaling. *J. Cell Physiol.* **2010**, *225*, 417–428. [[CrossRef](#)] [[PubMed](#)]
145. Fakhry, A.; Ratisoontorn, C.; Vedhachalam, C.; Salhab, I.; Koyama, E.; Leboy, P.; Pacifici, M.; Kirschner, R.E.; Nah, H.D. Effects of FGF-2/-9 in calvarial bone cell cultures: Differentiation stage-dependent mitogenic effect, inverse regulation of BMP-2 and noggin, and enhancement of osteogenic potential. *Bone* **2005**, *36*, 254–266. [[CrossRef](#)]
146. Homer-Bouthiette, C.; Doetschman, T.; Xiao, L.; Hurley, M.M. Knockout of Nuclear High Molecular Weight FGF2 Isoforms in Mice Modulates Bone and Phosphate Homeostasis. *J. Biol. Chem.* **2014**, *289*, 36303–36314. [[CrossRef](#)]
147. Yu, K.; Xu, J.; Liu, Z.; Susic, D.; Shao, J.; Olson, E.N.; Towler, D.A.; Ornitz, D.M. Conditional inactivation of FGF receptor 2 reveals an essential role for FGF signaling in the regulation of osteoblast function and bone growth. *Development* **2003**, *130*, 3063–3074. [[CrossRef](#)]
148. Jacob, A.L.; Smith, C.; Partanen, J.; Ornitz, D.M. Fibroblast growth factor receptor 1 signaling in the osteo-chondrogenic cell lineage regulates sequential steps of osteoblast maturation. *Dev. Biol.* **2006**, *296*, 315–328. [[CrossRef](#)]
149. Lee, C.-H.; Su, S.-Y.; Sittampalam, K.; Chen, P.C.-H.; Petersson, F.; Kao, Y.-C.; Carpenter, T.O.; Hsieh, T.-H.; Konishi, E.; Tsai, J.-W.; et al. Frequent overexpression of *klotho* in fusion-negative phosphaturic mesenchymal tumors with tumorigenic implications. *Mod. Pathol.* **2019**, *33*, 858–870. [[CrossRef](#)]
150. Fei, Y.; Xiao, L.; Doetschman, T.; Coffin, D.J.; Hurley, M.M. Fibroblast Growth Factor 2 Stimulation of Osteoblast Differentiation and Bone Formation Is Mediated by Modulation of the Wnt Signaling Pathway. *J. Biol. Chem.* **2011**, *286*, 40575–40583. [[CrossRef](#)]
151. Naganawa, T.; Xiao, L.; Abogunde, E.; Sobue, T.; Kalajzic, I.; Sabbieti, M.; Agas, D.; Hurley, M. In vivo and in vitro comparison of the effects of FGF-2 null and haplo-insufficiency on bone formation in mice. *Biochem. Biophys. Res. Commun.* **2006**, *339*, 490–498. [[CrossRef](#)]
152. Mansukhani, A.; Bellosta, P.; Sahni, M.; Basilico, C. Signaling by Fibroblast Growth Factors (Fgf) and Fibroblast Growth Factor Receptor 2 (Fgfr2)–Activating Mutations Blocks Mineralization and Induces Apoptosis in Osteoblasts. *J. Cell Biol.* **2000**, *149*, 1297–1308. [[CrossRef](#)]
153. Mansukhani, A.; Ambrosetti, D.; Holmes, G.; Cornivelli, L.; Basilico, C. Sox2 induction by FGF and FGFR2 activating mutations inhibits Wnt signaling and osteoblast differentiation. *J. Cell Biol.* **2005**, *168*, 1065–1076. [[CrossRef](#)]



154. Miraoui, H.; Oudina, K.; Petite, H.; Tanimoto, Y.; Moriyama, K.; Marie, P.J. Fibroblast growth factor receptor 2 promotes osteogenic differentiation in mesenchymal cells via ERK1/2 and protein kinase C signaling. *J. Biol. Chem.* **2009**, *284*, 4897–4904. [[CrossRef](#)]
155. Naganawa, T.; Xiao, L.; Coffin, J.; Doetschman, T.; Sabbieti, M.G.; Agas, D.; Hurley, M.M. Reduced expression and function of bone morphogenetic protein-2 in bones of Fgf2 null mice. *J. Cell. Biochem.* **2008**, *103*, 1975–1988. [[CrossRef](#)]
156. Xiao, G.; Jiang, D.; Gopalakrishnan, R.; Franceschi, R.T. Fibroblast Growth Factor 2 Induction of the Osteocalcin Gene Requires MAPK Activity and Phosphorylation of the Osteoblast Transcription Factor, Cbfa1/Runx2. *J. Biol. Chem.* **2002**, *277*, 36181–36187. [[CrossRef](#)]
157. Ai-Aql, Z.S.; Alagl, A.S.; Graves, D.T.; Gerstenfeld, L.C.; Einhorn, T.A. Molecular mechanisms controlling bone formation during fracture healing and distraction osteogenesis. *J. Dent Res.* **2008**, *87*, 107–118. [[CrossRef](#)] [[PubMed](#)]
158. Lemonnier, J.; Häy, E.; Delannoy, P.; Lomri, A.; Modrowski, D.; Caverzasio, J.; Marie, P.J. Role of N-Cadherin and Protein Kinase C in Osteoblast Gene Activation Induced by the S252W Fibroblast Growth Factor Receptor 2 Mutation in Apert Craniosynostosis. *J. Bone Miner. Res.* **2001**, *16*, 832–845. [[CrossRef](#)] [[PubMed](#)]
159. Debiais, F.; Lefèvre, G.; Lemonnier, J.; Le Mée, S.; Lasmoles, F.; Mascarelli, F.; Marie, P. Fibroblast growth factor-2 induces osteoblast survival through a phosphatidylinositol 3-kinase-dependent,  $\beta$ -catenin-independent signaling pathway. *Exp. Cell Res.* **2004**, *297*, 235–246. [[CrossRef](#)]
160. Montero, A.; Okada, Y.; Tomita, M.; Ito, M.; Tsurukami, H.; Nakamura, T.; Doetschman, T.; Coffin, J.D.; Hurley, M.M. Disruption of the fibroblast growth factor-2 gene results in decreased bone mass and bone formation. *J. Clin. Investig.* **2000**, *105*, 1085–1093. [[CrossRef](#)]
161. Xiao, L.; Sobue, T.; Eslinger, A.; Kronenberg, M.S.; Coffin, J.D.; Doetschman, T.; Hurley, M.M. Disruption of the Fgf2 gene activates the adipogenic and suppresses the osteogenic program in mesenchymal marrow stromal stem cells. *Bone* **2010**, *47*, 360–370. [[CrossRef](#)]
162. Coffin, J.D.; Florkiewicz, R.Z.; Neumann, J.; Mort-Hopkins, T.; Dorn, G.W.; Lightfoot, P.; German, R.; Howles, P.N.; Kier, A.; O'Toole, B.A. Abnormal bone growth and selective translational regulation in basic fibroblast growth factor (FGF-2) transgenic mice. *Mol. Biol. Cell* **1995**, *6*, 1861–1873. [[CrossRef](#)]
163. Sobue, T.; Naganawa, T.; Xiao, L.; Okada, Y.; Tanaka, Y.; Ito, M.; Okimoto, N.; Nakamura, T.; Coffin, J.; Hurley, M. Over-expression of fibroblast growth factor-2 causes defective bone mineralization and osteopenia in transgenic mice. *J. Cell. Biochem.* **2005**, *95*, 83–94. [[CrossRef](#)]
164. Berthiaume, F.; Maguire, T.J.; Yarmush, M.L. Tissue Engineering and Regenerative Medicine: History, Progress, and Challenges. *Annu. Rev. Chem. Biomol. Eng.* **2011**, *2*, 403–430. [[CrossRef](#)]
165. Langer, R.; Vacanti, J.P. Tissue engineering. *Science* **1993**, *260*, 920–926. [[CrossRef](#)]
166. Black, C.R.M.; Goriainov, V.; Gibbs, D.; Kanczler, J.M.; Tare, R.S.; Oreffo, R.O.C. Bone Tissue Engineering. *Curr. Mol. Biol. Rep.* **2015**, *1*, 132–140. [[CrossRef](#)]
167. Porter, J.R.; Ruckh, T.T.; Papat, K.C. Bone tissue engineering: A review in bone biomimetics and drug delivery strategies. *Biotechnol. Prog.* **2009**, *25*, 1539–1560. [[CrossRef](#)]
168. McCarthy, T.L.; Centrella, M.; Canalis, E. Effects of Fibroblast Growth Factors on Deoxyribonucleic Acid and Collagen Synthesis in Rat Parietal Bone Cells. *Endocrinology* **1989**, *125*, 2118–2126. [[CrossRef](#)]
169. Martin, I.; Muraglia, A.; Campanile, G.; Cancedda, R.; Quarto, R. Fibroblast growth factor-2 supports ex vivo expansion and maintenance of osteogenic precursors from human bone marrow. *Endocrinology* **1997**, *138*, 4456–4462. [[CrossRef](#)]
170. Gorin, C.; Rochefort, G.Y.; Bascetin, R.; Ying, H.; Lesieur, J.; Sadoine, J.; Beckouche, N.; Berndt, S.; Novais, A.; Lesage, M.; et al. Priming Dental Pulp Stem Cells With Fibroblast Growth Factor-2 Increases Angiogenesis of Implanted Tissue-Engineered Constructs Through Hepatocyte Growth Factor and Vascular Endothelial Growth Factor Secretion. *Stem Cells Transl. Med.* **2016**, *5*, 392–404. [[CrossRef](#)] [[PubMed](#)]
171. Hasegawa, T.; Chosa, N.; Asakawa, T.; Yoshimura, Y.; Asakawa, A.; Ishisaki, A.; Tanaka, M. Effect of fibroblast growth factor-2 on periodontal ligament cells derived from human deciduous teeth in vitro. *Exp. Ther. Med.* **2010**, *1*, 337–341. [[CrossRef](#)] [[PubMed](#)]
172. Wu, J.; Huang, G.T.-J.; He, W.; Wang, P.; Tong, Z.; Jia, Q.; Dong, L.; Niu, Z.; Ni, L. Basic Fibroblast Growth Factor Enhances Stemness of Human Stem Cells from the Apical Papilla. *J. Endod.* **2012**, *38*, 614–622. [[CrossRef](#)]
173. Novais, A.; Lesieur, J.; Sadoine, J.; Slimani, L.; Baroukh, B.; Saubaméa, B.; Schmitt, A.; Vital, S.; Poliard, A.; Hélarly, C.; et al. Priming Dental Pulp Stem Cells from Human Exfoliated Deciduous Teeth with Fibroblast Growth Factor-2 Enhances Mineralization Within Tissue-Engineered Constructs Implanted in Craniofacial Bone Defects. *Stem Cells Transl. Med.* **2019**, *8*, 844–857. [[CrossRef](#)] [[PubMed](#)]
174. Debiais, F.; Hott, M.; Graulet, A.M.; Marie, P.J. The Effects of Fibroblast Growth Factor-2 on Human Neonatal Calvaria Osteoblastic Cells Are Differentiation Stage Specific. *J. Bone Miner. Res.* **1998**, *13*, 645–654. [[CrossRef](#)] [[PubMed](#)]
175. Hurley, M.M.; Adams, U.J.; Wang, L.; Jiang, X.; Burt, P.M.; Du, E.; Xiao, L. Accelerated fracture healing in transgenic mice overexpressing an anabolic isoform of fibroblast growth factor 2. *J. Cell. Biochem.* **2016**, *117*, 599–611. [[CrossRef](#)] [[PubMed](#)]
176. Cowan, C.M.; Quarto, N.; Warren, S.M.; Salim, A.; Longaker, M.T. Age-related changes in the biomolecular mechanisms of calvarial osteoblast biology affect fibroblast growth factor-2 signaling and osteogenesis. *J. Biol. Chem.* **2003**, *278*, 45040. [[CrossRef](#)]
177. Shimabukuro, Y.; Ueda, M.; Ozasa, M.; Anzai, J.; Takedachi, M.; Yanagita, M.; Ito, M.; Hashikawa, T.; Yamada, S.; Murakami, S. Fibroblast Growth Factor-2 Regulates the Cell Function of Human Dental Pulp Cells. *J. Endod.* **2009**, *35*, 1529–1535. [[CrossRef](#)] [[PubMed](#)]

178. He, H.; Yu, J.; Liu, Y.; Lu, S.; Liu, H.; Shi, J.; Jin, Y. Effects of FGF2 and TGFbeta1 on the differentiation of human dental pulp stem cells in vitro. *Cell Biol. Int.* **2008**, *32*, 827–834. [[CrossRef](#)] [[PubMed](#)]
179. Nakao, K.; Itoh, M.; Tomita, Y.; Tomooka, Y.; Tsuji, T. FGF-2 potently induces both proliferation and DSP expression in collagen type I gel cultures of adult incisor immature pulp cells. *Biochem. Biophys. Res. Commun.* **2004**, *325*, 1052–1059. [[CrossRef](#)] [[PubMed](#)]
180. Hakki, S.S.; Hakki, E.E.; Nohutcu, R.M. Regulation of matrix metalloproteinases and tissue inhibitors of matrix metalloproteinases by basic fibroblast growth factor and dexamethasone in periodontal ligament cells. *J. Periodontol. Res.* **2009**, *44*, 794–802. [[CrossRef](#)]
181. Shimabukuro, Y.; Ueda, M.; Ichikawa, T.; Terashi, Y.; Yamada, S.; Kusumoto, Y.; Takedachi, M.; Terakura, M.; Kohya, A.; Hashikawa, T.; et al. Fibroblast Growth Factor-2 Stimulates Hyaluronan Production by Human Dental Pulp Cells. *J. Endod.* **2005**, *31*, 805–808. [[CrossRef](#)]
182. Nauman, E.A.; Sakata, T.; Keaveny, T.M.; Halloran, B.P.; Bikle, D.D. bFGF Administration Lowers the Phosphate Threshold for Mineralization in Bone Marrow Stromal Cells. *Calcif. Tissue Int.* **2003**, *73*, 147–152. [[CrossRef](#)]
183. Nakamura, S.; Nambu, M.; Ishizuka, T.; Hattori, H.; Kanatani, Y.; Takase, B.; Kishimoto, S.; Amano, Y.; Aoki, H.; Kiyosawa, T.; et al. Effect of controlled release of fibroblast growth factor-2 from chitosan/fucoidan micro complex-hydrogel on in vitro and in vivo vascularization. *J. Biomed. Mater. Res. Part. A* **2007**, *85*, 619–627. [[CrossRef](#)]
184. Tabata, Y.; Yamada, K.; Hong, L.; Miyamoto, S.; Hashimoto, N.; Ikada, Y. Skull bone regeneration in primates in response to basic fibroblast growth factor. *J. Neurosurg.* **1999**, *91*, 851–856. [[CrossRef](#)]
185. Kwan, M.D.; Sellmyer, M.A.; Quarto, N.; Ho, A.M.; Wandless, T.J.; Longaker, M.T. Chemical Control of FGF-2 Release for Promoting Calvarial Healing with Adipose Stem Cells. *J. Biol. Chem.* **2011**, *286*, 11307–11313. [[CrossRef](#)]
186. Nakahara, T.; Nakamura, T.; Kobayashi, E.; Inoue, M.; Shigeno, K.; Tabata, Y.; Eto, K.; Shimizu, Y. Novel Approach to Regeneration of Periodontal Tissues Based on in Situ Tissue Engineering: Effects of Controlled Release of Basic Fibroblast Growth Factor from a Sandwich Membrane. *Tissue Eng.* **2003**, *9*, 153–162. [[CrossRef](#)] [[PubMed](#)]
187. Anzai, J.; Nagayasu-Tanaka, T.; Terashima, A.; Asano, T.; Yamada, S.; Nozaki, T.; Kitamura, M.; Murakami, S. Long-term Observation of Regenerated Periodontium Induced by FGF-2 in the Beagle Dog 2-Wall Periodontal Defect Model. *PLoS ONE* **2016**, *11*, e0158485. [[CrossRef](#)]
188. Lee, J.-H.; Kang, K.-J.; Jang, Y.-J. Functional efficacy of human recombinant FGF-2s tagged with (His) 6 and (His-Asn) 6 at the N- and C-termini in human gingival fibroblast and periodontal ligament-derived cells. *Protein Expr. Purif.* **2017**, *135*, 37–44. [[CrossRef](#)] [[PubMed](#)]
189. Charles, L.F.; Woodman, J.L.; Ueno, D.; Gronowicz, G.; Hurley, M.M.; Kuhn, L.T. Effects of low dose FGF-2 and BMP-2 on healing of calvarial defects in old mice. *Exp. Gerontol.* **2015**, *64*, 62–69. [[CrossRef](#)]
190. Kigami, R.; Sato, S.; Tsuchiya, N.; Yoshimakai, T.; Arai, Y.; Ito, K. FGF-2 Angiogenesis in Bone Regeneration Within Critical-Sized Bone Defects in Rat Calvaria. *Implant. Dent.* **2013**, *22*, 422–427. [[CrossRef](#)] [[PubMed](#)]
191. Song, K.; Rao, N.-J.; Chen, M.-L.; Huang, Z.-J.; Cao, Y.-G. Enhanced bone regeneration with sequential delivery of basic fibroblast growth factor and sonic hedgehog. *Injury* **2011**, *42*, 796–802. [[CrossRef](#)] [[PubMed](#)]
192. Wang, B.; Mastrogiacomo, S.; Yang, F.; Shao, J.; Ong, M.M.A.; Chanchareonsook, N.; Jansen, J.A.; Walboomers, X.F.; Yu, N. Application of BMP-Bone Cement and FGF-Gel on Periodontal Tissue Regeneration in Nonhuman Primates. *Tissue Eng. Part C Methods* **2019**, *25*, 748–756. [[CrossRef](#)] [[PubMed](#)]
193. Collignon, A.-M.; Lesieur, J.; Anizan, N.; Ben Azzouna, R.; Poliard, A.; Gorin, C.; Letourneur, D.; Chaussain, C.; Rouzet, F.; Rochefort, G.Y. Early angiogenesis detected by PET imaging with <sup>64</sup>Cu-NODAGA-RGD is predictive of bone critical defect repair. *Acta Biomater.* **2018**, *82*, 111–121. [[CrossRef](#)]
194. Grosso, A.; Burger, M.G.; Lunger, A.; Schaefer, D.J.; Banfi, A.; Di Maggio, N. It Takes Two to Tango: Coupling of Angiogenesis and Osteogenesis for Bone Regeneration. *Front. Bioeng. Biotechnol.* **2017**, *5*, 68. [[CrossRef](#)]
195. Atlas, Y.; Gorin, C.; Novais, A.; Marchand, M.F.; Chatzopoulou, E.; Lesieur, J.; Bascetin, R.; Binet-Moussy, C.; Sadoine, J.; Lesage, M.; et al. Microvascular maturation by mesenchymal stem cells in vitro improves blood perfusion in implanted tissue constructs. *Biomaterials* **2021**, *268*, 120594. [[CrossRef](#)]
196. Kitamura, M.; Akamatsu, M.; Machigashira, M.; Hara, Y.; Sakagami, R.; Hirofuji, T.; Hamachi, T.; Maeda, K.; Yokota, M.; Kido, J.; et al. FGF-2 stimulates periodontal regeneration: Results of a multi-center randomized clinical trial. *J. Dent. Res.* **2011**, *90*, 35–40. [[CrossRef](#)]
197. Santana, D.B.R.; de Santana, C.M. Human intrabony defect regeneration with rhFGF-2 and hyaluronic acid—A randomized controlled clinical trial. *J. Clin. Periodontol.* **2015**, *42*, 658–665. [[CrossRef](#)] [[PubMed](#)]
198. Cochran, D.; Oh, T.-J.; Mills, M.; Clem, D.; McClain, P.; Schallhorn, R.; McGuire, M.; Scheyer, E.; Giannobile, W.; Reddy, M.; et al. A Randomized Clinical Trial Evaluating rh-FGF-2/ $\beta$ -TCP in Periodontal Defects. *J. Dent. Res.* **2016**, *95*, 523–530. [[CrossRef](#)] [[PubMed](#)]
199. Khoshkam, V.; Chan, H.L.; Lin, G.H.; Mailoa, J.; Giannobile, W.V.; Wang, H.L.; Oh, T.J. Outcomes of regenerative treatment with rhPDGF-BB and rhFGF-2 for periodontal intra-bony defects: A systematic review and meta-analysis. *J. Clin. Periodontol.* **2015**, *42*, 272–280. [[CrossRef](#)] [[PubMed](#)]
200. Li, F.; Yu, F.; Xu, X.; Li, C.; Huang, D.; Zhou, X.; Ye, L.; Zheng, L. Evaluation of Recombinant Human FGF-2 and PDGF-BB in Periodontal Regeneration: A Systematic Review and Meta-Analysis. *Sci. Rep.* **2017**, *7*, 1–10. [[CrossRef](#)]

201. Gronowicz, G.; Jacobs, E.; Peng, T.; Zhu, L.; Hurley, M.; Kuhn, L.T. Calvarial Bone Regeneration Is Enhanced by Sequential Delivery of FGF-2 and BMP-2 from Layer-by-Layer Coatings with a Biomimetic Calcium Phosphate Barrier Layer. *Tissue Eng. Part. A* **2017**, *23*, 1490–1501. [[CrossRef](#)]
202. Ou, G.; Charles, L.; Matton, S.; Rodner, C.; Hurley, M.; Kuhn, L.; Gronowicz, G. Fibroblast Growth Factor-2 Stimulates the Proliferation of Mesenchyme-Derived Progenitor Cells From Aging Mouse and Human Bone. *J. Gerontol. Ser. A Boil. Sci. Med. Sci.* **2010**, *65*, 1051–1059. [[CrossRef](#)] [[PubMed](#)]
203. Sukarawan, W.; Nowwarote, N.; Kerdpon, P.; Pavasant, P.; Osathanon, T. Effect of basic fibroblast growth factor on pluripotent marker expression and colony forming unit capacity of stem cells isolated from human exfoliated deciduous teeth. *Odontology* **2013**, *102*, 160–166. [[CrossRef](#)]
204. Ludwig, T.E.; Levenstein, M.E.; Jones, J.M.; Berggren, W.T.; Mitchen, E.R.; Frane, J.L.; Crandall, L.J.; Daigh, C.A.; Conard, K.R.; Piekarczyk, M.S.; et al. Derivation of human embryonic stem cells in defined conditions. *Nat. Biotechnol.* **2006**, *24*, 185–187. [[CrossRef](#)] [[PubMed](#)]
205. Varkey, M.; Kucharski, C.; Haque, T.; Sebald, W.; Uludağ, H. In Vitro Osteogenic Response of Rat Bone Marrow Cells to bFGF and BMP-2 Treatments. *Clin. Orthop. Relat. Res.* **2006**, *443*, 113–123. [[CrossRef](#)]
206. Dunstan, C.R.; Boyce, R.; Boyce, B.F.; Garrett, I.R.; Izbicka, E.; Burgess, W.H.; Mundy, G.R. Systemic Administration of Acidic Fibroblast Growth Factor (FGF-1) Prevents Bone Loss and Increases New Bone Formation in Ovariectomized Rats. *J. Bone Miner. Res.* **1999**, *14*, 953–959. [[CrossRef](#)] [[PubMed](#)]
207. Kato, T.; Kawaguchi, H.; Hanada, K.; Aoyama, I.; Hiyama, Y.; Nakamura, T.; Kuzutani, K.; Tamura, M.; Kurokawa, T.; Nakamura, K. Single local injection of recombinant fibroblast growth factor-2 stimulates healing of segmental bone defects in rabbits. *J. Orthop. Res.* **1998**, *16*, 654–659. [[CrossRef](#)] [[PubMed](#)]
208. Okazaki, H.; Kurokawa, T.; Nakamura, K.; Matsushita, T.; Mamada, K.; Kawaguchi, H. Stimulation of bone formation by recombinant fibroblast growth factor-2 in callotasis bone lengthening of rabbits. *Calcif. Tissue Int.* **1999**, *64*, 542–546. [[CrossRef](#)]
209. Nakamura, T.; Hara, Y.; Tagawa, M.; Tamura, M.; Yuge, T.; Fukuda, H.; Nigi, H. Recombinant Human Basic Fibroblast Growth Factor Accelerates Fracture Healing by Enhancing Callus Remodeling in Experimental Dog Tibial Fracture. *J. Bone Miner. Res.* **1998**, *13*, 942–949. [[CrossRef](#)]
210. Kawaguchi, H.; Nakamura, K.; Tabata, Y.; Ikada, Y.; Aoyama, I.; Anzai, J.; Nakamura, T.; Hiyama, Y.; Tamura, M. Acceleration of fracture healing in nonhuman primates by fibroblast growth factor-2. *J. Clin. Endocrinol. Metab.* **2001**, *86*, 875–880. [[CrossRef](#)]
211. Radomsky, M.L.; Aufdemorte, T.B.; Swain, L.D.; Fox, W.C.; Spiro, R.C.; Poser, J.W. Novel formulation of fibroblast growth factor-2 in a hyaluronan gel accelerates fracture healing in nonhuman primates. *J. Orthop. Res.* **1999**, *17*, 607–614. [[CrossRef](#)] [[PubMed](#)]
212. Kamo, K.; Miyakoshi, N.; Kasukawa, Y.; Sasaki, H.; Shimada, Y. Effects of single and cyclical local injections of basic fibroblast growth factor on cancellous bone defects in rabbits. *J. Orthop. Sci.* **2009**, *14*, 811–819. [[CrossRef](#)] [[PubMed](#)]
213. Nakamura, T.; Hanada, K.; Tamura, M.; Shibunishi, T.; Nigi, H.; Tagawa, M.; Fukumoto, S.; Matsumoto, T. Stimulation of endosteal bone formation by systemic injections of recombinant basic fibroblast growth factor in rats. *Endocrinology* **1995**, *136*, 1276–1284. [[CrossRef](#)]
214. Nagai, H.; Tsukuda, R.; Mayahara, H. Effects of basic fibroblast growth factor (bFGF) on bone formation in growing rats. *Bone* **1995**, *16*, 367–373. [[CrossRef](#)]
215. Mayahara, H.; Ito, T.; Nagai, H.; Miyajima, H.; Tsukuda, R.; Taketomi, S.; Mizoguchi, J.; Kato, K. In Vivo Stimulation of Endosteal Bone Formation by Basic Fibroblast Growth Factor in Rats. *Growth Factors* **1993**, *9*, 73–80. [[CrossRef](#)] [[PubMed](#)]
216. Lane, N.E.; Yao, W.; Kumer, J.; Breuning, T.; Wronski, T.; Modin, G.; Kinney, J.H. Basic fibroblast growth factor partially restores trabecular bone architecture in osteopenic ovariectomized rats. *Osteoporos. Int.* **2003**, *4*, 374–382.
217. Abbaspour, A.; Takata, S.; Sairyo, K.; Katoh, S.; Yukata, K.; Yasui, N. Continuous local infusion of fibroblast growth factor-2 enhances consolidation of the bone segment lengthened by distraction osteogenesis in rabbit experiment. *Bone* **2008**, *42*, 98–106. [[CrossRef](#)]
218. Benington, L.; Rajan, G.; Locher, C.; Lim, L.Y. Fibroblast Growth Factor 2—A Review of Stabilisation Approaches for Clinical Applications. *Pharmaceutics* **2020**, *12*, 508. [[CrossRef](#)]
219. Kuroda, Y.; Kawai, T.; Goto, K.; Matsuda, S. Clinical application of injectable growth factor for bone regeneration: A systematic review. *Inflamm. Regen.* **2019**, *39*, 20. [[CrossRef](#)]
220. Arakawa, T.; Prestrelski, S.J.; Kenney, W.C.; Carpenter, J.F. Factors affecting short-term and long-term stabilities of proteins. *Adv. Drug Deliv. Rev.* **2001**, *46*, 307–326. [[CrossRef](#)]
221. Tabata, Y.; Hijikata, S.; Ikada, Y. Enhanced vascularization and tissue granulation by basic fibroblast growth factor impregnated in gelatin hydrogels. *J. Control. Release* **1994**, *31*, 189–199. [[CrossRef](#)]
222. Kawaguchi, H.; Jingushi, S.; Izumi, T.; Fukunaga, M.; Matsushita, T.; Nakamura, T.; Mizuno, K.; Nakamura, T.; Nakamura, K. Local application of recombinant human fibroblast growth factor-2 on bone repair: A dose-escalation prospective trial on patients with osteotomy. *J. Orthop. Res.* **2007**, *25*, 480–487. [[CrossRef](#)] [[PubMed](#)]
223. Kawaguchi, H.; Oka, H.; Jingushi, S.; Izumi, T.; Fukunaga, M.; Sato, K.; Matsushita, T.; Nakamura, K. For the TESK Group A local application of recombinant human fibroblast growth factor 2 for tibial shaft fractures: A randomized, placebo-controlled trial. *J. Bone Miner. Res.* **2010**, *25*, 2735–2743. [[CrossRef](#)]

224. Nakamura, S.; Kanatani, Y.; Kishimoto, S.; Nakamura, S.I.; Ohno, C.; Horio, T.; Masanori, F.; Hattori, H.; Tanaka, Y.; Kiyosawa, T.; et al. Controlled release of FGF-2 using fragmin/protamine microparticles and effect on neovascularization. *J. Biomed. Mater. Res. A* **2009**, *91*, 814–823. [[CrossRef](#)]
225. Mammadov, R.; Mammadov, B.; Guler, M.O.; Tekinay, A.B. Growth Factor Binding on Heparin Mimetic Peptide Nanofibers. *Biomacromolecules* **2012**, *13*, 3311–3319. [[CrossRef](#)] [[PubMed](#)]
226. Kim, S.H.; Kiick, K.L. Heparin-mimetic sulfated peptides with modulated affinities for heparin-binding peptides and growth factors. *Peptides* **2007**, *28*, 2125–2136. [[CrossRef](#)]
227. Tardieu, M.; Gamby, C.; Avramoglou, T.; Jozefonvicz, J.; Barritault, D. Derivatized dextrans mimic heparin as stabilizers, potentiators, and protectors of acidic or basic FGF. *J. Cell. Physiol.* **1992**, *150*, 194–203. [[CrossRef](#)]
228. Liekens, S.; Leali, D.; Neyts, J.; Esnouf, R.; Rusnati, M.; Dell’Era, P.; Maudgal, P.C.; De Clercq, E.; Presta, M. Modulation of Fibroblast Growth Factor-2 Receptor Binding, Signaling, and Mitogenic Activity by Heparin-Mimicking Polysulfonated Compounds. *Mol. Pharmacol.* **1999**, *56*, 204–213. [[CrossRef](#)]
229. Nguyen, T.H.; Paluck, S.J.; McGahran, A.J.; Maynard, H.D. Poly(vinyl sulfonate) Facilitates bFGF-Induced Cell Proliferation. *Biomacromolecules* **2015**, *16*, 2684–2692. [[CrossRef](#)] [[PubMed](#)]
230. Arunkumar, P.; Dougherty, J.A.; Weist, J.; Kumar, N.; Angelos, M.G.; Powell, H.M.; Khan, M. Sustained Release of Basic Fibroblast Growth Factor (bFGF) Encapsulated Polycaprolactone (PCL) Microspheres Promote Angiogenesis In Vivo. *Nanomaterials* **2019**, *9*, 1037. [[CrossRef](#)]
231. Ali, Z.; Islam, A.; Sherrell, P.; Le-Moine, M.; Lolas, G.; Syrigos, K.; Rafat, M.; Jensen, L.D. Adjustable delivery of pro-angiogenic FGF-2 by alginate:collagen microspheres. *Biol. Open* **2018**, *7*, bio027060. [[CrossRef](#)] [[PubMed](#)]
232. Murahashi, Y.; Yano, F.; Nakamoto, H.; Maenohara, Y.; Iba, K.; Yamashita, T.; Tanaka, S.; Ishihara, K.; Okamura, Y.; Moro, T.; et al. Multi-layered PLLA-nanosheets loaded with FGF-2 induce robust bone regeneration with controlled release in critical-sized mouse femoral defects. *Acta Biomater.* **2019**, *85*, 172–179. [[CrossRef](#)]
233. Gromolak, S.; Krawczenko, A.; Antończyk, A.; Buczak, K.; Kiełbowicz, Z.; Klimczak, A. Biological Characteristics and Osteogenic Differentiation of Ovine Bone Marrow Derived Mesenchymal Stem Cells Stimulated with FGF-2 and BMP-2. *Int. J. Mol. Sci.* **2020**, *21*, 9726. [[CrossRef](#)]
234. Akita, S.; Fukui, M.; Nakagawa, H.; Fujii, T.; Akino, K. Cranial bone defect healing is accelerated by mesenchymal stem cells induced by coadministration of bone morphogenetic protein-2 and basic fibroblast growth factor. *Wound Repair Regen.* **2004**, *12*, 252–259. [[CrossRef](#)]
235. Hanada, K.; Dennis, J.E.; Caplan, A.I. Stimulatory Effects of Basic Fibroblast Growth Factor and Bone Morphogenetic Protein-2 on Osteogenic Differentiation of Rat Bone Marrow-Derived Mesenchymal Stem Cells. *J. Bone Miner. Res.* **1997**, *12*, 1606–1614. [[CrossRef](#)]
236. Fujimura, K.; Bessho, K.; Okubo, Y.; Kusumoto, K.; Segami, N.; Iizuka, T. The effect of fibroblast growth factor-2 on the osteoinductive activity of recombinant human bone morphogenetic protein-2 in rat muscle. *Arch. Oral Biol.* **2002**, *47*, 577–584. [[CrossRef](#)]
237. Takita, H.; Tsuruga, E.; Ono, I.; Kuboki, Y. Enhancement by bFGF of osteogenesis induced by rhBMP-2 in rats. *Eur. J. Oral Sci.* **1997**, *105*, 588–592. [[CrossRef](#)]
238. Nakamura, Y.; Tensho, K.; Nakaya, H.; Nawata, M.; Okabe, T.; Wakitani, S. Low dose fibroblast growth factor-2 (FGF-2) enhances bone morphogenetic protein-2 (BMP-2)-induced ectopic bone formation in mice. *Bone* **2005**, *36*, 399–407. [[CrossRef](#)] [[PubMed](#)]
239. Berendsen, A.D.; Olsen, B.R. Bone development. *Bone* **2015**, *80*, 14–18. [[CrossRef](#)]
240. Breeland, G.; Menezes, R.G. *Embryology, Bone Ossification*; StatPearls Publishing LLC.: Treasure Island, FL, USA, 2019.
241. Deckers, M.M.; Van Bezooijen, R.L.; Van Der Horst, G.; Hoogendam, J.; van Der Bent, C.; Papapoulos, S.E.; Löwik, C.W. Bone morphogenetic proteins stimulate angiogenesis through osteoblast-derived vascular endothelial growth factor A. *Endocrinology* **2002**, *143*, 1545–1553. [[CrossRef](#)] [[PubMed](#)]
242. Wan, M.; Cao, X. BMP signaling in skeletal development. *Biochem. Biophys. Res. Commun.* **2005**, *328*, 651–657. [[CrossRef](#)] [[PubMed](#)]
243. Peng, H.; Usas, A.; Olshanski, A.; Ho, A.M.; Gearhart, B.; Cooper, G.M.; Huard, J. VEGF Improves, Whereas sFlt1 Inhibits, BMP2-Induced Bone Formation and Bone Healing Through Modulation of Angiogenesis. *J. Bone Miner. Res.* **2005**, *20*, 2017–2027. [[CrossRef](#)]
244. Valdimarsdottir, G.; Goumans, M.-J.; Rosendahl, A.; Brugman, M.; Itoh, S.; Lebrin, F.; Sideras, P.; Dijke, P.T. Stimulation of Id1 Expression by Bone Morphogenetic Protein Is Sufficient and Necessary for Bone Morphogenetic Protein-Induced Activation of Endothelial Cells. *Circulation* **2002**, *106*, 2263–2270. [[CrossRef](#)] [[PubMed](#)]
245. Salazar, V.S.; Gamer, L.W.; Rosen, V. BMP signalling in skeletal development, disease and repair. *Nat. Rev. Endocrinol.* **2016**, *12*, 203–221. [[CrossRef](#)]
246. Carano, R.A.; Filvaroff, E.H. Angiogenesis and bone repair. *Drug Discov. Today* **2003**, *8*, 980–989. [[CrossRef](#)]
247. Chen, G.; Deng, C.; Li, Y.P. TGF-beta and BMP signaling in osteoblast differentiation and bone formation. *Int. J. Biol. Sci.* **2012**, *8*, 272–288. [[CrossRef](#)]
248. Yang, E.Y.; Moses, H.L. Transforming growth factor beta 1-induced changes in cell migration, proliferation, and angiogenesis in the chicken chorioallantoic membrane. *J. Cell Biol.* **1990**, *111*, 731–741. [[CrossRef](#)] [[PubMed](#)]

249. Sabbieti, M.G.; Marchetti, L.; Gabrielli, M.G.; Menghi, M.; Materazzi, S.; Menghi, G.; Raisz, L.G.; Hurley, M.M. Prostaglandins differently regulate FGF-2 and FGF receptor expression and induce nuclear translocation in osteoblasts via MAPK kinase. *Cell Tissue Res.* **2004**, *319*, 267–278. [[CrossRef](#)]
250. Sobue, T.; Gravelly, T.; Hand, A.; Min, Y.K.; Pilbeam, C.; Raisz, L.G.; Zhang, X.; Larocca, D.; Florkiewicz, R.; Hurley, M.M. Regulation of fibroblast growth factor 2 and fibroblast growth factor receptors by transforming growth factor beta in human osteoblastic MG-63 cells. *J. Bone Miner. Res.* **2002**, *17*, 502–512. [[CrossRef](#)]
251. Fiedler, J.; Röderer, G.; Günther, K.-P.; Brenner, R.E. BMP-2, BMP-4, and PDGF-bb stimulate chemotactic migration of primary human mesenchymal progenitor cells. *J. Cell. Biochem.* **2002**, *87*, 305–312. [[CrossRef](#)]
252. Edwards, J.R.; Nyman, J.S.; Lwin, S.T.; Moore, M.M.; Esparza, J.; O’Quinn, E.C.; Hart, A.J.; Biswas, S.; Patil, C.A.; Lonning, S.; et al. Inhibition of TGF-beta signaling by 1D11 antibody treatment increases bone mass and quality in vivo. *J. Bone Miner. Res.* **2010**, *25*, 2419–2426. [[CrossRef](#)] [[PubMed](#)]
253. Tang, Y.; Wu, X.; Lei, W.; Pang, L.; Wan, C.; Shi, Z.; Zhao, L.; Nagy, T.R.; Peng, X.; Hu, J.; et al. TGF-beta1-induced migration of bone mesenchymal stem cells couples bone resorption with formation. *Nat. Med.* **2009**, *15*, 757–765. [[CrossRef](#)] [[PubMed](#)]
254. Hollinger, J.O.; Hart, C.E.; Hirsch, S.N.; Lynch, S.; Friedlaender, G.E. Recombinant Human Platelet-Derived Growth Factor: Biology and Clinical Applications. *J. Bone Jt. Surg. Am. Vol.* **2008**, *90*, 48–54. [[CrossRef](#)] [[PubMed](#)]
255. Vu, T.H.; Shipley, J.; Bergers, G.; Berger, J.E.; Helms, J.A.; Hanahan, D.; Shapiro, S.D.; Senior, R.M.; Werb, Z. MMP-9/Gelatinase B Is a Key Regulator of Growth Plate Angiogenesis and Apoptosis of Hypertrophic Chondrocytes. *Cell* **1998**, *93*, 411–422. [[CrossRef](#)]
256. Stickens, D.; Behonick, D.J.; Ortega, N.; Heyer, B.; Hartenstein, B.; Yu, Y.; Fosang, A.J.; Schorpp-Kistner, M.; Angel, P.; Werb, Z. Altered endochondral bone development in matrix metalloproteinase 13-deficient mice. *Development* **2004**, *131*, 5883–5895. [[CrossRef](#)] [[PubMed](#)]
257. Ramasamy, S.K.; Kusumbe, A.P.; Wang, L.; Adams, R.H. Endothelial Notch activity promotes angiogenesis and osteogenesis in bone. *Nature* **2014**, *507*, 376–380. [[CrossRef](#)]