



**HAL**  
open science

# HELMINTHS OF APODEMUS SYLVATICUS (MURIDAE) DISTRIBUTED ON THE SOUTHERN EUROPEAN BORDER (ITALIAN PENINSULA)

C. Milazzo, G. Aloise, M. Cagnin, C. Di Bella, F. Geraci, C. Feliu, J. C.  
Casanova

► **To cite this version:**

C. Milazzo, G. Aloise, M. Cagnin, C. Di Bella, F. Geraci, et al.. HELMINTHS OF APODEMUS SYLVATICUS (MURIDAE) DISTRIBUTED ON THE SOUTHERN EUROPEAN BORDER (ITALIAN PENINSULA). *Vie et Milieu / Life & Environment*, 2005, pp.45-51. hal-03219008

**HAL Id: hal-03219008**

<https://hal.sorbonne-universite.fr/hal-03219008v1>

Submitted on 6 May 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## HELMINTHS OF *APODEMUS SYLVATICUS* (MURIDAE) DISTRIBUTED ON THE SOUTHERN EUROPEAN BORDER (ITALIAN PENINSULA)

C. MILAZZO<sup>1-3</sup>, G. ALOISE<sup>1</sup>, M. CAGNIN<sup>1</sup>, C. DI BELLA<sup>2</sup>, F. GERACI<sup>2</sup>,  
C. FELIU<sup>3</sup>, J. C. CASANOVA<sup>3</sup>

<sup>1</sup>Dipartimento di Ecologia, Università della Calabria, Via P. Bucci s/n., 87036 Rende (Cosenza), Italy  
<sup>2</sup>Istituto Zooprofilattico Sperimentale della Sicilia "A. Mirri," Via G. Marinuzzi 3, 90129 Palermo, Italy

<sup>3</sup>Laboratori de Parasitologia, Facultat de Farmàcia, Universitat de Barcelona,  
Avda Diagonal s/n 08028 Barcelona, Spain  
Corresponding author: dibella@pa.izs.it

HELMINTHES  
*APODEMUS SYLVATICUS*  
ITALIE  
MURIDAE

RÉSUMÉ. – Cinq aires du sud de l'Italie ont été examinées et ont révélé la présence d'*Apodemus sylvaticus*. Au total, 14 espèces d'Helminthes intestinaux ont été identifiées : Trematoda (*Corrigia vitta*), Cestoda (larves de *Taenia taeniaeformis*, *Gallegoides arfaai*, *Hymenolepis* sp. et *Rodentolepis straminea*), et Nematoda (*Trichuris muris*, *Aonchotheca muris-sylvatici*, A. sp., *Mastophorus muris*, *Rictularia proni*, *Heligmosomoides polygyrus*, *Syphacia stroma*, *S. frederici* et *Aspicularis tetraptera*). L'intensité d'infection par les Helminthes intestinaux est plus importante chez les adultes et chez les mâles. Le nombre d'espèces rencontrées en Calabre s'est révélé moins important que celui de la communauté de Vers identifiés en Europe du Nord et dans les aires continentales. La richesse spécifique, en effet, semble être identique à celle observée pour les vertébrés: dans une péninsule, le nombre d'espèces diminue parce que l'environnement réduit progressivement sa capacité à être colonisé par des vertébrés.

HELMINTHS  
*APODEMUS SYLVATICUS*  
ITALY  
MURIDAE

ABSTRACT. – Five areas populated by *Apodemus sylvaticus* in South of Italy have been analyzed. In total, fourteen species of intestinal helminths belonging to trematodes (*Corrigia vitta*), cestodes (*Taenia taeniaeformis* larvae, *Gallegoides arfaai*, *Hymenolepis* sp. and *Rodentolepis straminea*), and nematodes (*Trichuris muris*, *Aonchotheca muris-sylvatici*, A. sp., *Mastophorus muris*, *Rictularia proni*, *Heligmosomoides polygyrus*, *Syphacia stroma*, *S. frederici* and *Aspicularis tetraptera*) were found. Mean intensities of intestinal helminths were higher in adults and males. The number of parasite species found in Calabria was lower compared to helminth communities of the northern part of the European peninsulas and of other continental areas. The specific richness was similar to that observed in vertebrate taxa: in a peninsula, the number of species decreases due to habitat barriers that progressively reduce the possibility of colonization.

### INTRODUCTION

When describing the biodiversity of an area, parasites and infectious diseases are to be considered as "the forgotten half of biodiversity" (Dobson 1999). According to Dobson, there are ten species of parasites associated with any population of living mammals and each individual host contains a whole community of parasites. The helminth faunas of European *Apodemus* sp. have been well documented from extensive research mainly carried out in continental and insular areas. More than thirty helminth species have been reported parasitizing *A. sylvaticus*, *A. flavicollis*, *A. agrarius*, *A. microps* and *A. mystacinus* (Feliu *et al.* 1987, 1997, Montgomery & Montgomery 1990, Asakawa &

Tenora 1996, Behnke *et al.* 1999, Abu-Madi *et al.* 2000). Part of these studies provide ecological data on host sex and/or host age compositions, seasonal compositions or habitat influences. From these studies, it is hypothesized that species diversity should be lower on islands than on the mainland (Goüy de Bellocq *et al.* 2003). Very few reports of *Apodemus* sp. helminths in the European peninsulas are available. Thus, the study of parasite communities in the southern parts of Europe (Iberian and Italian peninsulas) should be of great interest since these were subject to variations due to the fact that these areas established a shelter for faunas during the Quaternary period (Hewitt 1996, 1999). Data are scarce on parasitic helminths from rodents in Italy (Macchioni 1967, Virga *et al.* 1995, Goüy de Bellocq *et al.* 2003, Milazzo *et al.* 2003a, b). In

contrast, parasitic helminths of murine rodents, mostly of the woodmice, are well documented in different countries in Europe (Goüy de Bellocq *et al.* 2002). Helminths of Italian *Apodemus* sp. are almost unknown. Therefore, the aim of this research is to provide the first description of helminth communities of *A. sylvaticus* in the southern Italian Peninsula (Calabria). The species richness in *A. sylvaticus* from Calabria will be discussed, taking into account the biogeography of rodents in Italy.

## METHODS

A total of 697 *A. sylvaticus* were studied for intestinal helminths. The wood mice were trapped in five different areas of Calabria (South of Italy): Crati River Valley (N 16: 33, E 39: 21) (years 1990-91; n= 168; monthly captured from February 1990 to February 1991), Sila Grande (N 16: 35, E 39: 23) (years 1994-95; n= 88; monthly from June 1991 to July 1992), Sila Greca (N 16: 42, E 39: 29) (years 1994-95; n= 163; monthly from April 1994 to March 1995), Northern Jonian rivers ("fiumare") (Avena river: N 16: 33, E 39: 55, Satanasso river: N 16: 31, E 30: 52) (years 1993-94; n= 204; monthly July 1992 to July 1993) and Catena Costiera (N 16: 06, E 39: 22) (years 1988-02; n= 74; may 1989 to January 1990).

**Crati River Valley:** The valley, shaped by the main river of Calabria, is about 91 km long. The landscape shows a mosaic of biotopes consisting of small patches of residual *Quercus virgiliana* woods, olive groves, cereal farming, abandoned fields in which rudimentary vegetation prevails.

**Sila Grande:** This is an extensive undulating plateau ranging from a minimum of 800 m to a maximum of 1930 m in altitude in the northern part of the Calabrian Apennines. Between 600 and 1100 m of woodland is covered by chestnut (*Castanea sativa*). Some areas consist of deciduous woodland of different types of oak (*Quercus* spp.) and maple (*Acer* spp.). Above 1100 m, the forest is covered by the Calabrian black pine (*Pinus laricio*), beech (*Fagus sylvatica*), white fir (*Abies alba*), hazel (*Corylus avellana*) and poplar (*Populus tremula*).

**Sila Greca:** The massif is located in the north eastern part of Calabria, ranging from 40 to 1480 m. In the lower part prevails the sclerophyll vegetation with Mediterranean maquis and *Quercus ilex* woods. Instead, in the upper part, the deciduous woods prevail (*Quercus pubescens*, *Q. cerris*, *Q. frainetto*, *Q. virgiliana*).

**Catena Costiera:** This is a mountain range of the north western part of Calabria, with few peaks higher than 1400 m. This area is covered by *Quercus virgiliana* and *Castanea sativa* woods in the lower part and *Fagus sylvatica* woods in the upper part.

**Northern Jonian rivers ("fiumare"):** Rodents were collected in the two "fiumare" Avena and Satanasso, located in the north eastern coast of Calabria. The "fiumare" are temporary rivers characterized by a short and rapid route whose final part is wide and stony. The area's climate is that of a typical dry Mediterranean cli-

mate. Vegetation is scarce downstream river and it is mainly dominated by *Chenopodium bothryos*, *Helicrisum italicum* and *Artemisia variabilis*. On the beside of the river we can find wood of *Quercus ilex*, *Pistacia lentiscus* or *Pinus halepensis*.

Animals were trapped using pitfall traps, according to a standardized procedure that allows to compare the relative density of animals at different stations (Cagnin *et al.* 1998) and these were stored in a 70% ethanol solution for a later examination. We examined 382 males and 314 females (1 specimen remained undetermined for sex). These animals were classified into two different categories of age as a result of the four age classes established by Adamcewska-Andrejewska (1976): juveniles (Class I and II = younger than two months; 263 specimens), adults (Class III and IV = older than two months; 428 specimens) (6 specimens remain undetermined for age). The stomach, the intestine, the liver and the pancreas were split and examined for any presence of helminths. Parasitic worms were treated for permanent mounting in Canada Balsam (cestodes and trematodes previously stained with Semichon acetocarmine) or temporarily examined with Amann lactophenol (nematodes). For specific identification of helminths the works of Verster (1969), Tenora (1963), Tenora & Meszaros (1975), Tenora & Mas Coma (1977), Meszaros & Murai (1979), Moravec (1982), Moravec *et al.* (1987), Casanova *et al.* (2001), Feliu *et al.* (2000) were consulted.

The characteristics of helminth faunas were analyzed in terms of prevalence (P) and the Average Intensity (AI) according to Bush *et al.* (1997). The diversity index of Gini-Simpson were calculated according to Gini (1912).

## RESULTS

Out of a total of 697 specimens of *A. sylvaticus* examined (Table I), 311 were infested (prevalence: P = 44.6) by 15790 parasites (average intensity AI = 51). The parasites belonged to 14 species (Table I) of intestinal helminths: one trematode (*Corrigia vitta* Dujardin, 1845), four cestodes (*Taenia taeniaeformis* (Batsch, 1786) larvae, *Gallegoides arfaai* (Mobedi et Ghadirian, 1977), *Hymenolepis* sp. and *Rodentolepis straminea* (Goeze, 1782) and nine nematodes (*Trichuris muris* (Schrunk, 1788), *Aonchotheca muris-sylvatici* (Dujardin, 1843), *Aonchotheca* sp., *Mastophorus muris* (Gmelin, 1790), *Rictularia proni* Seurat, 1915, *Heligmosomoides polygyrus* (Dujardin, 1845), *Syphacia stroma* (Von Linstow, 1884), *S. frederici* Roman, 1945 and *Aspiculuris tetraptera* (Nitsh, 1821). Only one species of trematode was collected in the pancreatic ducts of ten animals. Three species of cestodes are intestinal worms whereas *H. taeniaeformis* was found in the liver of the larval stage from one host. All species of nematodes are intestinal worms except *M. muris* and *R. proni* that are stomachal parasites. General helminth infections in *A. sylvaticus* are low not exceeding half of the studied animals (Table I). Only two species

Table I. – Value of prevalence (P%) and mean intensity (MI) for each recorded parasite. N = number of hosts; 95% l.c.l.–u.c.l. = lower and upper confidential limits at 95% of significance;  $s^2$  = variance; Max-Min = maximum and minimum number of parasites;  $N_p$  = number of parasites.

	N	P% (95% l.c.l.-u.c.l.)	MI	$\sigma^2$	Max-Min	$N_p$
<b>TREMATODA</b>						
<i>C.vitta</i>	10	1.4 (0.7-2.7)	5.1	17.8	14-1	51
<b>CESTODA</b>						
<i>T. taeniaeformis</i> (l)	1	0.1 (0.0-0.9)	1.0	0.0	1-1	1
<i>G. arfaai</i>	9	1.3 (0.6-2.5)	2.4	3.5	6-1	22
<i>Hymenolepis</i> sp.	2	0.3 (0.0-1.2)	1.0	0.0	1-1	2
<i>R. straminea</i>	1	0.1 (0.0-0.9)	1.0	0.0	1-1	1
<b>NEMATODA</b>						
<i>T. muris</i>	10	1.4 (0.7-2.7)	4.3	18.6	14-1	43
<i>A. muris-sylvatici</i>	13	1.9 (1.0-3.3)	6.5	59.6	24-1	84
<i>Aonchotheca</i> sp.	3	0.4 (0.1-1.4)	3.3	10.3	7-1	10
<i>M. muris</i>	1	0.1 (0.0-0.9)	1.0	0.0	1-1	1
<i>R. proni</i>	30	4.3 (3.0-6.2)	5.8	114.4	49-1	173
<i>H. polygyrus</i>	193	27.7 (24.4-31.2)	34.4	11,103.8	1350-1	6,643
<i>S. stroma</i>	50	7.2 (5.4-9.4)	39.6	3,426.5	310-1	1,981
<i>S. frederici</i>	142	20.4 (17.5-23.6)	43.7	7,356.3	782-1	6,212
<i>A. tetraptera</i>	26	3.7 (2.5-5.5)	21.8	2,909.1	255-1	566
<b>TOTAL</b>	697	44.6 (40.9-48.4)	51	12,444.9	1374-1	15,790

*H. polygyrus* and *S. frederici* show prevalences higher than 10% and could be considered as members of the component community of the woodmouse in Calabria. Differences in the composition of the community were found between males and females and between juveniles and adults. General prevalences and number of species in males and females are similar (Table II). In general all species found in both sexes show low prevalences and only *H. polygyrus* and *S. frederici* could be considered in the component community of these animals (Table II). Young individuals are

found parasitised by 11 worm species versus 13 found in adult animals (Table III). General prevalence and all prevalences of individual species in young host were lower than in adults (Table III). Also, both in adult and young animals only *H. polygyrus* and *S. frederici* show prevalences higher than 10%. The helminth communities structure was tested with the diversity index Gini-Simpson which shows to be fairly similar between adults and juveniles (0.38 versus 0.34) and quite different between males and females (0.31 versus 0.43).

Table II. – Value of prevalence and mean intensity for every recorded parasite inside sex variable. Abbreviations: see table I.

	P%		MI		$\sigma^2$		Max-Min		N <sub>p</sub>	
	(95% l.c.l.-u.c.l.)		F	M	F	M	F	M	F	M
	F (N = 314)	M (N = 382)								
<b>TREMATODA</b>										
<i>C. vitta</i>	1.6 (0.6-3.9)	1.3 (0.5-3.2)	4.4	5.8	12.3	26.7	10-1	14-1	22	29
<b>CESTODA</b>										
<i>T. taeniaeformis</i>	-	0.3 (0.0-1.7)	-	1.0	-	0.0	-	1-1	-	1
<i>G. arfaai</i>	0.6 (0.1-2.5)	1.8 (0.8-3.9)	1.0	2.9	0.0	3.8	1-1	6-1	2	20
<i>Hymenolepis</i> sp.	0.3 (0.0-0.2)	0.3 (0.0-1.7)	1.0	1.0	0.0	0.0	1-1	1-1	1	1
<i>R. straminea</i>	0.3 (0.0-2.0)	-	1.0	-	0.0	-	1-1	-	1	-
<b>NEMATODA</b>										
<i>T. muris</i>	1.0 (0.2-3.0)	1.8 (0.8-3.9)	1.3	5.6	0.3	21.6	2-1	14-1	4	39
<i>A. muris-sylvatici</i>	1.0 (0.2-3.0)	2.6 (1.3-4.9)	9.0	5.7	169.0	39.1	24-1	20-1	27	57
<i>Aonchotheca</i> sp.	0.3 (0.0-2.0)	0.5 (0.1-2.1)	7.0	1.5	0.0	0.5	7-7	2-1	7	3
<i>M. muris</i>	0.3 (0.0-2.0)	-	1.0	-	0.0	-	1-1	-	1	-
<i>R. proni</i>	3.8 (2.1-6.8)	4.7 (2.9-7.5)	5.7	5.8	107.1	125.9	37-1	49-1	68	105
<i>H. polygyrus</i>	25.8 (21.1-31.1)	29.3 (24.9-34.2)	36.7	32.8	22,598.2	2,913.0	1,350-1	279-1	2,974	3,669
<i>S. stroma</i>	5.4 (3.3-8.7)	8.6 (6.1-12.0)	30.9	44.1	2,092.2	4,139.2	183-1	310-1	525	1,456
<i>S. frederici</i>	15.0 (11.3-19.5)	24.9 (20.7-29.6)	28.4	51.3	3,294.6	9,246.8	299-1	782-1	1,336	4,876
<i>A. tetraptera</i>	2.5 (1.2-5.2)	4.7 (2.9-7.5)	6.0	28.8	51.4	4,087.9	18-1	255-1	48	518
<b>TOTAL</b>	40.4 (35.0-46.1)	48.2 (43.1-53.3)	39.4	58.5	16,674.3	9,451.7	1374-1	795-1	5016	10,774

## DISCUSSION

The helminth fauna of *A. sylvaticus* in Calabria is composed of common species of *Apodemus* spp. in Europe but highest species richness has been detected in continental Europe (Tenora 1962, Asakawa & Tenora 1996, Gouÿ de Bellocq *et al.* 2002, 2003). Peninsular studies on helminths of *A. sylvaticus*, mostly carried out in the Iberian Peninsula, show that wood mice have a richer fauna than other rodents in the Peninsula, with a number of species similar to continental Europe (Feliu *et*

*al.* 1997). Insular helminth populations parasitizing *A. sylvaticus* show a species richness that ranged from typical configurations in the British Isles (9 species from Montgomery & Montgomery 1990; eight species reported by Behnke *et al.* 1999) to those configurations reached in the Balearic Islands (see Mas Coma & Feliu 1984). In Calabria, the rodents' fauna is composed of a reduced number of species (31) compared to the rodents' fauna of northern regions, which is probably due to a "peninsular effect" (Simpson 1964, Cagnin *et al.* 1998, Baquero & Telleria 2001). To date known helminth faunas of rodents in Calabria seem also

Table III. – Value of prevalence and mean intensity for every recorded parasite within age variable. Abbreviations: see table I.

	P% (95% l.c.l.-u.c.l.)		MI		$\sigma^2$		Max-Min		N <sub>p</sub>	
	J (N = 263)	A (N = 428)	J	A	J	A	J	A	J	A
<b>TREMATODA</b>										
<i>C. vitta</i>	0.4 (0.0-2.1)	2.1 (1.0-4.1)	1.0	5.5	0.0	17.7	1-1	14-1	1	50
<b>CESTODA</b>										
<i>T. taeniaeformis</i>	-	0.2 (0.0-1.5)	-	1.0	-	0.0	-	1-1	-	1
<i>G. arfaai</i>	0.8 (0.1-2.7)	1.6 (0.7-3.5)	1.0	2.9	0.0	3.8	1-1	6-1	2	20
<i>Hymenolepis</i> sp.	-	0.5 (0.1-1.9)	-	1.0	-	0.0	-	1-1	-	2
<i>R. straminea</i>	0.4 (0.0-2.1)	-	1.0	-	0.0	-	1-1	-	1	-
<b>NEMATODA</b>										
<i>T. muris</i>	0,8 (0.1-2.7)	1,9 (0.9-3.8)	5,5	4	24.5	20.0	9-2	14-1	11	32
<i>A. muris-sylvatici</i>	1.5 (0.4-3.8)	2.1 (1.0-4.1)	3.5	7.8	19.0	75.9	10-1	24-1	14	70
<i>Aonchotheca</i> sp.	0.4 (0.0-2.1)	0.5 (0.1-1.9)	7.0	1.5	0.0	0.5	7-7	2-1	7	3
<i>M.muris</i>	-	0.2 (0.0-1.5)	-	1.0	-	0.0	-	1-1	-	1
<i>R. proni</i>	1.9 (0.6-4.4)	5.8 (3.9-8.6)	2.0	6.5	3.0	134.2	5-1	49-1	10	163
<i>H. polygyrus</i>	17.5 (13.1-22.6)	34.1 (29.7-38.8)	13.5	41.2	626.9	14,315.8	129-1	1350-1	620	6,020
<i>S. stroma</i>	4.6 (2.4-7.8)	8.9 (6.4-12.1)	34.3	41.3	2,486.3	3,786.3	178-1	310-1	411	1,570
<i>S. frederici</i>	13.3 (9.4-18.0)	25.0 (21.0-29.4)	24.3	50.1	1,558.2	9,119.1	200-1	782-1	849	5,363
<i>A. tetraptera</i>	1.5 (0.4-3.8)	5.1 (3.3-7.8)	3.0	25.2	3.3	3,383.4	5-1	255-1	12	554
<b>TOTAL</b>	27.8 (22.4-33.6)	55.4 (50.5-60.1)	26.5	58.8	1,625.7	15,710.1	200-1	1374-1	1,938	13,849

affected by the “peninsula effect” that influenced both prevalences and infection intensities of helminth parasites. The helminth fauna of *C. glareolus* was well studied in Calabria (Milazzo *et al.* 2003 b). The results of this study show a helminth community with a low species richness with a range of prevalences between 0,5 to 5,1%. The only data previously available on rodent parasitic helminths in Calabria comes from the study of Di Bella *et al.* (1997) which shows that *A. sylvaticus* and *A. flavicollis* are parasitized by the same and lower number of species. In Göyü de Bellocq *et al.* (2003), 12 helminth species were

cited in *A. sylvaticus* from Italy, all of these present in Calabria. No data are available on the prevalence or infection intensities of these species in the host population. In our study a general prevalence of 44,6% was found but only two species *H. polygyrus* and *S. frederici* show a prevalence higher than 10%. The rest of the species are not present in the component community of *A. sylvaticus* in Calabria. This fact is rare in the known helminth communities of this host in Europe. Normally specific species of *Apodemus* spp. are in the component communities of this host in Europe and, in several islands, also less specific species show

prevalences higher than 10%. In Sicily, despite the close proximity to Calabria, the number of species found in *A. sylvaticus* is higher than in Calabrian populations of this host. In another study of Goüy de Bellocq *et al.* (2003) all the species found are members of the component community of woodmouse. Probably, the non specific parasites of rodents such as *C. vitta*, *T. muris*, *A. dujardini* and *Mesocestoides* sp. are found in high prevalences in Sicilian woodmouse by the absence of other habitual host of these parasites as different species of Arvicolinae (i.e. *Clethrionomys glareolus*, habitual host of these worms in Europe mainland) (Feliu *et al.* 1997, Milazzo *et al.* 2003 b). Unfortunately, the rodent helminth faunas in Sicily are poorly known. To date only three rodent species in Sicily are studied helminthologically, *A. sylvaticus*, *Mus domesticus* and *Rattus rattus* and only three common species are found in these hosts, *H. diminuta*, *M. muris* and *T. muris* (Goüy de Bellocq *et al.* 2003, Milazzo *et al.* 2003a, Virga *et al.* 1995). In Calabria, the helminth faunas of *C. glareolus* and *M. domesticus* are known and are also characterised by a low helminth richness with several species (*C. vitta*, *M. muris*) parasitizing Murinae and Arvicolinae and *H. polygyrus* and *A. tetraptera* present in Murinae (*M. domesticus*, *A. sylvaticus*, *A. flavicollis*) (Di Bella *et al.* 1997, Milazzo *et al.* 2003, 2004). Prevalences of these common species in the different hosts are always lower than 10% except for *H. polygyrus* in *A. sylvaticus* found in our study (Di Bella *et al.* 1997, Milazzo *et al.* 2004). Concerning the continental populations of *A. sylvaticus*, the results found in Calabria are surprising. In the study of Behnke *et al.* (1999) in England nine helminth species were found, all of these except *A. muris sylvatici*, *H. taeniaeformis* and *Brachylaima recurva* with prevalences higher than 8,2%.

Different factors could be contributes to increase helminth richness in the different European populations of the woodmouse. In Iberia, *A. sylvaticus* is parasitized by endemic species which are mostly found in the Pyrenean mountains that separate the peninsula from the continental areas (Feliu *et al.* 1987). Most of these species are digenetic Trematodes with two intermediate hosts (in most cases also endemic species) with a distribution influenced by paleobiogeographic factors. The absence of several species in the helminth fauna of *A. sylvaticus* in Calabria could be explained by the historical biogeography of this host because the Italo-Balkan populations of woodmouse are characterised by a lower genetic diversity (Michaux *et al.* 2003). Sicily, on the contrary, as it is demonstrated in Michaux *et al.* (2003) represents a genetic hot spot of *A. sylvaticus* and probably this could explain the high helminth richness in this host (Goüy de Bellocq *et al.* 2003).

Our results are in accordance with the reports showing a lesser role of sex in the structure of the

helminth communities in several small mammals (Kisielewska 1970, Abu-Madi *et al.* 2000, Milazzo *et al.* 2003 a, b). After Haukisalmi *et al.* (1988), Montgomery & Montgomery (1990), Gregory (1992), Behnke *et al.* (1999) host age play a significant role in the structure of the component communities. Our results about *A. sylvaticus* in Calabria suggest that the host age is not an important factor determining the structure of the helminth community since species richness are similar in young and adult animals.

ACKNOWLEDGEMENTS. – The authors wish to thank the Comando Forestale of the Sila Grande Mountain for allowing the trapping in the Sila Grande area. This study was supported by two grants from the MIUR-Italy (ex 40%: local coordinator Prof Brandmayr; ex 60%: local coordinator Dr M Cagnin) and by the Comisionat per la Recerca i Universitats de la Generalitat de Catalunya 2001SGR00088.

## REFERENCES

- Abu-Madi MA, Behnke JM, Lewis JW, Gilbert FS 2000. Seasonal and site specific variation in the component community structure of intestinal helminths in *Apodemus sylvaticus* from three contrasting habitats in South East of England. *J Helminthol* 74: 7-15.
- Adamcewska-Andrejewska KA 1976. Age reference model for *Apodemus flavicollis* (Melchior, 1834). *Ekol Pol Ser A* 15 (41): 788-790.
- Asakawa M, Tenora F 1996. A checklist of Nematode Parasite of the Genus *Apodemus* (Murinae: Rodentia) throughout the World excluding Japan. *J Rakuno Gakuen Univ* 20 (2): 181-213.
- Baquero RA, Telleria JL 2001. Species richness, rarity and endemicity of European mammals: a biogeographical approach. *Biodiver Conserv* 10: 29-44.
- Behnke JM, Lewis JW, Mohd Zain SN, Gilbert FS 1999. Helminth infections in *Apodemus sylvaticus* in Southern England: interactive effects of host age, sex, and year on the prevalence and abundance of infections. *J Helminthol* 73: 31-44.
- Bush AO, Lafferty KD, Lotz JM, Shostak AW 1997. Parasitology meets ecology on its own terms: Margolis *et al.* revisited. *J Parasitol* 83: 575-583.
- Cagnin M, Moreno S, Aloise G, Garofalo G, Villafuerte R, Cristaldi M 1998. Comparative study of Spanish and Italian terrestrial small mammals of different biotopes in the Mediterranean Peninsular Tips. *J Biogeogr* 25: 1105-1113.
- Casanova JC, Santalla F, Durant P, Vaucher C, Feliu C, Renaud F 2001. Morphological and genetic differentiation of *Rodentolepis straminea* (Goeze, 1752) and *Rodentolepis microstoma* (Dujardin, 1845) (Hymenolepididae). *Parasitol Res* 87 (6): 439-444.
- Di Bella C, Milazzo C, Cagnin M 1997. Helminth parasites of *Apodemus sylvaticus* and *Apodemus flavicollis* from different biotopes of the Trionto river basin (N-E Calabria). *S It E Atti* 18: 125-126.

- Dobson A 1999. The role of parasite in ecological systems. In Farina A ed., *Perspective in Ecology*. Backhuys Publishers, Leiden: 51-64.
- Feliu C, Gracenea M, Torregrosa M 1987. Consideraciones ecológicas sobre la helmintofauna de *Apodemus sylvaticus* (Linnaeus, 1758) (Rodentia: Muridae) en le Pireneo Oriental Espanol. In Sans-Coma V, Mas-Coma S & Gosálbez J eds, *Mamíferos y Helmintos*. Ketres, Barcelona: 175-181.
- Feliu C, Spakulova M, Casanova JC, Renaud F, Morand S, Hugot JP, Santalla F, Durant P 2000. Genetic and morphological heterogeneity in small rodent white worms in south western Europe: characterization of *Trichuris muris* and description of *Trichuris arvicolae* n. sp. *J Parasitol* 86 (3): 442-449.
- Feliu C, Renaud F, Catzeffli F, Hugot JP, Durand P, Morand S 1997. A comparative analysis of parasite species richness of Iberian rodents. *Parasitology* 115: 453-466.
- Gini C. 1912. Variabilità e mutabilità. Studi Econ Giur Fac Giurisprudenza Univ Cagliari, anno III, parte II.
- Gregory R.D. 1992. On the interpretation of host-parasite ecology: *Heligmosomoides polygyrus* (Nematoda) in wild wood mouse (*Apodemus sylvaticus*) populations. *J Zoology* London 226: 109-121.
- Goüy De Bellocq J, Morand S, Feliu C 2002. Patterns of parasite species richness of western palearctic micro-mammals: island effects. *Ecology* 25: 173-183.
- Goüy De Bellocq J, Sarà M, Casanova JC, Feliu C, Morand S 2003. A comparison of the structure of helminth communities in the woodmouse, *Apodemus sylvaticus*, on islands of the western Mediterranean and continental Europe. *Parasitology Research* 90 (1): 64-70.
- Haukisalml V, Henttonen H, Tenora F 1988. Population dynamics of common and rare helminths in cyclic vole populations. *J Animal Ecology* 57: 807-825.
- Hewitt GM 1996. Some genetic consequences of ice ages, and their role in divergence and speciation. *Biol J Linn Soc* 58: 247-276.
- Hewitt GM 1999. Post-glacial re-colonization of European biota. *Biol J Linn Soc* 68: 87-112.
- Kisielewska K 1970. Ecological organization of intestinal helminth groupings in *Clethrionomys glareolus* (Schreb.) (Rodentia). IV. Spatial structure of a helminth grouping within the host population. *Acta Parasitol Pol* 18: 177-196.
- Macchioni G 1967. Indagine sulla elmintofauna dei muridi in Italia. *Ann Fac Med Vet Pisa* 20: 40-51.
- Mas-Coma S, Feliu C 1984. Helminth fauna from small mammals (Insectivores and Rodents) on the Pityusic Islands In *Biogeography and Ecology of the Pityusic Island*; Monographiae Biologicae. Dr W Junk Publ, The Hague, 469-525.
- Meszáros F, Murai E 1979. Contribution to the Knowledge of rodents' helminths in Rumania. *Parasitol Hung* 12: 55-70.
- Michaux JR, Magnanou E, Paradis E, Nieberding C, Li-bois R 2003. Mitochondrial phylogeography of the woodmice (*Apodemus sylvaticus*) in the western Palearctic region. *Mol Ecol* 12: 685-697.
- Milazzo C, Goüy De Bellocq J, Cagnin M, Casanova JC, Di Bella C, Morand S, Feliu C, Fons R, Morand S, Santalla F 2003a. Helminths and Ectoparasites of *Rattus rattus* and *Mus musculus* from Sicily, Italy. *Comp Parasitol* 70 (2): 99-104.
- Milazzo C, Casanova JC, Aloise G, Ribas Alexis, Cagnin M 2003b. Helminths of the bank vole *Clethrionomys glareolus* (Rodentia, Arvicolinae) in Southern Italy. *Ital J Zool* 70: 333-337.
- Milazzo C, Casanova JC, Ribas A, Aloise G, Cagnin M 2004. Helminths of *Mus domesticus* (Rodentia, Muridae) in Calabria (South Italy). S Mas Coma, MD Bargues, JG Esteban, MA Valero Eds. IX European Multicolloquium of Parasitology, Valencia, Spain 18-23 July: 588.
- Montgomery SSJ, Montgomery WI 1990. Structure, stability and species interactions in helminth communities of wood mice, *Apodemus sylvaticus*. *Int J Parasitol* 20 (2): 225-242.
- Moravec F 1982. Proposal of a new systematic arrangement of nematodes of the Capillariidae family. *Folia Parasitol* 29: 119-132.
- Moravec F, Prokopic P, Shilkas AV 1987. The biology of nematodes of the family Capillariidae Neveu-Lemaire, 1936. *Folia Parasitol* 34: 39-56.
- Simpson GG 1964. Species density of North American recent mammals. *Syst Zool* 13: 57-63.
- Tenora F 1962. Parasites in field-mice of the genus *Apodemus* in the CSSR. Thesis-chair of Zoology.
- Tenora F 1963. Review of parasitic worms in rodents of genus *Apodemus* in the Czech Republic. *Zool Listy* 12: 331-336.
- Tenora F, Mas-Coma S 1977. Records of *Gallegoides arfaai* (Mobedi et Ghadirian, 1977) n. comb. (Cestoda: Anoplocephalidae) in *Apodemus sylvaticus* L. from Western Europe. Proposition of *Gallegoides* n. gen. *Saugetierk Mitt* 40 (3): 222-226.
- Tenora F, Meszaros F 1975. Nematodes of the genus *Symphacia* Seurat, 1916 (Nematoda) Rodents' parasites (Rodentia) in the Czech Republic and Hungary. *Acta Univ Agric* 23 (3): 537-554.
- Verster A 1969. A taxonomic revision of the genus *Taenia* Linnaeus, 1758 s. str. *Onderstepoort. J Vet Res* 36: 3-58.
- Virga A, Canestri Trotti G, Nobile L, Pampiglione S 1995. Elminti intestinali di roditori selvatici della Sicilia Occidentale. *Atti Soc Ital Sci Vet* 49: 751-752.

Reçu le 15 janvier 2004; received January 15, 2004  
 Accepté le 7 mai 2004; accepted May 7, 2004