



HAL
open science

PICOPHYTOPLANKTON: BOTTOM-UP AND TOP-DOWN CONTROLS ON ECOLOGY AND EVOLUTION

J A Raven, Z V Finkel, A J Irwin

► **To cite this version:**

J A Raven, Z V Finkel, A J Irwin. PICOPHYTOPLANKTON: BOTTOM-UP AND TOP-DOWN CONTROLS ON ECOLOGY AND EVOLUTION. *Vie et Milieu / Life & Environment*, 2005, pp.209-215. hal-03219059

HAL Id: hal-03219059

<https://hal.sorbonne-universite.fr/hal-03219059v1>

Submitted on 6 May 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

PICOPHYTOPLANKTON: BOTTOM-UP AND TOP-DOWN CONTROLS ON ECOLOGY AND EVOLUTION

J. A. RAVEN¹, Z. V. FINKEL², A. J. IRWIN³

¹Plant Research Unit, University of Dundee at SCRI, Scottish Crop Research Institute, Invergowrie, Dundee DD2 5DA, UK

²Environmental Science Program, Mount Allison University, Sackville, NB, Canada E4L 1A7

³Department of Mathematics and Computer Science, Mount Allison University, Sackville, NB, Canada E4L 1E6
Corresponding author: j.a.raven@dundee.ac.uk

CHLOROPHYTA
CYANOBACTERIA
EUKARYA
GRAZING
PARASITISM
RESOURCE ACQUISITION
SEDIMENTATION
VIRUSES

ABSTRACT. – Picophytoplankton organisms were derived from larger ancestors in both cyanobacteria and (polyphyletically) in eukarya. There are a number of putative advantages in the acquisition of photosynthetically active radiation and nutrient solutes from resource-limited habitats, and probably of maximum specific growth rate, for very small cells relative to the situation for larger phytoplankton cells. However, there are also putative disadvantages for such small cells with respect to bottom-up factors (i.e. those limiting biomass production), including an increased potential for solute leakage and an increased metabolic cost of screening out damaging UV-B. Among top-down factors (i.e. those removing biomass), picophytoplankton may be at an advantage relative to larger phytoplankton cells in avoiding damage from eukaryotic parasites, and losses from sedimentation. However, viruses and (small) grazers can attack picophytoplankton, just as viruses and (larger) grazers can attack larger phytoplankton. Picophytoplankton may be at a disadvantage relative to larger phytoplankton in environments with temporally variable resource supply.

CHLOROPHYTES
CYANOBACTÉRIES
EUCARYOTES
BROUTAGE
PARASITISME
EXPLOITATION DES RESSOURCES
SÉDIMENTATION
VIRUS

RÉSUMÉ. – Les organismes du picophytoplancton dérivent d'ancêtres de plus grande taille parmi les cyanobactéries et les eucaryotes (polyphylétiquement). Ces organismes ont un nombre d'avantages potentiels dans l'acquisition de la radiation active photosynthétiquement et des éléments nutritifs solubles des habitats limités en ressources, et probablement dans le taux maximum de croissance spécifique, pour de petites cellules par rapport à la situation de cellules du phytoplancton plus grandes. Cependant, il y a aussi des désavantages potentiels pour de si petites cellules par rapport aux facteurs de croissance "bottom-up" (c.-à-d. ceux qui limitent la production de biomasse), y compris une augmentation du potentiel pour l'écoulement des substances solubles et un coût métabolique élevé pour filtrer les rayons UV-B qui causent des dommages. Parmi les facteurs « herbivores » ("top-down") (c.-à-d. ceux qui suppriment la biomasse), le picophytoplancton pourrait avoir l'avantage, par rapport aux cellules plus grandes du phytoplancton d'éviter les dommages causés par les parasites eucaryotes, et les pertes dues à la sédimentation. Cependant, les virus et les (petits) herbivores peuvent attaquer le picophytoplancton, de même que les virus et les herbivores (plus grands) peuvent attaquer le phytoplancton plus gros. Le picophytoplancton peut-être désavantagé par rapport au phytoplancton de plus forte taille dans les environnements aux ressources variables dans le temps.

INTRODUCTION

Picophytoplankton are defined here as planktonic photosynthetic organisms which are not retained by a 2 µm pore diameter filter. Molecular phylogenetic analyses show that these very small planktonic photolithotrophs were derived from larger ancestors, that, at least among eukarya, the picophytoplankton condition is polyphyletic, and

that miniaturization of genomes and cells can increase the rate of evolution (Raven 1998, Moreira & López-García 2002, Vaulot *et al.* 2002, Dufresne *et al.* 2005, Giovannoni *et al.* 2005).

This paper aims to examine the possible ecological and evolutionary advantages and disadvantages of very small size for phytoplankton by comparison with larger photosynthetic plankton. These possible costs and benefits are discussed first for bottom-up factors, then for top-down factors.

BOTTOM-UP FACTORS

Non-scalable components

This heading includes any environmental factor which decreases the growth of phytoplankton. Examples include the restricted or excessive availability of photosynthetically active radiation (PAR) or of nutrient solutes, and growth rate-inhibiting fluxes of UV-B. The proportion of cell volume taken up by non-scalable components, such as the minimal suite of genes needed for photolithotrophic growth and biological membranes of constant thickness, increases as cell and genome size is reduced, potentially forcing the displacement of some protein catalysts of growth processes leading to increased specialization and a reduction in average growth rate over a range of environmental conditions (Raven 1986, 1998). In general there is an increase in maximum specific growth rate (μ_m ; biomass increase per unit biomass per unit time) of organisms with decreasing body size of the form $\mu_m = a \cdot \text{biomass}^{-b}$, where a is a taxon-specific 'constant'. While b is held to have a taxon-independent value of 0.25, it may not be possible to reject the possibility of lower values of b for the members of some Phyla or Classes (Raven 1998). If the non-scalable factors restricting μ_m become significant within the size range observed for picophytoplankton, then the value of μ_m for the smallest picophytoplankton should be lower than

predicted by the scaling relationship. While the smallest known photolithotrophic eukaryote, *Ostreococcus tauri*, has very high μ_m values (Fouilland *et al.* 2004), the smallest O_2 -evolver, *Prochlorococcus*, has a lower μ_m than many rather larger cyanobacteria (Sullivan *et al.* 2005). Based on available observations of maximum growth rates normalized to cell volume, bigger cells appear to have higher intrinsic maximum growth rates (Table I) although there is a great deal of taxonomic and experimental variation in estimates of μ_m .

Absorption of PAR and UV-B

Turning to resource-limited growth rates, there are sound physical arguments for more effective acquisition of PAR and of nutrient solutes by smaller than by larger organisms (Fogg 1986, Raven 1986, Chisholm 1992, Raven 1998). For PAR there is less package effect in smaller than in otherwise similar larger cells, so that each pigment molecule is more effective at absorbing photons, and it takes less time for a pigment-protein complex to absorb enough photons from a given radiation field to recoup the energy cost of synthesizing the complex, in smaller cells (Raven 1984, 1998, Finkel *et al.* 2004). It is thus predicted, and observed, that the allometric coefficient b is smaller (more negative) in light-limited than in resource-saturated growth of phytoplankton organisms (Finkel & Irwin 2000, Finkel 2001, Finkel *et al.* 2004).

Table I. – Measured maximum growth rates for picophytoplankton and growth rates normalized to a cell volume of $1 \mu\text{m}^3$.

| Size and Taxonomic grouping | Cell diameter μm | Measured μ_{max} day^{-1} | Estimated μ_{max} for $1 \mu\text{m}^3$ cell | References | |
|------------------------------------|--------------------------------|--|--|-------------|---|
| Prokaryotes (Bacteria) | | | | | |
| <i>Prochlorococcus</i> spp. | Cyanobacteria | ~0.7 | 0.99 | 0.7 | Shalapyonok <i>et al.</i> (1988) |
| <i>Synechococcus</i> spp. | Cyanobacteria | ~1 | 1.97 | 1.8 | Kana and Glibert (1987) |
| Eukaryotes | | | | | |
| <i>Ostreococcus tauri</i> | Prasinophyceae | 0.8-1.1 | 1.1,2.4,8* | 0.9,2.4,6.5 | Courties <i>et al.</i> (1998); Fouilland <i>et al.</i> (2004); Rodriguez <i>et al.</i> (2005) |
| <i>Micromonas pusilla</i> | Prasinophyceae | 1.4-19 | 0.9,3.5* | 1.1,5.0 | Thronsen (1976); DuRand <i>et al.</i> (2002) |
| <i>Aureococcus anophagefferens</i> | Pelagophyceae | 1.5-2 | 0.9,2.3* | 1.1,3.0 | Pustazzi <i>et al.</i> (2004); Caron <i>et al.</i> (2004) |
| <i>Pycnococcus provasplii</i> | Prasinophyceae | 2.7 | 0.7 | 1.2 | Ho <i>et al.</i> (2003) |
| <i>Nannochloris atomus</i> | Trebouxiophyceae | 3 | 0.6 | 1.2 | Ho <i>et al.</i> (2003) |
| <i>Chaetoceros cf. tenuissimus</i> | Bacillariophyceae | 4.0 | 1.6 | 3.9 | Doblin <i>et al.</i> (1999) |
| <i>Thalassiosira</i> spp. | Bacillariophyceae | ~4 | 3* | 7.2 | Furnas (1991) |
| Small pennate spp. | Bacillariophyceae | ~4 | 3.5* | 8.5 | Furnas (1991) |
| Small Gymnodiniaceae | Dinophyceae | ~4 | 1.0* | 2.5 | Furnas (1991) |
| <i>Emilinia luxleyi</i> | Prymnesiophyceae | ~4-6 | 1.3,1.9 | 3.7,5.5 | Brand and Guillard (1981); Rhodes <i>et al.</i> (1995) |
| <i>Skeletonema costatum</i> | Bacillariophyceae | ~8 | 5.9* | 23.9* | Furnas (1990) |
| Medium Gymnodiniaceae | Dinophyceae | ~10 | 1.0* | 4.6 | Furnas (1991) |

*data from field study, all other data from laboratory experiments

†computed assuming allometric scaling with an exponent of -0.25

Cell size data predominantly from references indicated and Vaultot *et al.* (2004).

For Furnas (1990 and 1991) size data were not provided; the size estimates are based on an interpretation of the term "small" and "medium" for diatoms and dinoflagellates.

Furthermore, the smaller package effect in picophytoplankton than in larger organisms means that the spectral diversity among photosynthetic pigments is expressed to a greater extent in the *in vivo* absorption spectrum in the smaller organisms, which indeed have a greater diversity of photosynthetic pigments than in larger organisms (Raven 1998, Larkum & Kühl 2005, Miller *et al.* 2005). This spectral diversity of pigments is of ecological and evolutionary significance in niche partitioning among picophytoplankton species (Stomp *et al.* 2005) even if this diversity is not as important on larger evolutionary scales (Falkowski *et al.* 2004a,b).

Restriction of growth rate by UV-B radiation resembles photoinhibition by high PAR rather than limitation by low PAR. However, there is an implication of small cell size for the effectiveness of soluble intracellular UV-B screening compounds in restricting UV-B access to targets such as DNA (Raven 1998). The smaller intracellular optical path length in picophytoplankton means that a certain concentration of soluble UV-B-absorbing compounds absorbs a smaller fraction of the UV-B incident on the cell than would be the case for a larger cell, so a higher UV-B flux reaches targets (Raven 1998). Other possibilities of avoiding UV-B damage, and variations in the potential to repair UV-B damage, mean that the prediction is not obeyed universally (Raven 1998, Day & Neale 2002, Sommaruga *et al.* 2005).

Solute acquisition and loss

Smaller cells have an enhanced potential for nutrient solute influxes from low bulk phase concentrations through the diffusion boundary layers and the plasmalemma relative to the requirement for growth, granted the allometry of the potential growth rate as a function of organism size (Raven 1986, Chisholm 1992, Raven 1998). There is, however, also more potential for the loss of solutes from the smaller cells (Raven 1986, 1998). Such increased leakage can reduce the energetic efficiency of photosynthetic inorganic carbon concentrating mechanisms (CCMs) by increasing the rate constant for efflux of accumulated CO₂ (Raven 1986, 1998, Giordano *et al.* 2005). The problem is exacerbated for cyanobacteria by the low CO₂ affinity, and low CO₂/O₂ selectivity, of both the Form IA and Form IB ribulose biphosphate carboxylase-oxygenases or Rubiscos (Horken & Tabita 1999, Badger & Price 2003, Giordano *et al.* 2005). The effectiveness of the CCM in *Synechocystis* in suppressing the oxygenase activity of Rubisco is seen by the absence of effect on growth in air-equilibrium solutions of the deletion of glycine decarboxylase, an enzyme of the photorespiratory carbon oxidation cycle which is required to consume

glycolate (Hagemann *et al.* 2005). Another potential problem for resource acquisition, in this case N₂ fixation, is that of keeping oxygen away from, the nitrogenase – nitrogenase reductase complex. Even with nitrogen fixation limited to the dark phase there is a higher energy cost of removing oxygen per unit nitrogenase activity in picocyanobacteria (and other picoplankton) than in larger diazotrophs. Despite this, picophytoplanktonic cyanobacteria in which nitrogen fixation has been demonstrated, or in which have the genetic potential for nitrogen fixation, occur in the ocean (Zehr *et al.* 1998 Falcon *et al.* 2005).

Implications of the streamlining of genome

The smallest O₂-evolving photolithotrophs (*Prochlorococcus* strains) have lost a number of functions, e.g. the ability to use certain oxidized nitrogen sources (Hess 2004). Such gene loss, with genome streamlining, characterizes not only picophytoplanktonic but also the picochemo-organotrophic bacteria of the open ocean (Bryant 2002, Giovannoni *et al.* 2005). This reduction in the size of the genome offsets considerations of scalability, but ultimately the minimum size of cell and genome would mean a greater fraction of biomass is occupied by DNA, with a corresponding decrease in cellular C:N and C:P relative to the Redfield Ratio (see Table I of Geider & La Roche 2002). An increased fraction of plasmalemma in smaller cells would, as a result of the high protein and phospholipid content of the membrane, also decrease C:N and C:P relative to the Redfield Ratio (Geider & La Roche 2002). However, the observation is that C:N and C:P ratios in these very small cells may be higher than the Redfield Ratio average for larger phytoplankton cells (Geider & La Roche 2002), with the additional organic C helping to further increase the surface area per unit N or P in the small cells (reviewed by Raven *et al.* 2005, Thingstad *et al.* 2005). A further possibility for increasing surface area per unit N or P is to have a more dilute cytoplasm (see Raven *et al.* 2005). The option of vacuolation is not used by the smallest phytoplankton cells (Raven 1998), with their resource storage role taken by the essentially particulate polymers rather than dissolved monomers, with an order of magnitude smaller volume required to store unit N or P (Table II). Overall (Raven 1998, Table 6), there is the potential for a more effective use of already-acquired resources in obtaining further resources in picophytoplankton than in larger phytoplankton cells both at resource saturation and when resources are limiting. There is, however, the proviso that at the lowest sizes of picophytoplankton the fraction of the cell taken up by non-scalable components may decrease the effectiveness of already-acquired resources in obtaining further resources. Small and large cells

Table II. – Volume needed per mol P or N stored in vacuoles isosmotic with seawater, or as particulate polyphosphate or polypeptide.

| Element stored, chemical form | m ³ mol ⁻¹ | References |
|--|----------------------------------|---------------------------|
| P as KH ₂ PO ₄ | 1.54 | Weast (1969/1970) |
| P as polyphosphate, assuming density is identical to that of solid calcium pyrophosphate | 0.041 | Weast (1969/1970) |
| P as polyphosphate at the highest concentration measured in acidocalcisomes | 0.125 | Docampo and Moreno (2001) |
| N as KNO ₃ | 1.52 | Weast (1969/1970) |
| N as cyanophycin, the polypeptide N storage compound (1 arginine:1 aspartate) of cyanobacteria, assuming the same density as protein | 0.042 | Boyd and Gradmann (2002) |

alike may be able to store nutrients when they are supplied in excess of need, despite using different storage strategies, but the characteristics of the temporal pulse will favour one size over others; small cells with insufficient biomass-normalized storage ability will not be able to take full advantage of large nutrient pulses.

TOP-DOWN FACTORS

Sinking

Sinking of live cells out of the euphotic zone is one factor removing biomass from a phytoplankton population. Stokes' Law shows that, if a 50 µm radius spherical cell with density 50 kg m⁻³ greater than the surrounding water sinks at 26 m day⁻¹ relative to the surrounding water, an otherwise similar cell of 0.5 µm radius cell would only sink 2.6 mm day⁻¹ (Raven 1998). We shall return to sinking in the context of parasitism. Coagulation of particles can dramatically change the size distribution and sinking fluxes, probably increasing the flux due to small particles above a Stokes' law prediction (Stemmann *et al.* 2004a).

Biophagy: eukaryotic parasitoids

Raven (1998) points out that picophytoplankton organisms are very unlikely to support eukaryotic parasites (parasitoids): see Raven & Waite (2004). The smallest known eukaryote is the picophytoplankton organism *Ostreococcus tauri*, with a

volume of 5.24·10⁻¹⁹ m³; the largest spherical picophytoplankton cell has a volume of 4.19·10⁻¹⁸ m³. A hypothetical parasitoid with the same volume as *Ostreococcus* infecting the largest picophytoplankton organism as host could produce only four new parasitoid cells per infecting parasitoid if only half of the host biomass is converted into new parasitoids (the rest being unusable, or respired). No such parasitoid is known. Clearly no eukaryotic parasitoid, even as small as the hypothetical example used above, could have the smallest picophytoplankton cyanobacteria as hosts.

Biophagy: viruses

Viruses are widespread top-down factors in cyanobacterial (Sullivan *et al.* 2005) and eukaryotic (Wilson *et al.* 2005) phytoplankton, including picophytoplankton. This view is supported by the recent characterisation of two viruses of diatoms, a major phytoplankton taxon for which there had been no previous characterisation of viruses (Nagasaki *et al.* 2004, 2005). Viruses may be involved in bloom termination in cyanobacterial and eukaryotic picophytoplankton (Evans *et al.* 2003). While viruses are much smaller than eukaryotic parasitoids, viral reproduction requires significant quantities of P and the cell quota of P is very low in P-starved *Prochlorococcus* cells (Bertilsson *et al.* 2003). Furthermore, Wilson *et al.* (1996) showed that viruses infecting a *Synechococcus* culture which was starved of P had substantially reduced burst sizes. With half of the cell quota of 0.36 fg P in the 1.65 Mbp genome, the burst size of

cyanomyophage P-SSM2, with a 0.252 Mbp genome (Sullivan *et al.* 2005), can only be 13, while if a phage was as large as the 0.407 Mbp coccolithovirus (Wilson *et al.* 2005) the burst size would only be 8. Could this be a constraint on the size of cyanophages infecting the smallest picophytoplankton cells? A similar suggestion, with much more extensive documentation, has been made independently by C Brown and collaborators (personal communication; manuscript submitted).

The top-down factor of viral infection may interact with bottom-up effects independently through the recycling of nutrients by cell lysis. The *Prochlorococcus* phages P-SSM2 and P-SSM4 each have copies of two cyanobacterial genes (*phoH* and *pstS*) that are expressed under P deficiency, while the *Synechococcus* phage S-PM2 has, in addition, *phoH*; could this be related to their infection of P-depleted cells (Sullivan *et al.* 2005)? *Prochlorococcus* (Sullivan *et al.* 2005) and *Synechococcus* (Mann *et al.* 2005) phages also contain genes related to photosynthesis, another possible interaction between top-down and bottom-up factors.

There is the possibility that increased sinking rates of phytoplankton damaged by parasitoid or virus infection could be a means of purging healthy surface-dwelling populations by the faster sinking of infected organisms (Lawrence & Suttle 2004, Raven & Waite 2004). Even for larger phytoplankton cells the evolution and operation of such a mechanism has several constraints, e.g. host specificity of the parasitoid or virus, and the hydrodynamic regime of the upper mixed layer (Lawrence & Suttle 2004, Raven & Waite 2004), while the sinking rate of picophytoplankton is so small as to eliminate this mechanism for removing infected cells (Waite *et al.* 1997, Raven 1998, Raven & Waite 2004).

Biophagy: grazers

Finally we address the impact of grazers. Picophytoplankton escape grazing by larger grazers, but can be consumed by smaller grazers (see Raven 1998, Chrisaki *et al.* 1999, Fouilland *et al.* 2004); there are now known to be grazers of picoplankton size which have higher maximum specific growth rates than *Prochlorococcus*, and so could exert control over this picocyanobacterium (Gouillou *et al.* 1999). There can be significant discrimination among picophytoplankton by grazers, e.g. between *Prochlorococcus* and *Synechococcus* (Christaki *et al.* 1999, Worden *et al.* 2004). There is thus the frequent interposition of another trophic level, i.e. ciliate and flagellate grazers, between the primary producers and the many larger zooplankton grazers for picophytoplankton but not for larger phytoplankton, possibly causing a reversal

in the direction of top-down effects on different size categories of phytoplankton by, for example, changes in the populations of the organisms consuming the larger phytoplankton.

CONCLUSIONS

Many of the ecological and evolutionary aspects of picophytoplankton can be related to their small size. However, 'biology' complicates almost all of the arguments made purely on the basis of cell size (Table I).

ACKNOWLEDGEMENTS. – Work in JAR's laboratory on phytoplankton is supported by BBSRC and NERC. ZVF and AJI are supported by NSERC. We are grateful for comments on the manuscript from Dr D Campbell and two anonymous reviewers, and for translation of the abstract into French by M Robertson.

REFERENCES

- Badger MR, Price GD 2003. CO₂ concentrating mechanisms in cyanobacteria: molecular components, their diversity and evolution. *J Exp Bot* 54(383): 609-622.
- Bertilsson S, Berglund O, Karl DM, Chisholm SW 2003. Elemental composition of marine *Prochlorococcus* and *Synechococcus*: implications for the ecological stoichiometry of the sea. *Limnol Oceanogr* 48(5): 1721-1731.
- Boyd CM, Gradmann D 2002. Impact of osmolytes of marine phytoplankton. *Mar Biol* 141(4): 605-618.
- Brand L E, Guillard RRL 1981. The effects of continuous light and light intensity on the reproduction rates of twenty-two species of marine phytoplankton. *J Exp Mar Biol Ecol* 50(2-3): 119-132.
- Bryant DA 2003. The beauty in small things revealed. *Proc Natl Acad Sci USA* 100(17): 9647-9649.
- Caron DA, Gobler CJ, Lonsdale DJ, Cerato RM, Schaffer RA, Rose JM, Buck NJ, Taylor G, Boissenneault KR, Mehran R 2004. Microbial herbivory on the brown tide alga *Aureococcus anophagefferans*: results from natural ecosystems, mesocosm and laboratory experiments. *Harmful Algae* 3(4): 439-457.
- Chisholm SW 1992. Phytoplankton size. In Falkowski PG & Woodhead AD eds, *Primary Productivity and Biogeochemical Cycles in the Sea*. Plenum Press, New York: 213-237.
- Chistaki U, Jacquet S, Dolan JR, Vaulot D, Rassoulzadegan F 1999. Growth and grazing on *Prochlorococcus* and *Synechococcus* by two marine ciliates. *Limnol Oceanogr* 44(1): 52-61.
- Courties C, Perasso R, Chretien-Dinet MJ, Gouillou L, Troussellier M 1998. Phylogenetic analysis and genome size of *Ostreococcus tauri* (Chlorophyta, Prasinophyceae). *J Phycol* 34(5): 844-849.
- Day TA, Neale PJ 2002. Effects of UV-B radiation on terrestrial and aquatic primary producers. *Annu Rev Ecol Systematics* 33: 371-396.

- Doblin MA, Blackburn SI, Hallgraeaf GM 1999. Comparative study of selenium requirements of three phytoplankton species: *Gymnodinium catenatum*, *Alexandrium minutum* (Dinophyta) and *Chaetoceros cf. tenuissimus*. *J Plankt Res* 24(6): 1153-1169.
- Docampo R, Moreno SNJ 2001. The acidocalcisome. *Molec Biochem Parasitol* 114(3): 151-159.
- Dufresne A, Garzarek L, Partensky F 2005. Accelerated evolution associated with genome reduction in a free-living prokaryote. *Genome Biol* 6(2): R14.1-R14.10.
- Durand MD, Green CJ, Sosik MM, Olson RJ 2002. Diel variations in optical properties of *Micromonas pusilla* (Prasinophyceae). *J Phycol* 38(6): 1132-1142.
- Evans C, Archer SD, Jacquet S, Wilson WH 2003. Direct estimates of viral lysis and microzooplankton grazing to the decline of a *Micromonas* spp. population. *Aquat Microbial Ecol* 30(3): 207-219.
- Falcon LI, Pluvinage S, Carpenter EJ 2005. Growth kinetics of marine unicellular N₂-fixing cyanobacterial isolates in continuous culture in relation to phosphorus and temperature. *Mar Ecol Progr Ser* 285: 3-9.
- Falkowski PG, Katz ME, Knoll AH, Quigg A, Raven JA, Schofield O, Taylor FJR 2004b. The evolutionary history of eukaryotic phytoplankton. *Science* 305(5682): 354-360.
- Falkowski PG, Schofield O, Katz ME, Van de Schootbrugge B, Knoll AH 2004. Why is the land green and the ocean red? In Theirstein H & Young J eds, *Coccolithophores – from molecular processes to global impact*. Elsevier, Amsterdam: 429-453.
- Finkel ZV, Irwin AJ, Schofield O 2004. Resource allocation alters the ¾ size scaling of metabolic rates in phytoplankton. *Mar Ecol Progr Ser* 273: 269-279.
- Finkel ZV, Irwin AJ 2000. Modelling size-dependent photosynthesis: light absorption and the allometric rule. *J Theoret Biol* 204(3): 361-369.
- Finkel ZV 2001. Light absorption and size scaling of light-limited metabolism of marine diatoms. *Limnol Oceanogr* 46(1): 86-94.
- Fogg GE 1986. Picoplankton. *Proc Roy Soc Lond B* 228(1250): 1-30.
- Fouilland E, Descolas-Gros C, Courties C, Collos Y, Vaquer A, Gasc A 2004. Productivity and growth of the smallest free-living eukaryote under nitrogen deficiency and sufficiency. *Microbial Ecol* 48(1): 103-110.
- Furnas MJ 1990. *In situ* growth rates of marine phytoplankton: approaches to measurement, community and species growth rates. *J Plankt Res* 12(6): 1117-51.
- Furnas MJ 1991. Net *in situ* growth rate of phytoplankton in an oligotrophic, tropical shelf ecosystem. *Limnol Oceanogr* 36(1): 13-29.
- Geider RJ, La Roche J 2002. Redfield revisited: variability of C: N: P in marine microalgae and its biochemical basis. *Eur J Phycol* 37(1): 1-17.
- Giordano M, Beardall J, Raven JA 2005. CO₂ concentrating mechanisms in algae: mechanisms, environmental modulation and evolution. *Annu Rev Plant Biol* 56: 641-658.
- Giovannoni SJ, Tripp HJ, Givan S, Podar M, Vergin KL, Baptista D, Bibbs L, Eads J, Richardson TH, Nooreweir M, Rappé MS, Short JM, Carrington JC, Mathur EJ 2005. Genome streamlining in a cosmopolitan oceanic bacterium. *Science* 309(5738): 1242-1245.
- Gouillou L, Chrétiennot-Dinet MJ, Boulben S, Moonvan der Staay SY, Vaultot D 1999. *Symbiomonas scintillans* and *Picophagus flagellatus* gen et sp nov (Heterokonta): two new heterotrophic flagellates of picoplanktonic size. *Protist* 150(4): 383-398.
- Hagemann M, Vinnemeir J, Oberpichler I, Boldt R, Bauwe H 2005. The glycine decarboxylase complex is not essential for a cyanobacterium *Synechocystis* sp. strain PCC 6803. *Plant Biol* 7(1): 15-22.
- Hess WR 2004. Genome analysis of marine photosynthetic microbes and their global role. *Curr Opin Biotech* 15(3): 191-198.
- Ho TY, Quigg A, Finkel Z, Milligan AJ, Wyman K, Falkowski PG, Morel FMM 2003. The elemental composition of some marine phytoplankton. *J Phycol* 39(6): 1145-1159.
- Horken KM, Tabita FR 1999. Closely related Form I ribulose biphosphate carboxylase/oxygenase molecules that possess different CO₂/O₂ substrate specificities. *Arch Biochem Biophys* 361(2): 183-194.
- Kana TM, Glibert PM 1987. Effect of irradiances up to 2000 µmol m⁻² s⁻¹ on marine *Synechococcus* WH7803. 1. Growth, pigmentation, and cell composition. *Deep Sea Res Part A Oceanogr Res Papers* 34(4) 479-495.
- Larkum AWD, Kühl M 2005. Chlorophyll *d*: the puzzle resolved. *Trends Plant Sci* 10(8): 355-357.
- Lawrence JE, Suttle CA 2004. Effect of viral infection on sinking rates of *Heterosigma akashiwo* and its implication for bloom termination. *Aquat Microbial Ecol* 37(1): 1-7.
- Mann NH, Clokie MRJ, Millard A, Cook A, Wilson WH, Wheatley PJ, Letarov A, Krisch HM 2005. The genome of S-PM2, a “photosynthetic” T4-type bacteriophage that infects marine *Synechococcus* strains. *J Bacteriol* 187(9) 3188-3200.
- Miller SR, Augustine S, Olson TL, Blankenship RE, Selker J, Wood AM 2005. Discovery of a free-living chlorophyll *d*-producing cyanobacterium with a hybrid proteobacterial-cyanobacterial small-subunit rRNA gene. *Proc Nat Acad Sci USA* 102(3): 850-855.
- Moreira D, López-García P 2002. The molecular ecology of eukaryotes unveils a hidden world. *Trends Microbiol* 10(1): 31-38.
- Nagasaki K, Tomaru Y, Katanozaka N, Shirai Y, Nishida K, Itakura S, Yamaguchi M 2004. Isolation and characterization of a novel single-stranded RNA virus infecting the bloom-forming diatom *Rhizosolenia setigera*. *Appl Env Microbiol* 70(2): 704-711.
- Nagasaki K, Tomura Y, Takao Y, Nishida K, Shirai Y, Suzuki H, Nagumo T 2005. Previously unknown virus infects marine diatom. *Appl Env Microbiol* 71(2): 3528-3535.
- Pustazzi F, MacIntyre H, Warren ME, Hutchins DA 2004. Interactions of nitrogen source and light intensity in the growth and photosynthesis of the brown tide alga *Aureococcus anophagefferens*. *Harmful Algae* 3(4): 343-360.
- Raven JA, Andrews JA, Quigg A 2005. The evolution of oligotrophy: implications for the breeding of crop plants for low input agricultural systems. *Ann Appl Biol* 146(3): 261-280.
- Raven JA, Waite A 2004. Tansley review. The evolution of silicification in diatoms: inescapable sinking and sinking as escape? *New Phytol* 162(1): 45-61.

- Raven JA 1984. A cost-benefit analysis of photon absorption by photosynthetic unicells. *New Phytol* 98(4): 593-625.
- Raven JA 1986. Physiological consequences of extremely small size for autotrophic organisms in the sea. In Platt T & Li WKW eds, Photosynthetic Picoplankton. *Can Bull Fish Aquat Sci* 214: 1-70.
- Raven JA 1998. The twelfth Tansley lecture. Small is beautiful: the picophytoplankton. *Funct Ecol* 12(4): 503-513.
- Rhodes LL, Peake BM, MacKenzie AL, Marvuch S 1995. Coccolithophores *Gephyrocapsa oceanica* and *Emiliana huxleyi* (Prymnesiophyceae equals Haptophyceae) in New Zealand coastal waters – characteristics of blooms and growth in laboratory culture. *N Z J Mar Freshwater Res* 29(3): 345-357.
- Rodriguez F, Derelle E, Guillou L, LeGall F, Vaulot D, Moreau H 2005. Ecotype diversity in the marine picoeukaryote *Ostreococcus* (Chlorophyta, Prasinophyceae). *Env Microbiol* 7(6): 853-859.
- Shalapyonok A, Olson RJ, Shalapyonok LS 1998. Ultra-dian growth in *Prochlorococcus* spp. *Appl Env Microbiol* 64(3): 1066-1069.
- Sommaruga R, Hofer JS, Alonso-Sáez L, Gasol JM 2005. Differential sunlight sensitivity of picophytoplankton from surface Mediterranean coastal waters. *Appl Env Microbiol* 71(4): 2154-2157.
- Stemmann L, Jackson GA, Ianson D 2004. Vertical model of particle size distributions and fluxes in the mid-water column that includes biological and physical process – Part I. Model formulation. *Deep Sea Res I* 51(7): 856-84.
- Stomp M, Huisman J, de Jongh F, Veraart A, Gerlia D, Rijkboer M, Ibelings BW, Wollfenzien UIA, Stal LJ 2004. Adaptive divergences in pigment composition promotes phytoplankton biodiversity. *Nature* 432(7013): 104-107.
- Sullivan MB, Coleman ML, Weigele P, Rohwer F, Chisholm SW 2005. Three *Prochlorococcus* cyanophage genomes: signature features and ecological interpretations. *PLoS Biology* 3(5): 0790-0806.
- Thingstad TF, Øvreås L, Egge JK, Løvdal T, Heldal M 2005. Use of non-limiting substrates to increase size; a generic strategy to simultaneously optimize uptake and minimize predation in pelagic osmotrophs? *Ecol Lett* 8(7): 675-682.
- Thronsdren J 1976. Occurrence and productivity of small marine flagellates. *Norw J Bot* 23(4): 269-293.
- Vaulot D, Romani K, Not F 2002. Are autotrophs less diverse than heterotrophs in marine picoplankton? *Trends Microbiol* 10(6): 266-267.
- Waite A, Fisher A, Thompson PA, Harrison PJ 1997. Sinking rate versus cell volume relationships illuminate sinking rate control mechanisms in marine diatoms. *Mar Ecol Progr Ser* 157: 97-108.
- Weast RC 1969/1970. Handbook of Chemistry and Physics. 50th Edition. Chemical Rubber Company, Cleveland, Ohio.
- Wilson WH, Carr NG, Mann NH 1996. The effect of phosphate status on the kinetics of cyanophage infection in the oceanic cyanobacterium *Synechococcus* sp. WH7803. *J Phycol* 32(4): 506-516.
- Wilson WH, Schroeder DC, Allen MJ, Holden MTG, Parkhill J, Barrell BG, Churcher C, Hamlin N, Mungall K, Norbertczak H, Quail MA, Price C, Rabbino-witsch E, Walker D, Craigon D, Roy D, Ghazal P 2005. Complete genome sequence and lytic phase transcription profile of a *Coccolithovirus*. *Science* 309(5737): 1090-1092.
- Worden AZ, Nolan JK, Palenik B 2004. Assessing the dynamics and ecology of marine picophycoplankton: The importance of the eukaryotic component. *Limnol Oceanogr* 49(1): 168-179.
- Zehr JP, Mellon MT, Zarn S 1998. New nitrogen-fixing microorganisms detected in oligotrophic oceans by amplification of nitrogenase (nifH) genes. *Appl Env Microbiol* 64(9): 3444-3450.

Reçu le 7 octobre 2005; received October 7, 2005

Accepté le 12 novembre 2005; accepted November 12, 2005