Diversity and taxonomic revision of tribes Rhipileae and Rhipiliopsidae (Halimedaceae, Chlorophyta) based on molecular and morphological data

Laura Lagourgue, Claude E Payri

► To cite this version:

Laura Lagourgue, Claude E Payri. Diversity and taxonomic revision of tribes Rhipileae and Rhipiliopsidae (Halimedaceae, Chlorophyta) based on molecular and morphological data. *Journal of Phycology*, In press, 10.1111/jpy.13186 . hal-03232042

HAL Id: hal-03232042

<https://hal.sorbonne-universite.fr/hal-03232042>

Submitted on 21 May 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

DIVERSITY AND TAXONOMIC REVISION OF TRIBES RHIPILEAE AND
RHIPILOPSIDEAE (HALIMEDACEAE, CHLOROPHYTA) BASED ON MOLECULAR
AND MORPHOLOGICAL DATA ⁽¹⁾

Laura Lagourgue ⁽²⁾

Sorbonne Universités, UPMC Univ Paris 06, IFD, 4 Place Jussieu, 75252 Paris Cedex 05,
France

UMR ENTROPIE (IRD, UR, UNC, Ifremer, CNRS), Institut de Recherche pour le
Développement, B.P. A5 Nouméa Cedex, Nouvelle-Calédonie, 98848, France.

Claude E. Payri

UMR ENTROPIE (IRD, UR, UNC, Ifremer, CNRS), Institut de Recherche pour le
Développement, B.P. A5 Nouméa Cedex, Nouvelle-Calédonie, 98848, France.

⁽²⁾ Corresponding Author: laura.lagourgue@ird.fr, (+687) 518710

RUNNING TITLE: Rhipileae and Rhipiliopsidae diversity and taxonomy

ABSTRACT:

Genera and species of the tribes Rhipileae and Rhipiliopsidae are abundant in most coral reef ecosystems worldwide. However, the group has been largely overlooked, and very little genetic data is available to accurately assess its diversity, phylogenetic relationships, and geographical distribution. Our study provided an in-depth reassessment of tribes Rhipileae and Rhipiliopsidae based on a species-rich dataset and the combination of molecular species

delimitation, multilocus phylogenetic analyses (*tufA*, *rbcL* and 18S rDNA), and morpho-anatomical observations. Our results revealed an unexpected diversity of 38 morphologically-validated species hypotheses, including 20 new species, two of which are described in this paper and one resurrected species (*Rhipilia diaphana*). Based on our phylogenetic results we proposed to redefine the genera *Rhipilia* and *Rhipiliopsis* and described two new genera, *Kraftalia* gen. nov. (Rhipileae) and *Rhipiliospina* gen. nov. (Rhipiliopsidae). Finally, we validated *Rhipiliella* Kraft and included it in tribe Rhipileae. Although *Rhipilia* and *Rhipiliopsis* have a pantropical distribution, none of the species studied here appeared cosmopolitan; instead, they have restricted distributions.

KEYWORDS: Chlorophyta; *Kraftalia* gen. nov.; macroalgae; phylogeny; Rhipiliaceae; *Rhipiliospina* gen. nov; siphonous; species delimitation.

38 *Abbreviations:* ABGD, automatic barcode gap discovery; AIC, akaike information criterion;
39 BEAST, bayesian evolutionary analysis sampling trees; bGMYC, bayesian general mixed
40 yule coalescent; BI, bayesian inference; bs, bootstraps; ESS, effective sample size; GMYC,
41 general mixed yule coalescent; GTR, general time reversible; K80, kimura model; MCCT,
42 maximum clade credibility tree; MCMC, markov monte carlo chain; ML, maximum
43 likelihood; mPTP, multi-rate poisson pree process; nov., nova/novum; PP, posterior
44 probabilities; PSH, primary species hypothesis; PTP, poisson tree process; RAXML,
45 randomized axelerated maximum likelihood; SSH, secondary species hypothesis; sp., species;
46 s.s., *sensu stricto*; tufA, elongation factor Tu.

INTRODUCTION

In algae, traditional taxonomy has long been based on morphological characters with, as a corollary, a multitude of poorly defined taxa or *nomina dubia* and a classification that only partially reflects the natural relationships among taxa (De Clerck et al., 2013; Leliaert et al., 2014). In current works, the contributions of DNA sequence data combined with morphological and often geographical criteria have made it possible to revise taxonomic ambiguities (*e.g.*, Vieira et al., 2014; Caragnano et al., 2018; Hughey et al., 2019). The siphonous green macroalgae Bryopsidales are a good example of a group for which morphologically based taxonomy has led to several problems and has been revised in several works, including the resurrection of old unused species names (*e.g.*, *Tydemania gardineri*, Lagourgue et al, 2020), the synonymy of others (*e.g.*, in *Codium*, Verbruggen et al., 2007), or the description of new taxa in response to the cryptic diversity revealed by DNA analyses (*e.g.*, whole order (Verbruggen et al., 2009a), Udoteaceae (Lagourgue and Payri, 2020) or *Halimeda* (Cremen et al., 2016)). Sequence-based species delimitation approaches are recognized as powerful tools to study species diversity (Luo et al., 2018). Many methods have been developed, either based on genetic distances or on phylogenetic trees. The species delimitation process can be used independently for the purpose of referencing genetic diversity, or as part of a broader integrative taxonomic approach to assist in both delimitation and species identification (*e.g.*, Bond and Stockman, 2008; Hotelling et al., 2016; Mason et al., 2016). Species delimitation approaches have been demonstrated as the best tool to assess macroalgal diversity (*e.g.*, Leliaert et al., 2014), and within the green algae, these tools have been successfully used for groups such as *Chlorella*-like species (Zou et al., 2016), *Boodlea* (Leliaert et al., 2009), the Udoteaceae (Lagourgue et al., 2018; Lagourgue and Payri, 2020) or Ulvophyceae (Sauvage et al., 2016). Species delimitation methods have also proved successful to detect cryptic species or, conversely, phenotypic plasticity (*e.g.*, Vieira et al.,

2014), which is critical for taxonomic baseline data, biodiversity inventories, or to better understand ecological, physiological or evolutionary processes. Through phylogenies and character state mapping, DNA sequence data are also essential for classifications to reflect natural relationships and for studying the evolution of morpho-anatomical characters across lineages. In particular, comparative phylogenetic methods (PCMs) are designed to study how an organism's morpho-anatomical characters or traits have changed over time and which have influenced speciation or extinction events. Although these methods are very powerful, the evolution of morphological characters has been inferred on phylogenies only in a few studies of Bryopsidales (*e.g.*, Verbruggen et al. (2007) on *Codium*; Lagourgue and Payri (2020) on Udoteaceae; Verbruggen et al. (2009b) on *Halimeda*; and Payri and Verbruggen (2009) on *Pseudocodium*). Finally, phylogenetic inference has also been used to decipher biogeographical history, using distribution data to estimate the lineages evolution in space and time (*e.g.*, Vieira et al., 2017, Leliaert et al., 2018, or Vieira et al., 2021).

Rhipileae and Rhipiliopsidae species are siphonous green macroalgae whose geographical distribution is mainly tropical and associated with coral reef ecosystems. These species inhabit a wide variety of habitats from the surface to 150 m depth (Eiseman and Earle, 1983). They are found in seagrass meadows, lagoons, reef patches, reef slopes, and some endolithic species are even found in coral skeletons (Marcellino and Verbruggen, 2016).

Except for two species, *Rhipilia tomentosa* and *Johnson-sea-linkia profunda*, recorded from the Caribbean region, most species are distributed in the Indo-Pacific region.

The family Rhipiliaceae was merged with the family Halimedaceae by Cremen et al. (2019) and its species transferred to two tribes: Rhipileae and Rhipiliopsidae. The former, described initially by Hillis-Collinvaux (1984), was emended by Cremen et al. (2019) and now includes species of *Rhipilia*, the monospecific genus *Johnson-sea-linkia*, *Pseudochlorodesmis* sp., *Boodleopsis pusilla*, and *Boodleopsis* sp. Cremen et al. (2019) also proposed the new tribe

Rhipiliopsidae to accommodate two species: *Rhipiliopsis peltata* and *Callipsygma wilsonis*. The Rhipiliaceae was initially proposed by Dragastan et al. (1997) to distinguish the genera *Rhipilia*, *Rhipiliopsis* and *Rhipiliella*, and the fossil genus *Baratangia*, from other members of the Udoteaceae. Molecular phylogenetic analyses confirmed that *Rhipilia* and *Rhipiliopsis* are genetically distinct from Udoteaceae (Verbruggen et al., 2009c), while in the absence of genetic data, *Rhipiliella* was maintained within the Udoteaceae. Additionally, phylogenetic studies, including representative *Rhipilia* and *Rhipiliopsis*, revealed that none of these genera was monophyletic (Verbruggen et al., 2009a, c; Cremen et al., 2019). Cremen et al. (2019) also showed that Rhipiliaceae was polyphyletic, as *Rhipilia* and *Rhipiliopsis* do not form a monophyletic clade, and *Rhipiliopsis* rather branches as a sister lineage to *Halimeda* and *Callipsygma*. They resurrected *Johnson-sea-linkia* to accommodate *Rhipiliopsis profunda* and resolved the polyphyly of *Rhipiliopsis*. Tribes Rhipileae and Rhipiliopsidae are not as well-known as the closely related Udoteae, Halimedae or Caulerpaceae, for which unexpected species diversity has been revealed (Verbruggen et al., 2005a, b; Sauvage et al., 2013; Lagourgue and Payri, 2020). Indeed, most of the Rhipileae and Rhipiliopsidae species have been described from morphological characters only, and the DNA sequence data available for these lineages is limited to five species of *Rhipilia*, two species of *Rhipiliopsis* and one species of each *Johnson-sea-linkia* and *Callipsygma*, most of which are represented by a single marker.

Morphologically, species of the former family Rhipiliaceae are non-calcified and they consist of an erect cylindrical stipe, sometimes very small (or even indistinct), anchored to the substratum by a rhizomatous base and topped with siphonous filaments (*i.e.*, siphons). These siphons are either free or joined into a flabelliform, peltate or cyathiform frond. Initially, the family was characterized by the presence of particular secondary structures that allow the adjacent siphons to adhere more or less firmly to each other and known as tenacula

in *Rhipilia* and papillae in *Rhipiliopsis*. *Rhipilia* includes 12 currently recognized species (Guiry and Guiry, 2020) and is morphologically diverse, ranging from fronds composed of free siphons (*e.g.*, *R. penicilloides* or *R. coppejansii*), to more or less fan- or funnel-shaped (infundibuliform) blades that can be thin or compact (*e.g.*, *R. tomentosa*/ *R. orientalis*). The tenacula of *Rhipilia* species can be of various shapes (forked, pronged, hook-shaped, bent, or discoid) and are observed throughout the frond or only at the base in species with free siphons. *Rhipiliopsis* currently includes 19 species (including *Johnson-sea-linkia profunda*, Guiry and Guiry, 2020) that are much smaller in size and more delicate than *Rhipilia* species. *Rhipiliopsis* species consist of a mono- or multisiphonous stipe and a mono- or pluristromatic blade (flabellate, peltate, or cyathiform). The papillae are less developed than the tenacula of *Rhipilia* but give a cohesive and net-like appearance to the blade. Four types of lateral cohesion have been described by Coppejans et al. (1999): papillae with or without a thickening ring, direct longitudinal contact between the siphons or adhesion by differentiated apices of siphons. Finally, *Rhipiliella* was proposed by Kraft (1986) to accommodate specimens with deciduous blades. The only species, *Rhipiliella verticillata*, is characterized by whorls of abscission scars left on the stipe by successively lost deciduous blades.

To date, these lineages are poorly documented genetically, likely because of their small size or their ecology, as they preferred habitats like cracks or crevices that are difficult to access (particularly *Rhipiliopsis* and *Rhipiliella*). The main objective of our study was to reassess the diversity and systematics of Rhipileae and Rhipiliopsidae using a combined morphological and molecular approach applied to a large specimen dataset, and to meet the different objectives of a multidisciplinary approach, using integrative taxonomy. A rich collection of specimens collected from most of the geographical range of the relevant species was used to acquire new molecular and morphological data. Using several methods, including molecular species delimitation, multilocus phylogenetic analyses (*tufA*, *rbcL* and 18S rDNA),

and morpho-anatomical observations, we aimed to (1) explore species diversity, (2) analyze species phylogenetic relationships, and, where necessary, (3) resolve taxonomic ambiguities within these lineages.

MATERIAL AND METHODS

Sampling

A total of 587 Rhipileae and Rhipiliopsidae samples were included in this study. They were collected by the authors and several collaborators using SCUBA at various localities in the Indo-Pacific region (Table S1 in Supplementary Information). Vouchers were pressed-dried on herbarium sheets and mainly housed at NOU, GENT, MEL, and PERTH (herbarium abbreviations follow Thiers (2021), continuously updated). Subsamples were preserved in 95 % ethanol and silica gel for DNA analyses, and in a formaldehyde solution (5% in seawater) for morpho-anatomical studies.

DNA extraction, amplification, and sequencing

Extractions were conducted using the Plant mini Kit (Qiagen Inc, Valencia, CA, USA) for *Rhipilia* and CTAB protocol for all other genera. Two chloroplast markers, *tufA* and *rbcL*, and the 18S rDNA nuclear gene were sequenced using previously published primers (Kooistra, 2002; Lam and Zechman, 2006; Verbruggen et al., 2009c; Händeler et al., 2010) (see Table S2 in Supplementary Information). In some instances, the *rbcL* and 18S rDNA genes were amplified in two fragments (*rbcL*5' and *rbcL*3'; 18S5' and 18S3'). PCR reactions were conducted in a final volume of 25 μ L including 1X of AmpliTaq Gold 360 Master Mix (Applied Biosystems), 0.4 μ M of each primer, 3 % of dimethylsulfoxide (DMSO), 0.4 μ g. μ L⁻¹ of bovine serum albumin (BSA) and 1 ng. μ L⁻¹ of DNA. PCR programs follow Lagourgue et al. (2018), and the Sanger sequencing reaction was carried out by Genoscreen

(Lille, France). Sequences were edited with Geneious version 7.1.9 (<http://www.geneious.com>, Kearse et al., 2012). Additional sequences were retrieved from GenBank (18 *tufA*, 16 *rbcL*, and one 18S rDNA) and added to our dataset. All sequences were aligned for each marker separately using the MUSCLE algorithm available in the Geneious software. The CLUSTAW algorithm was also used for DNA regions that were difficult to align (*e.g.*, 18S rDNA gene). Species delimitation methods were performed on the two chloroplast datasets independently, while phylogenetic reconstructions were performed on a multilocus (*tufA*, *rbcL* and 18S rDNA) concatenated matrix.

Phylogenetic reconstructions

Phylogenetic reconstructions for species delimitation analyses were performed for each marker individually, selecting only distinct haplotypes in each dataset, and using maximum likelihood (ML) and Bayesian inference (BI) for ultrametric trees. The datasets were analyzed with Partition Finder v1.1.0 to determine the most suitable evolutionary models according to the Akaike information criterion (AIC). For the evaluation of partition schemes, *rbcL* was tested both as one entire marker and as two distinct datasets (*rbcL*5' and *rbcL*3'; *i.e.*, the two-fragment sequencing scheme) because of differences in sequencing success and sampling sizes. ML trees were reconstructed in RAXML (Stamatakis, 2014) on the CIPRES web portal (Miller et al., 2010) (see Table S3 in Supplementary Information for more details and analyses parameters). Bayesian ultrametric trees were computed using BEAST (Drummond et al., 2012). The global clock hypothesis was rejected (Likelihood ratio test in MEGA 6, Tamura et al., 2013), and the two analyses were performed under a relaxed lognormal molecular clock associated with a coalescent constant size tree prior, as recommended by Monaghan et al. (2009). For each run, the convergence of the Markov Chains Monte Carlo (MCMC), and the effective sample sizes (> 200) were checked in Tracer v.1.6 (Rambaut and

Drummond, 2007). Runs were then combined using Log Combiner without the first 10% generations, removed as burn-in. The Maximum Clade Credibility Tree (MCCT) was then calculated using Tree Annotator (included in the BEAST package).

For the final phylogenetic analyses, ML and BI reconstructions were performed on multilocus matrices (*tufA*, *rbcL*, and 18S). The first dataset (*i.e.*, dataset #1 in Table S3) included several representative members of the suborder Halimedineae (data detailed in Table S1 in Supplementary Information) to assess the taxonomic position and composition of the tribes. Two other datasets were created to represent the Rhipileae (dataset #2) and Rhipiliopsidae (dataset #3) tribes, including only one specimen per species for supra-generic level analyses. Finally, for analyses at the genus level, datasets with several representatives per species were assembled (datasets #4 to 7), provided that sequences were available for at least two of the three markers - except for *Rhipilia tomentosa* and *Rhipiliopsis reticulata*, which were not present in our collection, and for which only one sequence each was available on GenBank (*rbcL* and 18S, respectively). *Boodleopsis* and *Pseudochlorodesmis* were excluded from our analyses since both filamentous genera are unresolved (cf. Cremen et al., 2019). Outgroup species, partition schemes, evolutionary models used, and reconstruction parameters for ML and BI trees are detailed in Table S3 (Supplementary Information) for each analysis.

Bayesian phylogenetic analyses were performed in MrBayes v.3.2 (Ronquist and Huelsenbeck, 2003) through the CIPRES web portal. The effective sample size (ESS>200) values and the Markov chain Monte Carlo (MCMC) convergence were checked in TRACER v.1.5 (Rambaut and Drummond, 2007) before computing a consensus topology and posterior probabilities. ML reconstructions were conducted in RAXML (Stamatakis, 2014) also through the CIPRES web portal.

Species delimitation

Five species delimitation methods were used in combination to assess species boundaries. They included four tree-based methods: the General Mixed Yule Coalescent (GMYC) (Pons et al., 2006), its Bayesian implementation: bGMYC (Reid and Carstens, 2012), the Poisson tree process model (hPTP, Zhang et al., 2013) and the Multi-rate version, mPTP (Kapli et al., 2017); and a distance-based method: the Automatic Barcode Gap Discovery (ABGD, Puillandre et al., 2012a). We chose to combine several methods because each is based on different assumptions and models, which allows balancing the biases specific to each of them. Indeed, searching for congruence between the results of each method and between markers allows converging towards the most robust species hypotheses (Carstens and Knowles, 2007; Dupuis et al., 2012; Puillandre et al., 2012b; Carstens et al., 2013; Leliaert et al., 2014; Rannala, 2015). The delimitation methods allowed us to define primary species hypotheses (PSHs), while searching for congruence between markers and methods led us to select secondary species hypotheses (SSHs), which were then confirmed or not using morpho-anatomical information. Besides, comparing molecular-based hypotheses to non-genetic data (*e.g.*, morpho-anatomical, ecological) is recommended to corroborate species boundaries (Carstens and Knowles, 2007; Wiens, 2007; Fujita et al., 2012; Carstens et al., 2013; Talavera et al., 2013).

Before applying species delimitation methods, datasets were treated using the Collapsetypes v4.6 perl script (Chesters, 2013) to prevent potential bias linked to identical haplotypes, as recommended by Pons et al. (2006) and Reid and Carstens (2012). Species delimitation methods were then applied as follows:

The ABGD method was applied directly to each marker sequence alignments. The *tufA* marker was analyzed using the single distance method, with parameter X (relative minimum gap width) set at 0.8. For *rbcL*, two sets of data were analyzed, the *rbcL*5' and *rbcL*3' fragments, taking into account the imbalance in the amplification performance of the two

markers and the sensitivity of the methods to missing data. The Kimura and the Single Distance methods were applied to the *rbcL5'* and *rbcL3'* datasets, respectively; parameter X was set to 0.8.

GMYC analyses were performed in R (R Development Core Team, 2019) using the "splits" package on the MCCTs obtained with BEAST for each marker. The bGMYC method was applied to a subsample of 100 trees from the same analyses. After exploratory tests, the *tufA* analysis was run for 30M generations and sampled every 100 generations with a burn-in of 10,000 generations. The *rbcL* analysis was run for 20 M generations and sampled every 100 generations with a burn-in of 5 M generations.

The hPTP method was implemented on the online server (<http://sco.hits.org/exelixis/web/software/PTP/index.html>) using ML trees and 500,000 generations sampled every 100 generations, for both markers. The mPTP analyses were performed on both the ML and MCCT trees for both markers and via the website (<http://mPTP.hits.org>) using default settings.

Morphology

Species identification and observation of morpho-anatomical characters were based on the most relevant literature reference for the group: the work of Gepp and Gepp (1911), including several *Rhipilia* species and one *Rhipiliopsis* species; the monograph of Millar and Kraft (2001) as well as the work of N'Yeurt and Keats (1997), and Verbruggen and Schils (2012), among others, for *Rhipilia*; and mainly the works of Kraft (1986 and 2000), Farghaly and Denizot (1979), Eiseman and Earle (1983), Norris and Olsen (1991) and Coppejans et al. (1999) for *Rhipiliopsis*. The morpho-anatomical characters observed included the shape of the thallus, the frond, the stipe (*Rhipilia*) or stalk (*Rhipiliopsis*) and the stolon (for *Rhipilia*); the diameter and appearance of the siphons and the type of dichotomies and constrictions; the

shape, size, frequency, and position of the tenacula (*Rhipilia*) or papillae (*Rhipiliopsis*) on the siphons.

RESULTS AND DISCUSSION

Species delimitation analyses

A total of 906 sequences (*tufA*: 440 sequences; *rbcL*: 363 sequences; 18S rDNA: 103 sequences) were successfully produced, to which we added sequences available in GenBank (*i.e.*, for 25 additional specimens). The list of sequences and corresponding specimens and accession numbers are presented in Table S1 (Supplementary Information). The variability of datasets is reported in Table S4 (Supplementary Information). The phylogenetic analyses of the multilocus matrix (*tufA*, *rbcL*, and 18S rDNA) at the suborder level (Figure S1) led us to consider three clades for the species delimitation approach: a group corresponding to tribe Rhipileae (including specimens of *Rhipilia*, *Rhipiliopsis* and *Rhipiliella*), a “Rhipiliopsidae group 1” (including specimens of *Rhipiliopsis* and *Callipsygma*), and a “Rhipiliopsidae group 2” (including specimens of *Rhipiliopsis*). We have followed this architecture in subsequent analyses, but it is important to note that the relationships among the different tribes are only weakly supported.

Exploratory species delimitation analyses of tufA and rbcL datasets: For the *tufA* dataset, all lineages combined, a total of 14 PSHs were common to all five methods. A summary of the species delimitation results for the *tufA* marker for all lineages is presented in Table 1. The detailed results for each lineage (Rhipileae, “Rhipiliopsidae group 1” and “Rhipiliopsidae group 2”) are presented in Supplementary Information (Appendix S1 and Figures S2 to S4). The support values and *a posteriori* probabilities (PP) associated with the partitions delimited

by hPTP and bGMYC, respectively, are detailed in Appendix S2 and Table S5 (Supplementary Information).

For the *rbcL* dataset and all groups included, a total of 14 SSHs were common to all five methods (see Appendix S3 and Figure S5 and S6 for detailed results on Rhipileae and Rhipiliopsidae lineages). The summary of species delimitation results are presented in Table 1. The hPTP partitions support values, and the PP of the bGMYC partitions are detailed in Appendix S4 and Table S6 (Supplementary Information).

SSH definition and species name assignment: At the level of markers, a significant number of PSHs were common to all species delimitation methods (Table 1). Thirty-six SSHs were unambiguously defined based on the PSHs common to both markers. Three additional SSHs were more difficult to define due to discrepancies between the two markers. The resolution process is detailed in Table S7.

In total, 39 SSHs were delimited within the Rhipileae and Rhipiliopsidae, of which only 16 could be unambiguously named using morpho-anatomical observations. Two additional SSHs still await confirmation before final species name assignment (SSH20: *R. sp1* cf. *mortensenii*; SSH29: *R. sp14* cf. *echinocaulos*). Twenty SSHs could not be assigned to current species and probably represent new species. One SSH (SSH21) was only represented by GenBank sequences and could not be morphologically analyzed. Details about the species assignment of SSHs are available in Figures S2 to S6 (Supplementary Information).

Marker variability and the need to combine them: The chloroplast markers, *tufA* and *rbcL*, were used in the species delimitation approach due to their variability and discriminatory power at the species level, as recognized in previous studies (Verbruggen et al., 2009c; Saunders and Kucera, 2010; Leliaert et al., 2014). In this study, both markers proved to be

effective in providing species hypotheses and discriminating between species, in addition to being good substitutes for barcodes (*i.e.*, for species delimitation and identification, respectively, *sensu* Collins and Cruickshank, 2013). The nuclear 18S rDNA marker was more conserved than chloroplast markers and, therefore, not appropriate for species delineation analyses. However, this marker, which has been used in previous studies at various taxonomic levels (Kooistra et al. (2002) for *Halimeda*, Kazi et al. (2013) for *Caulerpa*, Lagourgue et al. (2018) for Udoteaceae, Verbruggen et al. (2009a, c) for the Bryopsidales), remained relevant for phylogenetic analyses, in combination with other markers.

The performance of species delimitation methods: The performance of the species delimitation methods depends on the context, particularly the dataset analyzed (Knowles and Carstens, 2007), since all statistical methods are sensitive to a lack of information on intraspecific variability (Puillandre et al., 2012b; Kekkonen and Hebert, 2014). Adding samples or genetic markers leads to improve species signatures (Knowles and Carstens, 2007), helps to resolve ambiguous cases or conversely, may reveal different partition schemes. However, some methods may be biased toward species discrimination and not recognize the phylogenetic signature of speciation, particularly in cases of rapid and recent diversification events (or adaptive radiations), as revealed by short terminal branches in phylogenetic trees (such as GMYC in Kubatko and Degnan (2007) and Luo et al. (2018)). In this study, these biases were observed in the results of (m)PTP and PTP methods when analyzing tribe Rhipileae, and with the PTP method for the analysis of “Rhipiliopsidae group 1”. Conversely, GMYC produced a higher number of partitions than the other methods, but, for *tufA*, GMYC (and bGMYC) led to species hypotheses that were the closest to those morphologically identified.

Species delimitation results also directly depend on the selection of genetic markers and their variability. In this study, species delimitation analyses were conducted independently on the two markers, following Kubatko and Degnan's (2007) recommendations. The combination of several methods in our study revealed a significant congruence between them. In addition, when methods based on genetic distances were found to be congruent with those based on phylogenetic trees, as observed several times during this study, the robustness of SSHs was increased (Ross et al., 2010; Fujita et al., 2012). However, using several methods was necessary, as none of them alone was able to delimit species defined *a posteriori*. Comparing different methods is important to counterbalance the bias of starting hypotheses, concepts or models, and to overcome the limits inherent in each method, and finally, to define the most likely species hypotheses (Carstens and Knowles, 2007; Dupuis et al., 2012; Puillandre et al., 2012b; Carstens et al., 2013; Leliaert et al., 2014; Rannala, 2015). In addition, taking into account data of different types makes possible the identification of possible differences in the evolutionary signal and is therefore particularly recommended to compare molecular results to non-genetic data, such as the morphological observations used here (Carstens and Knowles, 2007; Knowles and Carstens, 2007; Wiens, 2007; Fujita et al., 2012; Carstens et al., 2013; Talavera et al., 2013).

Morphology remains essential: In our study, morpho-anatomical observations were successfully used to unambiguously validate and assign 16 SSHs to known species and 20 SSHs to new entities and to document the morphological diversity of genera. The identification of SSHs was hampered by the limited genetic data available and erroneous species assignments in published sequences. Direct examination of the sequenced specimens and access to morphological studies were essential to detect misidentifications and to assign SSHs to the correct species. However, besides being time-consuming, the morpho-anatomical

approach involves particular best practices, such as the availability of type specimens to ensure correct species assignment, or the study of a large number of specimens to accurately document within species polymorphism (see Wiens and Servedio, 2000). Hence, the success of concomitant morphological and molecular approaches presupposes that the morphological and anatomical characters are sufficiently documented.

Phylogenetic relationships

Global-scale phylogeny (Suborder Halimedineae): Our phylogenetic reconstructions at the scale of the suborder Halimedineae (dataset #1; cf. Table S3) confirmed that the former family Rhipiliaceae does not form a monophyletic group (Figure S1), corroborating earlier results by Cremen et al. (2019). However, our analysis, which included more samples and species than in previous studies, resulted in three major lineages (Figure S1), and not two as in Cremen et al.'s study, corresponding to: (1) tribe Rhipileae (bs: 100; PP:1) containing the type genus of the tribe: *Rhipilia*; (2) "Rhipiliopsidae group 1" (bs: 100; PP:1) including the genus *Callipsygma* (bs: 100; PP:1) and a group of *Rhipiliopsis*-like specimens; and (3) "Rhipiliopsidae group 2" (bs: 98; PP:1) also containing *Rhipiliopsis*-like species and branching as a sister lineage to the genus *Halimeda* (although not well supported: bs: 72; PP: 0.90). These results indicate that the tribe Rhipiliopsidae, erected by Cremen et al. (2019), is likely polyphyletic.

Additionally, *Rhipiliopsis* species (or at least *Rhipiliopsis*-like specimens) were found in all three lineages, with the type-species for the genus, *R. peltata*, included in "Rhipiliopsidae group 2". The polyphyly of the genus was already shown by Cremen et al. (2019), who reinstated the genus *Johnson-sea-linkia profunda*, the basionym of *Rhipiliopsis profunda* (tribe Rhipileae), in an attempt to solve the *Rhipiliopsis* polyphyly. With a more extensive selection of specimens and taxa, our results point out the need for more taxonomic revisions.

At least another five *Rhipiliopsis*-like species were found in tribe Rhipileae, 7 in “Rhipiliopsidae group 1”, and 9 in “Rhipiliopsidae group 2”.

Phylogeny of the Rhipileae lineage: Our phylogenetic reconstructions based on the Rhipileae multilocus matrix including one specimen per species (dataset #2; cf. Table S3) indicated that the tribe can be subdivided into two groups (Figure 1). The first (group 1, bs: 47; PP: 0.98, Fig. 1) is further divided into a strongly supported subclade (A) containing the type and six other species of *Rhipilia* on the one hand, and (B) four sequences referring to four genera (*Rhipilia*, *Rhipiliopsis*, *Johnson-sea-linkia* and *Rhipiliella*) on the other hand. The second group consists of several *Rhipiliopsis* and *Rhipilia* species (group 2, bs: 99; PP: 1, Fig. 1). Although the polyphyly of the genus *Rhipilia* was shown previously (Verbruggen et al., 2009a and c), the extent of it is more significant in our study. We found *Rhipilia* species in three different sections of the tree. One clade contains the type species, *R. tomentosa*. A second clade is composed of five *Rhipilia* species clustering with four *Rhipiliopsis* species (group 2, Fig. 1), and *R. pusilla* represents the third clade.

Following this result, we consider the clade containing the type species, *Rhipilia tomentosa*, as representing the genus *Rhipilia* (Group1, B in Figure 1). Species clustering in the second and third sections thus need to be revised and transferred to other genera. Millar and Kraft (2001) proposed various subdivisions for *Rhipilia* based on the abundance and shape of tenacula, but this classification is not compatible with our results. Indeed, species with both rare (e.g., *R. penicilloides*) and abundant tenacula (e.g., *R. tomentosa*) have been found in the same group. Similarly, the grouping of species according to the shape of the fronds (blade or free siphons) did not produce monophyletic groups in our phylogeny. Although species with free siphon fronds, such as *R. penicilloides* or *R. coppejansii*, were mainly found in the first group, they also clustered with fan-shaped frond species (*R. sp1* and *R. tomentosa*). In our

study, simple forms (with free siphons, and no or few and poorly developed tenacula) did not appear as ancestral characters, such as hypothesized by Womersley (1984) or Millar and Kraft (2001). We observed that species with simple morphologies appear independently throughout the Rhipileae lineage, instead of forming a single clade. This observation is the same as for the evolution of the morpho-anatomical characters of the tribe Udoteae, which does not follow a "from simple to complex" scenario; rather, complex character states have been estimated at the family's ancestral node and then, complex and simple morphologies are found at random in the various lineages of the tribe (see Lagourgue and Payri, 2020). Other similar evolutionary examples are known in macroalgae, including several life history traits in brown algae that do not follow a simple to complex scenario (*e.g.*, heteromorphy/isomorphy, numbers of plastids, fertilization, growth or the macroscopic thallus architecture in brown algae crown radiation (BACR) orders (Silberfeld et al., 2010; Bringloe et al., 2020)), or the current uni- and multicellularity representations among the Ulvophyceae (Del Cortona et al., 2019).

Species in group 2 (*Rhipilia* and *Rhipiliopsis*-like species) also require taxonomic revision as the type species for both *Rhipilia* and *Rhipiliopsis* belong to other clades (Fig. 1 and 2). We propose the new genus *Kraftalia* gen. nov. to accommodate the nine species of group 2. Our results also confirmed that *Rhipiliella* should be included in tribe Rhipileae as proposed by Dragastan et al. (1997; as family Rhipiliaceae). Finally, we maintain the taxonomic status of the monospecific genera *Rhipiliella* and *Johnson-sea-linkia*.

Phylogeny of the “Rhipiliopsidae” lineages: Our phylogenetic reconstructions of the Rhipiliopsidae lineages (from the multilocus matrix including one specimen per species; dataset #3 cf. Table S3) produced two well-supported non-sister clades (bs: 100; PP: 1, Figure 2) containing both *Rhipiliopsis*-like species. “Rhipiliopsidae group 1” was further subdivided

into two moderately to strongly supported subclades, one representing *Callipsygma* (bs: 83; PP: 0.97) and the other one clustering seven *Rhipiliopsis*-like species (bs: 100; PP: 1). “Rhipiliopsidae group 2” formed a well-supported clade sister to *Halimeda* (bs: 72; PP: 0.9) and contained nine *Rhipiliopsis*-like species, including the *Rhipiliopsis* type species (Figure 2).

Considering the polyphyly of *Rhipiliopsis*, we propose the following taxonomic solutions: (i) to redefine *Rhipiliopsis* and include only species clustering with its type-species, *Rhipiliopsis peltata* (i.e., “Rhipiliopsidae group 2”); and (ii) to describe *Rhipiliospina* gen. nov., to accommodate the *Rhipiliopsis*-like species of “Rhipiliopsidae group 1”.

We also considered two options to solve the polyphyly of the Rhipiliopsidae lineages: (i) maintain the tribe for its type genus, *Rhipiliopsis*, and describe a new tribe to accommodate species of “Rhipiliopsidae group 1” (i.e., *Callipsygma* and *Rhipiliospina*); or (ii) merge all three genera into the monogeneric tribe Halimedae. For the time being, we believe the genera should remain in tribe Rhipiliopsidae until more data is collected and more reliable node supports are obtained to demonstrate whether the tribe is monophyletic and taxonomically valid or polyphyletic and requires taxonomic revision.

It is interesting to note that here again, genera with complex morpho-anatomy, such as *Halimeda* or *Udotea*, are phylogenetically more related to genera with much simpler and more delicate forms, such as *Callipsygma* or *Chlorodesmis*, than to each other. The morphological contrast between closely related taxa appears as a recurrent phylogenetic pattern in the suborder Halimedineae (cf. Fig. S1 and examples given above).

Systematic revision and diversity of the various genera

The diversity and the necessary taxonomic revisions of the various Rhipileae and Rhipiliopsidae genera included in this study (existing, revised, and new ones) are discussed below based on molecular, morphological, and phylogenetic results.

Rhipilia (tribe *Rhipileae*): Our multilocus phylogeny (several representatives per species; dataset # 4 cf. Table S3) indicated that *Rhipilia* should be revised to include seven species only (and not 12 as currently recorded in Guiry and Guiry, 2020) (Figure 3). Three of them are currently accepted species: *R. tomentosa* (the type species), *R. penicilloides*, and *R. coppejansii*, to which we add *R. diaphana* (resurrected here), and three other undescribed species, *R. sp1*, *R. sp2*, and *R. sp3* (Figure 3). The resolution of ambiguous species hypotheses in the delimitation analyses, and the morphological verification of some GenBank specimens, could reveal additional species. For instance, GenBank sequences identified as *R. nigrescens* and *R. orientalis* clustered with our specimens of *R. diaphana* (Figures 3, S2 and S3). Additional genetic data and careful morphological analyses could help to make the correct taxonomic decision regarding these specimens.

Rhipilia diaphana is currently regarded a synonym of *R. orientalis* (Millar and Kraft, 2001), but both species appear genetically distinct. The latter was confirmed from specimens collected in Papua New-Guinea, which fully matched the original diagnosis (Gepp and Gepp, 1911; type locality: Fau Island, Eastern Indonesia). *Rhipilia diaphana* was also identified in our collection, among specimens from the Solomon Islands and Fiji, particularly from deep habitats (60 and 70 m), which are similar to those of the type locality (Bikini Island, Marshall Is., samples dredged from 50 m). Specimens also morphologically matched the diagnosis of Taylor (1950). We thus propose to resurrect *R. diaphana*. The latter can be distinguished from *R. orientalis* by its longer stipe, broader and thinner blades and its soft green color (Taylor, 1950). *Rhipilia orientalis* is generally smaller in size, with a thicker

blade, darker in color, and blackens as it dries (Taylor, 1950). We also found that the two species are anatomically distinguished by numerous tenacula and the presence of basal constrictions in *R. diaphana*, whereas tenacula are rare and not constricted in *R. orientalis*. Although *Rhipilia* includes species with widely diverging morphologies, its species have several characters in common, including the presence of a stolon (although *R. tomentosa* has been observed without stolon), dichotomies with a subdichotomous bulge and supra-dichotomous constrictions, and simple tenacula (2-3 prongs, up to four in *R. sp1*). *Rhipilia* has a pantropical distribution extending from the Indo-Pacific to the northwestern Atlantic (Caribbean). In our study, none of the species was present in all three oceans. Most species appeared restricted to small geographical areas, such as *R. sp1* or *R. sp3*, which were collected only in the southwest Pacific, whereas *R. coppejansii* was found throughout the Indo-Pacific. *Rhipilia tomentosa*, described from the Caribbean (Antigua), was found only at this locality during our study. Records from the Pacific (*e.g.*, Carolina Islands (Tsuda, 1972), Australia (Millar and Kraft, 2001), the Philippines (Ang et al., 2014)) and in the Indian Ocean (Seychelles (Titlyanova et al., 1992)), which are based on morphological observations only, should be confirmed with DNA sequences. Indeed, we assigned several of our specimens from the Chesterfield Islands to *R. tomentosa* based on the morphological description of specimens from Australia by Millar and Kraft (2001). However, our DNA analyses revealed that they actually belong to the new genus *Kraftalia*, and that *R. tomentosa* is probably restricted to the Atlantic. Any record of *R. tomentosa* from outside this region is a possible misidentification. Overall, in the absence of combined DNA analysis and in-depth morpho-anatomical observations, *Rhipilia* species can be easily confused, which could partly explain overestimated geographical ranges.

The new genus *Kraftalia* (tribe *Rhipileae*): The results of our phylogeny (Fig. 1; from dataset #2) indicated the need to describe a new genus for nine *Rhipilia* and *Rhipiliopsis*-like species clustering in a strongly supported subclade of tribe *Rhipileae* (bs: 99; PP: 1, Fig. 1). *Kraftalia* gen. nov is proposed to accommodate the four species *Rhipilia orientalis*, *Rhipilia crassa*, *Rhipiliopsis yaeyamensis*, and *Rhipiliopsis gracilis*, as well as five other undescribed species. *Kraftalia* is characterized by a fan-shaped frond, the absence of stolon, relatively thin siphon diameters ($< 100\ \mu\text{m}$), and the cohesion of siphons by one or more particular types of structures (direct longitudinal contact, papillae, differentiated siphons or tenacula). *Kraftalia* is found in the Indo-Pacific with species restricted to specific geographical areas (Western Indian Ocean, West Pacific) (Figure 4). Only *K. crassa* occurs both in the Indian Ocean and the West Pacific. In our study, *K. orientalis* was collected only in the Indian Ocean and the Coral Triangle. There is no specimen from the Pacific or Atlantic oceans corresponding to this species, which raises questions about published records (Guiry and Guiry, 2020; as “*Rhipilia orientalis*”). For example, records of *K. orientalis* in southern Japan (Itono, 1986; as *Rhipilia orientalis*) could be *Rhipilia diaphana*, which is morphologically very similar and has a predominantly Pacific distribution. Again, verification is needed for GenBank sequences to confirm the correct geographical distribution of these species.

Rhipiliella (tribe *Rhipileae*): Our study provides the first genetic record of the monospecific *Rhipiliella*, containing only *R. verticillata*. Our species delimitation analyses, however, indicated that there is possibly more than one species, although these species hypotheses require confirmation with additional specimens from a more extensive geographical range. *Rhipiliella* is monophyletic and well-supported (bs: 100; PP: 1; Figure S7; from dataset #5). *Rhipiliella* is morphologically distinct from other related genera by the presence of scars from deciduous blades along the monosiphonous stipe, its monostromatic blade, and the presence

of papillae (Kraft, 1986). The specimens in our collection come from two different localities in New Caledonia (Grande Terre and Surprise Is.), which is not far from the type locality on the Australian Great Barrier Reef (Wistari Reef). It also perfectly matched the original diagnosis (Kraft, 1986). To date, the geographical distribution of *Rhipiliella* is restricted to the southwestern Pacific (Figure S7).

Other Rhipileae species: Additional species are clustering in tribe Rhipileae but their phylogenetic positions are not well supported and/or species richness is not enough documented to proceed with taxonomic revisions (*e.g.*, species such as *Pseudochlorodesmis* sp. or *Boodleopsis* sp. were not included) (Figure 1). Indeed, some species are represented by only one or a few specimens (*R. profunda* or *R. pusilla*), or by specimens from a single geographical area (New Caledonia for *Rhipiliella* and *R. cf. mortensenii*).

Rhipilia pusilla is one of them. It is sister to *Johnson-sea-linkia profunda* (Figures 1 and S7). *Rhipilia pusilla* is distinguished by a frond with free siphons, anisomorphic dichotomies and rare tenacula (Ducker, 1967; Womersley, 1984), while *J. profunda* is characterized by intersecting (“criss-cross”) siphons visible on the blade, and does not have scars left by deciduous blades along the stipe (Eiseman and Earle, 1983). In the absence of stronger phylogenetic support, and because of the limited morphological similarities to justify the grouping of these two species, we prefer to maintain the genus *Johnson-sea-linkia* and leave the status of *R. pusilla* in question.

The last Rhipileae species that branches separately is “*Rhipiliopsis*” cf. *mortensenii* (Figures 1 and S7). It is interesting to note that *R. mortensenii* was the type species of the genus *Geppella* (family Codiaceae) before the genus became a synonym of *Rhipiliopsis*. Here, the position of the species outside *Rhipiliopsis* raises the question of the resurrection of the genus *Geppella* (although the other ex-*Geppella* species do not cluster with

R. mortensenii). However, the weak node supports and lack of genetic data to correctly assess the species richness of the possible genus prevented us from reliably concluding about its taxonomic status.

These three species are geographically restricted. *Johnson-sea-linkia profunda* is only found in the Caribbean, *R. pusilla* in Southern Australia, and *R. cf. mortensenii* in New Caledonia and surrounding islands (Figure S7). Additional phylogenetic analyses on geographically larger datasets are needed to resolve the phylogenetic relationships within this set of species.

Rhipiliopsis (tribe *Rhipiliopsidae*): “*Rhipiliopsidae* group 2” corresponds to *Rhipiliopsis* sensu stricto (Figure 2; N.B.: although the position of the type species, *R. peltata*, is not the same in all trees (Fig. 2 and 5), maybe due to differences in sample size, it still represents “*Rhipiliopsidae* group2”). The results of our species delimitation analyses (Figure 5; dataset #6) indicate that the revised genus consists of nine species. They include *R. peltata* (the type-species), *R. corticata*, *R. reticulata*, *R. papuensis*, and five additional species, which have yet to be described: *R. sp5*, *R. sp6*, *R. sp7*, *R. sp8*, and *R. sp9*.

Based on our data, the genus *Rhipiliopsis* s.s. is characterized by the following: a strongly corticated stipe (ascending siphons or protuberances), supra-dichotomous constrictions and two types of adhesions between the siphons, *i.e.*, papillae of type I (bilateral contact in H structure) or II (unilateral). Interestingly, morphologically similar species can occur in very distant localities, *e.g.*, *R. corticata* from New Caledonia and *R. sp5* from Madagascar; or *R. reticulata* from the Caribbean and its sister species *R. sp7* from Madagascar (Figure 5).

According to our distribution map (Figure 5), each species of *Rhipiliopsis* s.s. is restricted geographically. Still, the genus has a cosmopolitan distribution, with *R. reticulata* occurring

in the Atlantic and other species in the Indo-Pacific (four in the Western Indian Ocean and five in the West Pacific).

However, our dataset for this group is relatively limited (*e.g.*, only one sequence for some species and limited node support for others), and a more comprehensive sampling is needed to better document species diversity, phylogenetic relationships, and geographical distribution.

The new genus Rhipiliospina (tribe Rhipiliopsidae): We propose *Rhipiliospina* gen. nov. to accommodate *Rhipiliopsis*-like species clustering in clade “Rhipiliopsidae group 1” (bs: 100; PP: 1; Figure 2). According to our delimitation analyses, the new genus includes seven species (Figure 6; dataset #7). Six require formal description (the type species, *R. stellifera* sp. nov. is described in Taxonomic Treatment section), and one requires further investigation (*R. sp5* cf. *Rhipiliopsis echinocaulos*). Each species is strongly supported (bs>98; PP: 1) except *Rhipiliospina* sp4 (bs: 89; PP: 1).

Rhipiliospina gen. nov. is characterized by a monosiphonous and corticated stipe with very remarkable spines (hence the genus name), which are simple or forked. Besides, all species have broad dichotomies without subdichotomous bulge, but with marked supra-dichotomous constrictions. Siphons adhere to each other by papillae of type I (bilateral contact in H structure) or II (unilateral).

Based on our data, the genus has a strict Indo-Pacific distribution. In this study, we found that these species have geographically restricted ranges and could be endemic to them. For instance, *R. sp2* has only been collected from the Isle of Pines in New Caledonia, and *R. sp6* is so far only known from the Chesterfield and Surprise islands in New Caledonia.

Rhipiliospina sp1 has the widest distribution and is found both in southern Japan and Papua New-Guinea (Figure 6).

The genus Callipsygma (tribe Rhipiliopsidae): *Callipsygma* is currently known as a monospecific genus and is reported only from Australia. In our analyses, specimens of the genus *Callipsygma* formed a well-supported clade (bs: 92; PP: 1, Fig. 6) branching as a sister lineage to the new genus *Rhipiliospina*. The results of our species delimitation analyses indicate that it consists of two species, including the type-species *Callipsygma wilsonis* and a new species, *Callipsygma brevis* sp. nov. (Figure 6).

The genus *Callipsygma* is characterized by an upper vegetative part composed of free siphons adhering together by lateral ramifications (Gepp and Gepp, 1911). The two species *C. wilsonis* and *C. brevis* can be distinguished from each other by the smaller size of thallus and stipe length in *C. brevis* and the diameter of their siphons, which is more than twice as large in the type species. They also have distinct geographical distributions, with the type species known only from Australia and the new species so far being collected only from northern Madagascar. The addition of the latter to *Callipsygma* thus extends the geographical distribution of the genus to the Western Indian Ocean (Figure 6).

Using different tools to better understand taxonomy and diversity

Our integrative taxonomic approach used a combination of tools to explore the diversity, phylogeny and systematics of the tribes Rhipileae and Rhipiliopsidae. They included species delimitation methods, based on genetic data, and morpho-anatomical observations. The species delimitation approach was used as a first step in a comprehensive integrative taxonomy approach to map species diversity (and not only genetic diversity) and resolve taxonomic ambiguities. The phylogenetic approach was also used to study and assess the diversity of the different genera and their evolutionary relationships within the two tribes. Our results underline the systematic value of morpho-anatomical characters in an integrative taxonomy approach, as already pointed out by several authors (Cianciola et al., 2010; Vieira

et al., 2014, Lagourgue et al., 2018) and the importance of combining morphological and genetic data. Without proper molecular-based species delimitation analyses, some of the species would not have been distinguished using morphological analyses alone. Similarly, without morpho-anatomical observations, most of the SSHs defined by the species delimitation analyses could not have been assigned to correct species due to the lack of available valid genetic data for most species (*e.g.*, *Rhipilia diaphana*). Also, a number of species hypotheses would not have been verified and confirmed. In phycology, most taxonomic studies are based on morphology resulting in an inestimable amount of information. The morphological characters recorded in the literature are critical to identify species, but their relevance and diagnostic robustness need to be verified, particularly in the context of taxonomic revision. The combination of morphological and molecular approaches has proven relevant, if not essential, to assess specific diversity accurately and provide correct species identifications.

It is by combining all these complementary and relevant tools and methods that we have been able to provide a significant taxonomic update about the diversity and phylogenetic relationships among members of tribes Rhipileae and Rhipiliopsidae.

TAXONOMIC TREATMENT

Rhipilia Kützing emend.

Description emended from Kützing (1858) and Gepp and Gepp (1911): Thallus uncalcified, green, stipitate or sessile, with a stolon, and with a frond of variable form, flabellate, cuneate, peltate, infundibuliform or composed of free siphons, sometimes zonate. Siphons cylindrical, straight, bent or tortuous, 20-320 µm in diameter, very laxly interwoven and dichotomously branched. Dichotomies have a subdichotomous bulge and supra-dichotomous

666 constrictions, with often a cell-wall thickening. Blade siphons (sometimes only basal siphons)
667 have at least one of the four types of adhesion structures: 1) tenacula with 2-3(4) prongs, often
668 with basal constrictions; 2) discoid tenacula; 3) hook-shaped tenacula; 4) differentiated bent
669 siphons apices.

670 *Distribution (confirmed by DNA sequences):* Atlantic Ocean: Mexico (Lam and Zechman,
671 2006); Indian Ocean: Madagascar (This study), Mayotte (This study), Western Australia
672 (Scott Reef) (Verbruggen et al., 2009; Verbruggen and Schils, 2012); Pacific Ocean: New
673 Caledonia (Chesterfield Is., Surprise Is., Grande Terre) (This study), Fiji (This study), Guam
674 (Verbruggen and Schils, 2012), Papua New-Guinea (This study), Solomon Is. (This study),
675 Australia (Queensland: Heron Is.; Masthead Is.) (Verbruggen and Schils, 2012); Tuvalu (This
676 study); Southwestern Asia: Philippines (Verbruggen et al., 2009).

677 *Type species:* *R. tomentosa* Kützinger; *Type locality:* Antigua, Antilles, West Indies; *Lectotype:*
678 MEL 14088 (and 13 isoelectotype specimens).

679 *List of other species (confirmed by DNA sequences in this study):* *R. penicilloides*, *R.*
680 *coppejansii*, *R. diaphana*, and three undescribed species.

681

682 We also propose the resurrection of:

683 *Rhipilia diaphana* W.R.Taylor 1950: 72, 205, pl. 37

684 *Type locality:* Bikini Atoll, Marshall Is.

685 *Type:* Holotype: Taylor, 13.iv.1946, MICH 1306664 (=WRT 46-195), dredged from 57 m.

686 *Description emended from Taylor (1950):* Uncalcified thalli, composed of a creeping stolon, a
687 simple or compound stipe, from which arise a flabellate frond. The frond is large, thin and
688 diaphanous, green in color, and zonate. Siphons are visible at the surface, they are tortuous,
689 subparallel, rarely interwoven, 30-55 μm (up to 50-60 μm) in diameter; Siphons are
690 dichotomously divided with isomorphic and lax dichotomies, subdichotomous bulges, and

691 symmetrical supra-dichotomous constrictions with cell-wall thickening. Siphons have many
692 adhesion structures that are found all along the blade, and which correspond either to two-
693 pronged tenacula (150-300 μm long) with basal constriction or spines.

694 *Distribution (confirmed by DNA sequences in this study*)*: Pacific Ocean: Fiji*, Solomon
695 Is.*, Marshall Is. (type locality, no DNA data).

696 *List of vouchers (limited to two per locality)*: Fiji, 2007: NOU 204022, NOU 204069;
697 Solomon Is., 2006: NOU 087399, NOU 087400

698

699 ***Kraftalia* Lagourgue & Payri gen. nov., Figure 7.**

700 *Type species: Kraftalia orientalis* (A. Gepp and E.S. Gepp) Lagourgue and Payri **comb. nov.**;

701 *Basionym: Rhipilia orientalis* A. Gepp and E.S. Gepp 1911: 57, 140, pl. XVI: figs. 134-136

702 *Description*: Uncalcified thalli, anchor system (no stolon), a corticated or uncorticated stalk,
703 which can be mono or multisiphonous, and a fan-shaped blade, which can be mono or
704 pluristromatic. Siphons are dichotomously divided with or without supra-dichotomous
705 constrictions. Siphons diameter are $< 100 \mu\text{m}$. Cohesion between siphons is due to one or
706 several types of adhesion structures (direct longitudinal contact, differentiated siphons,
707 papillae or tenacula).

708 *Etymology*: The name honors Dr. Gerald T. Kraft, who described three of the nine species
709 included in the genus.

710 *Distribution (confirmed by DNA sequences*)*: Indian Ocean: Madagascar* (This study), Juan
711 de Nova* (This study), Western Australia* (Scott Reef) (Verbruggen et al. 2009), Mayotte*
712 (This study); Southwestern Asia: Indonesia (Bunaken*) (This study), Philippines*
713 (Verbruggen et al., 2009); Pacific Ocean: Australia (Heron Is.)(type locality, no DNA data),
714 Papua New-Guinea* (This study), Tuvalu* (This study), Fiji (This study), New Caledonia*
715 (Chesterfield Is., Surprise Is., Grande Terre) (This study), Japan* (Sauvage et al., 2016).

716 *Species included in the genus (confirmed by molecular data in this study): Kraftalia*
717 *orientalis*, *K. crassa*, *K. gracilis*, *K. yaeyamensis*, and five undescribed species.

718

719 We propose the following new combinations for the transfer of selected *Rhipilia* species to
720 the new genus *Kraftalia*:

721 *Kraftalia orientalis* (A. Gepp and E.S. Gepp) Lagourgue & Payri **comb. nov.**

722 *Basionym:* *Rhipilia orientalis* A. Gepp and E.S. Gepp 1911: 57, 140, pl. XVI: figs. 134-136

723 *Syntypes localities:* Fau Island, Malay Archipelago; Pulu Sebangkatan, Borneo Bank

724 *Type:* n°334; L 3997222 (holotype); fig. 134a of Gepp and Gepp 1911, ex L 937, 279...308 =
725 MELU A235, and MICH 21873 (lectotypes); MICH 23026 (isotype)

726 *Description emended from Gepp and Gepp (1911) and Millar and Kraft (2001), see also Fig.*
727 *7A, 7F, and 7K:* Plants uncalcified, brownish-green to yellow-green (blackening when dried),
728 small (6-10 cm in length), without stolon, stipitate with simple or compound narrow and short
729 stipes (up to 1 cm long, 0.1—0.2 cm thick), expanding above into the frond. Frond
730 flabelliform to infundibuliform or peltate, small and thick (mostly 1—3 (up to 6 cm) cm-long,
731 1—2.5 (rarely 4) cm-wide), soft and finely meshed, almost like brown-stained muslin, not or
732 rarely zonate and with rounded to lacerate margins. Frond siphons (22-) 30-36 (-55) µm in
733 diameter, straight or slightly bent, interwoven, with a recurved, rounded or swollen apex.
734 Siphons are dichotomously divided with asymmetrical supra-dichotomous constrictions and
735 slight cell-wall thickening. Cohesion between siphons are due to either (i) simple, short and
736 stubby pronged-tenacula (2-3 prongs, variable in length: (70) 170-500 µm-long) without basal
737 constriction; (ii) hook-shaped tenacula without basal constriction; or (iii) differentiated
738 siphons (adhesion by rounded apex). Adhesion structures are rare.

739 *Distribution (confirmed by DNA sequences in this study*)*: South-east Asia: Indonesia
740 (Borneo Bank, Fau Is.) (type locality, no DNA data); Indian Ocean: Mayotte* (this study);
741 Pacific Ocean: Papua New-Guinea* (this study).

742 *List of vouchers (limited to two per locality)*: Mayotte, 2010: NOU 204163, NOU 204170;
743 2016: NOU 203544, NOU 203569; Papua New-Guinea, Madang, 2012: NOU 203532, NOU
744 204123; Papua New-Guinea, Kavieng, 2014: NOU 203350, NOU 203353

745

746 *Kraftalia crassa* (A.J.K. Millar and Kraft) Lagourgue & Payri **comb. nov**
747 *Basionym*: *Rhipilia crassa* A.J.K. Millar and Kraft 2001: 32, figs 37-40, 53-58
748 *Type locality*: Heron Island, Capricorn Group, Great Barrier Reef, Australia
749 *Type*: MELU A37571 (holotype); MELU A35070 and A37569-74 (isotypes).

750 *Description*: see Millar and Kraft (2001; see also Fig. 7B, 7G, and 7L

751 *Distribution (confirmed by DNA sequences*)*: Indian Ocean: Madagascar* (This study), Juan
752 de Nova* (This study), Western Australia* (Scott Reef) (Verbruggen et al. 2009);
753 Southwestern Asia: Indonesia* (Bunaken) (this study), Philippines* (Verbruggen et al.,
754 2009); Pacific Ocean: Japan* (Sauvage et al., 2016); Australia (Heron Island) (type locality,
755 no DNA data).

756 *List of vouchers (limited to two per locality)*: Madagascar, 2016: NOU 203728, NOU 203731;
757 Juan de Nova, 2013: NOU 204191; Indonesia, Bunaken, 2014: NOU 203475, NOU 203483.

758

759 *Kraftalia gracilis* (Kraft) Lagourgue & Payri **comb. nov.**
760 *Basionym*: *Rhipiliopsis gracilis* Kraft 1986: 55, figs 17-21
761 *Type locality*: Heron Island, Capricorn Group, Great Barrier Reef, Australia
762 *Type*: MELU K16136 (holotype); MELU KI5568 and MELU KI6161 (isotypes).

763 *Description*: see Kraft (1986); see also Fig. 7C, 7H, and 7M.

764 *Distribution (confirmed by DNA sequences*)*: Pacific Ocean: New Caledonia* (Chesterfield,
 765 Grande Terre, Surprises Is.) (This study); Australia (Heron Is.) (type locality, no DNA data)
 766 *List of vouchers (limited to two per locality)*: New Caledonia, Chesterfield, 2015: NOU
 767 203281, NOU 203320; New Caledonia, Grande Terre, 2017: NOU 203756, NOU 203866;
 768 New Caledonia, Surprise Is., 2017: NOU 203949, NOU 203963.
 769
 770 *Kraftalia yaeyamensis* (Tanaka) Lagourgue & Payri **comb. nov.**
 771 *Basionym*: *Geppella yaeyamensis*, Tanaka 1963: 65, figs 2, 3
 772 *Type locality*: Iriomotejima, Funauke, Ryukyu Island, Japan
 773 *Type*: T. Tanaka, 2.xi.1959, 20m deep (holotype)
 774 *Synonym*: *Rhipiliopsis yaeyamensis* (Tanaka) Kraft 1986: 71
 775 *Description*: see Tanaka (1963) and Kraft (1986); see also Fig. 7D, 7I, and 7N.
 776 *Distribution (confirmed by DNA sequences*)*: Pacific Ocean: New Caledonia* (Grande Terre,
 777 Surprises); Japan (type locality, no DNA data)
 778 *List of Vouchers (limited to two per locality)*: New Caledonia, Grande Terre, 2017: NOU
 779 203750, NOU 203762; New Caledonia, Isle of Pines, 2013: NOU 203405, NOU 203406;
 780 New Caledonia, Surprise Is., 2017: NOU 203903, NOU 203915.
 781
 782 *Rhipiliopsis* s.s. A. Gepp and E.S. Gepp 1911: 57, 140, pl. XVI: figs. 134-136
 783 *Description*: see Gepp and Gepp (1911).
 784 *Distribution (confirmed by DNA sequences*)*: Atlantic: Antilles (type locality, no DNA data);
 785 Panama* (Kooistra, 2002); Indian Ocean: Maldives Is.* (This study), Madagascar* (This
 786 study); Pacific Ocean: Australia* (Victoria) (Cremen et al., 2019); Lord Howe Is. (type
 787 locality, no DNA data); New Caledonia* (Chesterfield, Grande Terre, Surprise Is.) (This
 788 study), Papua New-Guinea* (This study).

789 *Type species: R. peltata* (J. Agardh) A. Gepp and E.S. Gepp

790 *Type:* Agardh, LD 15800 (BM)

791 *Type locality:* Port Phillip Heads, Victoria, Australia

792 *Basionym:* *Udotea peltata* J. Agardh.

793 *Other species included in the genus (as a result of the present study):* *Rhipiliopsis corticata*,

794 *R. reticulata*, *R. papuensis*, and five undescribed species.

795

796 ***Rhipiliospina* Lagourgue & Payri gen. nov.**

797 *Type species: Rhipiliospina stellifera* Lagourgue & Payri **sp. nov.**

798 *Description:* Uncalcified thalli composed of a monosiphonous and corticated stipe with very

799 remarkable spines, simple or forked, and a flabelliform or cyathiform frond, mono or

800 pluristromatic. Siphons dichotomously divided and < 50 µm in diameter. Broad dichotomies,

801 with deep supra-dichotomous constrictions. Adhesion of the siphons by papillae of type I

802 (bilateral contact in H structure) or II (unilateral).

803 *Etymology:* The name refers to its resemblance to the genus *Rhipiliopsis* and the presence of

804 remarkable spines on the stipe.

805 *Distribution (confirmed by DNA sequences):* Indian Ocean: Madagascar (This study); Pacific

806 Ocean: New Caledonia (Iles of Pines, Chesterfield, Grande Terre, Surprise Is.) (This study),

807 Papua New-Guinea (Madang) (This study); Japan (Sauvage et al., 2016; as *Rhipiliaceae* sp.)

808 *List of species:* *Rhipiliospina stellifera* and six undescribed species.

809

810 ***Rhipiliospina stellifera* Lagourgue & Payri sp. nov., Figure 8**

811 *Holotype:* NOU203095

812 *Type locality:* Ouen Islet, Canal Woodin, New Caledonia.

813 *Description:* Uncalcified thalli composed of a monosiphonous and corticated stipe (150 µm in
814 diameter), with forked and complex spines, including star-shaped spines, and a pluristromatic
815 rounded and flabelliform frond that is also thin and zonate. Siphons dichotomously divided,
816 tortuous, 10-30 µm in diameter, entangled in a disorganized network. Broad dichotomies with
817 a square or trapezoid shape, and symmetrical supra-dichotomous constrictions, with or
818 without cell-wall thickening. Siphon adhesion is provided by numerous and proximate
819 papillae of type I (bilateral contact in H structure) or II (unilateral) with a ring of cell-wall in
820 the contact zone. Papillae also adhere to siphons in different layers, giving a “3D” cortication
821 aspect.

822 *Etymology:* The name refers to the star-shaped spines on the stipe.

823 *Distribution confirmed by molecular data:* Pacific Ocean: New Caledonia (This study).

824 *List of vouchers and representative species sequences:* New Caledonia, Western lagoon, Voh,
825 2017: NOU 203758 (*tufA*: MT782677, *rbcL*: MT783058; 18S: MT782551); NOU 203761
826 (*tufA*: MT782798; *rbcL*: MT783164; 18S: MT782606); NOU 203764 (*tufA*: MT782722;
827 *rbcL*: MT783101); New Caledonia, Southern Lagoon, Ouen Isle, 2015: NOU 203095 (*tufA*:
828 MT782684; *rbcL*: MT783065; 18S: MT782553), NOU 203096 (*tufA*: MT782673).

829

830 ***Callipsygma brevis* Lagourgue & Payri sp. nov., Figure 9**

831 *Holotype:* NOU203608

832 *Type locality:* Madagascar, South, Diego Suarez Bay

833 *Description:* Uncalcified thalli, green, with a multisiphonous stipe and a tufted frond
834 composed of free siphons weakly adhering to each other by lateral ramifications, which form
835 a cohesive, feather-like whole. Stipe siphons with protuberances and deformed lateral
836 branches. Frond siphons lightly tortuous, thin, 50-70 µm in diameter, dichotomously divided,
837 and with rounded apices. Dichotomies (45°) with subdichotomous bulges and symmetrical

838 constrictions above, with a ring of cell-wall thickening almost occlusive; Adhesion between
839 siphons with a few circular, uni- or bilateral papillae.

840 *Etymology*: In reference to the size of the stipe and thallus, which are shorter than the type
841 species (*C. wilsonis*).

842 *List of vouchers and representative species sequences*: Madagascar, South, Diego Suarez
843 Bay, 2016: NOU 203608 (*tufA*: MT782750; *rbcL*: MT783124); NOU 203609 (*tufA*:
844 MT782810; *rbcL*: MT783174).

ACKNOWLEDGMENTS

This work was supported by the DUNE Labex-CORAIL and ENTROPIE funds. The authors are grateful to Heroen Verbruggen and Chiela Cremen for providing additional sequences. Many thanks to Francesca Benzoni for her help with latin names, and Lydiane Mattio for her critical review of the manuscript. The English language has been verified by blue[c]weed. Samples have been collected during several campaigns: Guadeloupe, 2014, Onema and MNHN fieldtrip; Lesser Antilles, 2015: R/V ANTEA, Pacotilles, [doi](#); Bunaken, 2014 : INDESO project (research permit 133/SIP/FRP/SM/V/2015 and 918/BLITBANKKP/II/2016); Fiji, 2007: R/V Alis, BSM-Fidji, [doi](#); Papua New-Guinea, Madang, 2012: NUIGUINI campaign, [doi](#); Papua New-Guinea, Kavieng, 2014: [doi](#); Madagascar, 2016: R/V Antea, MAD, [doi](#); Maldives Islands, 2009 : « Programme Maldives 2009 »; Mayotte, 2010 : TARA; 2016: SIREME; New Caledonia, 2005 : BSM-LOYAUTE, [doi](#); 2008 : CORALCAL2, [doi](#); 2012: CORALCAL4, [doi](#); Iles of Pines and Surprises Islands, 2013, LOF ; Iles of Pines and Chesterfield Islands, 2015: R/V ALIS, CHEST, [doi](#); Grande Terre, and Surprises Islands, 2017: R/V ALIS PostBlanco1 and TARA-NC ; Scattered Islands, Glorioso Is., 2012 and Juan de Nova, 2013: BIORECIE; Solomon Islands, 2004: R/V ALIS, BSM-Salomon; Tuvalu, 2014: “Programme SARGASSES”.

No conflict of interest.

REFERENCES

- Ang, P. O., Leung, S. M. & Choi, M. M. 2014. A verification of reports of marine algal species from the Philippines. *Philipp. J. Sci.* 142:5-49.
- Bond, J. E. & Stockman, A. K. 2008. An integrative method for delimiting cohesion species: Finding the population-species interface in a group of Californian trapdoor spiders with extreme genetic divergence and geographic structuring. *Syst. Biol.* 57:628–46.
- Bringloe, T. T., Starko, S., Wade, R. M., Vieira, C., Kawai, H., De Clerck, O., Cock, J. M., Coelho, S. M., Destombe, C., Valero, M., Neiva, J., Pearson, G. A., Faugeron, S., Serrão, E. A. & Verbruggen, H. 2020. Phylogeny and evolution of the brown algae. *Crit. Rev. Plant Sci.* 39:281–321.
- Caragnano, A., Foetisch, A., Maneveldt, G. W., Millet, L., Liu, L. C., Lin, S. M., Rodondi, G., & Payri, C. E. 2018. Revision of Corallinaceae (Corallinales, Rhodophyta): recognizing *Dawsoniolithon* gen. nov., *Parvicellularium* gen. nov. and Chamberlainoideae subfam. nov. containing *Chamberlainium* gen. nov. and *Pneophyllum*. *J. Phycol.* 54:391–409.
- Carstens, B. C. & Knowles, L. L. 2007. Estimating species phylogeny from gene-tree probabilities despite incomplete lineage sorting : An example from *Melanoplus* grasshoppers. *Syst. Biol.* 56:400–11.
- Carstens, B. C., Pelletier, T. A., Reid, N. M. & Satler, J.D. 2013. How to fail at species delimitation. *Mol. Ecol.* 22:4369–83.
- Chesters, D. 2013. Collapsetypes.pl. Available at: <https://sourceforge.net/projects/collapsetypes/> (last accessed 23 March 2020).
- Collins, R. A., & Cruickshank, R. H. 2013. The seven deadly sins of DNA barcoding. *Mol. Ecol. Resour.* 13:969–75.
- Coppejans, E., De Clerck, O. & Leliaert, F. 1999. New records of the green-algal genus

889 *Rhipiliopsis* (Udoteaceae, Caulerpales) from Papua New Guinea and the Seychelles Islands,
890 including the description of *Rhipiliopsis papuensis* Coppejans, De Clerck & Leliaert, sp. nov.
891 *Bot. Mar.* 42:377–82.

892 Cremen, C. M., Huisman, J. M., Marcelino, V. R., & Verbruggen, H. 2016. Taxonomic
893 revision of *Halimeda* (Bryopsidales , Chlorophyta) in south-western Australia. *Aust. Syst.*
894 *Bot.* 29:41–54.

895 Cremen, C. M., Leliaert, F., West, J., Lam, D. W., Shimada, S., Lopez-Bautista, J. M. &
896 Verbruggen, H. 2019. Reassessment of the classification of Bryopsidales (Chlorophyta) based
897 on chloroplast phylogenomic analyses. *Mol. Phylogenet. Evol.* 130:397–405.

898 De Clerck, O., Guiry, M. D., Leliaert, F., Samyn, Y., & Verbruggen, H. 2013. Algal
899 Taxonomy: A Road to Nowhere? *J. Phycol.* 49:215–25.

900 Dragastan, O., Richter, D. K., Kube, B., Popa, M., Sarbu, A. & Ciugulea, I. 1997. A new
901 family of paleo-mesozoic calcareous green siphons-algae (Order Bryopsidales, Class
902 Bryosidophyceae, Phylum Siphonophyta). *Rev. Esp. Micropaleontol.* 29:69–135.

903 Drummond, A. J., Suchard, M. A., Xie, D. & Rambaut, A. 2012. Bayesian phylogenetics with
904 BEAUti and the BEAST 1.7. *Mol. Biol. Evol.* 29:1969–73.

905 Ducker, S. 1967. The genus *Chlorodesmis* (Chlorophyta) in the Indo-Pacific region. *Nov.*
906 *Hedwigia* 13:145–82.

907 Dupuis, J. R., Roe, A. D. & Sperling, F. A. H. 2012. Multi-locus species delimitation in
908 closely related animals and fungi: one marker is not enough. *Mol. Ecol.* 21:4422–36.

909 Eiseman, N. J. & Earle, S. A. 1983. *Johnson-sea-linkia profunda*, a new genus and species of
910 deep-water Chlorophyta from the Bahama Islands, *Phycologia* 22:1-6.

911 Farghaly, M. S. & Denizot, M. 1979. A propos de trois especes nouvelles d’Udoteacees
912 (Caulerpales, Chlorophycees) considerees comme appartenant au genre *Rhipiliopsis* . *Rev.*
913 *Algol.* 14:71–72.

914 Fujita, M. K., Leache, A. D., Burbrink, F. T., McGuire, J. A. & Moritz, C. 2012. Coalescent-
 915 based species delimitation in an integrative taxonomy. *Trends Ecol. Evol.* 27:480–8.
 916 Gepp, A. & Gepp, E. S. 1911. The Codiaceae of the Siboga Expedition, including a
 917 monograph of Flabellarieae and Udoteaceae. *Siboga-Expeditie* 62:1–150.
 918 Guiry, M. D. & Guiry, G. M. 2020. AlgaeBase. World-wide electronic publication, National
 919 University of Ireland, Galway. <http://www.algaebase.org> (last accessed 20 January 2020)
 920 Händeler, K., Wägele, H., Wahrmund, U., Rüdinger, M. & Knoop, V., 2010. Slugs’ last
 921 meals: molecular identification of sequestered chloroplasts from different algal origins in
 922 Sacoglossa (Opisthobranchia, Gastropoda). *Mol. Ecol. Resour.* 10:968–78.
 923 Hotaling, S., Foley, M. E., Lawrence, N. M., Bocanegra, J., Blanco, M. B., Rasoloarison, R.,
 924 Kappeler, P. M., Barrett, M. A., Yoder, A. D., & Weisrock, D. W. 2016. Species discovery
 925 and validation in a cryptic radiation of endangered primates: coalescent-based species
 926 delimitation in Madagascar’s mouse lemurs. *Mol. Ecol.* 25:2029–45.
 927 Hughey, J. R., Maggs, C. A., Mineur, F., Jarvis, C., Miller, K. A., Shabaka, S. H., &
 928 Gabrielson, P. W. (2019). Genetic analysis of the Linnaean *Ulva lactuca* (Ulvales,
 929 Chlorophyta) holotype and related type specimens reveals name misapplications, unexpected
 930 origins, and new synonymies. *J. Phycol.* 55: 503–8
 931 Itono, H. 1986. New records of marine algae from southern parts of Japan. *Japanese J.*
 932 *Phycol.* 34:72–82.
 933 Kapli, P., Lutteropp, S., Zhang, J., Kobert, K., Pavlidis, P., Stamatakis, A. & Flouri, T. 2017.
 934 Multi-rate poisson tree processes for single-locus species delimitation under maximum
 935 likelihood and markov chain Monte Carlo. *Bioinformatics* 33:1630–8.
 936 Kazi, M. A., Reddy, C. R. K., & Jha, B. 2013. Molecular phylogeny and barcoding of
 937 *Caulerpa* (Bryopsidales) based on the *tufA*, *rbcL*, 18S rDNA and ITS rDNA genes. *PLoS*
 938 *ONE* 8:e82438.

939 Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton, S.,
 940 Cooper, A., Markowitz, S., Duran, C., Thierer, T., Ashton, B., Meintjes, P. & Drummond, A.
 941 2012. Geneious Basic: An integrated and extendable desktop software platform for the
 942 organization and analysis of sequence data. *Bioinformatics* 28:1647–49.

943 Kekkonen, M., & Hebert, P. D. N. 2014. DNA barcode-based delineation of putative species:
 944 efficient start for taxonomic workflows. *Mol. Ecol. Resour.* 14:706–15.

945 Knowles, L. L., & Carstens, B. C. 2007. Delimiting Species without Monophyletic Gene
 946 Trees. *Syst. Biol.* 56:887–95.

947 Kooistra, W. H. C. F. 2002. Molecular phylogenies of Udoteaceae (Bryopsidales,
 948 Chlorophyta) reveal nonmonophyly for *Udotea*, *Penicillus* and *Chlorodesmis*. *Phycologia*
 949 41:453–62.

950 Kooistra, W. H. C. F., Coppejans, E. G. G., & Payri, C. E. 2002. Molecular systematics,
 951 historical ecology, and phylogeography of Halimeda (Bryopsidales). *Mol. Phylogenet. Evol.*
 952 24:121–38.

953 Kraft, G. T. 1986. The green algal genera *Rhipiliopsis* A. & E.S. Gepp and *Rhipiliella* gen.
 954 nov. (Udoteaceae, Bryopsidales) in Australia and the Philippines. *Phycologia* 25:47–72.

955 Kraft, G. T. 2000. Marine and estuarine benthic green algae (Chlorophyta) of Lord Howe
 956 Island, south-western Pacific. *Aust. Syst. Bot.* 13:509–648.

957 Kubatko, L. S., & Degnan, J. H. 2007. Inconsistency of phylogenetic estimates from
 958 concatenated data under coalescence. *Syst. Biol.* 56:17–24.

959 Kützing, F. T. 1858. Tabulae phycologicae. oder, Abbildungen der Tange VIII, i–ii, 1–48 pp,
 960 100 pls.

961 Lagourgue, L., Puillandre, N. & Payri, C. E. 2018. Exploring the Udoteaceae diversity
 962 (Bryopsidales, Chlorophyta) in the Caribbean region based on molecular and morphological
 963 data. *Mol. Phylogenet. Evol.* 127:758–69.

964 Lagourgue, L., Verbruggen, H., Ampou, E. E., & Payri, C. E. 2020. One hundred years later,
 965 resurrection of *Tydemanian gardineri* A. Gepp & E. Gepp (Udoteaceae, Chlorophyta) based on
 966 molecular and morphological data. *Eur. J. Phycol.* 55:89–99.

967 Lagourgue, L. & Payri, C. E. 2020: Large scale diversity reassessment, evolutionary history,
 968 and taxonomic revision of the green macroalgae family Udoteaceae (Bryopsidales,
 969 Chlorophyta). *J. Syst. Evol.* 00:1-27.

970 Lam, D. W. & Zechman, F. W. 2006. Phylogenetic analyses of the Bryopsidales
 971 (Ulvophyceae, Chlorophyta) based on Rubisco large subunit gene sequences. *J. Phycol.*
 972 42:669–78.

973 Leliaert, F., Verbruggen, H., Wysor, B., & Clerck, O. De. 2009. DNA taxonomy in
 974 morphologically plastic taxa: Algorithmic species delimitation in the *Boodlea* complex
 975 (Chlorophyta: Cladophorales). *Mol. Phylogenet. Evol.* 53:122–33

976 Leliaert, F., Verbruggen, H., Vanormelingen, P., Steen, F., López-Bautista, J.M., Zuccarello,
 977 G. C. & De Clerck, O. 2014. DNA-based species delimitation in algae. *Eur. J. Phycol.* 49:
 978 179–96

979 Leliaert, F., Payo, D. A., Gurgel, C. F. D., Schils, T., Draisma, S. G. A., Saunders, G. W.,
 980 Kamiya, M., Sherwood, A. R., Lin, S.-M., Huisman, J. M., Le Gall, L., Anderson, R. J.,
 981 Bolton, J. J., Mattio, L., Zubia, M., Spokes, T., Vieira, C., Payri, C. E., Coppejans, E., ... De
 982 Clerck, O. 2018. Patterns and drivers of species diversity in the Indo-Pacific red seaweed
 983 *Portieria*. *J. Biogeogr.* 45:2299–313.

984 Luo, A., Ling, C., Ho, S. Y. W., Zhu, C. D., & Mueller, R. 2018. Comparison of methods for
 985 molecular species delimitation across a range of speciation scenarios. *Syst. Biol.* 67:830–846.

986 Mason, V. C., Li, G., Minx, P., Schmitz, J., Churakov, G., Doronina, L., Melin, A. D.,
 987 Dominy, N. J., Lim, N. T.-L., Springer, M. S., Wilson, R. K., Warren, W. C., Helgen, K. M.,
 988 & Murphy, W. J. 2016. Genomic analysis reveals hidden biodiversity within colugos, the

989 sister group to primates. *Science Advances* 2:e1600633.

990 Millar, A. J. K. & Kraft, G. T. 2001. Monograph of the green macroalgal genus *Rhipilia*
 991 (Udoteaceae, Halimedaes), with a description of *R. crassa* sp. nov. from Australia and the
 992 Philippines. *Phycologia* 40:21–34.

993 Miller, M. A., Pfeiffer, W. & Schwartz, T. 2010. Creating the CIPRES Science Gateway for
 994 inference of large phylogenetic trees, in: 2010 *Gateway Computing Environments Workshop*
 995 (*GCE*). IEEE, 1–8.

996 Monaghan, M. T., Wild, R., Elliot, M., Fujisawa, T., Balke, M., Inward, D. J. G., Lees, D. C.,
 997 Ranaivosolo, R., Eggleton, P., Barracough, T. G. & Vogler, A. P. 2009. Accelerated species
 998 inventory on Madagascar using coalescent-based models of species delineation. *Syst. Biol.*
 999 58:298–311.

1000 Montecinos, A. E., Couceiro, L., Peters, A. F., Desrut, A., Valero, M., & Guillemin, M. L.
 1001 2017. Species delimitation and phylogeographic analyses in the *Ectocarpus* subgroup
 1002 *siliculosi* (Ectocarpales, Phaeophyceae). *J. Phycol.*, 53:17–31.

1003 N’Yeurt, A. D. R. & Keats, D. W. 1997. *Rhipilia penicilloides* sp. nov. (Udoteaceae,
 1004 Chlorophyta) from Fiji. *Phycologia* 36:172–8.

1005 Norris, J. N. & Olsen, J. L. 1991. Deep-watter green algae from the Bahamas, including
 1006 *Cladophora Vandenhoekii* sp.nov (Cladophorales). *Phycologia* 30:315–28.

1007 Payri, C. E., & Verbruggen, H. 2009. *Pseudocodium mucronatum*, a new species from New
 1008 Caledonia, and an analysis of the evolution of climatic preferences in the genus (Bryopsidales,
 1009 Chlorophyta). *J. Phycol.* 45:953–61.

1010 Pons, J., Barracough, T. G., Gomez-zurita, J., Cardoso, A., Duran, D. P., Hazell, S., Kamoun,
 1011 S., Sumlin, W. D. & Vogler, A. P. 2006. Sequence-Based Species Delimitation for the DNA
 1012 Taxonomy of Undescribed Insects. *Syst. Biol* 55:595–609.

1013 Puillandre, N., Lambert, A., Brouillet, S. & Achaz, G. 2012a. ABGD, Automatic Barcode

Gap Discovery for primary species delimitation. *Mol. Biol. Evol.* 21:1864–77.

Puillandre, N., Modica, M. V, Gustave, O., Place, L. L. & West, C. P., 2012b. Large-scale species delimitation method for hyperdiverse groups. *Mol. Ecol.* 21:2671–91.

R Development Core Team, 2019. R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from <https://www.r-project.org/>

Rambaut, A., Drummond, A., 2007. Tracer version 1.5. Available at: <http://beast.bio.ed.ac.uk/Tracer> (Last accessed 18 February 2020).

Rannala, B. 2015. The art and science of species delimitation. *Curr. Zool.* 61:846–53

Reid, N. M. & Carstens, B. C. 2012. Phylogenetic estimation error can decrease the accuracy of species delimitation : a Bayesian implementation of the general mixed Yule-coalescent model. *BMC Evol. Biol.* 12:196.

Ronquist, F. & Huelsenbeck, J. P. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19:1572–4.

Ross, K. G., Gotzek, D., Ascunce, M. S., & Shoemaker, D. D. 2010. Species delimitation: A case study in a problematic ant taxon. *Syst. Biol.* 59:162–84.

Saunders, G. W., & Kucera, H. 2010. An evaluation of *rbcL*, *tufA*, UPA, LSU and ITS as DNA barcode markers for the marine green macroalgae. *Cryptogamie Algol.* 31:487–528.

Sauvage, T., Payri, C., Draisma, S. G., Prud'Homme van Reine, W. F., Verbruggen, H., Belton, G. S., Gurgel, F. D., Gabriel D., Sherwood A. R. & Fredericq S., 2013. Molecular diversity of the *Caulerpa racemosa*-*Caulerpa peltata* complex (Caulerpaceae, Bryopsidales) in New Caledonia, with new Australasian records for *C. racemosa* var. *cylindracea*. *Phycologia* 52:6–13.

Sauvage, T., Schmidt, W. E., Suda, S., & Fredericq, S. 2016. A metabarcoding framework for facilitated survey of endolithic phototrophs with *tufA*. *BMC Ecol.* 16:1–21.

1039 Silberfeld, T., Leigh, J. W., Verbruggen, H., Cruaud, C., de Reviers, B., & Rousseau, F. 2010.
 1040 A multi-locus time-calibrated phylogeny of the brown algae (Heterokonta, Ochrophyta,
 1041 Phaeophyceae): Investigating the evolutionary nature of the “brown algal crown radiation.”
 1042 *Mol. Phylogenet. Evol.* 56:659–74.

1043 Stamatakis, A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
 1044 large phylogenies. *Bioinformatics* 30:1312–13

1045 Talavera, G., Dinc, V. & Vila, R. 2013. Factors affecting species delimitations with the
 1046 GMYC model : insights from a butterfly survey, *Methods Ecol. Evol.* 4:1101–10.

1047 Tamura, K., Stecher, G., Peterson, D., Filipski, A. & Kumar, S. 2013. MEGA6: Molecular
 1048 Evolutionary Genetics Analysis version 6.0. *Mol. Biol. Evol.* 30:2725–9.

1049 Tanaka, T. 1963. Studies on some marine algae from Southern Japan. IV. *Mem. Fac. Fish.*
 1050 Kagoshima Univ. 12:1–6.

1051 Taylor, W. R. 1950. Plants of Bikini and other northern Marshall Islands. Ann Arbor:
 1052 University of Michigan Press. 227 pp, 79 pls.

1053 Thiers, B., 2021. Index Herbariorum: A Global Directory of Public Herbaria and Associated
 1054 Staff. New York Botanical Garden’s Virtual Herbarium.

1055 Titlyanova, T. V., Perestenko, L. P. & Kalugina-Gutnik, A. 1992. Catalogus algarum
 1056 benthicarum ac berbarium Insularum Seychellarum (Oceanus Indicus). *Nov. Sist.* 28:40–47.

1057 Tsuda, R. T. 1972. Some marine benthic algae from Truk and Kuop, Caroline Islands. *Atoll*
 1058 *Res. Bull.* 155:1–10.

1059 Verbruggen, H., De Clerck, O., Cocquyt, E., Kooistra, W. H. C. F. & Coppejans, E. 2005a.
 1060 Morphometric taxonomy of siphonous green algae: a methodological study within the genus
 1061 *Halimeda* (Bryopsidales). *J. Phycol.* 41:126–39.

1062 Verbruggen, H., De Clerck, O., Kooistra, W. H. C. F. & Coppejans, E. 2005b. Molecular and
 1063 morphometric data pinpoint species boundaries in *Halimeda* section *Rhipsalis* (Bryopsidales,

Chlorophyta). *J. Phycol.* 41:606–21.

Verbruggen, H., Leliaert, F., Maggs, C. A., Shimada, S., Schils, T., Provan, J., Booth, D.,
Murphy, S., De Clerck, O., Littler, D. S., Littler, M. M., & Coppejans, E. 2007. Species
boundaries and phylogenetic relationships within the green algal genus *Codium*
(Bryopsidales) based on plastid DNA sequences. *Mol. Phylogenet. Evol.* 44:240–54.

Verbruggen, H., Vlaeminck, C., Sauvage, T., Sherwood, A. R., Leliaert, F. & De Clerck, O.
2009a. Phylogenetic analysis of *Pseudochlorodesmis* strains reveals cryptic diversity above
the family level in the siphonous green algae (Bryopsidales, Chlorophyta). *J. Phycol.* 45:726–
31.

Verbruggen, H., Tyberghein, L., Pauly, K., Vlaeminck, C., Nieuwenhuyze, K. Van, Kooistra,
W. H. C. F., Leliaert, F., & Clerck, O. De. 2009b. Macroecology meets macroevolution:
evolutionary niche dynamics in the seaweed *Halimeda*. *Global Ecol. Biogeogr.* 18:393–405.

Verbruggen, H., Ashworth, M., LoDuca, S. T., Vlaeminck, C., Cocquyt, E., Sauvage, T.,
Zechman, F. W., Littler, D. S., Littler, M. M., Leliaert, F. & De Clerck, O. 2009c. A multi-
locus time-calibrated phylogeny of the siphonous green algae. *Mol. Phylogenet. Evol.* 50:642–
53.

Verbruggen, H. & Schils, T. 2012. *Rhipilia coppejansii*, a new coral reef-associated species
from Guam (Bryopsidales, Chlorophyta). *J. Phycol.* 48:1090–8.

Vieira, C., D’hondt, S., De Clerck, O., & Payri, C. E. 2014. Toward an inordinate fondness
for stars, beetles and *Lobophora* ? Species diversity of the genus *Lobophora* (Dictyotales,
Phaeophyceae) in New Caledonia. *J. Phycol.* 50:1101–19.

Vieira, C., Camacho, O., Sun, Z., Fredericq, S., Leliaert, F., Payri, C., & De Clerck, O. 2017.
Historical biogeography of the highly diverse brown seaweed *Lobophora* (Dictyotales,
Phaeophyceae). *Mol. Phylogenet. Evol.*, 110:81–92.

Vieira, C., Steen, F., D’hondt, S., Bafort, Q., Tyberghein, L., Fernandez-García, C., Wysor,

1089 B., Tronholm, A., Mattio, L., Payri, C., Kawai, H., Saunders, G., Leliaert, F., Verbruggen, H.,
 1090 & De Clerck, O. 2021. Global biogeography and diversification of a group of brown seaweeds
 1091 (Phaeophyceae) driven by clade-specific evolutionary processes. *J. Biogeogr.*, 48:703–15.
 1092 Wiens, J. J., & Servedio, M. R. 2000. Species delimitation in systematics : inferring
 1093 diagnostic differences between species. *Proc. R. Soc. Lond. B-Biol. Sci.* 267:631–6.
 1094 Wiens, J. J. 2007. Species delimitation: new approaches for discovering diversity. *Syst. Biol.*
 1095 56:875–8.
 1096 Will, K. W., Mishler, B. D., & Wheeler, Q. D. 2005. The perils of DNA barcoding and the
 1097 need for integrative taxonomy. *Syst. Biol.* 54:844–51.
 1098 Womersley, H. B. S. 1984. The marine benthic flora of southern Australia. Part I.,
 1099 Government Printer, Adelaide, 329 pp
 1100 Zhang, J., Kapli, P., Pavlidis, P. & Stamatakis, A. 2013. A general species delimitation
 1101 method with applications to phylogenetic placements. *Bioinformatics* 29:2869–76.
 1102 Zou, S., Fei, C., Song, J., Bao, Y., He, M., & Wang, C. 2016. Combining and comparing
 1103 coalescent, distance and character-based approaches for barcoding microalgae: a test with
 1104 *Chlorella*-like species (Chlorophyta). *Plos One* 11:e0153833.
 1105

TABLES

Table 1: Summary of the number of hypotheses delimited for each method applied to the *tufA* and *rbcL* datasets (alternative (b)GMYC partitions are indicated between brackets), including the number of singletons and summary of the number of PSHs common to all methods for each marker.

Methods		GMYC	bGMYC	hPTP	mPTP	ABGD
Number of delimited	<i>tufA</i>	48 7	(33)43 4	31 7	37 5	39 3
PSHs number of singletons	<i>rbcL</i>	47 10	37 8	31 12	41 12	42 9
PSHs in common (<i>tufA</i> <i>rbcL</i>)	bGMYC	38 30				
	hPTP	20 19	19 22			
	mPTP	32 28	32 30	19 23		
	ABGD	32 27	36 33	16 21	30 30	

1112 FIGURES LEGENDS:

1113 **Figure 1:** ML phylogeny of tribe Rhipileae obtained from the multilocus matrix (*tufA*, *rbcL*,
1114 and 18S rDNA), with bootstraps and posterior probabilities indicated at the nodes (bs/PP).

1115 Species of the same genus (as recognized by Guiry and Guiry (searched on the 10th of
1116 December 2019)) are noted using the same color. The type species of *Rhipilia* is indicated in
1117 bold. Outgroup species: *Caulerpa taxifolia*, *Caulerpa cupressoides* and *Caulerpa verticillata*.

1118 **Figure 2:** ML phylogeny of "Rhipiliopsideae group 1" and "Rhipiliopsideae group 2"
1119 obtained from the multilocus matrix (*tufA*, *rbcL*, and 18S rDNA), with bootstraps and
1120 posterior probabilities indicated at the nodes (bs/PP). Species of the same genus (as
1121 recognized by Guiry and Guiry (searched on the 10th of December 2019) are noted using the
1122 same color: Outgroup species: *Caulerpa taxifolia*, *Caulerpa sertularioides* and *Caulerpa*
1123 *verticillata*.

1124 **Figure 3:** Bayesian phylogeny of *Rhipilia* obtained from the multilocus matrix (*tufA*, *rbcL*,
1125 and 18S rDNA), with bootstraps and posterior probabilities indicated at the nodes (bs/PP).
1126 Species delimitation results obtained using the five methods applied to *tufA* and *rbcL* markers
1127 are shown in the middle section, with species names and illustrations. Distribution of species
1128 (from molecular data + type localities) is shown on the map to the right (A= *Rhipilia*
1129 *penicilloides*; C= *R. sp1*; D= *R. diaphana*; F= *R. sp3*). Outgroup species: *Rhipiliella*
1130 *verticillata*, *Kraftalia gracilis* and *Kraftalia orientalis*. Image rights: Payri C.E.; * from Littler
1131 and Littler (2000).

1132 **Figure 4:** ML phylogeny of *Kraftalia* gen. nov. obtained from the multilocus matrix (*tufA*,
1133 *rbcL*, and 18S rDNA), with bootstraps and posterior probabilities indicated at the nodes
1134 (bs/PP). Species delimitation results obtained using the five methods applied to *tufA* and *rbcL*
1135 markers are shown in the middle section, with species names and illustrations. Distribution of
1136 the species (from molecular data + type localities) is shown on the map to the right (A =

1137 *Kraftalia crassa*; B = *K. sp1*; C = *K. sp2*; D = *K. orientalis*; E = *K. sp3*; H = *K. yaeyamensis*;
1138 I = *K. sp4*; J = *K. gracilis*; K = *K. sp5*). Outgroup species: *Caulerpa taxifolia*, *Caulerpa*
1139 *verticillata*, *Rhipilia penicilloides*, *R. coppejansii*, *R. sp1* and *R. sp3*. Image rights: Payri C.E.

1140 **Figure 5:** ML phylogeny of *Rhipiliopsis* obtained from the multilocus matrix (*tufA*, *rbcL*, and
1141 18S rDNA), with bootstraps and posterior probabilities indicated at the nodes (bs/PP). Species
1142 delimitation results obtained using the five methods applied to *tufA* and *rbcL* markers are
1143 shown to the right, with species names and illustrations. Distribution of the species (from
1144 molecular data + type localities) is shown on the map at the bottom (A = *R. sp5*; B = *R. sp7*).
1145 Outgroup taxa: *Rhipilia penicilloides*, *Kraftalia orientalis* and *Rhipiliella verticillata*. Images
1146 rights: Payri, C.E.; * from Algaebase; ** from Littler and Littler (2000).

1147 **Figure 6:** ML phylogeny of *Rhipiliopsis* gen. nov. and *Callipsyigma* obtained from the
1148 multilocus matrix (*tufA*, *rbcL*, and 18S rDNA) with bootstraps and posterior probabilities
1149 indicated at nodes (bs/PP). Species delimitation results obtained using the five methods
1150 applied to *tufA* and *rbcL* markers are shown in the middle section, with species names and
1151 illustrations. Distribution of the species (from molecular data + type localities) is shown on
1152 the map to the right (C = *Rhipiliopsis* sp6; D = *R. sp2*; E = *R. sp1*; F = *R. sp3*; G = *R. sp4*; I = *R.*
1153 *sp7*). Outgroup species: *Caulerpa taxifolia*, *Caulerpa verticillata*, *Caulerpa sertularioides*.
1154 Image rights: Payri, C.E.; *: from Cremen and al. (2019).

1155 **Figure 7:** *Kraftalia* gen. nov. **A-E:** Species external habit, **A:** *K. orientalis* (NOU 204095), **B:**
1156 *K. crassa* (NOU 203593), **C:** *K. gracilis* (NOU 203756), **D:** *K. yaeyamensis* (NOU 203801),
1157 **E:** *K. sp5* (NOU 203798); **F:** Siphons disposition in *K. orientalis* (NOU 204123); **G:**
1158 Dichotomies with bulge and constrictions in *K. crassa* (NOU 203483); **H:** Siphons
1159 disposition in *K. gracilis* (NOU 203320); **I:** Siphons disposition in *K. yaeyamensis* (NOU
1160 203816); **J:** Siphons disposition in *K. sp5* (NOU 203798); **K-O:** Adhesion structures between
1161 siphons: **K and L :** Tenacula in *K. orientalis* (NOU 204123) and *K. crassa* (NOU 203593),

1162 respectively; **M**: Differentiated bent siphon apices on one of the two dichotomous branches
 1163 (circles); arising from unconstricted dichotomies (arrows) in *K. gracilis* (NOU 203320), **N**:
 1164 Direct contact between siphons in *K. yaeyamensis* (NOU 203816), **O**: Unilateral papillae in
 1165 *K. sp5* (NOU 203798); **Scale bars**: A: 1.5 cm ; B: 6.5 mm; C: 1 mm; D: 900 μ m; E: 1.5 mm;
 1166 F: 115 μ m; G: 55 μ m; H: 180 μ m; I: 200 μ m ; J: 250 μ m; K: 40 μ m; L: 60 μ m; M: 40 μ m; N:
 1167 40 μ m; O: 33 μ m.

1168 **Figure 8: *Rhipiliospina stellifera* sp. nov. A-C :** Habit of the plant, **A**: NOU 203095, **B**: NOU
 1169 203758, **C**: NOU 203764; **D-G**: Stipe with spinous or star-shaped cortication, **D and F**: NOU
 1170 203095, **E and G**: NOU 203758; **H**: Spinous protuberances in siphons from the basal part of
 1171 the blade (NOU 203095); **I**: Net-like aspect of the blade (NOU 203095); **J**: Tortuous siphons
 1172 dichotomously divided and adhering to each other with papillae (NOU 203758); **H**:
 1173 Dichotomies with symmetrical constrictions and adhesion between siphons with bilateral
 1174 papillae forming H structures (NOU 203758). **Scale bars**: A: 1 mm; B: 1.5 mm; C: 125 mm;
 1175 D: 100 μ m; E: 115 μ m; F: 50 μ m; G: 40 μ m; H: 50 μ m; I: 300 μ m; J: 130 μ m; K: 25 μ m.

1176 **Figure 9: *Callipsygma brevis* sp. nov. A:** Habitat of the species (in Madagascar); **B**: External
 1177 habit of the species *in-situ*; **C-D**: External habit of the species *ex-situ*; **E**: Stipe siphons with
 1178 protuberances and deformed lateral branches; **F**: Dichotomies with symmetrical constrictions
 1179 and ring of cell-wall thickening; **G**: Cohesion between siphons with uni- or bilateral papillae;
 1180 **H**: Overview of siphons dichotomously divided and adhering by papillae. **Scale bars**: A:
 1181 4cm; B: 1.25 cm; C: 0.8 cm; D: 1.25 cm; E: 85 μ m; F: 45.5 μ m; G: 100 μ m; H: 100 μ m.

1182 SUPPLEMENTARY TABLES AND FIGURES LEGENDS

1183

1184 **Appendix S1:** Species delimitation analyses of the *tufA* datasets

1185 **Appendix S2:** Supports (ML) of hPTP partitions for the *tufA* datasets

1186 **Appendix S3:** Species delimitation analyses of the *rbcL* datasets

1187 **Appendix S4:** Supports (ML) of hPTP partitions for the *rbcL* datasets

1188 **Figure S1:** Phylogenetic relationships among suborder Halimedineae obtained from the
1189 concatenated multilocus matrix (*tufA*, *rbcL*, 18S rDNA), and position of the former
1190 Rhipiliaceae lineages (light green). Values at nodes indicate bootstraps and posterior
1191 probabilities (bs/PP) obtained from ML and BI reconstructions, respectively. Type species
1192 appear in red. Outgroup species: *Codium duthieae*, *Codium platylobium*, and *Bryopsis*
1193 *plumosa*.

1194 **Figure S2:** Species delimitation results for tribe Rhipileae obtained with the five methods
1195 (ABGD, GMYC, bGMYC, PTP and mPTP) on the *tufA* dataset. The tree represented is MCCT
1196 tree from the BEAST analysis. Partitions retained as SSHs following the majority rule are
1197 indicated by black bars. Blue bars represent the partition retained as SSHs, although not in the
1198 majority rule, while grey bars are the different partitions not retained. The defined SSHs
1199 (= clades) are indicated in the right column, together with species assignments obtained from
1200 morpho-anatomical observations.

1201 **Figure S3:** Species delimitation results for “Rhipiliopsidae group 1” obtained with the five
1202 methods (ABGD, GMYC, bGMYC, PTP and mPTP) on the *tufA* dataset. Partitions retained as
1203 SSHs following the majority rule are indicated by black bars, while grey bars are the different
1204 partitions not retained. The defined SSHs (= clades) are indicated in the right column, together
1205 with species assignments obtained from morpho-anatomical observations.

Figure S4: Species delimitation results for “Rhipiliopsidae group 2” obtained with the five methods (ABGD, GMYC, bGMYC, PTP and mPTP) on the *tufA* dataset. Partitions retained as SSHs following the majority rule are indicated by black bars, while grey bars are the different partitions not retained. The defined SSHs (= clades) are indicated in the right column, together with species assignments obtained from morpho-anatomical observations.

Figure S5: Species delimitation results for tribe Rhipileae obtained with the five methods (ABGD, GMYC, bGMYC, PTP and mPTP) on the *rbcL* dataset. Partitions retained as SSHs following the majority rule are indicated by black bars, while grey bars are the different partitions not retained. The defined SSHs (= clades) are indicated in the right column, together with species assignments obtained from morpho-anatomical observations.

Figure S6: Species delimitation results for Rhipiliopsidae lineages (group 1 & 2) obtained with the five methods (ABGD, GMYC, bGMYC, PTP and mPTP) on the *rbcL* dataset. The tree represented is MCCT tree from the BEAST analysis. Partitions retained as SSHs following the majority rule are indicated by black bars. Blue bars represent the partition retained as SSHs, although not in the majority rule, while grey bars are the different partitions not retained. The defined SSHs (= clades) are indicated in the right column, together with species assignments obtained from morpho-anatomical observations.

Figure S7 : ML phylogeny of other Rhipileae species, including *Rhipiliella verticillata*, obtained from the multilocus matrix (*tufA*, *rbcL* and 18S rDNA), with bootstraps and posterior probabilities indicated at the nodes (bs/PP). The species delimitation results obtained using the five methods applied to *tufA* and *rbcL* markers are shown in the middle section, with species names and illustrations. The distribution of the species (from molecular data + type localities) is shown on the map to the right (A = *Rhipiliella verticillata*; B = *Rhipiliopsis* cf. *mortensenii*). Outgroup species: *Caulerpa taxifolia*, *Caulerpa verticillata*, *Rhipilia*

1230 *penicilloides*, *R. coppejansii*, *R. sp1* and *R. sp3*. Image rights: * from Littler and Littler
1231 (2000); ** from Womersley, 1984.

1232 **Table S1:** List of specimens with sample ID, species identification, location of sampling,
1233 GenBank accession numbers (or BOLD sequence ID for those not submitted), and the
1234 sequences used in the species delimitation approach and the corresponding SSH number.

1235 **Table S2:** Primers used for the amplification of *tufA*, *rbcL*, and 18S rDNA markers

1236 **Table S3:** Details of phylogenetic analysis for both ML and BI reconstructions according to the
1237 various datasets.

1238 **Table S4:** Variability of the datasets.

1239 **Table S5:** *A posteriori* probabilities (PP) of the partitions defined by the bGMYC method on
1240 the *tufA* marker for Rhipileae and Rhipiliopsidae lineages.

1241 **Table S6:** *A posteriori* probabilities (PP) of the partitions defined by the bGMYC method on
1242 the *rbcL* marker for Rhipileae and Rhipiliopsidae lineages.

1243 **Table S7:** Details of the incongruence resolution process and species assignment of the SSHs.

1244

1245

1246