



HAL
open science

Novel cytokinetic ring components drive negative feedback in cortical contractility

Kathryn Rechain Bell, Michael Werner, Anusha Doshi, Daniel Cortes, Adam Sattler, Thanh Vuong-Brender, Michel Labouesse, Amy Shaub Maddox

► **To cite this version:**

Kathryn Rechain Bell, Michael Werner, Anusha Doshi, Daniel Cortes, Adam Sattler, et al.. Novel cytokinetic ring components drive negative feedback in cortical contractility. *Molecular Biology of the Cell*, 2020, 31 (15), pp.1623-1636. 10.1091/mbc.E20-05-0304) . hal-03244176

HAL Id: hal-03244176

<https://hal.sorbonne-universite.fr/hal-03244176v1>

Submitted on 1 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Novel cytokinetic ring components drive negative feedback in cortical contractility

Kathryn Rehai Bell^{a,b,†}, Michael E. Werner^{a,†}, Anusha Doshi^a, Daniel B. Cortes^a, Adam Sattler^a, Thanh Vuong-Brender^c, Michel Labouesse^c, and Amy Shaub Maddox^{a,b,*}

^aDepartment of Biology and ^bCurriculum in Genetics and Molecular Biology, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599; ^cInstitut de Biologie Paris-Seine, Sorbonne Université, INSERM, 75005 Paris, France

ABSTRACT Actomyosin cortical contractility drives many cell shape changes including cytokinetic furrowing. While positive regulation of contractility is well characterized, counterbalancing negative regulation and mechanical brakes are less well understood. The small GTPase RhoA is a central regulator, activating cortical actomyosin contractility during cytokinesis and other events. Here we report how two novel cytokinetic ring components, GCK-1 (germinal center kinase-1) and CCM-3 (cerebral cavernous malformations-3), participate in a negative feedback loop among RhoA and its cytoskeletal effectors to inhibit contractility. GCK-1 and CCM-3 are recruited by active RhoA and anillin to the cytokinetic ring, where they in turn limit RhoA activity and contractility. This is evidenced by increased RhoA activity, anillin and nonmuscle myosin II in the cytokinetic ring, and faster cytokinetic furrowing, following depletion of GCK-1 or CCM-3. GCK-1 or CCM-3 depletion also reduced RGA-3 levels in pulses and increased baseline RhoA activity and pulsed contractility during zygote polarization. Together, our results suggest that GCK-1 and CCM-3 regulate cortical actomyosin contractility via negative feedback. These findings have implications for the molecular and cellular mechanisms of cerebral cavernous malformation pathologies.

Monitoring Editor

William Bement
University of Wisconsin,
Madison

Received: May 14, 2020
Accepted: May 22, 2020

This article was published online ahead of print in MBoc in Press (<http://www.molbiolcell.org/cgi/doi/10.1091/mboc.E20-05-0304>) on June 3, 2020.

The authors declare no competing financial interests.

Author contributions: K.R.B. and M.E.W. contributed to conceptualization, data curation, formal analysis, investigation, supervision, visualization, and writing of the original draft. M.E.W. also contributed methodology and writing (editing). A.D. contributed to investigation and visualization. D.B.C. contributed software and writing (review). A.S. contributed investigation and data curation. T.V.-B. contributed a resource. M.L. contributed to funding acquisition and supervision. A.S.M. contributed to conceptualization, funding acquisition, supervision, visualization, and writing (editing).

[†]These authors contributed equally.

*Address correspondence to: Amy Shaub Maddox (asm@unc.edu).

Abbreviations used: *C. elegans*, *Caenorhabditis elegans*; CRISPR, clustered regularly interspaced short palindromic repeats; DIC, differential interference contrast; DNA, deoxyribonucleic acid; ERM, ezrin-radixin-moesin; F-actin, filamentous actin; GaAsP PMT, Gallium arsenide phosphide photo-multiplier tube; GAP, GTPase activating protein; GEF, guanine nucleotide exchange factor; GFP, green fluorescent protein; GTP, guanosine triphosphate; GTPase, guanosine triphosphate hydrolase; NA, numerical aperture; NMMII, nonmuscle myosin II; PAR, abnormal embryonic PARTitioning of cytoplasm; PCR, polymerase chain reaction; RFP, red fluorescent protein; RNA, ribonucleic acid; RNAi, RNA-mediated interference; sgRNA, single guide RNA; STRIPAK, striatin-interacting phosphatase and kinase.

© 2020 Bell et al. This article is distributed by The American Society for Cell Biology under license from the author(s). Two months after publication it is available to the public under an Attribution-Noncommercial-Share Alike 3.0 Unported Creative Commons License (<http://creativecommons.org/licenses/by-nc-sa/3.0/>).

“ASCB®,” “The American Society for Cell Biology®,” and “Molecular Biology of the Cell®” are registered trademarks of The American Society for Cell Biology.

INTRODUCTION

Following faithful replication and segregation of the genome, nascent daughter nuclei are partitioned into individual daughter cells during cytokinesis (Green et al., 2012; Pollard and O’Shaughnessy, 2019). The spatiotemporal coupling of this partitioning with chromosome segregation is achieved by signaling from the anaphase spindle to elicit a zone of active RhoA GTPase at the equatorial cortex (Basant and Glotzer, 2018). RhoA activity initiates a cascade of downstream effects ultimately resulting in the polymerization of filamentous actin (F-actin) and activation of the motor protein nonmuscle myosin II (NMM-II). F-actin and NMM-II form the structural basis of the cytokinetic ring along with many other components, including anillin (D’Avino et al., 2015). Anillin binds F-actin, NMM-II, RhoA, and other structural and regulatory ring components, thus acting as a scaffold (D’Avino, 2009; Piekny and Maddox, 2010). Once the ring is assembled, it constricts, drawing the plasma membrane into a furrow that partitions the cytoplasm (Cheffings et al., 2016). Complete cytokinetic ring closure is essential for cells to maintain proper ploidy. Failure results in the formation of a polyploid cell that can undergo apoptosis or cancerous transformation (Lacroix and Maddox, 2012).

RhoA activity that promotes cytokinesis as well as contractility in many other biological processes is regulated by the delicate balance of activating guanine nucleotide exchange factors (GEFs) and inhibiting GTPase activating proteins (GAPs) as well as positive

and negative feedback loops (Bement *et al.*, 2015; Goryachev *et al.*, 2016; Bischof *et al.*, 2017; Nishikawa *et al.*, 2017; Michaux *et al.*, 2018). The main RhoA GAP during cytokinesis, RGA-3/4, plays a central role in negative feedback regulation of pulsed contractions during polarization of the *Caenorhabditis elegans* (*C. elegans*) zygote, but the mechanism of negative feedback regulation during cytokinesis remains poorly understood (Schmutz *et al.*, 2007; Schonegg *et al.*, 2007; Mayer *et al.*, 2010; Zanin *et al.*, 2013; Naganathan *et al.*, 2014, 2018; Zhang and Glotzer, 2015; Michaux *et al.*, 2018).

To gain new insights into the mechanisms of cytokinetic ring constriction, we identified novel anillin-interacting proteins and found germinal center kinase-1 (GCK-1), a serine/threonine kinase related to budding yeast Ste20 (sterile-20). GCK-1 is the only *C. elegans* orthologue of the mammalian germinal center kinase III subfamily, which is implicated in apoptosis, proliferation, polarity, and cell motility (Schouest *et al.*, 2009; Zheng *et al.*, 2010; Yin *et al.*, 2012; Rehain-Bell *et al.*, 2017). One well-characterized interaction partner of GCK-1 is CCM-3 (cerebral cavernous malformations-3), which is thought to recruit GCK-1 to the striatin-interacting phosphatase and kinase (STRIPAK) complex (Hwang and Pallas, 2014). Together these proteins are thought to regulate endothelial integrity in part by negatively regulating RhoA (Borikova *et al.*, 2010; Zheng *et al.*, 2010; Richardson *et al.*, 2013). We and others characterized the roles of *C. elegans* GCK-1 and its cofactor CCM-3 (collectively, GCK-1/CCM-3) in maintaining the structural integrity of the oogenic syncytial germline (Pal *et al.*, 2017; Rehain-Bell *et al.*, 2017). GCK-1/CCM-3 promote the stability of the intercellular bridges that connect developing oocytes with a shared cytoplasm by limiting the abundance of proteins that promote contractility (including anillin and NMY-2 [NMY-2]) (Pal *et al.*, 2017; Rehain-Bell *et al.*, 2017). We proposed that GCK-1/CCM-3 suppress anillin (ANI-1) and NMY-2 localization by inhibiting RhoA, as their vertebrate homologues are known to do (Richardson *et al.*, 2013; Rehain-Bell *et al.*, 2017).

The identification of novel anillin-interacting proteins that limit contractility fit logically with their localization to stable intercellular bridges. However, we and others noted that GCK-1/CCM-3 also enrich on the dynamic, contractile cytokinetic ring in the *C. elegans* zygote (Pal *et al.*, 2017; Rehain-Bell *et al.*, 2017). Here, we report investigating the regulation of contractility in the *C. elegans* zygote by GCK-1/CCM-3. We found that on the contractile ring, GCK-1/CCM-3 limit the abundance of “contractility” proteins. Partial depletion of GCK-1/CCM-3 boosts contractility during cytokinesis and polarization of the zygote. GCK-1/CCM-3 localize to the cytokinetic ring and cortical pulses downstream of the master regulator RhoA (RHO-1 in *C. elegans*) and of anillin (ANI-1), and also influence the abundance of active RhoA, likely through promoting the cortical recruitment or retention of RGA-3. Therefore, we conclude that GCK-1/CCM-3 are novel components of negative feedback in the cytokinetic ring. These findings advance the growing body of work showing that contractile networks in cells not only are activated by positive regulation, but also contain structural “brakes” and regulatory time-delayed negative feedback important for turnover and dynamics (Bement *et al.*, 2015; Dorn *et al.*, 2016; Goryachev *et al.*, 2016; Bischof *et al.*, 2017; Nishikawa *et al.*, 2017; Khaliullin *et al.*, 2018; Michaux *et al.*, 2018).

RESULTS

GCK-1 and CCM-3 regulate each other's stability and cortical targeting

C. elegans GCK-1 and CCM-3 are known to interact, and their mammalian homologues form a heterodimer (Ceccarelli *et al.*, 2011; Xu *et al.*, 2013; Zhang *et al.*, 2013; Lant *et al.*, 2015). The domains that allow for heterodimerization of GCK III subfamily members and

CCM3 are conserved in *C. elegans* GCK-1 and CCM-3, indicating that these proteins heterodimerize as well (Ceccarelli *et al.*, 2011). We first tested the dynamics of GCK-1/CCM-3 localization and their interdependence during cytokinesis. We followed the localization of fluorescently tagged GCK-1 and CCM-3 expressed under the control of their own promoters (Pal *et al.*, 2017; Rehain-Bell *et al.*, 2017). Both are present in the cytoplasm and enrich in the cytokinetic ring during anaphase (Figure 1, A and B). We then depleted GCK-1 or CCM-3 by RNA-mediated interference (RNAi) to test the requirement of each for localization of the other to the cytokinetic ring. Levels of GCK-1 or CCM-3 in the cytokinetic ring were significantly reduced following depletion of CCM-3 or GCK-1, respectively (Figure 1, A, A', B, and B'). These results demonstrate that GCK-1 and CCM-3 are interdependent for their enrichment on the cytokinetic ring and support the idea that they act as a complex during cytokinesis.

We next tested whether GCK-1 and CCM-3 affect each other's localization to the cytokinetic furrow via active recruitment or effect on protein level. To do so, we assessed the abundance of cytoplasmic GCK-1 or CCM-3 following depletion of the other (Figure 1, C and C'). CCM-3 depletion did not significantly affect the cytoplasmic levels of GCK-1::GFP (Figure 1C). Thus, we concluded that CCM-3 is not required for GCK-1 protein stability. Depletion of GCK-1, however, significantly decreased cytoplasmic levels of CCM-3::mNeonGreen (Figure 1C'), suggesting that GCK-1 is required for CCM-3 stability. Consistently, we found that cytokinetic ring enrichment, the ratio of cytokinetic ring and cytoplasmic protein levels, of CCM-3::mNeonGreen increased slightly (~20%) following GCK-1 depletion compared with control, indicating approximately equal reduction of both the ring and cytoplasmic protein pools and little effect on ring targeting (Figure 1D'). Conversely, GCK-1::GFP enrichment on the cytokinetic ring relative to the cytoplasm was significantly decreased (~60% decrease) following CCM-3 depletion (Figure 1D). Together these results suggest that CCM-3 targets GCK-1 to the cytokinetic ring, while GCK-1 promotes CCM-3 ring localization at least partly by regulating the CCM-3 protein level.

GCK-1 and CCM-3 localize downstream of RhoA

RhoA is the master regulator necessary for recruitment and activation of cytokinetic ring components in animal cells (Piekny *et al.*, 2005; Jordan and Canman, 2012; Basant and Glotzer, 2018). To test whether GCK-1/CCM-3 localize downstream of RhoA (RHO-1 in *C. elegans*), we performed time-lapse imaging of GCK-1::GFP and CCM-3::mNeonGreen on the cell cortex during anaphase. Shortly following anaphase onset in control cells, both GCK-1 and CCM-3 localize to large cortical foci in the anterior of the embryo and the cell equator (Figure 2A and Supplemental Figure 1A). This localization pattern mirrors that of many known cytokinetic ring components such as active RHO-1, anillin (ANI-1), and NMY-2 (Maddox *et al.*, 2005, 2007; Motegi *et al.*, 2006; Schonegg *et al.*, 2007; Velarde *et al.*, 2007; Werner *et al.*, 2007). In fact, GCK-1 and CCM-3 colocalized with NMY-2 during anaphase (Figure 2A, Supplemental Movie 1, and Supplemental Figure 1A). When the levels of active RHO-1 were reduced by partially depleting its main activator during cytokinesis, the RhoGEF ECT-2, the levels of cortical GCK-1 or CCM-3 at the cell equator during anaphase were significantly reduced (Figure 2, B and C, Supplemental Movie 2, and Supplemental Figure 1, B and C). We conclude that GCK-1/CCM-3 depend on active RhoA for recruitment to the cytokinetic ring.

RhoA effector proteins, including Diaphanous-family formins, Rho-kinase, and anillin, bind RhoA-GTP directly and independently, whereas downstream factors such as F-actin, NMY-2, and septins depend on these effectors (Piekny *et al.*, 2005; Piekny and Maddox,




FIGURE 1: Mutual regulation of GCK-1 and CCM-3 enrichment on the cytokinetic ring. (A) Representative single-plane images of control and *ccm-3(RNAi)* embryos expressing GCK-1::GFP. (B) Representative single-plane images of control and *gck-1(RNAi)* embryos expressing CCM-3::mNeonGreen. (A', B') Quantitation of average fluorescence intensity per micron relative to the perimeter of the cytokinetic ring in control ($n = 7$ embryos) and depleted ($n = 9$ or 6 embryos [CCM3 or GCK-1 depleted, respectively]) conditions. (C, C') Cytoplasmic levels of GCK-1::GFP (C) and CCM-3::mNeonGreen (C') in control and depleted cells. (D, D') Enrichment of GCK-1::GFP (D) and CCM-3::mNeonGreen (D') on the cytokinetic ring relative to the cytoplasm in control and depleted cells. Scale bars are $10 \mu\text{m}$ in all images in all figures. Error bars are SE in all figures unless otherwise indicated.

2010; Heasman and Ridley, 2008). To determine whether GCK-1 localization is dependent on other cytokinetic ring components, we depleted ANI-1 or NMY-2 by RNAi and assessed GCK-1 and CCM-3 levels in the cytokinetic ring. Depletion of ANI-1, but not NMY-2, caused a significant reduction in GCK-1 and CCM-3 in the cell equator (Figure 2, B, D, and E, Supplemental Movies 3 and 4, and Supplemental Figure 1, B, D, and E). Taken together, these results demonstrate that GCK-1 and CCM-3 are recruited to the cytokinetic ring downstream of active RHO-1 and ANI-1 but not NMY-2.

GCK-1 and CCM-3 limit contractility in cytokinesis

We next tested whether, as in the stable intercellular bridges of the *C. elegans* germline, GCK-1 or CCM-3 limits contractility in the

highly contractile zygote. We began by assessing embryonic viability and found that depletion of GCK-1 or CCM-3 leads to 11% and 13% embryonic lethality, respectively (Figure 3A). To explore whether this embryonic lethality could be due to defects in cytokinesis in early (<100 cells) embryos, we strongly depleted GCK-1 or CCM-3 and quantified the prevalence of multinucleation. Similar percentages of GCK-1 (15%) and CCM-3 (20%) depleted embryos contained at least one multinucleated blastomere (Figure 3B). However, cytokinesis failure was rare; only about 0.5% of blastomeres were multinucleate (Figure 3C). As reported previously, strong depletion of GCK-1 or CCM-3 caused severe defects in germline organization, leading to smaller and fewer embryos (Schouest *et al.*, 2009; Green *et al.*, 2011; Pal *et al.*, 2017;




FIGURE 2: GCK-1 localization is dependent on active RHO-1 and ANI-1 but not NMY-2. (A) Representative images of GCK-1::GFP and NMY-2::mKate localization during anaphase. (B) Representative images of embryos expressing both GCK-1::GFP and NMY-2::mKate in control, *ect-2(RNAi)*, *ani-1(RNAi)*, and *nmy-2(RNAi)* embryos 60 s after anaphase onset. (C–E) Background-adjusted equatorial GCK-1::GFP levels normalized to that at anaphase onset in control, *ect-2(RNAi)* (C), *ani-1(RNAi)* (D), and *nmy-2(RNAi)* (E) embryos ($n = 6$ for all conditions).

Rehain-Bell *et al.*, 2017; Priti *et al.*, 2018). As such, embryos assessed for cytokinesis defects were significantly smaller (Supplemental Figure 2A) and a correlation between embryo size and frequency of multinucleation suggested that the latter could at least partially be a secondary effect of abnormal embryo size (Supplemental Figure 2B).

To study the role of GCK-1/CCM3 in cytokinesis while avoiding the confounding factor of abnormal embryo size, we performed

feeding RNAi for 24 h to partially (>83%) deplete embryos of CCM-3 or GCK-1, which maintained normal embryo size (Supplemental Figure 2C). Under these conditions, cytokinetic furrows ingressed to completion in all zygotes observed, and furrowing completed significantly faster than in controls, largely due to increased maximal ingression speed of the cytokinetic ring (Figure 3, D–G). This suggests that GCK1/CCM-3 attenuate contractility during cytokinetic ring constriction.

To determine how GCK-1/CCM-3 affect cytokinetic ring ingression kinetics, we examined how they affect recruitment of cytokinetic ring components to the cell equator. The levels of both NMY-2 and ANI-1 at the cell equator were significantly increased following GCK-1 or CCM-3 depletion (Figure 3, H and I). This result suggests that GCK-1 and CCM-3 attenuate ingression speed by limiting the amount of proteins that drive contractility in the cytokinetic ring.

Taken together, these results demonstrate that GCK-1/CCM-3 limit the amount of ANI-1 and NMY-2 at the cell equator and negatively regulate the kinetics of cytokinetic ring ingression.

GCK-1 and CCM-3 negatively regulate cortical contractility during polarization

Before the first mitotic division of the *C. elegans* zygote, a highly contractile actomyosin network spans the embryonic cortex and becomes progressively polarized to the embryo anterior. To test whether GCK-1/CCM-3 limit zygote contractility not only in cytokinesis but also more generally, we next examined the role of GCK-1/CCM-3 during polarization, when a transient feature called the pseudocleavage furrow ingresses at the boundary of the highly contractile anterior and the more compliant posterior cortex, and regresses. The positioning and regulation of the pseudocleavage furrow are distinct and independent from those of the cytokinetic furrow that ingresses about 10 min later (Cowan and Hyman, 2007; Werner and Glotzer, 2008) (Figure 4A). Consistent with a potential role before cytokinesis, we observed that GCK-1/CCM-3 enrich in cortical patches during polarization, co-localizing with other contractile components such as NMY-2 and ANI-1 (Figure 4B, Supplemental Movie 5, and Supplemental Figure 3, A and B). Similar to our observations during anaphase, this localization is dependent on ECT-2 and ANI-1 but not NMY-2 (Supplemental Figure 3, A–D).

To test whether GCK-1/CCM-3 affect contractility during polarization, we depleted either CCM-3 or GCK-1 by RNAi and measured the regression timing and depth of the pseudocleavage furrow, both of which reflect contractility during polarization (Schmutz *et al.*, 2007; Schonegg *et al.*, 2007; Tse *et al.*, 2012; Reymann *et al.*, 2016). The persistence of the pseudocleavage furrow was significantly increased in zygotes depleted of either CCM-3 or GCK-1 (Figure 4, C and D, and Supplemental Movie 5). To control for possible defects in cell cycle progression, we also measured the time from pronuclear meeting to anaphase onset and found no statistically significant difference among control and embryos depleted of GCK-1 or CCM-3, indicating that cell cycle timing is not affected (Supplemental Figure 4A). Similarly, the maximal depth of pseudocleavage furrow as a percentage of embryo width was significantly increased following GCK-1 or CCM-3 depletion, when compared with controls (Figure 4, C and E). An additional measure of cortical contractility during polarization is the size of the anterior polarity domain (Schonegg *et al.*, 2007; Tse *et al.*, 2012). The size of the anterior domain, defined by localization of PAR-6, was significantly reduced in embryos depleted of GCK-1 or CCM-3 when compared with controls (Figure 4F). This was not due to defects in polarity establishment or maintenance following depletion of CCM-3 or GCK-1, since asymmetric spindle positioning or the extent of asymmetric enrichment of anterior PAR proteins was not affected (Supplemental Figure 4, B–B’). The dependence of the anterior enrichment of GCK-1 and CCM-3 on PARs (Supplemental Figure 4, C–G’) further demonstrated that GCK-1 and CCM-3 are downstream of PAR-driven polarity. Taken together, these findings support the conclusion that GCK-1 and CCM-3 limit cortical contractility during

polarization, in addition to during cytokinesis and on germline intercellular bridges.

GCK-1/CCM-3 limit RhoA activity in the cytokinetic ring and during pulsed contractility

GCK-1/CCM-3 homologues in cultured human endothelial cells and mouse neocortex negatively regulate RhoA (Borikova *et al.*, 2010; Zheng *et al.*, 2010; Louvi *et al.*, 2014), but we showed that they are recruited to the cytokinetic ring downstream of RHO-1 activity and ANI-1. We reasoned that this series of recruitment interdependencies and protein activities could constitute a negative feedback loop in which RHO-1 recruits ANI-1, which then recruits GCK-1/CCM-3, which in turn limits RHO-1 activity and contractility. To test this hypothesis, we first examined the pulsed cortical contractions that occur during embryo polarization. These cycles of quiescence, activation, contraction, and finally relaxation and disassembly are a powerful model to study feedback loops in the regulation of contractility (Reymann *et al.*, 2016; Nishikawa *et al.*, 2017; Michaux *et al.*, 2018; Naganathan *et al.*, 2018). RHO-1 activity in *C. elegans* has been assessed using a biosensor composed of the C-terminal third of ANI-1, which contains, among several functional elements, its RHO-1 binding domain (Sun *et al.*, 2015). Because GCK-1/CCM-3 and ANI-1 reciprocally affect the other’s localization and/or stability (Figures 2, B and D, and 3H), we avoided the use of the ANI-1-based biosensor and generated a fluorescently tagged Rho-kinase (LET-502) expressed from its endogenous locus to employ as a biosensor for RHO-1 activity. Rho-kinase is a well-characterized RhoA effector that is preferentially associated with the activated form of RhoA (Matsui *et al.*, 1996). To characterize this probe, we increased or decreased RHO-1 activity by depleting its inactivator RhoGAP RGA-3/4 or activator RhoGEF ECT-2, respectively. As expected, recruitment of LET-502 to the cell equator increased following RGA-3/4 depletion and decreased following ECT-2 depletion (Figure 5, A–C, and Supplemental Movies 6–8).

We next tested whether GCK-1/CCM-3 localization dynamics are consistent with a role in negative feedback during pulsed contractility. Consistent with previous observations, local enrichment of active RHO-1, as visualized via our LET-502 biosensor, precedes the localization of NMY-2, ANI-1, and RGA-3, all of which localize concurrently (Michaux *et al.*, 2018) (Figure 5, D, F, and H, and Supplemental Movie 9). We imaged cells coexpressing green fluorescent protein tagged (GFP-tagged) GCK-1 and red fluorescent protein tagged (RFP-tagged) NMY-2 with high temporal resolution and tracked cortical patches during polarization to determine the kinetics of GCK-1 localization relative to the accumulation of NMY-2. The maximal level of GCK-1 in cortical patches occurred approximately 3.5 s after that of NMY-2 and thus about 13 s after active RHO-1 recruitment, consistent with a role in time-delayed negative feedback regulation of RHO-1 and contractility (Figure 5, E, G, I, and J, and Supplemental Movie 10).

We predicted that if GCK1/CCM3 feed back negatively on RHO-1, depletion of GCK-1/CCM-3 would increase levels of active RHO-1. This prediction was substantiated by the positive control that depletion of RGA-3/4 increased the baseline RHO-1 activity level during pulsed contractility (Figures 5L and 6D). We tested the effects of depleting GCK-1/CCM-3 on the abundance of active RHO-1 in the cell equator and in cortical pulses during polarization. The abundance of LET-502 was significantly increased following GCK-1 or CCM-3 depletion, when compared with controls, in both the cell equator and cortical pulses (Figure 5, K and L). Together, our results support the conclusion that GCK-1/CCM3 are novel cytokinetic ring components recruited downstream of RHO-1 (via ANI-1) that in turn feed back negatively on RhoA activity (Figure 6G).




FIGURE 3: Depletion of GCK-1 or CCM-3 causes increased cytokinetic ingress speed and increases the amount of NMY-2 and ANI-1 in the furrow. (A) Quantification of embryonic lethality in control embryos or after depletion of *gck-1*(RNAi) and *ccm-3*(RNAi). (B) Percent of embryos with at least one multinucleated cell in control embryos or after thorough depletion of *gck-1*(RNAi), *ccm-3*(RNAi), or *rga-3*(RNAi) ($n > 80$ embryos for all conditions). (C) Quantification of the amount

GCK-1/CCM-3 regulate RhoA activity by promoting RGA-3/4 recruitment

What is the molecular mechanism of RhoA inhibition downstream of GCK-1/CCM-3? In zebrafish and human cultured cells, homologues of GCK-1 phosphorylate moesin, an ezrin-radixin-moesin (ERM) family protein, leading to a reduction of RhoA activity (Zheng *et al.*, 2010). To determine whether GCK-1 regulates RHO-1 activity by a similar mechanism in the *C. elegans* zygote, we tested whether depletion of the only *C. elegans* ERM protein, ERM-1, affects cytokinetic furrow ingression kinetics or contractility during polarization (Gobel *et al.*, 2004; Van Furden *et al.*, 2004). Neither cytokinetic furrow kinetics nor pseudocleavage regression timing was significantly different between control and ERM-1-depleted cells, and the pseudocleavage furrow was only slightly deeper following ERM-1 depletion, suggesting that ERM-1 is not the main target of GCK-1/CCM-3 during polarization or cytokinesis (Figure 6, A–C).

Another potential mediator of RHO-1 inhibition by GCK-1/CCM-3 is the RhoGAP RGA-3/4, a known negative regulator of RhoA activity during both polarization and cytokinesis that mediates negative feedback regulation during pulsed contractions (Schmutz *et al.*, 2007; Schonegg *et al.*, 2007; Zanin *et al.*, 2013; Michaux *et al.*, 2018). Severe reduction or elimination of RGA-3/4 levels following thorough RNAi or in homozygous *rga-3Δ*, *rga-4Δ* double mutants leads to ectopic contractility, cytokinetic furrow positioning defects, and undetectable active RHO-1 pulses, confounding quantitative comparisons with the contractile phenotypes following GCK-1/CCM-3 depletion (Schmutz *et al.*, 2007; Schonegg *et al.*, 2007; Zanin *et al.*, 2013; Michaux *et al.*, 2018). Interestingly, in heterozygous *rga-3/4 (+Δ, +Δ)* double mutants cortical contractility still increases but pulses of active RHO-1 persist (Michaux *et al.*, 2018). We used partial RGA-3/4 depletion to quantify RHO-1 pulses when RHO-1 activity is increased. We observed an increase in baseline active RhoA levels in cortical pulses without a significant effect on the length of the pulse (Figure 6D). Hence, weak RGA-3/4 depletion enhanced contractility during both cytokinesis and polarization in ways that closely resembled the defects observed following GCK-1/CCM-3 depletion (Figures 3E, 4, D and E, and 6, A–C).

We next tested whether GCK-1/CCM-3 depletion affected cortical abundance of RGA-3/4. Indeed, depletion of GCK-1 or CCM-3 significantly reduced GFP::RGA-3 levels in cortical pulses (Figure 6, E and F). These results support the idea that GCK-1/CCM-3 regulate RhoA activity by promoting the cortical recruitment of the RhoGAP RGA-3. Together, our results define a negative feedback loop among RHO-1, ANI-1, GCK-1/CCM-3, and RGA-3/4 (Figure 6G).

DISCUSSION

We characterized two novel cytokinetic ring proteins and their roles in regulating actomyosin contractility during cytokinesis. We found that GCK-1/CCM-3 are recruited downstream of active RHO-1 and the scaffold protein ANI-1, suggesting that they are indirect RHO-1

effectors. However, GCK-1/CCM-3 also limit RHO-1 activity and actomyosin contractility during pulsed contractility, polarization, and cytokinesis, in agreement with what we and others showed for the stable actomyosin cortex of the *C. elegans* syncytial germline (Pal *et al.*, 2017; Rehai-Bell *et al.*, 2017; Priti *et al.*, 2018). We report a potential mechanism by which GCK-1/CCM-3 inhibit Rho: they help recruit or retain the RhoGAP RGA-3/4 at the cortex. Therefore, we propose that GCK-1/CCM-3 contribute to a negative feedback loop acting on the RHO-1-regulated actomyosin cytoskeleton.

Our conclusion that GCK-1/CCM-3 contribute to negative feedback on RHO-1 activity is further supported by the delayed timing of GCK-1/CCM-3 localization to pulsed contractile cortical patches, with respect to active RHO-1 and NMY-2 and, by extension ANI-1, recruitment. Contractile pulses are driven by oscillations of RHO-1 activity, which drives F-actin assembly, which in turn promotes accumulation of the RhoGAP RGA-3/4 (Nishikawa *et al.*, 2017; Michaux *et al.*, 2018), which feeds back to inactivate RHO-1. To test how GCK-1/CCM-3 contribute to pulse dynamics, we first measured how pulses respond to a reduction of RHO-1 activity, since reduction and elimination of RGA-3/4 have qualitatively distinct effects (Michaux *et al.*, 2018). To approximate the reduction in RGA-3/4 that we observed following GCK-1/CCM-3 depletion, we examined pulses following depletion of RGA-3/4. In such cells, pulse dynamics were indistinguishable from those of control cells, but baseline Rho activity level was higher following RGA-3/4 depletion (Figure 6D). Similarly, depletion of GCK-1/CCM-3 increased the amount of baseline RHO-1 activity in cortical patches, at least partly by limiting the initial size of cortical patches, but did not change pulse dynamics (Figure 5L and Supplemental Figure 5). Higher baseline RhoA activity is expected to push an excitable system toward stable RhoA activation (Goryachev *et al.*, 2016). We therefore predict that thorough loss of GCK-1/CCM-3, especially in combination with other RHO-1 activity gain-of-function perturbations, could cause constitutive cortical contractility and reduce the responsiveness of the cortex to spatial cues from the spindle.

Together, these observations suggest that GCK-1 and CCM-3 contribute to negative regulation of the cortical actomyosin cytoskeleton. This conclusion supports a growing consensus about negative feedback in pulsed contractility (Bement *et al.*, 2015; Dorn *et al.*, 2016; Goryachev *et al.*, 2016; Bischof *et al.*, 2017; Nishikawa *et al.*, 2017; Khaliullin *et al.*, 2018; Michaux *et al.*, 2018). Interestingly though, it has not to our knowledge been demonstrated that negative feedback on RhoA contributes to normal cytokinetic dynamics. Our observations that negative feedback built into the cytokinetic ring limits its speed may reflect an advantage of the above-mentioned responsiveness of the system.

RHO-1 activity is required for equatorial enrichment of GCK-1/CCM-3. Several observations suggest that this is mediated by ANI-1: depletion of ANI-1 prevents GCK-1/CCM-3 equatorial enrichment (Figure 2, B and D), ANI-1/anillin recruitment is downstream

of multinucleated cells as a percentage of total cells in embryos quantified in B. (D) Representative montage of HIS::GFP and mNeonGreen::PH in the furrow region over time in control, *gck-1(RNAi)*, and *ccm-3(RNAi)* embryos. Colored bars represent the timing of distinct phases of cytokinesis shown in the schematic below. Initiation: from anaphase onset to the onset of furrow ingression. Involution: from onset of ingression to the appearance of a doubled membrane. Furrowing: from the appearance of a doubled membrane to complete ring closure. Length scale bar = 10 μ m. (E) Quantification of the duration of interval between furrowing onset and ring closure. (F, G) Cytokinetic ring closure dynamics in control ($n = 9$), *gck-1(RNAi)* ($n = 7$), and *ccm-3(RNAi)* ($n = 6$) embryos measured by following closure dynamics (F) or subdividing cytokinesis timing into three distinct stages (G) using a strain expressing mNeonGreen::ANI-1 and NMY-2::mKate2. (H) Quantification of background-adjusted equatorial mNeonGreen::ANI-1 levels relative to equatorial levels 20 s prior to onset of furrow ingression in control ($n = 6$), *gck-1(RNAi)* ($n = 9$), or *ccm-3(RNAi)* ($n = 9$) embryos. (I) Quantification of background-adjusted equatorial NMY-2::GFP levels relative to equatorial levels 20 s prior to onset of furrow ingression in control ($n = 9$), *gck-1(RNAi)* ($n = 10$), and *ccm-3(RNAi)* ($n = 12$) embryos. Error bars are SD in A, E, and G.




FIGURE 4: GCK-1 localizes to cortical patches during polarization and inhibits cortical contractility. (A) Schematic representation of cellular morphology and distribution of relevant protein complexes in the *C. elegans* zygote. (B) Representative images of single-plane cortical imaging of embryos expressing GCK-1::GFP and NMY-2::mKate2, or

of active RHO-1/RhoA (Straight *et al.*, 2003; Hickson and O'Farrell, 2008; Piekny and Glotzer, 2008; Piekny and Maddox, 2010), and GCK-1 coimmunoprecipitates with ANI-1 (Rehain-Bell *et al.*, 2017). Anillins are large multidomain scaffold proteins with many binding partners and putative roles in cytokinesis (Piekny and Maddox, 2010). It is possible that the increase in furrowing speed observed following partial ANI-1 depletion (Descovich *et al.*, 2018) is due in part to decreased GCK-1/CCM-3 recruitment.

Thorough depletion of GCK-1/CCM-3 leads to severe germline defects and precludes the formation of normal embryos (Schouest *et al.*, 2009; Green *et al.*, 2011; Pal *et al.*, 2017; Rehain-Bell *et al.*, 2017; Priti *et al.*, 2018). The partial nature of our GCK-1/CCM-3 depletions likely explains the partial, but significant, effect on RGA-3/4 levels we observed (Figure 6, E and F). GCK-1/CCM-3 have been anecdotally implicated in zygote cytokinesis and polarity (Pal *et al.*, 2017). We quantitatively examined these events following various degrees of GCK-1/CCM-3 loss of function. Strong GCK-1/CCM-3 depletion increased embryonic lethality, with a very low incidence of cytokinesis failure (less than 0.5% of cells; Figure 3C). However, we could not separate this cytokinesis failure from defects in embryo size, since we observe an increase in the incidence of cytokinesis failure in embryos whose size is reduced after GCK-1/CCM-3 depletion, suggesting that cytokinesis failure could be secondary to abnormal embryo size. Cytokinetic furrow regression was never observed in zygotes, even in embryos that were significantly smaller than controls. We also never observed polarity defects following 80–90% GCK-1/CCM-3 depletion. Instead, our findings suggest that GCK-1/CCM-3 are downstream of PAR protein-based polarity and that the primary phenotype resulting from partial loss of GCK-1/CCM-3 during cytokinesis is an increase in equatorial contractility that translates into faster cytokinetic ring constriction. While in the cells we studied, faster cytokinetic furrowing does not translate to cytokinesis failure, this type of dysregulation could be detrimental in other contexts, such as in cells with long chromosomes with point centromeres or other specialized cell types. More generally, CCM proteins including CCM-3 have been implicated in contributing to endothelial integrity by promoting the formation and maintenance of tight and adherens junctions (Zheng *et al.*, 2010; Fischer *et al.*, 2013). Future work will be aimed at investigating the intriguing possibility that CCM-3 contributes to endothelial integrity by coordinating junctional dynamics and contractility in the cytokinetic furrow.

MATERIALS AND METHODS

C. *elegans* strains and culture

Worm strains (Table 1) were maintained using standard procedures at 20°C (Brenner, 1974). Clustered regularly interspaced short palindromic repeats (CRISPR) knock-in of GFP at the N-terminus of LET-502 was carried out using an asymmetric repair template consisting of 0.5 kb 3' to the start codon of *let-502* and 1.5 kb of the *let-502* coding region. The latter portion of this repair template was obtained from an existing plasmid containing *mlc-4p::GFP::LET-502* (pML1595) via removing the C-terminus of *let-502* using Fast Cloning (Li *et al.*, 2011), leaving only 1.5 kb downstream of the start

codon. Then 0.5 kb upstream of *let-502* start codon was amplified from genomic DNA (primers: CCTAATCGTTGTCTTTTGATCGGCA and GGCTGCAGCTCGATTTTCGT) to replace the *mlc-4p* using overlap extension polymerase chain reaction (PCR) (Bryksin and Matsumura, 2010). To target Cas9 to the *let-502* locus, the sequence of the single guide RNA (sgRNA) was CGCAGCTCATCCTGCTCCA, which was cloned into pDD162 (Dickinson *et al.*, 2013) and the plasmid (containing Cas9) was injected at a concentration of 50 ng/μl. The asymmetric repair template was injected at a concentration of 20 ng/μl, and genome editing was detected via single worm PCR (primers: GCTTGCCTGTCTTATTCATGC [genomic DNA] and TCCGTATGTTGCATCACCTTACC [GFP]).

MDX27 was generated by crossing LP306 (Heppert *et al.*, 2016) to JCC719 (gift from Julie Canman) (Columbia University Medical Center, Department of Pathology and Cell Biology) to introduce mCherry::his58. MDX33 was generated by crossing LP148 (Dickinson *et al.*, 2013) and LP274 (Heppert *et al.*, 2016), and MDX68 was generated by crossing MDX38 (Rehain-Bell *et al.*, 2017) and LP229 (Dickinson *et al.*, 2017). MDX69 was generated by crossing MDX38 and LP244 (Dickinson *et al.*, 2017). MDX74 was generated by crossing WD478 and LP244. MDX77 was generated by crossing WD478 and LP229. OD437 was kindly provided by Karen Oegema (Department of Cellular and Molecular Medicine, Ludwig Institute for Cancer Research, University of California).

RNA-mediated interference

Depletions were conducted by feeding worms bacteria, from the Ahringer collection, expressing double-strand RNA as described previously for 20–24 h in all conditions except for Figure 3, A–C, and Supplemental Figure 2, A and B, for which feeding was carried out for 48 h (Kamath and Ahringer, 2003; Kamath *et al.*, 2003). For weak *rga-3(RNAi)* experiments, the *rga-3(RNAi)* feeding culture was diluted fivefold using L4440 and worms were incubated on the resulting feeding plates for 24 h.

Quantification of embryonic lethality and frequency of cytokinesis failure

Embryonic lethality was determined by incubating L4 worms on RNAi feeding plates for 24 h and transferring them to a fresh RNAi plate before incubation for another 24 h. Parental worms were picked off the plate, and embryonic lethality was determined by quantifying the ratio of unhatched embryos to total number of worms on the plate. For determining cytokinesis failure, embryos from MDX27 worms were extruded onto an agarose pad and visually inspected for the presence of multinucleated cells based on DNA morphology. Embryo length was determined in ImageJ, measuring the distance from anterior to posterior pole.

C. *elegans* embryo sample prep and live imaging conditions

Embryos were dissected from gravid hermaphrodites and mounted on 2% agar pads except for Figure 3, D–I, for which embryos were mounted as inverted hanging drops. Images for Figures 1 and 4, C–F, and Supplemental Figure 4 were acquired with a CoolSNAP Hq camera (Photometrics) mounted on a DeltaVision Image Restoration

CCM-3::mNeonGreen and NMY-2::mKate2. (C) Representative fluorescent images of control, *gck-1(RNAi)*, and *ccm-3(RNAi)* embryos expressing HIS::GFP and mNeonGreen::PH (top) and corresponding differential interference contrast (DIC) images (bottom). Left: maximal pseudocleavage furrow ingression depth; right: at pronuclear meeting.

(D) Quantification of the time interval between pronuclear meeting and complete regression of the pseudocleavage furrow in control, *gck-1(RNAi)*, and *ccm-3(RNAi)* embryos. (E) Quantification of maximal pseudocleavage ingression as a percentage of cell width in control, *gck-1(RNAi)*, and *ccm-3(RNAi)* embryos. (F) Quantification of the size of the anterior PAR domain at metaphase in control, *gck-1(RNAi)*, and *ccm-3(RNAi)* embryos expressing PAR-6::mKate2. Error bars are SD.




FIGURE 5: GCK-1 localizes to cortical foci after NMY-2 and limits RHO-1 activity on the cortex. (A) Representative images of endogenous GFP::LET-502 at onset of ingress in control, *ect-2(RNAi)*, or *rga-3/4(RNAi)* embryos. (B) Background-adjusted total equatorial GFP::LET-502 fluorescence levels in control embryos ($n = 7$) and embryos with increased (*rga-3/4(RNAi)*) [$n = 6$] or decreased (*ect-2(RNAi)*) [$n = 6$] active RHO-1 levels. (C) Relative increase in background-adjusted total equatorial GFP::LET-502 fluorescence levels in control embryos and embryos with increased (*rga-3/4(RNAi)*) or decreased (*ect-2(RNAi)*) active RHO-1 levels. (D) Kymograph showing pulsed GFP::LET-502 and NMY-2::mRFP localization during embryo polarization. (E) Kymograph showing pulsed GCK-1::GFP and NMY-2::mKate2 localization during embryo polarization. (F) Magnified view of GFP::LET-502 and NMY-2::mRFP localization to a single focus. (G) Magnified view of GCK-1::GFP and NMY-2::mKate2 localization to a single focus. (H) Mean accumulation profile of GFP::LET-502 and NMY-2::mRFP in cortical patches. (I) Mean accumulation profile of GCK-1::GFP and NMY-2::mKate2 in cortical patches. (J) Time delay in maximal GCK-1::GFP accumulation relative to NMY-2::mKate2 localization. Data points = individual foci; $n = 236$ total foci from a total of five embryos for H and $n = 423$ total foci from a total eight embryos for I and J. (K) Quantification of background-adjusted equatorial GFP::LET-502 levels relative to equatorial levels 30 s prior to onset of furrow ingress in control ($n = 7$), *gck-1(RNAi)* ($n = 10$), and *ccm-3(RNAi)* ($n = 10$) embryos. (L) Mean fluorescence intensity profile of GFP::LET-502 in cortical patches from control and *gck-1(RNAi)* embryos. Error bars are SE except for panel J, where error bars are SD.




FIGURE 6: GCK1/CCM-3 contribute to negative feedback regulation of RHO-1 by promoting RGA-3 localization. (A) Quantification of the duration of interval between furrowing onset and ring closure in control, *erm-1*(RNAi), or weak *rga-3*(RNAi) embryos. (B) Time interval between pronuclear meeting and complete regression of the pseudocleavage furrow in control, *erm-1*(RNAi), or weak *rga-3*(RNAi) embryos. (C) Quantification of maximal pseudocleavage ingress as a percentage of cell width in control, *erm-1*(RNAi), or weak *rga-3*(RNAi) embryos. (D) Mean fluorescence intensity profile of GFP::LET-502 in cortical patches from control and weak *rga-3*(RNAi) embryos ($n = 9$ embryos each for a total of 277 and 304 patches, respectively). (E) Mean fluorescence intensity profile of RGA-3::GFP in cortical patches from control ($n = 9$ embryo; 241 patches total) and *gck-1*(RNAi) embryos ($n = 8$ embryos; 155 patches total). (F) Mean fluorescence intensity profile of RGA-3::GFP in cortical patches from control and *ccm-3* (RNAi) embryos ($n = 8$ embryos; 140 total patches). (G) Summary diagram of relationship of GCK1/CCM-3 to canonical cytokinetic contractility regulation network. Error bars are SD in A–C; all others are SE.

System (GE) with a 60×/1.42 NA Plan Apochromat objective. For Figure 1 and Supplemental Figure 4, B–D, we acquired z-stacks with 1 μm spacing through the entire embryo every 20 s and every 15 s for Figure 4, C–E. Data for Supplemental Figure 4, A and E–G', were

gathered by taking single images at the midplane of the embryo every 60 s.

All other data were acquired on a Nikon A1R microscope body with a 60 × 1.27 numerical aperture (NA) Nikon Water Immersion

Strain	Genotype	Publication
EM328	<i>let-502(mc74[GFP::let-502])</i> ; (<i>zuls151 [nmy-2::NMY-2-mRFP; unc-119(+)]</i>)	This study
LP162	<i>cp13[nmy-2::gfp + LoxP]</i> I.	Dickinson et al., 2013
MDX27	<i>cpls53[Pmex-5::GFP-C1::PLCδ-PH::tbb-2 3'UTR + unc-119 (+)]</i> II; <i>ltIs37 [pAA64; pie-1/mCherry::his-58; unc-119 (+)]</i> IV	This study
MDX33	<i>cpls45[Pmex-5::mNeonGreen::PLCδ-PH::tbb-2 3'UTR + unc-119(+)]</i> II; <i>his-72 (cp10 [his-72::gfp+LoxP unc-119(+)] LoxP)</i> III	This study
MDX38	<i>ccm-3(mon9[ccm-3::mNeonGreen^Flag])</i> II	Rehain-Bell et al., 2017
MDX40	<i>nmy-2(cp52[nmy-2::mkate2 + LoxP unc-119(+)] LoxP)</i> I; <i>ani-1(mon7[mNeonGreen^3xFlag::ani-1])</i> III	Rehain-Bell et al., 2017
MDX68	<i>nmy-2(cp52[nmy-2::mkate2 + LoxP unc-119(+)] LoxP)</i> I; <i>ccm-3(mon9[ccm-3::mNeonGreen^Flag])</i> II;	This study
MDX69	<i>par-6(cp60[par-6::mKate2::3xMyc + LoxP unc119(+)] LoxP)</i> I; <i>ccm-3(mon9[ccm-3::mNeonGreen^Flag])</i> II	This study
MDX74	<i>par-6(cp60[par-6::mKate2::3xMyc + LoxP unc119(+)] LoxP)</i> I; <i>gck-1(onIs15[GFP::3xFlag::gck-1])</i> III	This study
MDX77	<i>nmy-2(cp52[nmy-2::mkate2 + LoxP unc-119(+)] LoxP)</i> I; <i>gck-1(onIs15[GFP::3xFlag::gck-1])</i> III	This study
OD437	<i>rga-3(ok1889)</i> V; <i>unc-119(ed3)</i> III; <i>ltSi25[pOD928/EZ-36; prga-3::GFP::RGA-3; cb-unc-119(+)]</i> II	This study
WD478	<i>gck-1(onIs15[GFP::3xFlag::gck-1])</i> III	Pal et al., 2017

TABLE 1: Strains.

Objective (Figure 3, B–I, and Supplemental Figure 3, A and B) or 60 × 1.41 NA Nikon Oil Immersion Objective (all other figures) with a Gallium arsenide phosphide photo-multiplier tube (GaAsP PMT) detector using NIS-elements. For Figure 3, D–I, 40 confocal sections with 1 μm spacing through the entire embryo were acquired every 2.7 s using the resonance scanner. All other data acquired on the A1R were from single Z-sections at the embryonic cortex with a sampling frequency of 1 s using the Galvano point scanner.

Segmentation and analysis of fluorescence in the cytokinetic ring

For Figures 1, A and B, and 3F, the ImageJ-based software Fiji was used to analyze z-stacks of the full embryo (Schindelin et al., 2012; Schneider et al., 2012; Rueden et al., 2017). First, the cytokinetic ring was isolated. Then, an end-on reconstruction was sum-intensity projected into two dimensions. The signal in this end-on reconstruction was automatically segmented utilizing the Weka-segmentation plug-in. In MATLAB, an ellipsoid was fitted to the segmented signal at each time point, giving a measurement of ring perimeter. Fluorescence intensity on the ring at each time point was measured from sum projections of cytoplasmic-background corrected images where the signal was identified by Weka-segmentation.

For all other quantifications of equatorial fluorescence levels, single cortical planes (Figures 2, C–E, and 5, B, C, and K, and Supplemental Figure 1, C–E) or sum intensity projections of z-stacks (Figure 3, H and I) were analyzed in Fiji by performing a line scan with a width of 5 μm centered around the ingressing furrow. Mean intensity values were calculated and background adjusted using custom MATLAB scripts by subtracting the mean fluorescence intensity of a 5-μm-square region in the posterior half of the embryo from the mean intensity data at each time point.

Analysis of cytoplasmic protein levels and cytokinetic ring enrichment and RNAi depletion levels

Average fluorescence intensity in the cytoplasm was measured using Fiji by measuring the mean fluorescence intensity of a 5 μm ×

5 μm square region in the center of the embryo. Enrichment on the ring at ~50% closure was calculated by dividing ring average fluorescence by cytoplasmic average fluorescence at the same time point (Figure 1, C–D'). Depletion levels were calculated by performing a line scan in the equatorial cortex with a width of 5 μm centered around the ingressing furrow and background adjusting by subtracting the mean fluorescence intensity of a 5 μm × 5 μm square region in the cytoplasm.

Line scan analysis and calculation of anterior enrichment

Line scan analysis was performed using Fiji. Lines were drawn along the cortex from the anterior to posterior end of the embryo. The GraphPad software PRISM was then used to fit a sigmoidal line to the data and to calculate the LogIC50. The LogIC50 marks the inflection point where anterior enrichment ends. Anterior enrichment was calculated by determining the ratio of average fluorescence intensity of the most anterior 25% of the cortex to the average fluorescence intensity of the most posterior 25% of the cortex (Supplemental Figure 4).

Analysis of pseudocleavage furrow depth and persistence

In the MDX33 or MDX27 strains, which express both a membrane and DNA marker, we used Fiji to measure the distance between the two ends of the pseudocleavage furrow at the point of maximal ingression. This number was then subtracted from the total embryo width to determine the depth of the pseudocleavage furrow. To normalize between conditions, we converted the pseudocleavage depth to a percentage of embryo width (Figure 4E). To measure the persistence of the pseudocleavage furrow, the time from full regression of the pseudocleavage furrow to pronuclear meeting was calculated (Figure 4D).

Analysis of total ingression time

We imaged cytokinesis in the strain MDX33 (Figure 3E) or MDX27 (Figure 6A), which expresses both a membrane and DNA marker, with 2.7 s time resolution and calculated the time from initiation

defined as the first indentation of the membrane following anaphase onset to ~100% closure of the cytokinetic ring, defined as the point when the cytokinetic ring did not decrease in size in the next two time points.

Tracking and analyzing cortical foci

Kymographs were obtained by performing 2- μm -thick line scans along the entire A–P axis of the embryo. Cortical foci during polarization were identified and tracked using Fiji plug-in TrackMate (Tinevez et al., 2017). Prior to analysis using TrackMate, single cortical plane images acquired every second were normalized by processing to “enhance contrast” in Fiji. For multichannel image sequences, fluorescence intensities from both channels were summed for each time point to create a binary mask. The normalized and summed stack was combined with the raw data stacks into a hyperstack that was used in TrackMate. Foci were tracked using the normalized and summed stack, and values within the tracked areas on the raw data were used for quantitative analysis. We used the TrackMate LoG detector with an estimated foci diameter of 3 μm and median filter thresholding applied in TrackMate to detect foci and the LAP tracer function with maximum linking distance of 2 μm and “no gap closing distance” to track foci. Processing and analysis of foci properties determined using TrackMate were performed using custom scripts in MATLAB. Mean intensity values within the tracked region were used to determine foci intensity. Mean intensity values were background adjusted at each time point. Background values were obtained by measuring the mean intensity of a 5 \times 5 μm square area in the posterior of the embryo that was devoid of foci for each time point. Only foci persisting for more than 20 s were considered in the analysis. Background-adjusted intensity values for each track were smoothed using a gaussian filter and aligned at the maximum intensity peak. Mean intensity curves were obtained by averaging intensity values of all aligned foci for a given condition. To compare localization timing of different fluorescently tagged proteins to the same cortical foci, background-adjusted mean intensity values were processed individually for each channel for each track. Relative localization timing was calculated by comparing the timing of when the maximum mean fluorescence intensity for each protein was reached.

Figures and statistical analysis

Figures were generated using Microsoft Excel, MATLAB, or GraphPad PRISM software. Statistical significance was determined using a two-tailed Student's *t* test. Assumptions for normality and equal variance were met for all data analyzed. A *p* value of less than 0.05 from a two-tailed *t* test was considered significant. Results of the statistical analysis are shown in all figures. All error bars represent standard error unless stated otherwise in the figure legends. Sample size (*n*) and *p* values are given on each figure panel or in the figure legends.

ACKNOWLEDGMENTS

We thank Julie Canman, Brent Derry, Dan Dickinson, Jenny Hepert, Edwin Munro, and Karen Oegema for providing worm strains. We thank Stephan Grill and members of the Maddox labs, especially Paul Maddox, Jenna Perry, and Vincent Boudreau, for critical reading and discussion of the manuscript. K.R.B. was supported in part by a grant from the National Institute of General Medical Sciences under award 5T32 GM007092. This work was supported by European Research Council grant #294744 to M.L. and GM102390 from the National Institutes of Health and 1616661 from the National Science Foundation to A.S.M.

REFERENCES

- Basant A, Glotzer M (2018). Spatiotemporal regulation of RhoA during cytokinesis. *Curr Biol* 28, R570–R580.
- Bement WM, Leda M, Moe AM, Kita AM, Larson ME, Golding AE, Pfeuti C, Su KC, Miller AL, Goryachev AB, von Dassow G (2015). Activator-inhibitor coupling between Rho signalling and actin assembly makes the cell cortex an excitable medium. *Nat Cell Biol* 17, 1471–1483.
- Bischof J, Brand CA, Somogyi K, Majer I, Thome S, Mori M, Schwarz US, Lenart P (2017). A cdk1 gradient guides surface contraction waves in oocytes. *Nat Commun* 8, 849.
- Borikova AL, Dibble CF, Sciacy N, Welch CM, Abell AN, Bencharit S, Johnson GL (2010). Rho kinase inhibition rescues the endothelial cell cerebral cavernous malformation phenotype. *J Biol Chem* 285, 11760–11764.
- Brenner S (1974). The genetics of *Caenorhabditis elegans*. *Genetics* 77, 71–94.
- Bryksin AV, Matsumura I (2010). Overlap extension PCR cloning: a simple and reliable way to create recombinant plasmids. *Biotechniques* 48, 463–465.
- Ceccarelli DF, Laister RC, Mulligan VK, Kean MJ, Goudreault M, Scott IC, Derry WB, Chakrabarty A, Gingras AC, Sicheri F (2011). CCM3/PDCCD10 heterodimerizes with germinal center kinase III (GCKIII) proteins using a mechanism analogous to CCM3 homodimerization. *J Biol Chem* 286, 25056–25064.
- Cheffings TH, Burroughs NJ, Balasubramanian MK (2016). Actomyosin ring formation and tension generation in eukaryotic cytokinesis. *Curr Biol* 26, R719–R737.
- Cowan CR, Hyman AA (2007). Acto-myosin reorganization and PAR polarity in *C. elegans*. *Development* 134, 1035–1043.
- D'Avino PP (2009). How to scaffold the contractile ring for a safe cytokinesis—lessons from anillin-related proteins. *J Cell Sci* 122, 1071–1079.
- D'Avino PP, Giansanti MG, Petronczki M (2015). Cytokinesis in animal cells. *Cold Spring Harb Perspect Biol* 7, a015834.
- Descovich CP, Cortes DB, Ryan S, Nash J, Zhang L, Maddox PS, Nedelec F, Maddox AS (2018). Cross-linkers both drive and brake cytoskeletal remodeling and furrowing in cytokinesis. *Mol Biol Cell* 29, 622–631.
- Dickinson DJ, Schwager F, Pintard L, Gotta M, Goldstein B (2017). A single-cell biochemistry approach reveals PAR complex dynamics during cell polarization. *Dev Cell* 42, 416–434.e411.
- Dickinson DJ, Ward JD, Reiner DJ, Goldstein B (2013). Engineering the *Caenorhabditis elegans* genome using Cas9-triggered homologous recombination. *Nat Methods* 10, 1028–1034.
- Dorn JF, Zhang L, Phi T-T, Lacroix B, Maddox PS, Liu J, Maddox AS (2016). A theoretical model of cytokinesis implicates feedback between membrane curvature and cytoskeletal organization in asymmetric cytokinetic furrowing. *Mol Biol Cell* 27, 1286–1299.
- Fischer A, Zalvide J, Faurobert E, Albiges-Rizo C, Tournier-Lasserre E (2013). Cerebral cavernous malformations: from CCM genes to endothelial cell homeostasis. *Trends Mol Med* 19, 302–308.
- Gobel V, Barrett PL, Hall DH, Fleming JT (2004). Lumen morphogenesis in *C. elegans* requires the membrane-cytoskeleton linker erm-1. *Dev Cell* 6, 865–873.
- Goryachev AB, Leda M, Miller AL, von Dassow G, Bement WM (2016). How to make a static cytokinetic furrow out of traveling excitable waves. *Small GTPases* 7, 65–70.
- Green RA, Kao HL, Audhya A, Arur S, Mayers JR, Fridolfsson HN, Schulman M, Schloissnig S, Niessen S, Laband K, et al. (2011). A high-resolution *C. elegans* essential gene network based on phenotypic profiling of a complex tissue. *Cell* 145, 470–482.
- Green RA, Paluch E, Oegema K (2012). Cytokinesis in animal cells. *Annu Rev Cell Dev Biol* 28, 29–58.
- Heasman SJ, Ridley AJ (2008). Mammalian Rho GTPases: new insights into their functions from in vivo studies. *Nat Rev Mol Cell Biol* 9, 690–701.
- Heppert JK, Dickinson DJ, Pani AM, Higgins CD, Steward A, Ahringer J, Kuhn JR, Goldstein B (2016). Comparative assessment of fluorescent proteins for in vivo imaging in an animal model system. *Mol Biol Cell* 27, 3385–3394.
- Hickson GR, O'Farrell PH (2008). Rho-dependent control of anillin behavior during cytokinesis. *J Cell Biol* 180, 285–294.
- Hwang J, Pallas DC (2014). STRIPAK complexes: structure, biological function, and involvement in human diseases. *Int J Biochem Cell Biol* 47, 118–148.
- Jordan SN, Canman JC (2012). Rho GTPases in animal cell cytokinesis: an occupation by the one percent. *Cytoskeleton* 69, 919–930.

- Kamath RS, Ahringer J (2003). Genome-wide RNAi screening in *Caenorhabditis elegans*. *Methods* 30, 313–321.
- Kamath RS, Fraser AG, Dong Y, Poulin G, Durbin R, Gotta M, Kanapin A, Le Bot N, Moreno S, Sohrmann M, et al. (2003). Systematic functional analysis of the *Caenorhabditis elegans* genome using RNAi. *Nature* 421, 231–237.
- Khaliullin RN, Green RA, Shi LZ, Gomez-Cavazos JS, Berns MW, Desai A, Oegema K (2018). A positive-feedback-based mechanism for constriction rate acceleration during cytokinesis in *Caenorhabditis elegans*. *Elife* 7, e36073.
- Lacroix B, Maddox AS (2012). Cytokinesis, ploidy and aneuploidy. *J Pathol* 226, 338–351.
- Lant B, Yu B, Goudreault M, Holmyard D, Knight JD, Xu P, Zhao L, Chin K, Wallace E, Zhen M, et al. (2015). CCM-3/STRIPAK promotes seamless tube extension through endocytic recycling. *Nat Commun* 6, 6449.
- Li C, Wen A, Shen B, Lu J, Huang Y, Chang Y (2011). FastCloning: a highly simplified, purification-free, sequence- and ligation-independent PCR cloning method. *BMC Biotechnol* 11, 92.
- Louvi A, Nishimura S, Gunel M (2014). Ccm3, a gene associated with cerebral cavernous malformations, is required for neuronal migration. *Development* 141, 1404–1415.
- Maddox AS, Habermann B, Desai A, Oegema K (2005). Distinct roles for two C. elegans anillins in the gonad and early embryo. *Development* 132, 2837–2848.
- Maddox AS, Lewellyn L, Desai A, Oegema K (2007). Anillin and the septins promote asymmetric ingression of the cytokinetic furrow. *Dev Cell* 12, 827–835.
- Matsui T, Amano M, Yamamoto T, Chihara K, Nakafuku M, Ito M, Nakano T, Okawa K, Iwamatsu A, Kaibuchi K (1996). Rho-associated kinase, a novel serine/threonine kinase, as a putative target for small GTP binding protein Rho. *EMBO J* 15, 2208–2216.
- Mayer M, Depken M, Bois JS, Julicher F, Grill SW (2010). Anisotropies in cortical tension reveal the physical basis of polarizing cortical flows. *Nature* 467, 617–623.
- Michaux JB, Robin FB, McFadden WM, Munro EM (2018). Excitable RhoA dynamics drive pulsed contractions in the early *C. elegans* embryo. *J Cell Biol* 217, 4230–4252.
- Motegi F, Velarde NV, Piano F, Sugimoto A (2006). Two phases of astral microtubule activity during cytokinesis in *C. elegans* embryos. *Dev Cell* 10, 509–520.
- Naganathan SR, Furthauer S, Nishikawa M, Julicher F, Grill SW (2014). Active torque generation by the actomyosin cell cortex drives left-right symmetry breaking. *Elife* 3, e04165.
- Naganathan SR, Furthauer S, Rodriguez J, Fievet BT, Julicher F, Ahringer J, Cannistraci CV, Grill SW (2018). Morphogenetic degeneracies in the actomyosin cortex. *Elife* 7, e37677.
- Nishikawa M, Naganathan SR, Julicher F, Grill SW (2017). Controlling contractile instabilities in the actomyosin cortex. *Elife* 6, e19595.
- Pal S, Lant B, Yu B, Tian R, Tong J, Krieger JR, Moran MF, Gingras AC, Derry WB (2017). CCM-3 promotes *C. elegans* germline development by regulating vesicle trafficking cytokinesis and polarity. *Curr Biol* 27, 868–876.
- Piekny A, Werner M, Glotzer M (2005). Cytokinesis: welcome to the Rho zone. *Trends Cell Biol* 15, 651–658.
- Piekny AJ, Glotzer M (2008). Anillin is a scaffold protein that links RhoA, actin, and myosin during cytokinesis. *Curr Biol* 18, 30–36.
- Piekny AJ, Maddox AS (2010). The myriad roles of anillin during cytokinesis. *Semin Cell Dev Biol* 21, 881–891.
- Pollard TD, O’Shaughnessy B (2019). Molecular mechanism of cytokinesis. *Annu Rev Biochem* 88, 661–689.
- Priti A, Ong HT, Toyama Y, Padmanabhan A, Dasgupta S, Krajnc M, Zaidel-Bar R (2018). Syncytial germline architecture is actively maintained by contraction of an internal actomyosin corset. *Nat Commun* 9, 4694.
- Rehain-Bell K, Love A, Werner ME, MacLeod I, Yates JR 3rd, Maddox AS (2017). A sterile 20 family kinase and its co-factor CCM-3 regulate contractile ring proteins on germline intercellular bridges. *Curr Biol* 27, 860–867.
- Reymann AC, Staniscia F, Erzberger A, Salbreux G, Grill SW (2016). Cortical flow aligns actin filaments to form a furrow. *eLife* 5, 1–25.
- Richardson BT, Dibble CF, Borikova AL, Johnson GL (2013). Cerebral cavernous malformation is a vascular disease associated with activated RhoA signaling. *Biol Chem* 394, 35–42.
- Rueden CT, Schindelin J, Hiner MC, DeZonia BE, Walter AE, Arena ET, Eliceiri KW (2017). ImageJ2: ImageJ for the next generation of scientific image data. *BMC Bioinformatics* 18, 529.
- Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B, et al. (2012). Fiji: an open-source platform for biological-image analysis. *Nat Methods* 9, 676–682.
- Schmutz C, Stevens J, Spang A (2007). Functions of the novel RhoGAP proteins RGA-3 and RGA-4 in the germ line and in the early embryo of *C. elegans*. *Development* 134, 3495–3505.
- Schneider CA, Rasband WS, Eliceiri KW (2012). NIH Image to ImageJ: 25 years of image analysis. *Nat Methods* 9, 671–675.
- Schonegg S, Constantinescu AT, Hoege C, Hyman AA (2007). The Rho GTPase-activating proteins RGA-3 and RGA-4 are required to set the initial size of PAR domains in *Caenorhabditis elegans* one-cell embryos. *Proc Natl Acad Sci USA* 104, 14976–14981.
- Schouest KR, Kurasawa Y, Furuta T, Hisamoto N, Matsumoto K, Schumacher JM (2009). The germinal center kinase GCK-1 is a negative regulator of MAP kinase activation and apoptosis in the *C. elegans* germline. *PLoS One* 4, e7450.
- Straight AF, Cheung A, Limouze J, Chen I, Westwood NJ, Sellers JR, Mitchison TJ (2003). Dissecting temporal and spatial control of cytokinesis with a myosin II inhibitor. *Science* 299, 1743–1748.
- Sun L, Guan R, Lee IJ, Liu Y, Chen M, Wang J, Wu JQ, Chen Z (2015). Mechanistic insights into the anchorage of the contractile ring by anillin and Mid1. *Dev Cell* 33, 413–426.
- Tinevez JY, Perry N, Schindelin J, Hoopes GM, Reynolds GD, Laplantine E, Bednarek SY, Shorte SL, Eliceiri KW (2017). TrackMate: an open and extensible platform for single-particle tracking. *Methods* 115, 80–90.
- Tse YC, Werner M, Longhini KM, Labbe JC, Goldstein B, Glotzer M (2012). RhoA activation during polarization and cytokinesis of the early *Caenorhabditis elegans* embryo is differentially dependent on NOP-1 and CYK-4. *Mol Biol Cell* 23, 4020–4031.
- Van Furden D, Johnson K, Segbert C, Bossinger O (2004). The *C. elegans* ezrin-radixin-moesin protein ERM-1 is necessary for apical junction remodeling and tubulogenesis in the intestine. *Dev Biol* 272, 262–276.
- Velarde N, Gunsalus KC, Piano F (2007). Diverse roles of actin in *C. elegans* early embryogenesis. *BMC Dev Biol* 7, 142.
- Werner M, Glotzer M (2008). Control of cortical contractility during cytokinesis. *Biochem Soc Trans* 36, 371–377.
- Werner M, Munro E, Glotzer M (2007). Astral signals spatially bias cortical myosin recruitment to break symmetry and promote cytokinesis. *Curr Biol* 17, 1286–1297.
- Xu X, Wang X, Zhang Y, Wang DC, Ding J (2013). Structural basis for the unique heterodimeric assembly between cerebral cavernous malformation 3 and germinal center kinase III. *Structure* 21, 1059–1066.
- Yin H, Shi Z, Jiao S, Chen C, Wang W, Greene MI, Zhou Z (2012). Germinal center kinases in immune regulation. *Cell Mol Immunol* 9, 439–445.
- Zanin E, Desai A, Poser I, Toyoda Y, Andree C, Moebius C, Bickle M, Conradt B, Piekny A, Oegema K (2013). A conserved RhoGAP limits M phase contractility and coordinates with microtubule asters to confine RhoA during cytokinesis. *Dev Cell* 26, 496–510.
- Zhang D, Glotzer M (2015). The RhoGAP activity of CYK-4/MgcRacGAP functions non-canonically by promoting RhoA activation during cytokinesis. *Elife* 4, e08898.
- Zhang M, Dong L, Shi Z, Jiao S, Zhang Z, Zhang W, Liu G, Chen C, Feng M, Hao Q, et al. (2013). Structural mechanism of CCM3 heterodimerization with GCKIII kinases. *Structure* 21, 680–688.
- Zheng X, Xu C, Di Lorenzo A, Kleaveland B, Zou Z, Seiler C, Chen M, Cheng L, Xiao J, He J, et al. (2010). CCM3 signaling through sterile 20-like kinases plays an essential role during zebrafish cardiovascular development and cerebral cavernous malformations. *J Clin Invest* 120, 2795–2804.