Burrowing in blindsnakes: a preliminary analysis of burrowing forces and consequences for the evolution of morphology

Anthony Herrel, Aurélien Lowie, Aurélien Miralles, Philippe Gaucher, Nathan J Kley, John Measey, Krystal A Tolley

► To cite this version:

Anthony Herrel, Aurélien Lowie, Aurélien Miralles, Philippe Gaucher, Nathan J Kley, et al.. Burrowing in blindsnakes: a preliminary analysis of burrowing forces and consequences for the evolution of morphology. *The Anatomical Record: Advances in Integrative Anatomy and Evolutionary Biology*, 2021, 10.1002/ar.24686 . hal-03253463

HAL Id: hal-03253463

<https://hal.sorbonne-universite.fr/hal-03253463>

Submitted on 8 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Burrowing in blindsnakes: a preliminary analysis of burrowing forces and consequences for the evolution of morphology.

Anthony Herrel^{1,2}, Aurélien Lowie², Aurélien Miralles³, Philippe Gaucher⁴, Nathan J. Kley⁵, John Measey⁶, Krystal A. Tolley^{7,8}

1. UMR 7179 C.N.R.S/M.N.H.N., Département Adaptations du Vivant, Bâtiment d'Anatomie Comparée, 55 rue Buffon, 75005, Paris, France.

2. Department of Biology, Evolutionary Morphology of Vertebrates, Ghent University, K.L. Ledeganckstraat 35, 9000 Ghent, Belgium.

3. Institut de Systématique, Évolution, Biodiversité (ISYEB), Muséum national d'Histoire naturelle, CNRS, Sorbonne Université, EPHE, Université des Antilles, 57 rue Cuvier, CP 50, 75005 Paris, France.

4. Laboratoire Ecologie, Evolution, Interactions des Systèmes amazoniens Centre de Recherche de Montabo, BP 70620, 97334 Cayenne cédex, French Guyana.

5. Department of Anatomical Sciences, Stony Brook University, Stony Brook, NY, USA

6. Centre for Invasion Biology, Department of Botany & Zoology, Stellenbosch University, Stellenbosch, South Africa.

7. Kirstenbosch Research Centre, South African National Biodiversity Institute, Private Bag X7, Claremont 7735, Cape Town, South Africa.

8. School of Animal, Plant and Environmental Sciences, University of the Witwatersrand, P.O. Wits, 2050 Johannesburg, South Africa

Orcid AH: 0000-0003-0991-4434; AL: 0000-0003-0065-7152; AM: 0000-0002-2538-7710; JM: 0000-0001-9939-7615; KT: 0000-0002-7778-1963

pages: 16; # tables: 1; # figures: 4

Address for correspondence:

Anthony Herrel

UMR 7179 C.N.R.S/M.N.H.N.

Département Adaptations du Vivant, Bâtiment d'Anatomie Comparée

55 rue Buffon, 75005, Paris, France

e-mail: anthony.herrel@mnhn.fr

Abstract

Burrowing is a common behavior in vertebrates. An underground life-style offers many advantages but also poses important challenges including the high energetic cost of burrowing. Scolecophidians are a group of morphologically derived subterranean snakes that show great diversity in form and function. Although it has been suggested that leptotyphlopids and anomalepidids mostly use existing underground passageways, typhlopids are thought to create their own burrows. However, the mechanisms used to create burrows and the associated forces that animals may be able to generate remain unknown. Here, we provide the first data on push forces in scolecophidians and compare them with those in some burrowing alethinophidian snakes. Our results show that typhlopids are capable of generating higher forces for a given size than other snakes. The observed differences are not due to variation in body diameter or length, suggesting fundamental differences in the mechanics of burrowing or the way in which axial muscles are used. Qualitative observations of skull and vertebral shape suggest that the higher forces exerted by typhlopids may have impacted the evolution of their anatomy. Our results provide the basis for future studies exploring the diversity of form and function in this fascinating group of animals. Quantitative comparisons of the cranial and vertebral shape in addition to collecting functional and ecological data on a wider array of species would be particularly important to test the patterns described here.

Introduction

Burrowing is a common behavior among vertebrates that dates back at least to the Devonian (Benton, 1988; Kinlaw, 1999). The exploitation of the underground offers many advantages including shelter from predators, novel prey resources and an environment buffered from extreme fluctuations in temperature (Sumner et al., 2004). However, the underground environment also poses important challenges including low oxygen content (McNabb, 1966; Arieli, 1979), and the high energetic cost of creating burrows (Vleck, 1979; Navas et al., 2004). Tetrapods have radiated into the underground environment many times independently and have developed solutions to the constraints of burrowing and underground locomotion (i.e., substrate density; see Nevo, 1979). As the cost of creating tunnels is directly related to the diameter of the animal (Navas et al., 2004; Wu et al., 2015), it is no surprise that many limbless vertebrates, characterized by a reduced body diameter, have colonised the underground environment including caecilians, lizards and snakes (Gans, 1986). However, the mechanisms used to create, enlarge, or maintain burrows remain relatively poorly understood (but see Gaymer, 1971; Gans et al., 1978; Gasc, 1982; O'Reilly et al., 1997; Quillin, 2000; Dorgan et al., 2005; Dorgan, 2015).

Among limbless tetrapods, snakes occupy a special place as many forms are known to burrow (Davis, 1946; Young & Morain, 2003; Deuffel, 2017). Moreover, snakes have been suggested to have gone through a fossorial phase early-on in their evolutionary history (da Silva et al., 2018; Miralles et al., 2018). Despite extensive studies on locomotion in limbless lizards in general, and snakes in particular (e.g., Gans, 1973, 1986; Gasc, 1984; Gasc & Gans, 1990; Hohl et al., 2014; Newman and Jayne, 2018), surprisingly little is known about their burrowing mechanics. Even more surprisingly, not a single study has been devoted to the analysis of burrowing in scolecophidians, an entire 'clade' of burrowing snakes. To date, only a single study has described the anatomy of the muscles of the axial system in any detail (Gasc, 1981), rendering our understanding of how these animals may create burrows nearly nonexistent.

Scolecophidians are now divided into two superfamilies, the Typhlopoidea (blind snakes) and Leptotyphlopoidea (thread snakes). Based on their anatomy the Anomalepidae have long been regarded as a third group within Scolecophidia, but recent molecular phylogenies suggest they might actually be more closely related to Alethinophidians (cfr. Miralles et al., 2018). Scolecophidians diverged from other snakes (Alethinophidia) about 125 to 130 Mya in the Early Cretaceous (Zheng & Wiens, 2016; Schneider Fachini et al., 2020), and the divergence between Typhlopoidea and Leptotyphlopoidea probably followed soon thereafter, around 120 – 130 Mya (Pyron & Burbrink, 2012; Zheng & Wiens, 2016; Miralles et al., 2018; but see Vidal et al., 2010). Comparatively, most

alethinophidian snake families diverged beginning in the Eocene into the Oligocene between 40 – 60 Mya (see Pyron & Burbrink, 2012; Zheng & Wiens, 2016; Zaher et al., 2019). Thus, the scolecophidians have long been on a separate evolutionary trajectory from other snakes, but within the group itself the family-level lineages are characterized by long branch lengths and early divergences dating back to the Cretaceous (Adalsteinsson et al., 2009; Pyron & Burbrink, 2012; Miralles et al., 2018).

Alethinophidian snakes are a species-rich clade (~3,900 species) with at least one clade, the Endoglyptodonta (*sensu* Zaher et al., 2019, or Colubroidea *sensu* Pyron & Burbrink, 2012), contributing to the bulk of the richness. Synapomorphies for this clade are cranial features comprising a venom delivery system, and this key innovation may have allowed them to diversify so suddenly and so widely (Pyron & Burbrink, 2012). The scolecophidians are comparatively species-poor with just 459 species described (see Uetz et al., 2020), although it is very likely that there are numerous cryptic species (Thomas & Hedges, 2007; Busschau et al., 2021). The bulk of these species (60%) resides within one family, the Typhlopidae (Pyron & Wallach, 2014; Uetz et al., 2020). This family has also been noted to have an unusually high net diversification rate (Pyron & Burbrink, 2012) but the mechanisms that drove this elevated diversification rate are obscure. It is possible that the initial advancement of scolecophidians toward specialization was their capitalization on a new trophic niche that arose around 130 Mya; i.e., the divergence of ants from other Hymenoptera (see Brady et al., 2006) and their subsequent dominance of ecosystems, making them a reliable prey source. Furthermore, the increased speciation rate for Typhlopidae beginning around 50 Mya (Pyron & Burbrink, 2012) could have been a response to the increased speciation rate that also occurred around 50 Mya in several New World ant genera (see Moreau & Bell, 2013).

Although scolecophidians have a conserved morphology and have therefore been erroneously termed ‘primitive’ or ‘basal’, they are in fact extremely specialized with a phenotype that is well-adapted to a strictly fossorial life-style (da Silva et al., 2018; Miralles et al., 2018). They forage underground, primarily on the eggs, larvae, pupae, and adults of ants and termites, and unlike alethinophidians, they consume large numbers of small prey (Shine & Webb, 1990; Webb & Shine 1993; Webb et al., 2001). This requires them to move through existing underground passageways (in the case of leptotyphlopids) or to create new underground tunnels (in the case of typhlopids) to invade ant or termite colonies. They use chemoreception to locate their prey, but this does not appear to be restricted to the detection of only a single prey species (Watkins et al., 1967; Webb & Shine, 1992). Leptotyphlopids can avoid detection from ants and termites through chemical crypsis, and this is possibly achieved through sequestration of their prey’s defensive compounds with the

subsequent release of these compounds through glands (Watkins et al., 1969; Webb et al., 2000; Savitzky et al., 2012). Conversely, typhlopids probably withstand attacks by ants or termites through their more heavily armoured body (Webb & Shine, 1993). Thus, while the two superfamilies of scolecophidians have basic similarities in terms of life-history and are superficially similar in morphology, they probably have quite different adaptations allowing them to have a highly specialised ecological niche and different specializations towards burrowing.

Here, we provide new data on burrowing performance and cranial and vertebral morphology in scolecophidian snakes and compare them to data for some alethinophidian burrowers. Given that leptotyphlopids are thought to use existing burrows in contrast to typhlopids which construct their own burrows, we predict that typhlopids will be better burrowers for their size. As most alethinophidian burrowers utilize the relatively ‘soft’ top compartment of the soil we predict them to be more similar to leptotyphlopids and produce less force for a given size.

Materials and methods

Animals

Burrowing forces were measured in the field in Kenya (*Afrotyphlops angolensis*, *Rhinotyphlops unitaeniatus*), or South Africa (*Leptotyphlops scutifrons*, *Rhinotyphlops lalandei*), or in the lab (*Liotyphlops beui*, *Myriopholis algeriensis*). For comparative purposes we also recorded data on alethinophidian snakes in the field (Kenya: *Aparallactus guentheri*; French Guyana: *Anilius scytale*, *Oxyrhopus melanogenys*) and in the lab (*Eryx colubrinus*, *Farancia abacura*, *Loxocemus bicolor*). The number of individuals per species used is summarized in Table 1.

To better understand how the measured burrowing forces may impact the anatomy of the head and vertebrae we CT-scanned one specimen each of *Anilius unguirostris* (MNHN 1895.449; voxel size: 16.3 μm), *Epictia tenella* (MNHN 2011.319; voxel size: 11.2 μm), and *Typhlophis squamosus* (MNHN 1999.8306; voxel size: 8.81 μm) at the AST-RX platform at the Muséum national d’histoire naturelle on a v|tome|x machine. For comparative purposes we also used CT-scans of *Anilius scytale* (KUH 125976; voxel size: 23.7 μm), *Leptotyphlops nigricans* (LSUSM Z57237; voxel size: 3.5 μm), *Liotyphlops albirostris* (UMMZ 48173; voxel size: 10.8 μm) and *Rhinotyphlops lalandei* (UMMZ 61525; voxel size: 11.3 μm) that were downloaded from Morphosource.

Morphometrics

Animals captured in the field were weighed using an electronic balance (Ohaus, ± 0.1 g) or a Pesola spring scale (± 0.5 g). Body diameter was measured using a digital calliper (Mitutoyo, ± 0.1 mm). The snout-vent length was measured by stretching the animals along a ruler (± 1 mm). A summary of the morphometric data is provided in Table 1.

Force measurements

Measurements of peak push forces were made using a piezoelectric force platform (Kistler Squirrel force plate, ± 0.1 N, Kistler Inc., Switzerland) as described previously (Vanhooydonck et al., 2011; Le Guilloux et al., 2020; Fig. 1). In brief, the force platform was positioned on a metal base and connected to a charge amplifier (Kistler Charge Amplifier type 9865, Kistler Inc.). A Perspex block with 1 cm-deep holes of different diameters was mounted on the force plate, level with the front edge. One of the holes was loosely filled with soil. A Perspex tunnel with a diameter approximatively equal to the maximal body diameter of the test animal was mounted on the metal base in front of (but not touching) the force plate, and aligned with the soil-filled hole in the Perspex block. An animal was then introduced into the tunnel and allowed to move through it until reaching the soil-filled chamber. Next, the animal was stimulated to burrow into the soil by touching the end of the tail sticking out of the tunnel, or by prodding the animal inside the tunnel with the blunt end of a thin wooden stick. Forces were recorded during 60 s recording sessions at 500 Hz, and three trials were performed for each individual, with at least 1 h between trials. Forces were recorded in three dimensions using the Bioware software (Kistler Inc.; Fig. 1). For each individual we then extracted the highest peak resultant force across all trials as an indicator of that animal's maximal push force. A summary of the force data is provided in Table 1.

Statistical analyses

All data were Log₁₀-transformed before analyses to ensure normality and homoscedasticity. To explore which traits (snout-vent length, body mass or body diameter) best explained variation in push force data (resultant force) we ran a stepwise multiple regression across data for all individuals. Next, we tested whether differences between groups (Alethinophidia, Typhlopidae, Leptotyphlopidae, Anomalepididae) were observed in resultant force, using an ANCOVA with snout-vent length as our co-variate. To test which groups differed from one another, we next ran simple regressions on maximal push force (resultant), body diameter, and body mass with snout-vent length as our predictor and extracted unstandardized residuals. We then ran an ANOVA on residual resultant force coupled to Bonferroni post-hoc tests to explore which groups differed from one another. Finally, we ran Pearson correlations between residual data to explore whether body

diameter or body mass, independent of the effect of snout-vent length, explained variation in residual maximal resultant force and ran a MANOVA to test whether clades differed in residual body mass and residual body diameter. All analyses were run in IBM SPSS V. 26. Significance was set at $P < 0.05$.

Although the data set is composed of different species and sometimes contains several individuals within species, we decided not to use phylogenetic comparative methods on species means given the small sample sizes and incomplete taxon sampling. Future analyses on larger and more robust data sets could provide an assessment on whether the patterns described here are born out.

Results

Maximal push forces across the species studied ranged from 0.21 N for *Myriopholis algeriensis*, the smallest species in our data set, to 18.18 N for *Afrotyphlops angolensis*, the largest scolecophidian in our data set (Table 1). In all taxa, forward-directed forces made up between 50 and 65% of the total force, suggesting that all taxa are capable of generating forward-directed forces (Fig. 1). Lateral and dorso-ventral forces contributed roughly equally (between 16 and 25%) to the remainder of the forces generated. No clear differences were observed between groups in how forces were applied, however. Rather individuals within a species sometimes showed different strategies, with one *Rhinotyphlops lalandei* pushing only 34% in the forward direction, whereas in another individual 59% of the force was directed forward.

A multiple stepwise regression with snout-vent length, body mass and body diameter retained a significant model with only snout-vent length as the predictor ($R^2 = 0.78$; $P < 0.001$; slope = 1.93 ± 0.20 , intercept = -4.40 ± 0.49). Thus, longer snakes were able to produce higher resultant forces (Fig. 2A). The ANCOVA detected significant effects of snout-vent length ($F_{1,28} = 62.54$; $P < 0.001$) and phylogenetic group ($F_{3,28} = 70.70$; $P < 0.001$) on maximal push force, suggesting that individuals from different phylogenetic groups differed in their maximal push force irrespective of variation in snout-vent length. The ANOVA run on residual maximal push force was significant ($F_{3,29} = 8.87$; $P < 0.001$). Post-hoc tests indicated that Typhlopidae differed from all other groups (all $P < 0.018$), yet none of the other groups differed. The inspection of the marginal means showed that typhlopids produced higher maximal push forces for a given snout-vent length than other species (marginal means; Typhlopidae: 0.329; Leptotyphlopidae: -0.149; Anomalepididae: -0.181; Alethinophidia: -0.053). Finally, residual maximal push force was correlated with residual body diameter ($r = 0.44$; $P = 0.011$) but not residual body mass ($r = 0.33$; $P = 0.077$), showing that snakes with relatively wider bodies were able to push harder (Fig. 2B). Yet, the difference in push force was not explained by the

difference in body diameter as a MANOVA on residual body diameter and residual body mass detected no differences between groups (Wilks' lambda = 0.69; $F_{6,48} = 1.67$; $P = 0.15$; uni-variate ANOVAs: body diameter: $F_{3,25} = 2.25$; $P = 0.11$; body mass: $F_{3,25} = 2.49$; $P = 0.08$).

Inspection of the CT-scans showed interesting differences in cranial and vertebral anatomy between the different species examined (Fig. 3). Overall, all burrowing species had blunt and robust snouts in dorsal view with surprisingly unfused and rather simple sutures. However, in lateral view differences between species were striking. Aside from differences in the tooth-bearing elements associated with their radically different feeding modes (Kley & Brainerd, 1999; Kley, 2001, 2006; Rieppel et al., 2009; Strong et al., 2021), the shape of the anterior part of the cranium was very different with *Anilius* and *Rhinotyphlops* having more pointed shapes. The vertebral morphology was also rather different with the condyle and cotyle of *Anilius* being wide and tall compared to those in the other species (Fig. 4). Moreover, the facets of the pre- and postzygapophyses appear more laterally positioned and more robust in *Anilius* (Fig. 4). Note however, that *Anilius* was larger than the other species examined and as such that allometry may partly be responsible for the observed differences.

Discussion

The first ever data on maximal push forces in burrowing snakes suggest that significant interspecific differences exist, with typhlopids snakes being able to generate higher forces for a given body length. As typhlopids are considered active burrowers in contrast to leptotyphlopids, which are thought to mostly use existing underground passageways, this observation is in line with our predictions. Alethinophidian snakes also generated relatively low forces for their size which may correspond to the fact that they mostly use the softer top compartment of the soil for burrowing. Interestingly, although snout-vent length and residual body diameter are both correlated to the force generation capacity in the animals included in our data set, these variables did not explain why typhlopids snakes were able to generate higher forces. Longer snakes can be expected to have more overall muscle mass and an increase in the relative diameter should allow for the packing of muscles with a greater cross-sectional area (Gans, 1974). This is confirmed by our results where longer snakes do indeed have a greater diameter (Pearson correlation: $r = 0.86$; $P < 0.001$). Whereas this may explain the observed relationships between force, length and diameter, this does not explain differences between the different groups of snakes. It is likely that different burrowing snakes use different mechanisms for burrowing (as suggested for amphisbaenians with different head shapes for example; Gans, 1974), allowing them to recruit the axial muscles differently. This is translated in differences in overall proportions with leptotyphlopids being generally extremely narrow and long in contrast to typhlopids which are much more robust and have a greater diameter for a given snout-

vent length [note that exceptions exist with, for example, *Anilius grypus* showing an aspect ratio (total length divided by midbody diameter) of 130].

Although the exact mechanisms used by these snakes to generate burrowing force while in a tunnel remains unknown, the mostly forward-directed forces (very different from what has been described in burrowing scincid lizards, for example; see Vanhooydonck et al., 2011) suggest that they may use a mechanism similar to what has been described for caecilians (Gaymer, 1971; O'Reilly et al., 1997) and hypothesized to exist in uropeltid snakes (Gans et al., 1978): internal concertina. This type of burrowing involves bending followed by an extension of the vertebral column within the skin envelope, allowing animals to recruit most of their body wall muscles to generate forward-directed forces. In contrast, non-specialized snakes will use muscular forces to push laterally against the side of a straight-sided tunnel in a typical concertina movement to traverse smooth tunnels (Gray & Lissmann, 1950; Gans, 1974). X-ray video analysis is needed to be able to understand the movements of the skin and vertebral column and thus to test whether skin-vertebral independence exists in scolecophidian snakes in general and typhlopids snakes more specifically. Unfortunately, our understanding of the axial musculature in burrowing snakes, and especially in scolecophidian snakes, remains extremely fragmentary (Gasc, 1981). The musculature has been suggested to be convergent on that observed in amphisbaenian lizards, another group of very strong burrowers (Gasc, 1982; Navas et al., 2004; Hohl et al., 2014) but with a typical snake bauplan (Gasc, 1981). A remarkable feature of the muscular anatomy in typhlopids is the absence of the *m. costo-cutanei* resulting in an anatomical independence between the axial and cutaneous muscles (Gasc, 1981). However, these data are based on the dissection of a single specimen of *Afrotyphlops* (*A. punctatus*) and differ from what has been reported by Mosauer (1935), illustrating the need for further studies on the axial musculature. The small size of many leptotyphlopids and anomalepids has prevented direct dissections of the musculature. However, contrast-enhanced μ CT scans may provide a unique opportunity to better understand the musculature in these extremely small animals (Metscher, 2009).

Our qualitative anatomical comparisons of skull and vertebral morphology suggest that the higher forces generated by typhlopids snakes may be reflected in the shape of these structures. Whereas all burrowing species had blunt and robust snouts in dorsal view, the shape of the anterior part of the cranium was very different with *Anilius* and *Rhinotyphlops* having more pointed shapes possibly facilitating substrate penetration. Given the high forces encountered during burrowing it was unexpected to see little or no fusion of the cranial sutures, in contrast to what is observed in, for example, trogonophid amphisbaenians where the cranial sutures are highly interdigitated (Gans,

1974). Elongate and highly interdigitated sutures have been suggested to be a response to torsional stresses in the skull induced by oscillatory digging (Gans, 1974). The fact that the recorded forces were mostly directed forward may explain the difference in skull morphology between burrowing snakes and amphisbaenians. However, the snout-complexes in scolecophidian skulls are often characterized by overlapping skull bones and inspection of the CT-scans showed this to be the case in the species included in the present study. Moreover, qualitatively this appeared to be more striking in typhlopids compared to leptotyphlopids and anomalepids, but this needs to be tested quantitatively. If confirmed, this could provide a mechanism by which the snout is reinforced and may allow typhlopids to withstand the higher forces generated during burrowing. The vertebral morphology was also rather different with the condyle and cotyle of the typhloid *Anilius* being relatively wide and tall. Moreover, the facets of the pre- and postzygapophyses appear more laterally positioned and more robust in this species. Given the higher push forces recorded for typhloid snakes the larger surface areas in contact with the cranium may permit a better dissipation of the substrate reaction forces during burrowing. Similarly, the more robust pre- and postzygapophyses may help dissipate load in addition to stabilizing the cervical vertebrae when loaded in compression during burrowing. Finite element analyses of the cranium and vertebrae in these animals could be particularly useful to better understand whether the observed morphology is indeed a response to the forces encountered during burrowing. Additionally, exploring variation due to allometry would be important as the animals examined here differed considerably in size.

The use of 3D geometric morphometric approaches (e.g., Fabre et al., 2016; Segall et al., 2016) could be of interest to: 1) quantify possible differences in head, cranial and vertebral shape in typhlopids in comparison to other burrowing snakes, and 2) to explore whether cranial and vertebral shape covary with the push forces measured *in vivo*. These studies are ongoing and may provide better insights into the evolution of burrowing and the diversity of form and function in burrowing snakes. Finally, as our study included only a very small part of the diversity of burrowing snakes, scolecophidian or otherwise, future measurements on other species are likely to change our understanding of the evolution of diversity of form and function in burrowing snakes.

Acknowledgements

The authors would like to thank Jim O'Reilly for his insights and for allowing us to use data collected in his lab. This work was supported by a grant from Agence Nationale de la Recherche under the LabEx ANR-10-LABX-0003-BCDiv, in the program 'Investissements d'avenir' n°0001 ANR-11-IDEX-0004-02 and the National Research Foundation (NRF) of South Africa (South

304 African Biosystematics Initiative-SABI, Key International Science Collaboration-KISC). Field
305 research was conducted under provincial permits in South Africa (Northern Cape: FAUNA
306 1243/2008; Limpopo: 333-00015) and permit MOEST 13/0001/36C 183 in Kenya. Ethics
307 clearance for field work in South Africa was from SANBI (0010/08).

308 References

- 309 Adalsteinsson, S. A., Branch, W. R., Trape, S., Vitt, L. J. & Hedges, S. B. (2009). Molecular phylogeny,
310 classification, and biogeography of snakes of the family Leptotyphlopidae (Squamata, Scolecophidia).
311 *Zootaxa*, 2244, 1-50.
- 312 Arieli, R. (1979). The atmospheric environment of fossorial mole rat (*Spalax ehrenbergi*): effect of
313 season, soil texture, rain, temperature and activity. *Comparative Biochemistry and Physiology A*
314 *Comparative Physiology*, 63, 569-575.
- 315 Benton, M. J. (1988). Burrowing by vertebrates. *Nature*, 331, 17-18.
- 316 Brady, S. G., Schultz, T. R., Fisher, B. L., & Ward, P. S. (2006). Evaluating alternative hypotheses for the
317 early evolution and diversification of ants. *Proceedings of the National Academy of Sciences U.S.A.*,
318 103, 18172-18177.
- 319 Busschau, T., Conradie, W., & Daniels, S. R. (2021). One species hides many: Molecular and
320 morphological evidence for cryptic speciation in a thread snake (Leptotyphlopidae: *Leptotyphlops*
321 *syvicolus* Broadley & Wallach, 1997). *Journal of Zoological Systematics and Evolutionary Research*,
322 59, 195-221.
- 323 Da Silva, F. O., Fabre, A-C., Savriama, Y., Ollonen, J., Mahlow, K., Herrel, A., Müller, J., & Di-Poï, N.
324 (2018). The ecological origins of snakes as revealed by skull evolution. *Nature Communications*, 9,
325 376.
- 326 Davis, D. D. (1946). Observations on the burrowing behavior of the hog-nosed snake. *Copeia*, 1946,
327 75-78.
- 328 Deufel, A. (2017). Burrowing with a kinetic snout in a snake (Elapidae: *Aspidelaps scutatus*). *Journal of*
329 *Morphology*, 278, 1706-1715.

330 Dorgan, K. M. (2015). The biomechanics of burrowing and boring. *Journal of Experimental Biology*,
331 218, 176-183.

332 Dorgan, K. M., Jumars, P. A., Johnson, B., Boudreau, B. P., & Landis, E. (2005). Burrow extension by
333 crack propagation. *Nature*, 433, 475.

334 Fabre, A-C., Bickford, D., Segall, M., & Herrel, A. (2016). The impact of diet, habitat use, and behavior
335 on head shape evolution in homalopsid snakes. *Biological Journal of the Linnean Society*, 118, 634-
336 647.

337 Gans, C. (1973). Locomotion and burrowing in limbless vertebrates. *Nature*, 242, 414-415.

338 Gans, C. (1974). Biomechanics. An approach to vertebrate biology. University of Michigan Press. 261
339 Pp.

340 Gans, C. (1986). Locomotion of limbless vertebrates: pattern and evolution. *Herpetologica*, 42, 33-46.

341 Gans, C., Dessauer, H. C., & Baic, D. (1978). Axial differences in the musculature of uropeltid snakes:
342 the freight-train approach to burrowing. *Science*, 199, 189-192.

343 Gasc, J-P. (1981). Axial musculature. In *Biology of the Reptilia*, Vol. 11 Morphology F (Eds. C. Gans &
344 T. Parsons). Pp. 355-435. Academic Press: London.

345 Gasc, J-P. (1982). Le mécanisme du fouissage chez *Amphisbaena alba* (Amphisbaenidae, Squamata).
346 *Vertebrata Hungarica*, 21, 147-155.

347 Gasc, J-P. (1984). Analyse expérimentale du déplacement du saurien *Bachia cophias* (Schneider),
348 microtéliidé a membres réduits. Un modèle simplifié de reptation par ondulations. *Amphibia-Reptilia*,
349 5, 355-365.

350 Gasc, J-P., & Gans, C. (1990). Tests on locomotion of the elongate and limbless lizard *Anguis fragilis*
351 (Squamata: Anguidae). *Copeia*, 1990, 1055-1067.

352 Gaymer, R. (1971). New method of locomotion in limbless terrestrial vertebrates. *Nature*, 234, 150-
353 151.

354 Gray, J., & Lissmann, H. W. (1950). The kinetics of locomotion of the grass-snake. *Journal of*
355 *Experimental Biology*, 26, 354-367.

356 Hohl, L. S. L., Loguerico, M. F. C., Buendia, R. A., Almeida-Santos, M., Viana, L. A., Barros-Filho, J. D., &
 357 Rocha-Barbosa, O. (2014). Fossorial gait patterns and performance of a shovel-headed
 358 amphisbaenian. *Journal of Zoology*, 294, 234-240.

359 Kinlaw, A. (1999). A review of burrowing by semi-fossorial vertebrates in arid environments. *Journal*
 360 *of Arid Environments*, 41, 127-145.

361 Kley, N. J. (2001). Prey transport mechanisms in blindsnakes and the evolution of unilateral feeding
 362 systems in snakes. *American Zoologist*, 41, 1321-1337.

363 Kley, N. J. (2006). Morphology of the lower jaw and suspensorium in the Texas blindsnake,
 364 *Leptotyphlops dulcis* (Scoleophidia: Leptotyphlopidae). *Journal of Morphology*, 267, 494-515.

365 Kley, N. J., & Brainerd, E. (1999). Feeding by mandibular raking in a snake. *Nature*, 402, 369-370.

366 Le Guilloux, M., Miralles, A., Measey, J., Vanhooydonck, B., O'Reilly, J. C., Lowie, A., & Herrel, A.
 367 (2020). Trade-offs between burrowing and biting force in fossorial scincid lizards? *Biological Journal*
 368 *of the Linnean Society*, 130, 310-319.

369 McNab, B. K. (1966). The metabolism of fossorial rodents: a study of convergence. *Ecology*, 47, 712-
 370 733.

371 Metscher, B. D. (2009). MicroCT for comparative morphology: simple staining methods allow high-
 372 contrast 3D imaging of diverse non-mineralized animal tissues. *BMC Physiology*, 9, 11.

373 Miralles, A., Marin, J., Markus, D., Herrel, A., Hedges, S. B., & Vidal, N. (2018). Molecular evidence for
 374 the paraphyly of Scoleophidia and its evolutionary implications. *Journal of Evolutionary Biology*, 31,
 375 1782-1793.

376 Moreau, C. S., & Bell, C. D. (2013). Testing the museum versus cradle tropical biological diversity
 377 hypothesis: phylogeny, diversification, and ancestral biogeographic range evolution of the ants.
 378 *Evolution*, 67, 2240-2257.

379 Mosauer, W. (1935). The myology of the trunk region of snakes and its significance for ophidian
 380 taxonomy and phylogeny. University of California at Los Angeles Publications in Biological Sciences 1:
 381 81-120.

382 Navas, C. A., Antoniazzi, M. M., Carvalho, J. E., Chaui-Berlink, J. G., James, R. S., Jared, C., Kohlsdorf,
383 T., Pai-Silva, M. D., & Wilson, R. S. (2004). Morphological and physiological specialization for digging
384 in amphisbaenians, an ancient lineage of fossorial vertebrates. *Journal of Experimental Biology*, 207,
385 2433-2441.

386 Nevo, E. (1979). Adaptive convergence and divergence of subterranean mammals. *Annual Reviews*
387 *Ecology Systematics*, 10, 269-308.

388 Newman, S. J., & Jayne, B. C. (2018). Crawling without wiggling: muscular mechanisms and
389 kinematics of rectilinear locomotion in boa constrictors. *Journal of Experimental Biology*, 221,
390 jeb166199.

391 O'Reilly, J. C., Ritter, D. A., & Carrier, D. R. (1997). Hydrostatic locomotion in a limbless tetrapod.
392 *Nature*, 386, 269-272.

393 Pyron, R. A., & Burbrink, F. T. (2012). Extinction, ecological opportunity, and the origins of global
394 snake diversity. *Evolution*, 66, 163-178.

395 Pyron, R. A., and Wallach, V. (2014). Systematics of the blindsnakes (Serpentes: Scolecophidia:
396 Typhlopoidea) based on molecular and morphological evidence. *Zootaxa*, 3829, 1-081.

397 Quillin, K. J. (2000). Ontogenetic scaling of burrowing forces in the earthworm *Lumbricus terrestris*.
398 *Journal of Experimental Biology*, 203, 2757-2770.

399 Rieppel, O., Kley, N. J., & Maisano, J. (2009). Morphology of the skull of the white-nosed blindsnake,
400 *Liotyphlops albirostris* (Scolecophidia: Anomalepididae). *Journal of Morphology*, 270, 536-557.

401 Savitzky, A. H., Mori, A., Hutchinson, D. A., Saporito, R. A., Burghardt, G. M., Lillywhite, H. B., &
402 Meinwald, J. (2012). Sequestered defensive toxins in tetrapod vertebrates: principles, patterns, and
403 prospects for future studies. *Chemoecology*, 22, 141-158.

404 Schineider Facini, T., Onary, S., Palci, A., Lee, M. S. Y., Bronzati, M., & Schmaltz Hsiou A. (2020).
405 Cretaceous blind snake from Brazil fills major gap in snake evolution. *iScience*, 23, 101834.

406 Segall, M., Cornette, R., Fabre, A-C., Godoy-Diana, R., & Herrel, A. (2016). Does aquatic foraging
407 impact head shape evolution in snakes? *Proceeding of the Royal Society B*, 283, 20161645.

408 Shine, R. & Webb, J. K. (1990). Natural history of Australian typhlopoid snakes. *Journal of*
409 *Herpetology*, 24, 357-363.

410 Strong, C. R. C., Scherz, M. D. & Caldwell, M. W. (2021) Deconstructing the Gestalt: New concepts and
411 tests of homology, as exemplified by a re-conceptualization of “microstomy” in squamates. *Anat.*
412 *Rec.* DOI: 10.1002/ar.24630

413 Šumbera, R., Chitaukali, W. N., Elichová, M., Kubová J., & Burda, H. (2004). Microclimatic stability in
414 burrows of an Afrotropical solitary bathyergid rodent, the silvery mole-rat (*Heliophobius*
415 *argenteocinereus*). *Journal of Zoology*, 263, 409-416.

416 Thomas, R. & Hedges, B. S. (2007). Eleven new species of snakes of the genus *Typhlops* (Serpentes:
417 Typhlopidae) from Hispaniola and Cuba. *Zootaxa*, 1400, 1-26.

418 Uetz, P., Freed, P. & Hošek, J. (2020). The Reptile Database, <http://www.reptile-database.org>,
419 accessed 30/12/2020

420 Vanhooydonck, B., Boistel, R., Fernandez, V., & Herrel, A. (2011). Push and bite: trade-offs between
421 burrowing and biting in a burrowing skink (*Acontias percivali*). *Biological Journal of the Linnean*
422 *Society*, 101, 461-475.

423 Vidal, N., Marin, J., Morini, M., Donnelan, S., Branch, W. R., Thomas, R., Vences, M., Wynn, A.,
424 Cruaud, C., & Hedges, S. B. (2010). Blindsnake evolutionary tree reveals long history on Gondwana.
425 *Biology Letters*, 6, 558-561.

426 Vleck, D. (1979). The energy costs of burrowing by the pocket gopher *Thomomys bottae*.
427 *Physiological Zoology*, 52, 122-136.

428 Watkins, J. F., Gehlbach, F. R., & Baldrige, R. S. (1967). Ability of the blind snake, *Leptotyphlops*
429 *dulcis*, to follow pheromone trails of army ants, *Neivamyrmex nigrescens* and *N. opacithorax*. *The*
430 *Southwestern Naturalist*, 12, 455-462.

431 Watkins, J. F., Gehlbach, F. R., & Kroll, J. C. (1969). Attractant-repellent secretions of blind snakes
432 (*Leptotyphlops dulcis*) and their army ant prey (*Neivamyrmex nigrescens*). *Ecology*, 50, 1098-1102.

433 Webb, J. K., & Shine, R. (1992). To find an ant: trail-following in Australian blindsnakes (Typhlopidae).
434 *Animal Behaviour*, 43, 941-948.

435 Webb, J. K. & Shine, R. (1993). Prey-size selection, gape limitation and predator vulnerability in
 436 Australian blindsnakes (Typhlopidae). *Animal Behaviour*, 45, 1117-1126.

437 Webb, J. K., Branch, W. R. & Shine, R. (2001). Dietary habits and reproductive biology of typhloid
 438 snakes from southern Africa. *Journal of Herpetology*, 35, 558-567.

439 Webb, J. K., Shine, R., Branch, W. R. & Harlow, P. S. (2000). Life history strategies in basal snakes:
 440 reproduction and dietary habits of the African thread snake, *Leptotyphlops scutifrons* (Serpentes,
 441 Leptotyphlopidae). *Journal of Zoology*, 250, 321-327.

442 Wu, N. C., Alton, L. A., Clemente, C. J., Kearney, M. R., & White, C. R. (2015). Morphology and
 443 burrowing energetics of semi-fossorial skinks (*Liopholis* spp.). *Journal of Experimental Biology*, 218,
 444 2416-2426.

445 Young, B.A., & Morain, M. (2003). Vertical burrowing in the Saharan sand vipers (*Cerastes*). *Copeia*,
 446 2003, 131-137.

447 Zaher, H., Murphy, R. W., Arredondo, J. C., Graboski, R., Machado-Filho, P. R., Mahlow, K.,
 448 Montingelli, G. G., Bottallo Quadros, A., Orlov, N. L., Wilkinson, M., Zhang, Y-P., & Grazziotin F. G.
 449 (2019). Large-scale molecular phylogeny, morphology, divergence-time estimation, and the fossil
 450 record of advanced caenophidian snakes (Squamata: Serpentes). *PlosOne*, 14, e0216148.

451 Zheng, Y., & Wiens, J. J. (2016). Combining phylogenomic and supermatrix approaches, and a time-
 452 calibrated phylogeny for squamate reptiles (lizards and snakes) based on 52 genes and 4162 species.
 453 *Molecular Phylogenetics and Evolution*, 94, 537-547.

454 **Figure legends**

455 **Figure 1:** Top: set-up illustrating the force plate, tunnel and Perspex block during push force
456 measurements of an *Afrotyphlops angolensis* measured in Kenya. Bottom: example force trace
457 showing two pushes in a *Rhinotyphlops unitaeniatus* measured in South Africa. Note how the
458 forward-directed force (FX, red) is much greater than the lateral (FY, green) or dorso-ventral (FZ,
459 blue) forces, especially during the strongest push.

460 **Figure 2:** A) Scatterplot showing the relationship between snout-vent length and maximal push force
461 in snakes. Each point is the maximal force recorded for a given individual. On average typhlopids
462 generate higher forces for their snout-vent length than other snakes. Leptotyphlopids and
463 anomalepidids do not differ from alethinophidian snakes in the relationship between snout-vent
464 length and push force. B) Scatterplot illustrating the relationship between residual body diameter
465 and residual push force in snakes illustrating that snakes with relatively wider bodies push harder.
466 Interestingly, even for a given residual body diameter typhloid snakes still generate more force than
467 other snakes.

468 **Figure 3:** Cranial anatomy in burrowing snakes. Computed tomography (CT) scans of skulls in dorsal
469 (left) and left lateral (right) views. A) *Anilius scytale* - Aniliidae, B) *Liotyphlops albirostris* -
470 Anomalepididae, C) *Leptotyphlops nigricans* - Leptotyphlopidae, D) *Rhinotyphlops lalandei* -
471 Typhlopidae. Whereas all three scolecophidian species show robust and blunt crania in dorsal view,
472 the profile in lateral view is rather different with *Rhinotyphlops* having a more 'pointed' snout.

473 **Figure 4:** Fifth vertebra in cranial, left lateral, caudal, dorsal, and ventral views. Illustrated are
474 vertebrae for A) *Anilios unguirostris* - Typhlopidae, B) *Epictia tenella* - Leptotyphlopidae, and C)
475 *Typhlophis squamosus* - Anomalepididae. Note the robust condyle and corresponding cotyle and pre-
476 and postzygapophyses in *Anilios* compared to the other two species. CON, condyle; COT, cotyle; PA,
477 prezygapophysis; PO, postzygapophysis.

Table 1: Summary of the morphometric and force data for the species included in this study.

Genus	species	origin	N	force (N)	mass (g)	SVL (mm)	diameter (mm)
Alethinophidia							
<i>Anilius</i>	<i>scytale</i>	French Guyana	5	5.9 ± 3.3	26.3 ± 16.5	500.0 ± 93.5	8.8 ± 2.2
<i>Aparallactus</i>	<i>guentheri</i>	Kenya	1	1.6	5.1	385.0	6.0
<i>Eryx</i>	<i>colubrinus</i>	Pet trade	2	2.8 ± 1.1	77.0 ± 19.8	400.0 ± 28.3	16.0†
<i>Farancia</i>	<i>abacura</i>	Pet trade	2	11.8 ± 15.7	4.0*	600.0 ± 523.3	16.8 ± 15.0
<i>Loxocemus</i>	<i>bicolor</i>	Pet trade	2	14.1 ± 0.0	171.5 ± 2.1	745.0 ± 7.1	19.9†
<i>Oxyrhopus</i>	<i>melanogenys</i>	French Guyana	1	2.9	36.0	480.0	10.3
Typhlopidae							
<i>Afrotyphlops</i>	<i>angolensis</i>	Kenya	1	18.2	48.1	511.5	15.4
<i>Rhinotyphlops</i>	<i>lalandei</i>	South Africa	5	3.3 ± 1.6	6.5 ± 3.2	236.8 ± 78.0	6.1 ± 1.0
<i>Rhinotyphlops</i>	<i>unitaeniatus</i>	Kenya	1	8.1	12.7	415.0	6.7
Leptotyphlopidae							
<i>Leptotyphlops</i>	<i>scutifrons</i>	South Africa	6	0.4 ± 0.2	1.1 ± 0.4	140.7 ± 17.8	3.1 ± 0.5
<i>Myriopholis</i>	<i>algeriensis</i>	Pet trade	1	0.2	NA	137.2	1.2
Anomalepidae							
<i>Liotyphlops</i>	<i>beui</i>	Brazil	2	1.5 ± 0.7	2.9 ± 1.2	282.5 ± 24.8	4.0 ± 0.6

Table entries are means ± standard deviations. * Body mass data was available only for the smallest specimen. † body diameter was measured for only one of the specimens. N, number of individuals sampled; svl, snout-vent length. NA: not available.

A

B

C

D