



HAL
open science

Clarkson's Disease Episode or Secondary Systemic Capillary Leak-Syndrome

Marc Pineton de Chambrun, Jean-Michel Constantin, Alexis Mathian, Cyril Quemeneur, Victoria Lepere, Alain Combes, Charles-Edouard Luyt, Zahir Amoura

► **To cite this version:**

Marc Pineton de Chambrun, Jean-Michel Constantin, Alexis Mathian, Cyril Quemeneur, Victoria Lepere, et al.. Clarkson's Disease Episode or Secondary Systemic Capillary Leak-Syndrome. *Chest*, 2021, 159 (1), pp.441. 10.1016/j.chest.2020.07.084 . hal-03254103

HAL Id: hal-03254103

<https://hal.sorbonne-universite.fr/hal-03254103v1>

Submitted on 8 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Clarkson's Disease Episode or Secondary Systemic Capillary Leak-Syndrome



That Is the Question!

To the Editor:

We read with great interest the article by Case et al¹ titled "Systemic Capillary Leak Syndrome secondary to COVID-19." The authors described a fatal systemic capillary leak syndrome (SCLS) after severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection. They suggest SCLS could be a varied presentation of the multisystemic inflammatory syndromes (MIS) associated with the coronavirus disease 2019 (COVID-19).

We recently reported a SARS-CoV-2-induced crisis of a patient with a 7-year history of IgG Kappa Clarkson's disease,² and several articles lately highlighted the role of lung capillary leakage in the pathophysiologic condition of COVID-19.³

These data suggest that COVID-19 can induce both Clarkson's disease episodes (henceforth primary SCLS) and secondary SCLS. Several considerations must be addressed to determine the nature of the capillary-leak syndrome in the reported patient.

First, COVID-19-related MIS usually affects children or young adults⁴ while Clarkson's disease patients are typically >50 years old. Second, this case is highly typical of primary SCLS: very marked hemoconcentration (>20 g/dL), rapid clinical evolution, and 4-limb compartment syndrome. On the contrary, secondary SCLS usually exhibit incomplete capillary-leak syndrome with less obvious hematocrit level increase and evolves more chronically with unresolving anasarca. Third, although not reported by the authors, the presence of a monoclonal gammopathy is critical to differentiate primary or secondary SCLS.⁵ Fourth, inflammatory biomarkers could be useful to distinguish the two hypotheses: being very elevated in MIS but low in primary SCLS episodes (personal data). Fifth, 80% of patients with COVID-19-related MIS had a positive serologic assay,⁴ which suggests that MIS occurs in the late phase of COVID-19. However, the authors do not report how the diagnosis of SARS-CoV-2 was done. Last, therapeutic interventions (corticosteroids, IV immunoglobulins) seem to be effective in COVID-19-related MIS, although no treatment has shown its efficacy during primary SCLS severe episodes. Yet,

chronic treatment with IV immunoglobulins prevents relapse and improves survival of patients with Clarkson's disease.⁵

In conclusion, COVID-19 seems to be responsible for both primary and secondary SCLS. In our opinion, the type of capillary-leak syndrome that affected the reported patient remains unclear. Differentiating the two syndromes is crucial because it has very significant therapeutic implications.

Marc Pineton de Chambrun, MD
Jean-Michel Constantin, MD, PhD
Alexis Mathian, MD, PhD
Cyril Quemeneur, MD
Victoria Lepere, MD
Alain Combes, MD, PhD
Charles-Edouard Luyt, MD, PhD
Zahir Amoura, MD
Paris, France

AFFILIATIONS: From the Sorbonne Université, Assistance Publique-Hôpitaux de Paris, Institut de Cardiométabolisme et Nutrition (ICAN), Hôpital La Pitié-Salpêtrière, Service de Médecine Intensive-Réanimation (Drs Pineton de Chambrun, Combes, and Luyt); INSERM, UMRS_1166-ICAN, Institut de Cardiométabolisme et Nutrition (Drs Pineton de Chambrun, Combes, and Luyt); GRC 29, Assistance Publique-Hôpitaux de Paris, DMU DREAM, Hôpital La Pitié-Salpêtrière, Département d'Anesthésie et de Réanimation (Drs Constantin, Quemeneur, and Lepere); and Sorbonne Université, Assistance Publique-Hôpitaux de Paris, Hôpital de la Pitié-Salpêtrière, Service de Médecine Interne 2, Centre de Référence National Lupus et SAPL et Autres Maladies Auto-immunes et Systémiques Rares (Drs Pineton de Chambrun, Mathian, and Amoura).

FINANCIAL/NONFINANCIAL DISCLOSURES: None declared.

CORRESPONDENCE TO: Marc Pineton de Chambrun, MD, Service de Médecine Intensive Réanimation et Service de Médecine Interne 2, Hôpital La Pitié-Salpêtrière, 47-83, boulevard de l'Hôpital, 75651 Paris Cedex, France; e-mail: marc.dechambrun@gmail.com, marc.pinetondechambrun@aphp.fr

Copyright © 2020 American College of Chest Physicians. Published by Elsevier Inc. All rights reserved.

DOI: <https://doi.org/10.1016/j.chest.2020.07.084>

References

1. Case R, Ramaniuk A, Martin P, Simpson PJ, Harden C, Ataya A. Systemic capillary leak syndrome secondary to coronavirus disease 2019. *Chest*. 2020;158(6):e267-e268.
2. Pineton de Chambrun M, Cohen-Aubart F, Donker DW, et al. SARS-CoV-2 induces acute and refractory relapse of systemic capillary leak syndrome (Clarkson's Disease). *Am J Med*. 2020;133(11):e663-e664.
3. Wu MA, Fossali T, Pandolfi L, et al. COVID-19: the key role of pulmonary capillary leakage: an observational cohort study. *medRxiv*. 2020. 2020.05.17.20104877.
4. Dufort EM, Koumans EH, Chow EJ, et al. Multisystem inflammatory syndrome in children in New York State. *N Engl J Med*. 2020;283(4):347-358.
5. Pineton de Chambrun M, Gousseff M, Mauhin W, et al. Intravenous immunoglobulins improve survival in monoclonal gammopathy-associated systemic capillary-leak syndrome. *Am J Med*. 2017;130(10):1219.e19-1219.e27.