



# The first Palaeodictyoptera (Insecta) from the Carboniferous-Permian basin of Graissessac (France)

Antoine Logghe, Romain Garrouste, Jean-Sébastien Steyer, Jean-Marc Pouillon, André Nel

## ► To cite this version:

Antoine Logghe, Romain Garrouste, Jean-Sébastien Steyer, Jean-Marc Pouillon, André Nel. The first Palaeodictyoptera (Insecta) from the Carboniferous-Permian basin of Graissessac (France). *Historical Biology*, 2021, 34 (3), pp.486-492. 10.1080/08912963.2021.1929953 . hal-03257982

HAL Id: hal-03257982

<https://hal.sorbonne-universite.fr/hal-03257982>

Submitted on 11 Jun 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Historical biology

2

3 **The first Palaeodictyoptera (Insecta) from the Carboniferous-Permian basin  
4 of Graissessac (France)**

5

6 Antoine Logghe<sup>a</sup>, Romain Garrouste<sup>b</sup>, Jean-Sébastien Steyer<sup>a</sup>, Jean-Marc Pouillon<sup>c</sup> and André  
7 Nel<sup>b,\*</sup>

8

9 Running head

10 Palaeodictyopteran insect from Graissessac

11

12 <sup>a</sup>Centre de Recherches en Paléontologie de Paris, UMR 7202 – CNRS, MNHN, SU, EPHE,  
13 Muséum National d'Histoire Naturelle, 8 rue Buffon, CP38, F-75005 Paris, France. E-mails:  
14 [antoine.logghe@orange.fr](mailto:antoine.logghe@orange.fr), [jean-sebastien.steyer@mnhn.fr](mailto:jean-sebastien.steyer@mnhn.fr)

15 <sup>b</sup>Institut de Systématique, Evolution, Biodiversité (ISYEB), Muséum national d'Histoire  
16 naturelle, CNRS, Sorbonne Université, EPHE, Université des Antilles, CP 50, 57 rue Cuvier,  
17 F-75005 Paris, France. E-mails: [garroust@mnhn.fr](mailto:garroust@mnhn.fr), [anel@mnhn.fr](mailto:anel@mnhn.fr)

18 <sup>c</sup>179 Rue des Platières, 38300, Nivolas Vermelle, France. E-mail: [jmpdb@wanadoo.fr](mailto:jmpdb@wanadoo.fr)

19

20 Antoine Logghe: <https://orcid.org/0000-0002-2854-77>

21 Jean-Sébastien Steyer: <https://orcid.org/0000-0003-1835-7852>

22 Romain Garrouste: <https://orcid.org/0000-0002-0880-5781>

23 Jean-Marc Pouillon: <https://orcid.org/0000-0001-6016-4734>

24 André Nel: <https://orcid.org/0000-0002-4241-7651>

25

26 \* Corresponding author

27

28 ABSTRACT

29 A new dictyoneurid insect, *Dictyoneura goujonorum* n. sp. from the Latest Ghzelian – Asselian  
30 basin of Graissessac (Hérault, France) is described in details. It is represented by a well-  
31 preserved specimen with wings of 32-35 mm long and 13-14 mm wide and other peculiar  
32 diagnostic characters such a MP with four branches and a CuP with three branches. As all the  
33 other *Dictyoneura* species are known from the Namurian and/or the Wesphalian, *Dictyoneura*  
34 *goujonorum* n. sp. is the youngest representative of the genus. It is also the first record of the  
35 order Palaeodictyoptera from the Graissessac basin. The Carboniferous-Permian  
36 palaeodictyopterans are well-known to have lived in rather humid swamp forests. The global  
37 warming and drying of the climate during the Permian and/or the rise of potential predators  
38 may be responsible of their extinction.

39

40 KEYWORDS

41 Insecta; Palaeoptera; Carboniferous-Permian; Capitanian extinction; climatic changes; France.

42

43 **Introduction**

44 The Palaeodictyoptera are major herbivorous sucking insects known from the Late  
45 Carboniferous to the Early Permian. They were well-diversified, with at least known 50 genera  
46 in about 16 families (see Fossilworks database at <http://fossilworks.org>). However the  
47 palaeodictyopteran species are generally represented by very few specimens in comparison to  
48 other clades such as the Dictyoptera or the Archaeorthoptera (A.N. pers. obs.). Thus, any new  
49 fossil of these strange six-winged insects is welcome to improve our knowledge on their  
50 diversity and temporal range. Palaeodictyoptera are known of the swamp forests of the Late

51 Carboniferous, with a maximum of diversity in the well-known outcrop of Commentry  
52 (northern part of the French Massif Central) (Kukalová 1969a-b, 1970). They are generally  
53 associated with the warm and humid evergreen coal forests, together with some of their  
54 emblematic predators, the giant odonatopterans Meganeuridae called griffenflies. Some  
55 Palaeodictyoptera are also very large insects (e.g. *Homoioptera gigantea* Agnus, 1902 with a  
56 wing span of ca. 40 cm), but most of them were relatively small, with wing span ca. 10 cm.

57 The Latest Carboniferous – Earliest Permian Graissessac basin (Hérault, France) is a  
58 quite favorable area to find fossils of plants. Numerous geological researches, mainly linked to  
59 the exploitation of the coal, have been undertaken since the 19<sup>th</sup> century. The collect of fossil  
60 animals is decreasing since these last decades with the closure of these coal mines. The insect  
61 record is especially scarce. Thanks to the efforts of Mr and Mrs Goujon Claude and Monique,  
62 more fossils have been found recently. Here we describe in details the first Palaeodictyoptera  
63 from the Graissessac basin. It is also the first fossil insect properly described for this outcrop.  
64 It is compared with all the other dictyonurid genera. This new finding addresses our knowledge  
65 on the Carboniferous-Permian fauna of Graissessac. It is also the latest occurrence of this genus  
66 and therefore contribute to document the putative extinction of the clade before the Middle  
67 Permian.

68

## 69 **Material and methods**

70 The type specimen was collected in 2019 by Goujon and Monique Claude, in the dump issued  
71 of the open coal mine named ‘Carrière de Layrac’ (North East of the town of Graissessac). The  
72 Graissessac basin is situated in the southern part of the Massif Central, France. Its general  
73 orientation is west to east, forming a long synclinal of 30 km long and 2.5 km wide (Fig. 1)  
74 (Martín-Closas and Galtier 2005: fig. 1). Becq-Giraudon (1973) considered this basin as a  
75 channel of sedimentation flow and plant debris-flow alimenting a larger Carboniferous basin

76 located under the Permian sediments of Lodève. Seven to eight coal layers have been identified,  
77 one of them being up to six meters thick on average. They are interbedded between layers of  
78 sandstone and shale. The recent sedimentological study of these rocks allows to recognize a  
79 great diversity of depositional environments: torrential alluvial cones, fluvial environments of  
80 variable energy, flood plains and swamps in an intra-mountain sedimentary basin (Saint Martin  
81 1993; Rilliart 2013). These environments may have developed in a relatively low basin region  
82 framed by very high mountains (Becq-Giraudon and Van den Driessche 1993), forming  
83 possibly what we called here a “lost valley”. Indeed sedimentological data (exhaust craters,  
84 solifluxion flows, ice crystal ghosts) suggest the presence of periglacial phenomena and a rather  
85 cool paleoclimate for the Graissessac area (Becq-Giraudon and Van Den Driessche 1994; Becq-  
86 Giraudon et al. 1996).

87 Concerning the age of the Graissessac basin, Bruguier et al. (2003) estimated it around  
88  $295.5 \pm 5.1$  Ma, that-is-to-say Asselian (earliest Permian) based on zircon geochronology.  
89 However its plant assemblage suggests a late Gzhelian (latest Carboniferous) age (Poschmann  
90 et al. 2016). This Graissessac flora includes about 70 taxa of lycopsids, sphenopsids,  
91 pteridosperms, and prephanerograms (Grand'Eury 1877; Becq-Giraudon 1973; Doubinger,  
92 1983 Poschmann et al. 2016): these taxa are relatively well-known in the Carboniferous-  
93 Permian rocks of the other Massif Central and European basins (e.g. Doubinger et al. 1995).  
94 The presence of very large trunks of *Sigillaria* and *Calamites* suggests an environment of flood  
95 plains and swamps, also supported by Martín-Closas and Galtier (2005).

96 The animal fossil record is reduced. Mostly invertebrate have been recovered: Mollusca:  
97 *Carbonicola* sp. (Becq-Giraudon 1973); Arthropoda: xiphosuran *Euproops mariae* (Crônier  
98 and Courville, 2005), Chelicerata *Aenigmatarbus rasteli* Poschmann et al., 2016 and Scorpiones  
99 *Eoscorpius* sp., *Compsoscorpius* sp. (Poschmann et al. 2016); Insecta: Paoliida *Blattinopsis* sp.,  
100 a wing of a Polyneoptera ‘Prothortoptera’ (Becq-Giraudon 1972) and the endophytic

101 oviposition of an insect, possibly an Odonatoptera (Béthoux et al. 2004); Vertebrata: ‘fishes’  
102 have been mentioned but not described (Bergeron 1889; Louis 1954) and a branchiosaurid  
103 recently excavated is currently under description by us. Trace fossils have also been found in  
104 the Graissessac basin with the presence of trackways of the millipede-like *Arthropleura* sp.  
105 (Moreau et al. 2019). Compared with the faunas of the other Massif Central basins, this fauna  
106 is reduced in specimen number but well-diversified, with representatives of several major  
107 groups of organisms. This suggests that the biota of the Graissessac “lost valley” already  
108 presents complex trophic chains.

109 The photographs were taken using Nikon D800 camera with 60 mm 2,8 Micro Nikkor lens in  
110 the 3D Lab (ISYEB, MNHN), and the reconstruction of the venation was done under a  
111 binocular microscope Nikon SMZ 1500, with a camera lucida.

112 We follow the wing venation terminology of Kukalová-Peck (1991).

113 Institutional abbreviations.—MHNE, Muséum d’Histoire Naturelle et d’Ethnologie of Colmar,  
114 France.

115 Other abbreviations.—A anal veins; C costal vein; CuA cubitus anterior; CuP cubitus posterior;  
116 MA median anterior; MP median posterior; RA radius anterior; RP radius posterior; ScP  
117 subcostal posterior.

118 Nomenclatural acts.—This published work and the nomenclatural acts it contains, have been  
119 registered in urn:lsid:zoobank.org:pub:9220BD12-40CC-4181-8F90-798E433CDE55

120

121

## 122 Systematic palaeontology

123 Order: Palaeodictyoptera Goldenberg, 1877

124 Superfamily: Dictyoneuroidea Handlirsch, 1906a

125 Family: Dictyoneuridae Handlirsch, 1906a

- 126                   *Dictyoneura* Goldenberg, 1854
- 127        *Type species*
- 128        *Dictyoneura libelluloides* Goldenberg, 1854 (Westphalian D, Saar basin, Germany)
- 129
- 130        *Other species*
- 131        *Dictyoneura kemperi* Brauckmann and Koch, 1983 (Namurian, Hagen-Vorhalle, Germany),
- 132        *Dictyoneura nigra* Kliver, 1883 (Westphalian C, Saar basin), and *Dictyoneura goujonorum* n.
- 133        sp.
- 134
- 135                   *Dictyoneura goujonorum* n. sp.
- 136                   (Figure 2)
- 137        urn:lsid:zoobank.org:act:53F972EE-42A0-4F24-8A86-17D2A9073275
- 138        *Etymology*
- 139        Named after Goujon Claude and Monique who found the type specimen.
- 140
- 141        *Type material*
- 142        MHNE.2021.3.1 (imprint and counter-imprint of a fore- and a hind wing in life position with
- 143        fragments of abdomen and thorax), stored at the Muséum d'Histoire Naturelle et d'Ethnologie
- 144        of Colmar, France.
- 145
- 146        *Type locality*
- 147        Quarry of Layrac, Graissessac, Hérault, France (GPS coordinates available to qualified
- 148        researchers).
- 149
- 150        *Stratigraphic occurrence*

151 Latest Gzhelian to Asselian.

152

153 *Diagnosis*

154 Wings ca. 32.0-35.0 mm long, and 13.0-14.0 mm wide; ScP elongate; RP not pectinate; MA  
155 simple; MP with four branches; CuA simple; CuP with three branches; bases of CuA, MA and  
156 RP approximate.

157

158 *Description*

159 Wings without trace of coloration; forewing ca. 35.0 mm long, 13.0 mm wide; anterior margin  
160 not curved basally; costal area 1.0 mm wide; ScP at least 29.0 mm long, terminating on C well  
161 beyond midwing; RA simple; base of RP 11.0 mm from wing base; RP not pectinate, with two  
162 main branches, anterior one forked and posterior one with two-three branches; MA unbranched,  
163 weakly curved basally, separated from MP slightly basal to base of RP, 1.0 mm apart; MP with  
164 two main branches, each of them being forked again; CuA unbranched, weakly curved basally,  
165 separated from CuP slightly basal to base of MA, 2.0 mm apart; CuP with two-three branches;  
166 anal area 4.0 mm wide, anal veins poorly preserved; archaedictyon present but poorly visible.  
167 Hind wing ca. 32.0 mm long, 14.0 mm wide; triangular-shaped and with a broader anal area  
168 than in forewing; costal area as broad as that of forewing, 1.0 mm wide; venation identical to  
169 that of forewing; cubito-anal area broader than that of forewing, 6.0 mm wide.

170

171 **Discussion**

172 This fossil belongs to the Palaeodictyoptera rather than to the other orders of the  
173 Palaeodictyopterida because of the dense venation, with numerous branches of main veins, and  
174 the simple pattern of branching of RP and MA (Carpenter 1992). Riek (1976) proposed a key  
175 to palaeodictyopteran superfamilies, after which *Dictyoneura goujonorum* n. sp. would fall in

176 the Dictyoneuroidea, because of the following characters: CuA and MA simple; ScP separated  
177 from R; Archaeodictyon present. But after Riek (1976), the representatives of this superfamily  
178 would also have MP simple or three-branched, which is not the case for the type genus  
179 *Dictyoneura* and *Dictyoneura goujonorum* n. sp.

180 The current classification of the Palaeodictyoptera is not satisfactory. Sinitshenkova (2002:  
181 fig. 138) first proposed a phylogenetic hypothesis, but the absence of real outgroup(s) to  
182 basically polarize the character states, and the basal-most dichotomy established on the  
183 character ['wings wide basally' vs. 'wing base narrow'], with the two states supposedly  
184 supporting the two branches, while one should be plesiomorphic, prevent a total confidence to  
185 these results. Sroka et al (2015: fig. 11) proposed another hypothesis, better supported by true  
186 outgroups and computer treatment of the data, in which the order Palaeodictyoptera falls as a  
187 grade, sister group of the (Megasecoptera + (Permothemistidae + Diaphanopterodea)).

188 A comparison of this fossil to all the currently accepted families (as listed in the fossilworks  
189 database <http://fossilworks.org>) is necessary:

- 190 - the 'anterior margin of the wing not curved basally' excludes the Lithomanteidae  
191 Handlirsch, 1906;
- 192 - the posteriorly curved branches of M and Cu exclude the Megaptilidae Handlirsch 1906;
- 193 - the less elongated wings, with only three branches of MP instead of six, and the shorter  
194 abal area differs from *Archaemegaptillus kiefferi* Meunier, 1908 (Kukalová 1969: fig.  
195 46) and exclude the Archaemegaptilidae Handlirsch, 1919;
- 196 - the 'RP and MA not coalescent nor strongly approximate' excludes the Eugeronidae  
197 Handlirsch, 1906a;
- 198 - the elongate ScP excludes the Calvertiellidae Martynov, 1931 and the Stobbsiidae  
199 Handlirsch, 1908 (Laurentiaux and Laurentiaux-Vieira, 1951: fig. 5);
- 200 - the 'branched MP' excludes the Tchirkovaeidae Sinitshenkova, 1979;

- 201 - the ‘CuA simple’ excludes the Homoiopteridae Handlirsch, 1906a, Spilapteridae  
202 Handlirsch, 1906a, Fouqueidae Handlirsch, 1906a, Elmoboriidae Carpenter, 1976;  
203 - the ‘MA simple’ excludes the Homothetidae Scudder, 1885 based on a poorly known  
204 fossil;  
205 - the ‘CuA simple’ and the ‘MA simple’ exclude the Pteronidiidae Bolton, 1912,  
206 Mecynostomatidae Handlirsch, 1904, Straeleniellidae Laurentiaux-Vieira and  
207 Laurentiaux, 1986, and Eubleptidae Handlirsch, 1906a;  
208 - the ‘CuA simple’, and the broader and shorter wings and shorter cubito-anal area  
209 exclude the Peromapteridae Handlirsch, 1906a;  
210 - the ‘branched CuP and RP with more than three branches’ excludes the Psychroptilidae  
211 Riek, 1976 (originally considered as a Megasecoptera, but put in the Palaeodictyoptera  
212 in the fossilworks database <http://fossilworks.org>);  
213 - the ‘MA not arising in a distal position, and not on first branch of MP’ exclude’ the  
214 Lycocercidae Handlirsch, 1906a;  
215 - the ‘RP posteriorly pectinate’ excludes the Saarlandiidae Guthörl, 1930 and the  
216 Mongolianidae Özdkmen 2008 (replacement name for Mongolodictyidae  
217 Sinitshenkova, 1992), two poorly known families;  
218 - the ‘base of MA basal to that of RP’ excludes the Heolidae Handlirsch, 1906a and the  
219 Graphiptilidae Handlirsch, 1906a;  
220 Finally, The Jongmansiidae Laurentiaux, 1950 have a reduced RP. The Namuroningxiidae  
221 Prokop and Ren, 2007 have a richer venation, with bases of RP and MA at the same level, and  
222 much more branches of RP, MP, and CuP. The Polycreagridae Handlirsch, 1906b have much  
223 more branches of RP and MP and a forked CuA (Handlirsch 1906b). The Synarmogidae  
224 Handlirsch, 1910 have the median vein appressed to the radial one.

225        Affinities with the Breyeriidae Handlirsch, 1906a are more complicate to exclude. They  
226        generally have long crossveins between the main veins and their branches (viz. in *Breyeria* de  
227        Borre, 1875, *Jugobreyeria* Brauckmann et al., 1985, *Hasala* Brauckmann, 1995, *Megaptiloides*  
228        Handlirsch, 1906, and *Vermooija* Prokop et al., 2018), not present in *Dictyoneura goujonorum*  
229        n. sp. Thus this character would be sufficient to exclude affinities of *Dictyoneura goujonorum*  
230        n. sp. with the Breyeriidae. But in *Aviobreyeria* Prokop et al., 2013, these crossveins are mainly  
231        present in the distal half of the wing while there is an archedictyon in basal part of wing. The  
232        Breyeriidae can have a short ScP, ending at the level of first branch of anterior branch of RP  
233        (in *Breyeria*, *Jugobreyeria*) or an elongate one (in *Hasala*, *Aviobreyeria*, *Megaptiloides*, and  
234        *Vermooija*) (Laurentiaux and Laurentiaux-Vieira, 1951; Brauckmann et al., 1985; Brauckmann,  
235        1995; Prokop et al., 2013, 2018). *Hasala* and *Aviobreyeria* have a pectinate RP, unlike  
236        *Dictyoneura goujonorum* n. sp.

237            *Dictyoneura goujonorum* n. sp. has all the diagnostic characters of the Dictyoneuridae  
238        (as proposed by Carpenter 1992: 28). They share the absence of well-defined crossveins all  
239        over the wings (Guthörl 1934). It fits with the genus *Dictyoneura* (see diagnoses in Waterlot  
240        1934: 139 or Guthörl 1934), while it strongly differs from the other genera of this family in the  
241        combination of the following characters: RP not posteriorly pectinate, base of CuA nearly at  
242        the same level as those of RA/RP and MA/MP, base of MA slightly more basal than that of RP,  
243        CuP not simple, main veins posteriorly curved, ScP elongate, no trace of crossveins in costal  
244        area.

245            Brauckmann and Koch (1983), following Guthörl (1934), made a revision of  
246        *Dictyoneura*, with the three species *Dictyoneura libelluloides*, *Dictyoneura kemperi*, and  
247        *Dictyoneura nigra*. Waterlot (1934) included the two species *Dictyoneura rugosa* Handlirsch,  
248        1906a and *Dictyoneura sinuosa* Kliver, 1883 in this genus. Lastly *Dictyoneura higginsii*  
249        Handlirsch 1906a is a basal fourth of a wing, impossible to compare to the other species in the

250 genus. *Dictyoneura sinuosa* probably does not belong to this genus because its base of CuA and  
251 of RP are well basal to that of MA (Waterlot 1934: pl. 17, fig. 1). Also, *Dictyoneura rugosa*  
252 (based on a fragment of mid part of wing) probably does not belong to this genus because the  
253 base of CuA is well basal to that of MA (Waterlot 1934: text-fig. 43). *Dictyoneura kemperi* has  
254 five branches of MP, unlike four in the other species and *Dictyoneura goujonorum* sp. nov. But  
255 the value of this character remains uncertain because of the lack of information on the variations  
256 in the number of branches of the main veins in these insects. *Dictyoneura kemperi* has forewings  
257 ca. 43 mm long, 16 mm wide, and the hind wings ca. 39 mm long and 16 mm wide. *Dictyoneura*  
258 *libelluloides* has wings more than 55 mm long and 22 mm wide, after Waterlot (1934: 141),  
259 and/or wings more than 57 mm long and 22 mm wide, after Guthörl (1934: 52). *Dictyoneura*  
260 *nigra* has wings more than 44 mm long and 18 mm wide after Waterlot (1934: 142), and/or  
261 more than 35 mm long and 18 mm wide, after Guthörl (1934: 53). *Dictyoneura goujonorum* n.  
262 sp. is clearly a smaller insect with forewing ca. 35 mm long, 13 mm wide, and hind wing ca. 32  
263 mm long, 14 mm wide.

264

## 265 Conclusion

266 All the previously described *Dictyoneura* spp. are Pennsylvanian ('Namurian B' or  
267 'Westphalian C or D'), while *Dictyoneura goujonorum* n. sp. is latest Gzhelian to Asselian,  
268 thus the most recent known representative of the genus. The Namurian *Dictyoneura kempeli* is  
269 supposed to have lived in a moist area with rich vegetation along a coastal area in a Variscan  
270 foreland basin (Brauckmann 1988). *Dictyoneura libelluloides* was found in rocks formed in the  
271 Westphalian D swamp forests of the intramontane Saar basin (Uhl and Cleal, 2010), similarly  
272 to *Dictyoneura goujonorum* n. sp. *Dictyoneura nigra* was also found in the Westphalian C of  
273 the same area. Thus all these insects probably lived in similar environments, humid but possibly  
274 not very warm, even cool. This possible cool temperature could be linked with the proximity

275 of relatively high mountains, as suggested by Becq-Giraudon and Van den Driessche (1993)  
276 for the Graissessac basin.

277 The fossil record of the family Dictyoneuridae ranges between the ‘Namurian’ and the  
278 Artinskian (in Central Siberia and Northern China) (Sharov and Sinitshenkova 1977; Hong,  
279 1985). Thus they possibly disappeared because of the global warming which started during the  
280 Early Permian and continued after, during the whole Permian. The mean annual temperatures  
281 of the period during which they are currently recorded seem to have been under 18°C (Scotese  
282 et al. 2021: fig. 6). The floristic changes to a ‘seed plant-dominated world’, due to a supposed  
283 global drying, would have happened earlier, around the Permian-Carboniferous boundary  
284 (DiMichele et al. 2001). But it seems that the drying was not uniform during the Early Permian,  
285 much more serious in the Western part of Pangea than in North China (Yang et al. 2016). Also,  
286 the central Siberia was clearly less arid than the central Pangea during the Early to middle  
287 Permian (Fujimoto et al. 2012), possibly allowing the survival of the dictyoneurids in these  
288 regions at least till the end of the Early Permian. Nevertheless, it seems that the *Dictyoneura*  
289 spp. that were living in the humid (and possibly relatively cool) forests did not survive the  
290 earliest Permian. Another explanation for the apparent extinction of these insects could be the  
291 rise of some of their predators such as the reptiles, whose diversity increased during the  
292 Carboniferous-Permian. But, as some other palaeodictyopterans (e.g some Calvertiellidae;  
293 Béthoux et al. 2007) of the same range of body size survived till the end of the Middle Permian,  
294 this hypothesis is more unlikely.

295

## 296 Acknowledgements

297 We thank two anonymous referees for the very useful comments on the first version of the  
298 paper. We sincerely thank Claude and Monique Goujon who found the type specimen and gave  
299 it to Sylvie (borned Goujon) and Eric Monceret. We also sincerely thank Eric and Sylvie

300 Monceret for the donation of the holotype specimen to the Muséum d'Histoire Naturelle et  
301 d'Ethnologie of Colmar. This research was supported by several interdisciplinary programs  
302 called "RedPerm I" and "Redperm II", the "Actions Transversales du Muséum" (ATM) of the  
303 MNHN, The LabEx Research Program, and the "Projet Fédérateur MNHN" attributed to J-S.S.  
304 & R.G, and A.N. We thank the reviewers XXX and YYY for their constructive remarks.

305

### 306 **Disclosure statement**

307 No potential conflict of interest was reported by the authors.

308

### 309 **References**

- 310 Agnus A. 1902. Description d'un Neuroptère fossile nouveau. *Homoioptera gigantea*. Bulletin  
311 de la Société Entomologique de France 7:259–261.
- 312 Becq-Giraudon JF. 1972. Contribution à l'étude géologique du bassin houiller de Graissessac  
313 (Hérault). Unpublished Thèse de 3e cycle, Université de Paris-Sud:1–66.
- 314 Becq-Giraudon JF. 1973. Etude géologique du bassin houiller de Graissessac (Hérault).  
315 Bulletin du Bureau des Recherches Géologiques et Minières, (deuxième série), Section I 3:151–  
316 163.
- 317 Becq-Giraudon JF, Montenat C, Van Den Driessche J. 1996. Hercynian high-altitude  
318 phenomena in the French Massif Central: tectonic implications. Palaeogeography,  
319 Palaeoclimatology, Palaeoecology 122:227–241. doi.org/10.1016/0031-0182(95)00081-X
- 320 Becq-Giraudon JFVan Den Driessche J. 1994. Dépôts périglaciaires dans le Stéphano-Autunien  
321 du Massif central: témoin de l'effondrement gravitaire d'un haut plateau hercynien. Comptes  
322 Rendus de l'Académie des Sciences, Paris, (Série II) 318:675–682. pascal-  
323 francis.inist.fr/vibad/index.php?action=getRecordDetail&idt=4032519

- 324 Bergeron J. 1889. Etude géologique du massif ancien situé au Sud du Massif central. Thèse  
325 Paris. Annales de la Société Géologique de France 22:1–362.
- 326 Béthoux O, Galtier J, Nel A. 2004. Oldest evidence of insect endophytic oviposition. Palaios  
327 19:408–413. [jstor.org/stable/3515864](https://doi.org/10.1016/j.palaeo.2003.08.005)
- 328 Béthoux O, Nel A, Schneider JW, Gand G. 2007. *Lodetiella magnifica* nov. gen. and nov. sp.  
329 (Insecta: Palaeodictyoptera; Permian), an extreme situation in wing morphology of  
330 palaeopterous insects. Geobios 40:181–189. <https://doi.org/10.1016/j.geobios.2006.04.001>
- 331 Bolton H. 1912. Insect-remains from the Midland and South-Eastern coal measures. Quarterly  
332 Journal of the Geological Society 68:310–323. doi.org/10.1144/GSL.JGS.1912.068.01-04.25
- 333 Brauckmann C. 1988. Hagen-Vorhalle, a new important Namurian Insecta-bearing locality  
334 (Upper Carboniferous; FR Germany). Entomologia Generalis 14:73–79.  
335 [doi.org/10.1127/entom.gen/14/1988/73](https://doi.org/10.1127/entom.gen/14/1988/73)
- 336 Brauckmann C. 1995. Neue Insekten-Funde (Palaeodictyoptera, Breyeriidae) aus dem Ober-  
337 karbon von Osnabrück (Deutschland). Osnasbrucker Naturwissenschaften Mitteilungen, 20-21:  
338 157-166.
- 339 Brauckmann C, Koch L. 1983. Eine weitere neue Insektenart aus den Vorhalle-Schichten  
340 (Ober-Karbon, oberes Namurium B) von Hagen-Vorhalle. Dortmunder Beiträge zur  
341 Landeskunde, Naturwissenschaftliche Mitteilungen 17:3–8.
- 342 Brauckmann C, Koch L, Kemper M. 1985. Spinnentiere (Arachnida) und Insekten aus den  
343 Vorhalle-Schichten (Namurium B; Ober-Karbon) von Hagen-Vorhalle (West-Deutschland).  
344 Geologie und Paläontologie in Westfalen, Westfälisches Museum für Naturkunde 3: 1–132.
- 345 Bruguier O, Becq-Giraudon JF, Champenois M, Deloule E, Ludden J, Mangin D. 2003.  
346 Application of in situ zircon geochronology and accessory phase chemistry to constraining  
347 basin development during post-collisional extension: a case study from the French Massif  
348 Central. Chemical Geology 201:319–336. doi.org/10.1016/j.chemgeo.2003.08.005

- 349 Carpenter FM. 1976. The Lower Permian insects of Kansas: Part 12. Protorthoptera  
350 (continued), Neuroptera, additional Palaeodictyoptera & families of uncertain position. *Psyche*  
351 83:336–376. doi.org/10.1155/1976/932123
- 352 Carpenter FM. 1992. Superclass Hexapoda. In: Moore, R.C., and Kaesler, R.L. (eds). Treatise  
353 on Invertebrate Paleontology. The Geological Society of America and the University of Kansas,  
354 Boulder, Colorado, (R), Arthropoda 4, 3/4: xxii + 655 pp.
- 355 Crônier C, Courville P. 2005. New xiphosuran merostomata from the Upper Carboniferous of  
356 the Graissessac Basin (Massif Central, France). *Comptes Rendus Palevol* 4:123–133.  
357 [doi.org/10.1016/j.crpv.2004.11.002](https://doi.org/10.1016/j.crpv.2004.11.002)
- 358 de Borre PA. 1875. Notes sur des empreintes d'insectes fossiles découvertes dans les schistes  
359 houillers des environs de Mons. *Annales de la Société Entomologique de Belgique* 18:xxxix–  
360 xlvi.
- 361 DiMichele WA, Pfefferkorn HW, Gastaldo RA. 2001. Response of Late Carboniferous and  
362 Early Permian plant communities to climate change. *Annual Review of Earth and Planetary  
363 Sciences* 29:461–487. doi.org/10.1146/annurev.earth.29.1.461
- 364 Doubinger J, Feist M, Galtier J, Broutin J. 1983. Excursion dans le Paléozoïque et le  
365 Mésozoïque de l'Hérault. Montpellier: Livret Guide de la Conférence Européenne de  
366 Paléobotanique:1–24.
- 367 Doubinger, J., Vetter, P., Langiaux, J., Galtier, J., Broutin, J. 1995. La flore fossile du bassin  
368 houiller de Saint-Etienne. *Mémoires du Muséum National d'Histoire Naturelle* 164:1–357.
- 369 Fujimoto T, Otoh S, Orihashi Y, Hirata T, Yokoyama TD, Shimojo M, Kouchi Y, Obara H,  
370 Ishizaki Y, Tsukada K, Kurihara T, Nuramkhan M, Gonchigdor S. 2012. Permian Peri-glacial  
371 deposits from Central Mongolia in Central Asian orogenic belt: a possible indicator of the  
372 Capitanian cooling event. *Resource Geology* 62:408–422. doi.org/10.1111/j.1751-  
373 3928.2012.00204.x

- 374 Goldenberg F. 1854. Die fossilen Insecten der Kohlenformation von Saarbrücken.
- 375 *Palaeontographica* 4:17–38.
- 376 Goldenberg F. 1877. Die fossilen Thiere aus der Steinkohlenformation von Saarbrücken. *Fauna Saraepontana Fossilis* 2:i–iv + 1–54.
- 378 Grand'Eury FC. 1877. Mémoire sur la flore carbonifère du département de la Loire et du centre de la France. *Mémoires de l'Académie des Sciences et de l'Institut National de France* 24:1–624.
- 381 Guthörl P. 1930. Neue Insektenfunde aus dem Saarcarbon. *Neues Jahrbuch für Mineralogie, Geologie und Paläontologie Abhandlungen Abteilung B* 64:147–164.
- 383 Guthörl P. 1934. Die Arthropoden aus dem Karbon und Perm der Saar-Nahe-Pfalz-Gebiete. *Abhandlungen der Preussischen Geologischen Landesanstalt (Neue Folge)* 164:1–219.
- 385 Handlirsch A. 1904. Les insectes houillers de la Belgique. *Mémoires du Musée Royal d'Histoire Naturelle de Belgique* 3:1–20.
- 387 Handlirsch A. 1906a (-1908). Die fossilen Insekten und die Phylogenie der rezenten Formen. Ein Handbuch für Paläontologen und Zoologen. Engelmann, V.W. publ., Leipzig: 1430 pp.
- 389 Handlirsch A. 1910. Das erste fossile Insekt aus dem Oberkarbon Westfalens. *Verhandlungen der Kaiserlich-Königlichen Zoologisch-Botanischen Gesellschaft in Wien* 60:249–251.
- 391 Handlirsch A. 1906b. Revision of American Paleozoic insects. *Proceedings of the United States National Museum* 29:661–820. doi.org/10.5479/si.00963801.29-1441.661
- 393 Handlirsch A. 1919. Revision der Paläozoischen Insekten. *Denkschriften der Akademie Wissenschaften, Mathematisch-Naturwissenschaftliche Klasse* 96:511–592.
- 395 Hong Y-C. 1985. New fossil genera and species of Shanxi Formation in Xishan of Taiyuan. *Entomotaxonomia* 7:83-91. [in Chinese.]
- 397 Kliver M. 1883. Über einige neue Blattinarien, zwei *Dictyoneura* und zwei *Arthropleura*-Arten aus der Saarbrücker Steinkohlenformation. *Palaeontographica* 29:249–269.

- 399 Kukalová J. 1969a. Revisional study of the order Palaeodictyoptera in the Upper Carboniferous  
400 shales of Commentry, France. Part I. *Psyche* 76:163–215. doi.org/10.1155/1969/74019
- 401 Kukalová J. 1969b. Revisional study of the order Palaeodictyoptera in the Upper Carboniferous  
402 shales of Commentry, France. Part 2. *Psyche* 76:439–486. doi.org/10.1155/1969/20732
- 403 Kukalová J. 1970. Revisional study of the order Palaeodictyoptera in the Upper Carboniferous  
404 shales of Commentry, France. Part III. *Psyche* 77:1–44. doi.org/10.1155/1970/71691
- 405 Kukalová-Peck J. 1991. Chapter 6: Fossil history and the evolution of hexapod structures. pp.  
406 141–179. In: Naumann ID. (ed.). *The insects of Australia, a textbook for students and research*  
407 workers (2nd ed.), 1, (Melbourne University Press, Melbourne, CSIRO):542 pp.
- 408 Laurentiaux D. 1950. Les insectes houillers du Limbourg hollandais. *Mededelingen van de*  
409 *Geologische Stichling* 4:13–22.
- 410 Laurentiaux,D, Laurentiaux-Vieira F. 1951 (1952). Observations sur les Paléodictyoptères  
411 Breyeriidae. *Bulletin de la Société Géologique de France* (6) 1:585–596.
- 412 Laurentiaux-Vieira F, Laurentiaux D. 1986. Paléodictyoptère nouveau du Namurien belge.  
413 *Annales de la Société Géologique du Nord* 105:187–193.
- 414 Louis J. 1954. Tectonique des bassins houillers de la bordure orientale du Massif central. Thèse  
415 Lyon, A. Lémery ed.:173 pp.
- 416 Martín-Closas C, Galtier J. 2005. Plant taphonomy and paleoecology of Late Pennsylvanian  
417 intramontane wetlands in the Graissessac-Lodève basin (Languedoc, France). *Palaios* 20:249–  
418 265. doi.org/10.2110/palo.2003.p03-119
- 419 Martynov AV. 1931. On some remarkable Odonata from the Permian of Archangelsk District.  
420 *Izvestiya Akademii nauk SSSR* 1931:141-147.
- 421 Meunier F. 1908. Nouveaux Mégasécoptérides et un nouveau Paléodictyoptère de Commentry.  
422 *Bulletin du Museum National d'Histoire Naturelle* 14:172–175.

- 423 Moreau J-D, Gand G, Fara E, Galtier J, Aubert N, Fouché S. 2019. Trackways of *Arthropleura*  
424 from the Late Pennsylvanian of Graissessac (Hérault, southern France). Historical Biology,  
425 doi.org/10.1080/08912963.2019.1675055
- 426 Özdkmen H. 2008. Some nomenclatural changes for Blattodea and Dictyoneurida (=  
427 Palaeodictyoptera). Munis Entomology & Zoology 3:745–748.
- 428 Pfeifer LS, Soreghan GS, Pochat S, Van Den Driessche J, Thomson SN. 2018. Permian  
429 exhumation of the Montagne Noire metamorphic core complex recorded in provenance of upper  
430 Paleozoic clastic strata in the Graissessac-Lodève Basin, France. Basin Research 30:1–14.  
431 https://doi.org/10.1111/bre.12197
- 432 Poschmann M, Dunlop JA, Béthoux O, Galtier J. 2016. Carboniferous arachnids from the  
433 Graissessac Basin, Central Massif, France. Paläontologische Zeitschrift 90:33–48.  
434 /doi.org/10.1007/s12542-016-0295-7
- 435 Prokop J, Ren D. 2007. New significant fossil insects from the Upper Carboniferous of Ningxia  
436 in northern China (Palaeodictyoptera, Archaeorthoptera). European Journal of Entomology  
437 104:267–275. [doi.org/10.14411/eje.2007.041](https://doi.org/10.14411/eje.2007.041)
- 438 Prokop J, Pecharová M, Jarzembski EA, Ross AJ. 2018. New palaeodictyopterans from the  
439 Late Carboniferous of the UK (Insecta: Palaeodictyopterida). Earth and Environmental Science  
440 Transactions of the Royal Society of Edinburgh 107:99–107.  
441 doi.org/10.1017/S1755691017000408
- 442 Prokop J, Tippeltová S, Roques P, Nel A. 2013. A new genus and species of Breyeriidae and  
443 wings of immature stages from the Upper Carboniferous, Nord-Pas-de-Calais, France (Insecta:  
444 Palaeodictyoptera). Insect Systematics & Evolution 44:117–128. doi.org/10.1163/1876312X-  
445 44032098
- 446 Riek EF. 1976. Neosecoptera, a new insect suborder based on specimen discovered in the late  
447 Carboniferous of Tasmania. Alcheringa 1:227–234. doi.org/10.1080/03115517608619072

- 448 Rilliart P. 2013. Le Stéphanien du bassin de Graissessac. 28 septembre 2013, sortie dirigée par  
449 Philippe Rilliart avec la complicité de Danyck Gourdin et Bernard Guibbert:15 pp.
- 450 Saint Martin M. 1993. Evolution du bassin stéphanien de Graissessac en relation avec les  
451 tectoniques hercyniennes et tardi-hercyniennes de la Montagne Noire (Sud du Massif Central  
452 français). PhD Thesis, Université de Montpellier:89 pp.
- 453 Scotese CR, Song H-j, Mills BJW., van der Meer DG. 2021. Phanerozoic paleotemperatures:  
454 The earth's changing climate during the last 540 million years. Earth-Science Reviews 215  
455 (103503):1–47. doi.org/10.1016/j.earscirev.2021.103503
- 456 Scudder SH. 1885. 4. Classe Insecta. Insecten. Handbuch der Palaeontologie, I Abtheilung.  
457 Palaeozoologie 2:747–831.
- 458 Sharov AG, Sinitshenkova ND. 1977. New Palaeodictyoptera from the USSR. Paleontological  
459 Journal 11:44–59.
- 460 Sinitshenkova ND. 1979. A new family of the Palaeodictyoptera from the Carboniferous of  
461 Siberia. Paleontological Journal 13:192–205.
- 462 Sinitshenkova ND. 1992. Two new insect species (Insecta: Dictyoneurida = Palaeodictyoptera,  
463 Perlida = Plecoptera) from the Late Permian of South Mongolia, Trudy Sovmestnaya Sovetsko-  
464 Mongol'skaya Paleontologicheskaya Ekspeditsiya 4:98–100.
- 465 Sinitshenkova ND. 2002. Chapter 2.2.1.2.3. Superorder Dictyoneurida Handlirsch, 1906 (=  
466 Palaeodictyopteroidea). pp. 115–124. In: Rasnitsyn AP, Quicke DLJ. (eds). 2002. History of  
467 insects. Kluwer Academic Publishers, Dordrecht, Boston, London:xi + 517 pp.
- 468 Sroka P, Staniczek AH, Bechly, G. 2015. Revision of the giant pterygote insect *Bojophlebia*  
469 *prokopi* Kukalová-Peck, 1985 (Hydropalaeoptera: Bojophlebiidae) from the Carboniferous of  
470 the Czech Republic, with the first cladistic analysis of fossil palaeopterous insects. Journal of  
471 Systematic Palaeontology 13:963–982. doi.org/10.1080/14772019.2014.987958

472 Uhl D, Cleal CJ. 2010. Late Carboniferous vegetation change in lowland and intramontane  
473 basins in Germany. International Journal of Coal Geology 83:318–328.  
474 doi.org/10.1016/j.coal.2009.07.007

475 Waterlot G. 1934. Etude de la faune continentale du terrain houiller sarro-lorrain. In: Bassin  
476 houiller de la Sarre et de la Lorraine. 2. Faune fossile. Etude des Gites Minéraux de la France,  
477 Imprimerie L. Danel, Lille:1–317.

478 Yang J-h, Cawood PA, Du Y-s, Li W-q, Yan J-x. 2016. Reconstructing the Early Permian  
479 tropical climates from chemical weathering indices. Geological Society of America Bulletin  
480 128:739–751. [doi.org/10.1130/B31371.1](https://doi.org/10.1130/B31371.1)

481

482 **Figure 1.** Geological map of Carboniferous Graissessac basin (modified from Martín-Closas  
483 and Galtier 2005: fig. 1; Becq-Giraudon 1973; Saint Martin 1973; Pfeifer et al. 2018).

484

485 **Figure 2.** *Dictyoneura goujonorum* n. sp., holotype: MHNE.2021.3.1. Imprint. (A) forewing;  
486 (B) hind wing; (C) forewing under alcohol; (D) hind wing under alcohol; (E) drawing of  
487 forewing. Scale bars 10 mm.

488



