Is There an Association of Being Breastfed as an Infant and Fertility Status as an Adult?

Perrine Talla (*M.D.*)¹, Céline Faure (*Ph.D.*)², Virginie Rigourd (*M.D., Ph.D.*)^{3,4}, Sébastien Czernichow (*M.D., Ph.D.*)^{5,6}, Nathalie Sermondade (*M.D., Ph.D.*)¹, Rachel Lévy (*M.D., Ph.D.*)¹, Charlotte Dupont (*Pharm.D., Ph.D.*)¹ and ALIFERT collaborative group

Affiliations

¹Sorbonne Université, Saint Antoine Research center, INSERM équipe Lipodystrophies génétiques et acquises US938, Service de biologie de la reproduction-CECOS, AP-H, Hôpital Tenon, 4 rue de la Chine, 75020 Paris, France

²Service de biologie de la reproduction-CECOS, AP-H, Hôpital Tenon, 4 rue de la Chine, 75020 Paris, France

³Service de néonatalogie, Hôpital Necker Enfants Malades, 75015 Paris, France

⁴Banque de lait, Ile de France, Hôpital Necker Enfants Malades, 75015 Paris, France

⁵Université Paris Descartes, Paris, France

⁶APHP, Service de nutrition, Hôpital européen Georges-Pompidou, Paris, France.

ALIFERT collaborative group : **Isabelle Akinin**: Unité fonctionnelle de biologie de la reproduction, histologie – embryologie – cytogénétique, hôpital Nord, Saint-Étienne, France ; **Isabelle Cedrin-Durnerin**: Service de Médecine de la Reproduction, Hôpital Jean Verdier, APHP, Bondy, France ; **Steven Cens**, Centre d'AMP de PAU, Polyclinique de Navarre, Pau, France; **Serge Hercberg**: EREN, INSERM U557; INRA; CNAM; Université Paris 13, CRNH IdF, 93017 Bobigny, France; **Claude Uthurriague**, Centre d'AMP de PAU, Polyclinique de Navarre, Pau ; **Jean-Philippe Wolf**: Service d'Histologie-Embryologie-Biologie de la Reproduction, Hôpital Cochin, APHP, Paris, France.

Corresponding author:

Charlotte Dupont: charlotte.dupont@aphp.fr

Fax : 01.56.01.78.03 Phone : 01.56.01.78.01

Adresse : Service de biologie de la reproduction-CECOS, AP-H, Hôpital Tenon, 4 rue de la Chine, 75020 Paris, France

Word count for abstract: 227 words

Word count for text: 1611 words

Number of tables: 2

Abstract:

Background: Breastfeeding has many short-term and long-term health benefits for infants. Short-term benefits include protection against childhood infections and mortality in low income countries. The adult long-term effects usually emphasized are a reduction of excess weight and type 2 diabetes. However, there is a lack of available data on the impact of **having been** breastfed on adult fertility. Indeed, infertility **probably has** a multifactorial origin, including an environmental origin. The aim of this study was to investigate whether having been breastfed could be associated **with** unexplained infertility.

Materials and Methods: This research is an ancillary study of the case-control study ALIFERT, for which both fertile and infertile couples were recruited. Breastfeeding statuses, collected **from** childhood health records, were compared among fertile and infertile individuals. Anthropometrics parameters were also used for analysis.

Results: 65.6% of infertile women and 63.3% of fertile women were breastfed, and 69% of infertile men and 67.4% of fertile men were breastfed. There was no statistically significant difference between fertile and infertile groups. Nevertheless, infertile women who were not breastfed had a significantly higher BMI than those who were breastfed (25.8 kg/m² vs 23.2 kg/m²).

Conclusion: In our study, we did not observe any **association between having been breastfed** and fertility in adulthood. However, we observed that, in infertile women, **having not been** breastfed may **influence** weight in adulthood.

Trial registration: NCT01093378 ALIFERT. Registered: March 25, 2010.

Keywords: breastfeeding; BMI, fertility

Introduction

Several studies have reported that breastfeeding has many **short and long term** health benefits to infants. In 2001, World Health Organization (WHO) gave some worldwide recommendations on breastfeeding. Specifically, it encouraged exclusive breastfeeding until the age of 6 months¹. More evidence based on a systematic **literature** review **published** in 2007 confirmed these recommendations². These observations published by the WHO were recently confirmed in a meta-analysis published in the Lancet³. The short-term benefits highlighted were a protection against childhood infection such as diarrhoea and respiratory infections. Forty-six studies conducted in low income countries showed that breastfeeding is associated with a 68% reduction in malocclusions⁴. A reduction of the risk of death in high income countries was also highlighted⁵.

Long-term effects of breastfeeding were associated with a 13% reduction in adults becoming overweight or obese and a 35% reduction in the incidence of type 2 diabetes in adulthood³. **Having been breastfed** was also associated with increased performance in intelligence tests **during** childhood and adolescence, with a 3- to 4-point increase in intelligence quotient (IQ) points⁶.

Given the manifest health benefits of breastfeeding, we wondered about the possibility of an association between **having been breastfed** and fertility in adulthood. To our knowledge, there is no published research on this particular aspect of breastfeeding.

Few animal studies have been published **about** the influence of newborns overfeeding or underfeeding on their reproductive functions. Castellano demonstrated the influence of changes in early postnatal feeding on the timing of puberty and development of the hypothalamic kisspeptin system involved in the reproductive function⁷. These results have been confirmed by Caron's study⁸. They showed that neonatally undernourished and overnourished females

display perturbed development of neural projections from the arcuate nucleus to the preoptic region with adverse consequences on neural projection of kisspeptin and puberty onset.

These experiments underline the importance of early nutrition in the development of the reproductive system. Similarly, **having been** breastfed has largely been described as protective against several illnesses; thus, in this paper, we intended to evaluate if it may impact fertility in adulthood. In order to answer to this question, we compared the breastfeeding **status of** both fertile and infertile couples.

Materials and Methods

Couple recruitment

Data from patients recruited for the ALIFERT case-control study were analysed⁹. The purpose of the ALIFERT study was to **evaluate** the link between unexplained infertility and the patient's nutritional behaviour. Unexplained infertility is defined by a lack of diagnosis for couples that have failed to conceive after one years of unprotected sexual intercourse. Standard investigation protocol of unexplained infertility is somewhat limited and mainly involves **normal** ovulation and tubal **assessment** for women, **as well as normal** semen analysis for men.

Couples were recruited from September 2009 to December 2013 from 4 **fertility** centres in France (Jean Verdier Hospital in Bondy, Cochin Hospital in Paris, North Hospital in Saint Etienne, and Polyclinic Navarre in Pau).

The inclusion criteria for the infertile groups were: individuals who had experienced more than 12 months of unexplained infertility; **female or male ages** between 18 **and** 38, **or** 18 **and** 45 **years old, respectively**; and individuals being in possession of childhood health records.

The fertile couples were healthy volunteers recruited nearby these hospitals. The inclusion criteria for the fertile group were: **female or male age** between 18 **and** 38, **or** 18 to 45 **years old, respectively**; individuals who were the biological parent of a child under 2 years of age, spontaneously conceived in less than 12 months, and in possession of their childhood health records.

Data collection

To avoid reporting bias, only childhood health records completed by doctors were accepted as viable records of a participant's breastfeeding status. Each participant had his or her weight and height measured by the same trained investigator, using the same calibrated devices. The body mass index (BMI) of each participant was calculated as the weight in kilograms divided by the square of height in metres.

All participants gave their written informed consent. The ALIFERT study was approved by an ethics committee. (National biomedical research ID no. P071224; ethics committee approval ('Comité de Protection des Personnes') ID no. AOM 2009-A00256–51; NEudra CT ID no. 08180; clinicaltrials.gov ID no. NCT01093378).

Statistical analysis

Data was summarized using means and standard deviations. Statistical differences were analysed using unpaired Student's *t* test for the quantitative data; and the Chi 2 test for the qualitative data.

$P < 0.05$ was considered significant.

Results

Ninety-three infertile women, 98 fertile women, 87 infertile men and 95 fertile men were included in this ancillary study. All participants were born at term (gestational age was between 37 and 41 weeks of amenorrhea).

Age, BMI and breastfeeding status of the participants are presented in Table 1. Fertile and infertile men had comparable ages whereas fertile women were slightly older than the infertile women. The BMI of infertile men and women was significantly higher compared to fertile participants.

65.6% of infertile women and 63.3% of fertile women had been breastfed and 69% of infertile men and 67.4% of fertile men had been breastfed. The difference was not statistically significant between groups (respectively: $p=0.764$ and $p=0.874$).

We examined more specifically the BMI according to the different groups. We observed that infertile women who had not been breastfed had a significantly higher BMI than those who had been breastfed ($25.8 \text{ kg/m}^2 \pm 5.55$ vs $23.2 \text{ kg/m}^2 \pm 4.13$, $p=0.018$). We did not observe such differences in the other groups (Table 2).

Discussion

We did not observe any association between unexplained infertility in adulthood and **having been breastfed** in the neonate period, neither for women or men. Nevertheless, interestingly, among the infertile group, we noted that the non-breastfed women had significantly higher BMI than those who had been breastfed, with **a shift toward** the overweight BMI category.

Breastfeeding has many beneficial and protective effects on the short- and long-term health of individuals, as evidenced by the recommendations of the WHO on breastfeeding^{3,6,10}. Several studies have shown that long-term health programming mechanisms are established during the prenatal and first years of life¹¹. This concept is well-known as the "first 1000 days" of life

(including gestation and the first two years of life), a period of vulnerability in human development^{12,13}.

In the ALIFERT cohort, we had previously reported that an increased birth weight was a risk factor for unexplained infertility **both** in men¹⁴ and women¹⁵, **suggesting a** link between prenatal period and fertility in adulthood. In the present study, we aimed to examine the potential impact of early postnatal period on fertility **at** adulthood. **An association** between **having been** breastfed and fertility was not highlighted; but we observed a link between **having been** breastfed and **female weight in the infertile subgroup**.

Studies have indicated that nutrient imbalance in early life influences the risk of obesity later in life^{16,17}, suggesting that obesity may result from “developmental programming”. Breastfed newborns have a better regulation of the amount of milk ingested¹⁸ and they are significantly lighter at 9 months of age¹⁹. Some reports highlighted an association between **having been** breastfed and **a relative** protection against obesity later in life. Higher plasma-insulin concentrations in bottle-fed compared to breast-fed infants could stimulate fat deposition and lead to **an** early development of adipocytes²⁰. Although the origin of obesity is complex and multifactorial, rapid weight gain in early childhood has been clearly identified as a risk factor for the development of subsequent obesity and metabolic dysfunction²¹. Thus, breastfeeding is known to have a protective effect on the early rebound of adiposity in children, which is known to have deleterious effects on the onset of puberty and increases the risk of long-term obesity^{22,23}. Pubertal timing is an indicative marker for the neuroendocrine system, which regulates the development of reproductive system. A recent large-scale study showed that early pubertal timing was associated with a lower sperm concentration and negatively associated with estrogen levels²⁴. Both testicular somatic cells and germ cells are sources of estrogen in mammals. Exposure of testis to extra-estrogen contributes to lower sperm concentration²⁵.

178 Pubertal timing can therefore **be** used as an indicative marker for hormone levels in adult men,
179 and consequently for their fertility.

180 **Overweight and** obesity are **known** risk factors of infertility in both men ²⁶ and women ²⁷. We
181 assume that the lack of protective breastfeeding in early life, combined with an unhealthy
182 lifestyle in adulthood, could lead to obesity and therefore, by extension, **could** contribute to
183 infertility. An accumulation of risk factors could be envisaged, reinforcing our hypothesis that
184 unexplained infertility origin is multifactorial and may have a developmental origin (pre- and
185 post-natal). Studies have shown that infertile individuals are in poorer health than fertile
186 individuals ^{28,29}, and would **have been** more susceptible to unfavourable foetal or neonatal
187 programming. These hypotheses underline a possible indirect impact of **having been** breastfed
188 on the reproductive functions in adulthood

189 A direct effect of **having been** breastfed on fertility in adulthood may also be considered. Thus,
190 leptin is present in breast milk³⁰ and plays a critical role in the long-term protection against
191 obesity and metabolic disorders³¹. Leptin is also an essential factor for brain development and
192 neural projection³². A lack of leptin intake during the neonatal period could have consequences
193 on the development of the reproductive axis^{7,8,33} and, therefore, have consequences on fertility
194 in adulthood. Another theory is the potential role of epigenetics mechanisms through early
195 postnatal nutrition in the developmental programming. **Leptin may play a critical role** in the
196 DNA methylation patterns **establishment** and the response to dietary conditions in later life³⁴.
197 Furthermore, miRNAs **are** present in high concentration in breast milk ³⁵ and **could** influence
198 individual development.

199 The strengths of the study are the recruitment of two comparable groups of fertile and infertile
200 couples. The assessments on breastfeeding **status** were **registered** from health book completed
201 by a medical staff, **in order to** limit bias due to declarative information. However, our study

had some limitations, such as the lack of information concerning the duration of breastfeeding and the type of breastfeeding (exclusive or not). In some meta-analyses, the duration of exclusive breastfeeding is a protective factor for obesity in adulthood^{36,37}. The duration of breastfeeding may also have an impact on the age at which adiposity rebound occurs^{38,39}. We recognise that the limited sample size of the groups may decrease the **accuracy** of our study. Consequently, further studies are needed for meaningful conclusions; in particular, studies which take into account breastfeeding characteristics such as its duration and the type of breastfeeding (exclusive or not). It should be noted that animal studies could be useful in obtaining quick answers.

In conclusion, in our study, we did not observe **any association between having been breastfed** and fertility in adulthood. **However**, an association **was observed** between **having not been breastfed** and a high BMI **in the subgroup of infertile women, suggesting that not being breastfed could constitute** a factor contributing to the onset of infertility. Nevertheless, infertility may be multifactorial. **Although** further studies are needed to fully understand this phenomenon, breastfeeding should continue to be encouraged.

Acknowledgments:

The authors want to acknowledge all the participants involved in the study.

The authors would like to thank Clinical Research Unit Paris-centre coordinator Christelle Auger and CRA Deborah Rechard.

References

1. World Health Assembly 54. Global strategy for infant and young child feeding: the optimal duration of exclusive breastfeeding. *Annex: Expert consultation on the optimal duration of exclusive breastfeeding: conclusions and recommendations (Geneva, 28 to 30 March 2001)*. Published online 2001. Accessed December 6, 2019. <https://apps.who.int/iris/handle/10665/78801>
2. Horta BL, World Health Organization, Department of Child and Adolescent Health and Development. *Evidence on the Long-Term Effects of Breastfeeding*. WHO; 2007.
3. Victora CG, Bahl R, Barros AJD, et al. Breastfeeding in the 21st century:

- epidemiology, mechanisms, and lifelong effect. *The Lancet*. 2016;387(10017):475-490. doi:10.1016/S0140-6736(15)01024-7
4. Peres KG, Cascaes AM, Nascimento GG, Victora CG. Effect of breastfeeding on malocclusions: a systematic review and meta-analysis. *Acta Paediatr*. 2015;104(467):54-61. doi:10.1111/apa.13103
5. Ip S, Chung M, Raman G, et al. Breastfeeding and maternal and infant health outcomes in developed countries. *Evid Rep Technol Assess (Full Rep)*. 2007;(153):1-186.
6. Horta BL, Loret de Mola C, Victora CG. Breastfeeding and intelligence: a systematic review and meta-analysis. *Acta Paediatr*. 2015;104(467):14-19. doi:10.1111/apa.13139
7. Castellano JM, Bentsen AH, Sánchez-Garrido MA, et al. Early Metabolic Programming of Puberty Onset: Impact of Changes in Postnatal Feeding and Rearing Conditions on the Timing of Puberty and Development of the Hypothalamic Kisspeptin System. *Endocrinology*. 2011;152(9):3396-3408. doi:10.1210/en.2010-1415
8. Caron E, Ciofi P, Prevot V, Bouret SG. Alteration in neonatal nutrition causes perturbations in hypothalamic neural circuits controlling reproductive function. *J Neurosci*. 2012;32(33):11486-11494. doi:10.1523/JNEUROSCI.6074-11.2012
9. Foucaut A-M, Faure C, Julia C, et al. Sedentary behavior, physical inactivity and body composition in relation to idiopathic infertility among men and women. *PLOS ONE*. 2019;14(4):e0210770. doi:10.1371/journal.pone.0210770
10. Binns C, Lee M, Low WY. The Long-Term Public Health Benefits of Breastfeeding. *Asia Pac J Public Health*. 2016;28(1):7-14. doi:10.1177/1010539515624964
11. Barker DJP, Eriksson JG, Forsén T, Osmond C. Fetal origins of adult disease: strength of effects and biological basis. *Int J Epidemiol*. 2002;31(6):1235-1239. doi:10.1093/ije/31.6.1235
12. Why 1,000 Days. 1,000 Days. Accessed December 9, 2019. <https://thousanddays.org/why-1000-days/>
13. Victora CG, Adair L, Fall C, et al. Maternal and child undernutrition: consequences for adult health and human capital. *The Lancet*. 2008;371(9609):340-357. doi:10.1016/S0140-6736(07)61692-4
14. Faure C, Dupont C, Chavatte-Palmer P, Gautier B, Levy R. Are semen parameters related to birth weight? *Fertility and Sterility*. 2015;103(1):6-10. doi:10.1016/j.fertnstert.2014.11.027
15. Dupont C, Hulot A, Jaffrezic F, et al. Female ponderal index at birth and idiopathic infertility. *J Dev Orig Health Dis*. Published online July 16, 2019:1-5. doi:10.1017/S2040174419000394
16. Bouret S, Levin BE, Ozanne SE. Gene-environment interactions controlling energy and glucose homeostasis and the developmental origins of obesity. *Physiol Rev*. 2015;95(1):47-82. doi:10.1152/physrev.00007.2014
17. Lukaszewski M-A, Eberlé D, Vieau D, Breton C. Nutritional manipulations in the perinatal period program adipose tissue in offspring. *Am J Physiol Endocrinol Metab*. 2013;305(10):E1195-1207. doi:10.1152/ajpendo.00231.2013
18. Heinig MJ, Nommsen LA, Peerson JM, Lonnerdal B, Dewey KG. Intake and growth of breast-fed and formula-fed infants in relation to the timing of introduction of complementary foods: the DARLING study. Davis Area Research on Lactation, Infant Nutrition and Growth. *Acta Paediatr*. 1993;82(12):999-1006. doi:10.1111/j.1651-2227.1993.tb12798.x
19. Michaelsen KF, Petersen S, Greisen G, Thomsen BL. Weight, length, head circumference, and growth velocity in a longitudinal study of Danish infants. *Dan Med Bull*. 1994;41(5):577-585.
20. Lucas A, Blackburn AM, Aynsley-Green A, Sarson DL, Adrian TE, Bloom SR.

- BREAST vs BOTTLE: ENDOCRINE RESPONSES ARE DIFFERENT WITH FORMULA FEEDING. *The Lancet*. 1980;315(8181):1267-1269. doi:10.1016/S0140-6736(80)91731-6
21. Sacco MR, de Castro NP, Euclides VLV, Souza JM, Rondó PHC. Birth weight, rapid weight gain in infancy and markers of overweight and obesity in childhood. *Eur J Clin Nutr*. 2013;67(11):1147-1153. doi:10.1038/ejcn.2013.183
 22. Pietrobelli A, Agosti M, MeNu Group. Nutrition in the First 1000 Days: Ten Practices to Minimize Obesity Emerging from Published Science. *Int J Environ Res Public Health*. 2017;14(12). doi:10.3390/ijerph14121491
 23. Kang MJ. The adiposity rebound in the 21st century children: meaning for what? *Korean J Pediatr*. 2018;61(12):375-380. doi:10.3345/kjp.2018.07227
 24. Wang X, Zou P, Mo M, et al. Early pubertal timing is associated with lower sperm concentration in college students. *Oncotarget*. 2018;9(36):24178-24186. doi:10.18632/oncotarget.24415
 25. Handelsman DJ. Estrogens and falling sperm counts. *Reprod Fertil Dev*. 2001;13(4):317-324. doi:10.1071/rd00103
 26. Bellastella G, Menafrà D, Puliani G, Colao A, Savastano S. How much does obesity affect the male reproductive function? *Int J Obes Suppl*. 2019;9(1):50-64. doi:10.1038/s41367-019-0008-2
 27. Talmor A, Dunphy B. Female Obesity and Infertility. *Best Practice & Research Clinical Obstetrics & Gynaecology*. 2015;29(4):498-506. doi:10.1016/j.bpobgyn.2014.10.014
 28. Dupont C, Faure C, Daoud F, et al. Metabolic syndrome and smoking are independent risk factors of male idiopathic infertility. *Basic Clin Androl*. 2019;29:9. doi:10.1186/s12610-019-0090-x
 29. Broughton DE, Moley KH. Obesity and female infertility: potential mediators of obesity's impact. *Fertil Steril*. 2017;107(4):840-847. doi:10.1016/j.fertnstert.2017.01.017
 30. Miralles O, Sánchez J, Palou A, Picó C. A Physiological Role of Breast Milk Leptin in Body Weight Control in Developing Infants*. *Obesity (Silver Spring, Md)*. 2006;14:1371-1377. doi:10.1038/oby.2006.155
 31. Palou M, Picó C, Palou A. Leptin as a breast milk component for the prevention of obesity. *Nutr Rev*. 2018;76(12):875-892. doi:10.1093/nutrit/nuy046
 32. Stepan CM, Swick AG. A Role for Leptin in Brain Development. *Biochemical and Biophysical Research Communications*. 1999;256(3):600-602. doi:10.1006/bbrc.1999.0382
 33. Bouret SG, Simerly RB. Developmental programming of hypothalamic feeding circuits. *Clinical Genetics*. 2006;70(4):295-301. doi:10.1111/j.1399-0004.2006.00684.x
 34. Palou M, Picó C, McKay JA, et al. Protective effects of leptin during the suckling period against later obesity may be associated with changes in promoter methylation of the hypothalamic pro-opiomelanocortin gene. *Br J Nutr*. 2011;106(5):769-778. doi:10.1017/S0007114511000973
 35. Munch EM, Harris RA, Mohammad M, et al. Transcriptome Profiling of microRNA by Next-Gen Deep Sequencing Reveals Known and Novel miRNA Species in the Lipid Fraction of Human Breast Milk. *PLOS ONE*. 2013;8(2):e50564. doi:10.1371/journal.pone.0050564
 36. Horta BL, Bahl R, Martinés JC, Victora CG, Organization WH. *Evidence on the Long-Term Effects of Breastfeeding : Systematic Review and Meta-Analyses*. World Health Organization; 2007. Accessed February 27, 2020. <https://apps.who.int/iris/handle/10665/43623>
 37. Arenz S, Rückerl R, Koletzko B, Kries R von. Breast-feeding and childhood obesity—a systematic review. *International Journal of Obesity*. 2004;28(10):1247-1256. doi:10.1038/sj.ijo.0802758
 38. Chivers P, Hands B, Parker H, et al. Body mass index, adiposity rebound and early

feeding in a longitudinal cohort (Raine Study). *Int J Obes (Lond)*. 2010;34(7):1169-1176.
doi:10.1038/ijo.2010.61

39. Wu YY, Lye S, Briollais L. The role of early life growth development, the FTO gene and exclusive breastfeeding on child BMI trajectories. *Int J Epidemiol*. 2017;46(5):1512-1522. doi:10.1093/ije/dyx081

Conflicts of Interest

There are no potential conflicts of interest relevant to this article.

Funding

Disclosure statement: Financial support: this study was supported by French national biomedical research P071224 ALIFERT.

Availability of data and materials

Data are the property of the Public Assistance – Paris Hospitals [Assistance Publique – Hôpitaux de Paris (AP-HP)] that does not authorize as a promoter the sharing of data without a contract. Consultation by the editorial board or interested researchers may nevertheless be considered.

Authors' contributions

CD, CF participated in the study conception and design, in patient's recruitment, data acquisition, interpretation and analysis, and drafting of the manuscript. PT participated in study design, performed statistical analysis and participated in drafting of the manuscript. SC and NS participated in study conception and design, in patient recruitment and critical revisions of the manuscript for intellectual content. RL participated in study conception and design, interpretation of data, critical revision of the manuscript for intellectual content and supervised

the study. The collaborators of the ALIFERT collaborative group participated in study design and were involved in patients' recruitment. All authors read and approved the final manuscript.

Ethics approval and consent to participate

The ethics committee ("Comité de Protection des Personnes") approved the study. ALIFERT study (national biomedical research P071224/AOM 08180:NEudra CT 2009-A00256–51/clinical trials NCT01093378). All the participants signed a written informed consent. CPP Ile de France, Numéro de dossier: 2012-nov-13076.

	Infertile Women	Fertile Women	Infertile Men	Fertile Men
Total	93	98	87	95
Age	<i>30.9+/-4.25*</i>	<i>32.1+/-3.18*</i>	33.1+/-5.19	34.3+/-3.85
BMI (kg/m ²)	<i>24.1+/-4.77**</i>	<i>21.9+/-3.02**</i>	<i>26.3+/-4.17**</i>	<i>23.9+/-2.64**</i>
Breastfeeding (%)	61 (65.6%)	62 (63.3%)	60 (68.9%)	64(67.3%)

373 *p=0.02, **p<0.001

374 Table1: Age, BMI and breastfeeding status of the fertile and infertile women and men. Data are means ± standard deviations. Significant
375 differences are written in bold italic.

376

377

	Infertile Women		Fertile Women		Infertile Men		Fertile Men	
	Breastfed	Not breastfed	Breastfed	Not breastfed	Breastfed	Not breastfed	Breastfed	Not breastfed
Total	61	32	62	36	60	27	64	31
Weight (kg)	<i>63.4 +/- 11.8*</i>	<i>71 +/- 16*</i>	60.2 +/- 7.8	58.8 +/- 6.8	83.5 +/- 15.4	83.9 +/- 17.2	75.8 +/- 9.3	77.1 +/- 11.5
Height (m)	1.65 +/- 0.06	1.66 +/- 0.05	1.66 +/- 0.04	1.65 +/- 0.05	1.78 +/- 0.07	1.79 +/- 0.06	1.79 +/- 0.06	1.77 +/- 0.07
BMI (kg/m ²)	<i>23.2 +/- 4.1**</i>	<i>25.8 +/- 5.5**</i>	21.8 +/- 2.7	22 +/- 3.6	26.4 +/- 3.9	26.2 +/- 4.7	23.6 +/- 2.4	24.6 +/- 2.9

378 *p=0.012, **p=0.018

379 Table 2: BMI of infertile and fertile women and men. Data are means ± standard deviations. Significant differences are written in bold italic.

380