Grazoprevir/elbasvir for the immediate treatment of recently acquired HCV genotype 1 or 4 infection in MSM

Anders Boyd, Patrick Miallhes, Julie Chas, Marc-Antoine Valantin, Yazdan Yazdanpanah, Eric Rosenthal, Stephane Chevaliez, Lionel Piroth, Hayette Rougier, Gilles Peytavin, et al.

► To cite this version:

Anders Boyd, Patrick Miallhes, Julie Chas, Marc-Antoine Valantin, Yazdan Yazdanpanah, et al.. Grazoprevir/elbasvir for the immediate treatment of recently acquired HCV genotype 1 or 4 infection in MSM. *Journal of Antimicrobial Chemotherapy*, 2020, 75 (7), pp.1961-1968. 10.1093/jac/dkaa091 . hal-03272038

HAL Id: hal-03272038

<https://hal.sorbonne-universite.fr/hal-03272038>

Submitted on 28 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Grazoprevir/elbasvir for the immediate treatment of recently acquired hepatitis C virus genotype 1 or 4 infection in men having sex with men

Anders BOYD¹, Patrick MIALHES², Julie CHAS³, Marc-Antoine VALANTIN⁴, Yazdan YAZDANPANA⁵,
Eric ROSENTHAL⁶, Stephane CHEVALIEZ⁷, Lionel PIROTH⁸, Hayette ROUGIER⁹, Gilles PEYTAVIN¹⁰, Gilles
PIALOUX^{3,11}, Pierre-Marie GIRARD^{12,13}, Karine LACOMBE*^{12,13}

Institutional affiliations:

¹INSERM, Institut Pierre Louis d'Epidémiologie et de Santé Publique, F-75012, Paris, France;

²Hôpital de la Croix-Rousse, Hospices Civils de Lyon, Service de Maladies Infectieuses et Tropicales,
Lyon, France

³AP-HP, Hôpital Tenon, Service de Maladies Infectieuses et tropicales, Paris, France

⁴AP-HP, Groupe Hospitalier Pitié-Salpêtrière, Service de Maladies Infectieuses et tropicales, Paris,
France

⁵AP-HP, Hôpital Bichat-Claude Bernard, Service de Maladies Infectieuses et tropicales, Paris, France

⁶Hôpital de l'Archet, Service de médecine interne, Nice, France

⁷AP-HP, département de Virologie, Hôpital Henri Mondor; National Reference Center for Viral
Hepatitis B, C and delta; INSERM U955, Créteil, France

⁸Département d'infectiologie, CHU de Dijon, 21079 Dijon, France; Inserm CIC 1432, Université de
Bourgogne, 21079 Dijon, France

⁹Institut de Médecine et d'Epidémiologie Appliquée, Paris

¹⁰AP-HP, Hôpital Bichat-Claude Bernard, Laboratoire de Pharmacologie-Toxicologie and IAME, UMR
1137, Université Paris Diderot, Sorbonne Paris Cité and INSERM, Paris, France

¹¹Sorbonne Université, Paris, France

¹²Service de maladies infectieuses et tropicales Hôpital St Antoine, AP-HP, Paris, France

¹³Sorbonne Université, Inserm UMR-S1136, Institut Pierre Louis de Santé Publique, Paris, France

27

28 **Corresponding Author:**

29 Prof. Karine Lacombe, MD, PhD

30 Service de maladies infectieuses et tropicales

31 Hôpital Saint-Antoine

32 184 rue du Faubourg Saint-Antoine

33 75012 Paris – FRANCE

34 Tel: +33 1 49 28 24 38

35 Fax: +33 1 49 28 21 49

36 Email: karine.lacombe2@aphp.fr

37

38 **Running Title:** GZR/ EBR for acute HCV infection

SYNOPSIS

Background: Men having sex with men (MSM) belong to the most at-high risk populations that fuel the epidemic of recently acquired hepatitis C in Europe. Alternative associations of direct antiviral agents against hepatitis C virus (HCV) need to be assessed.

Patients and methods. In this pilot trial, MSM with recently acquired genotype 1-4 HCV infection were prospectively included and received 8 weeks of oral grazoprevir 100 mg and elbasvir 50 mg in a fixed-dose combination administered once daily. The primary endpoint was sustained virological response evaluated 12 weeks after end of treatment (SVR12). Secondary endpoints were the virological characterization of failures, the quality of life before, during and after treatment and the rate of reinfection.

Results. In a 15 months period 30 patients were enrolled, all of whom were men who have sex with men. Of the 29 patients completing follow-up, 28 (96%, 95%CI=82-99%) achieved SVR12. One patient interrupted follow-up (suicide) but had undetectable plasma HCV-RNA at the end of treatment. One patient with suboptimal adherence confirmed by plasma drug monitoring relapsed and developed NS3, NS5A and NS5B RASs at positions V36M, M28V, and S556G. The most common adverse events related to study drug were diarrhoea ($n=4$, 13%), insomnia ($n=2$, 7%), and fatigue ($n=2$, 7%), although no patient discontinued treatment. No HIV-RNA breakthrough was reported in the 28 patients with HIV co-infection. At week 48, reinfection was diagnosed in 3 patients.

Conclusions. Our data support the use of grazoprevir/elbasvir for immediate treatment against HCV in order to reduce HCV transmission in MSM.

INTRODUCTION

With the use of direct-acting antivirals (DAAs), clearance of hepatitis C virus (HCV) can be achieved with short-course treatment for all HCV-infected individuals.¹ This success has prompted the World Health Organization to target HCV elimination by 2030.²

In Western European countries, outbreaks of recently acquired HCV among HIV-positive MSM have begun since 1996³ and HCV incidence has failed to decline over the past decade.⁴ With the increased use of pre-exposure prophylaxis against HIV, clustering of recently acquired HCV infections has also been observed between HIV-positive and HIV-negative MSM.⁵ Although most settings delay treatment for 6 months in order to assess spontaneous HCV clearance, HIV-positive MSM with recently acquired HCV infection could still be engaging in at risk activity and continue to transmit HCV.⁶ Short-course DAA treatment during recently acquired HCV infection could therefore be beneficial to reduce HCV transmission. Modelling studies have indeed suggested that treating HCV early-on is an essential part of the interventions needed to curb HCV incidence and reach elimination.⁷⁻⁹

Few DAA combinations exploring diverse durations of treatment, mostly involving the NS5B inhibitor sofosbuvir, have been studied for individuals with acute HCV infection. In two one-arm, open label trials, sustained virological response 12 weeks after treatment (SVR12) was achieved with 6- or 8-week sofosbuvir/ledipasvir in 77% and 100% of HCV genotype 1-4 HIV-positive individuals.^{10, 11} In the SWIFT-C trial, a 12-week sofosbuvir/ribavirin regimen led to 59% with SVR12, which is clearly sub-optimal.¹² An 8-week course of the non-sofosbuvir based DAA regimen, paritaprevir/ritonavir/ombitasvir/dasabuvir±ribavirin, was examined in acute HCV genotype 1 infection, resulting in 97% with SVR12.¹³ These trials were limited in that patients with acute or early chronic HCV infection were combined.

Grazoprevir and elbasvir are potent NS3/4 and NS5A inhibitors for the treatment of chronic HCV genotype 1/4 infection and have resulted in high proportions of SVR12.^{14, 15} SVR12 of 96%-99% after 12-week grazoprevir/elbasvir have also been observed among HIV and chronic HCV co-infected patients harbouring HCV genotypes 1, 4 and 6 in both clinical trials^{16, 17} and real-life settings.¹⁸ Only one trial has evaluated the efficacy of this combination in individuals recently infected with hepatitis C genotype 1 or 4 in the Netherlands and after a shorter 8-week duration of treatment, reported an SVR12 of 99%.¹⁹ Further confirmation of these results are needed. We present herein the results of a clinical pilot trial conducted in France that included MSM with a recently acquired HCV genotype 1 or 4 infection undergoing eight-week grazoprevir/elbasvir.

PATIENTS AND METHODS

Study design

The SAHIV study is an open-label, single-arm, interventional, pilot trial evaluating the efficacy and safety of 8-week grazoprevir 100 mg and elbasvir 50 mg for the treatment of recently acquired genotype 1 or 4 HCV infection. Recruitment of study participants occurred at 6 university hospitals in France (Paris – Hôpital Saint-Antoine, Hôpital La Pitié-Salpêtrière, Hôpital Bichat, Hôpital Tenon; Lyon – Hôpital de la Croix-Rousse, Lyon; Nice – CHU de Nice). The protocol was approved in accordance with the Helsinki Declaration. The trial was registered at clinicaltrials.gov (identifier: NCT02886624).

Participants

Inclusion criteria were: age ≥ 18 years, recently acquired HCV genotype 1 or 4 infection (defined in the Supplementary methods), plasma HCV-RNA ≥ 1000 IU/mL, body weight 40-125 kg, and health

coverage under the French national health system. If HIV-positive, infection had to be confirmed with a positive western blot. HIV-positive individuals were allowed to either follow an ART combination not contraindicated with grazoprevir and elbasvir or remain untreated with ART while taking the study drug. All patients signed written informed consent prior to participating in the trial.

Non-inclusion criteria were: active or recent (<6 months) opportunistic infection; primary HIV infection; hepatitis B surface antigen-positive without treatment containing a third-generation nucleoside/nucleotide analogue [tenofovir, tenofovir alafenamide, entecavir] within 2 weeks of participating; confirmed cirrhosis prior to acute HCV diagnosis; any other known causes of acute hepatitis; pregnant or breast-feeding; liver transplant recipient; progressive malignancy; history of non-adherence; participation in another clinical trial with experimental treatment; under legal guardianship or incarcerated; haemoglobin <10 g/dL (female) or <11 g/dL (male); platelet count <50,000/mm³; neutrophil count < 750/mm³; or contraindication to grazoprevir and/or elbasvir or to any of the excipients.

Study procedures

Participants received an oral, fixed-dose, once-daily combination of grazoprevir (100 mg) and elbasvir (50 mg) for eight weeks. HCV genotype and HCV-RNA viral load were obtained either at screening visit or from a previous biological test performed 1-4 weeks before treatment initiation. RAS testing was not performed prior to treatment initiation.

After treatment initiation, patients were followed at week 4 (W4) and week 8 (W8) and post-treatment at week 4 (PTW4) and week 12 (PTW12). Blood samples were drawn at each visit to assess ALT, AST, total and conjugated bilirubin, γ -glutamyl transferase, alkaline phosphatase, full blood and platelet count, and creatinine levels. HCV-RNA viral load was measured using a commercially-

available PCR assay at the study centre (TaqMan 2.0 Assay Roche diagnostic or Abbott Realtime; lower limit of detection: 15 IU/mL and 12 IU/mL, respectively). Genotyping was performed using Versant® HCV Genotype 2.0 Assay (LiPA) (Siemens Healthcare Diagnostics Inc., Tarrytown, USA). For HIV-positive individuals, HIV-RNA viral load was measured using a commercially-available PCR assay. Patients were asked to fill in questionnaires on adherence at W4 and W8 and on quality of life [using the Short Form (SF)-12v2] at treatment initiation and PTW12.

If SVR12 was not achieved, drug concentrations were quantified on stored plasma samples obtained during treatment (W4 and W8) using UPLC coupled with tandem MS (Acquity-TQD, Waters Corporation; Milford, MA, USA).^{20,21,22}

For post-treatment study visits at which HCV-RNA levels were detectable, HCV drug resistance profiles were determined from stored serum samples at baseline and the study visit where failure was recorded^{23, 24} NS3, NS5A and NS5B sequencing analysis were performed by in-depth phylogenetic analysis. The phylogenetic trees were visualized using Dendroscope (version 3.5.9) and used to differentiate between late relapse of the same infection or reinfection with another HCV strain. HCV-RNA testing continued until 48 weeks post-treatment to assess HCV re-infection.

End-points

The primary outcome was SVR12, defined by undetectable plasma HCV-RNA twelve weeks after end of treatment (EOT). The secondary outcomes included virological failure at the end of follow-up, as determined by the number of patients harbouring HCV (NS5A, NS5B and NS3/4) resistance mutations, and reinfection. For HIV-positive individuals, the median levels of HIV-RNA and CD4+ T cell count at baseline and PTW12 were also evaluated.

Statistical analysis

Given the single-arm, non-comparative design, the sample size calculation was based on statistical precision and not on a comparative statistical test. Assuming an absolute precision of <30% between lower and upper bounds of the Clopper-Pearson 95% CIs for a binomial end-point and an SVR12 in ITT analysis at ≥80%, a total 30 patients were necessary.

In PP analysis, all available HCV-RNA data at PTW12 were used to calculate SVR12. In ITT analysis, all available HCV-RNA data were used to calculate SVR12, yet individuals lost to follow-up prior to PTW12 were considered as not achieving SVR12.

Scores from the SF-12 were calculated for each of the eight quality of life domains. All scores were transformed to range from 0 to 1 with higher scores indicating better mental or physical health. Scores were compared between treatment initiation and PTW12 using a fractional probit model with cluster-adjusted variance estimators.

Statistical analysis was conducted using STATA v15.0 (College Station, TX, USA). Significance was determined using a *p*-value <0.05.

Role of funding source

This was an investigator-initiated study funded by Merck Sharp & Dohme Corp. The funding source was not involved in any part of data collection, interpretation of data or writing of the manuscript.

RESULTS

Description of the study population

Between 31 May 2017 and 20 August 2018, of 33 individuals assessed for eligibility, 3 did not meet inclusion criteria (Figure 1). Thirty individuals were allocated to study treatment, all of whom received eight-week grazoprevir/elbasvir. The distribution of criteria sets used to define acute HCV infection at inclusion is provided in Table S1.

Patient characteristics at grazoprevir/elbasvir initiation are described in Table 1. Median time between suspected presentation of acute HCV symptoms and treatment initiation was 85 days (IQR=63-130). Illicit drug use over the past 12 months was found in 18 (75%) patients (Table S2). Ten patients (33%) were diagnosed as re-infected, either by change in HCV genotype from previous known infection ($n=4$) or positive HCV-RNA viral load from previously successful treatment ($n=6$). Eight (27%) patients had an STI within 12 months prior to inclusion, with the most common STIs being *Neisseria gonorrhoea* (5 infections) and syphilis (4 infections).

Median HCV-RNA was 5.71 log₁₀ IU/mL (IQR=5.08-6.44) and patients harboured genotype 1a ($n=15$), genotype 1b ($n=1$) or genotype 4 ($n=14$). Of note, 8 patients had an HCV-RNA viral load >800,000 IU/mL and genotype 1a. Median ALT and AST levels were 282 IU/L (IQR=100-202) and 128 IU/L (IQR=100-202), respectively, and 3 (10%) presented with ALT or AST levels $\leq 2\times$ upper limit of normal.

The large majority of patients were HIV-positive ($n=28$), all of whom were undergoing ART with either undetectable HIV-RNA ($n=26$, <40 copies/mL) or low-level replication ($n=2$, at 51 and 90 copies/mL). No patient was on an agent having possible drug interactions with study medication. Immunosuppression was largely mild with only 3 (11%) HIV-positive patients having a previous AIDS-defining illness and median CD4⁺ cell count at 580/mm³ (IQR=467-706).

Virological and biochemical response

216

217 Of the 30 included patients, all completed 8 weeks of grazoprevir/elbasvir. During post-treatment
218 follow-up, one patient died (suicide occurring 3 weeks after the W8 visit). In total, 29 completed the
219 PTW12 visit. The proportion of detectable HCV-RNA at each follow-up visit is given for both PP and
220 ITT analysis in Figure 2A. The proportion of patients achieving SVR12 was 93% (95%CI=78-99%) in ITT
221 analysis and 96% (95%CI=82-99%) in PP analysis. SVR12 was not different between individuals with or
222 without re-infection ($n=10/10$, 100% and $n=18/19$, 95%; respectively) or between individuals with
223 baseline HCV-RNA $\leq 800,000$ IU/mL or $>800,000$ IU/mL ($n=16/16$, 100% and $n=12/13$, 92%;
224 respectively).

225

226 As shown in Figure 2B, most patients were able to normalize transaminase levels at the PTW12 visit.
227 One patient had increasing ALT levels from 78 IU/L at baseline to 109 IU/L PTW4, then decreasing to
228 37 IU/L at PTW12. This patient had undetectable HCV-RNA at all visits during and post-treatment.
229 Two patients had ALT levels >40 IU/L at their PTW12 visit: one with undetectable HCV-RNA at all
230 visits after W4 and the other failing treatment (described below). No patient had a flare in ALT or AST
231 >250 IU/L during or post-treatment.

232

233 One patient with GT1a experienced virological failure. Virological and biological evolution of this
234 patient is described in Figure 3. At W4, both plasma GZR and EBR concentrations were undetectable
235 (<5 ng/mL) and concomitant plasma antiretroviral concentrations were low (FTC, 111 ng/mL; TDF, 48
236 ng/mL; RPV 78 ng/mL). At W8, plasma concentrations of study drug were normal (GZR, 144 ng/mL;
237 EBR, 105 ng/mL) including those of the antiretroviral regimen (FTC, 1074 ng/mL; TDF, 213 ng/mL;
238 RPV, 90 ng/mL). HCV-RNA level was undetectable at PTW4 but rebounded to $6.57 \log_{10}$ IU/mL at
239 PTW12. Phylogenetic analysis indicated the same HCV viral strain compared to baseline, ruling out
240 HCV re-infection. Genotypic resistance analysis at PTW12 revealed the V36M amino acid
241 substitution in NS3, M28V in NS5A, and S556G in NS5B regions of the HCV genome.

242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267

Self-reported adherence and quality of life

At the W4 and W8 visits, respectively, 1 (4%) and 2 (7%) patients declared missing grazoprevir/elbasvir within the last 4 days and 2 (7%) and 1 (3%) forgot to take grazoprevir/elbasvir at least once during the weekend. The level of grazoprevir/elbasvir adherence was comparable to that of ART, for which 4 (15%) and 3 (11%) patients reported missing ≥ 1 pills included in their regimen within the last 4 days and 1 (3%) and 1 (3%) forgetting to take their medication at least once during the weekend at the W4 and W8 visits, respectively.

Based on scores from the SF-12 (Table S3), significant improvement between treatment initiation and PTW12 were observed in role physical ($p=0.02$) and role emotional ($p=0.02$) domains. No significant differences between treatment initiation and PTW12 were observed in the other six domains: physical functioning, bodily pain, general health, vitality, social functioning, and mental health. At baseline, when compared to the overall average SF-12 across domains in the entire study population (0.61), the only patient failing treatment had an average SF-12 of 0.58 and the two patients with completed and attempted suicide, respectively, had an average SF-12 of 0.79 and 0.40.

Control of HIV-infection during follow-up

At PTW12, all 27 HIV-positive participants with available data had HIV-RNA < 50 copies/mL and median CD4⁺ T cell count was 655/mm³ (IQR=488-829). After treatment with grazoprevir/elbasvir, 3 patients switched from TDF/FTC/RAL ($n=2$) or TDF/FTC/DTG ($n=1$) to TAF/FTC/elvitegravir/COBI (unrelated to HIV virological failure or toxicity). No HIV virological failure was reported during treatment or post-treatment.

Adverse events and serious adverse events

During treatment and post-treatment, 60 clinical adverse events were reported and ≥ 1 clinical adverse event occurred in 23 (77%) participants. Only 3 serious clinical adverse events occurred: one patient had a panic attack with underlying suicidal ideation and another patient committed suicide. None of the serious clinical adverse events was related to study treatment. All other possible and probable drug-related adverse events are listed in Table 2.

Reinfection rate

No HCV re-infections were observed within 12 weeks after EOT. However, HCV-RNA was detected in 3 patients after PTW12. One patient with previous GT1a infection acquired a GT4d strain 48 weeks after PTW12, the second patient with a previous GT4 infection acquired an HCV GT1b strain 24 weeks after PTW12, and the third patient with a previous GT1a infection acquired a GT1a strain 28 weeks after PTW12, but phylogenetic analysis showed that it was a different strain from baseline. In total, the reinfection rate was 3/28 (10.7/100 patient-month). All have successfully been retreated with various DAA combinations for 8 to 16 weeks.

DISCUSSION

In this multicentre pilot study of MSM with recently acquired HCV infection, we observed that 96% (28/29) were able to achieve SVR12 after an eight-week course of treatment with grazoprevir/elbasvir. This proportion was 93% (28/30) assuming those lost to follow-up did not achieve SVR at the end of follow-up. These SVR rates are comparable to chronically HCV-infected individuals undergoing the same treatment combination for 12-weeks, regardless of HIV status.¹⁵⁻¹⁸ Of those completing the PTW12 visit, only one exhibited RAS associated with grazoprevir/elbasvir

294 treatment failure. These data provide strong evidence that eight-week grazoprevir/elbasvir is
295 appropriate for the treatment of acute HCV infection genotype 1 and 4, and particularly in HIV-
296 positive patients.

297
298 In chronic HCV infection, the C-WORTHY trial showed that higher SVR12 rates were achieved in
299 patients treated with 12-week versus 8-week grazoprevir/elbasvir.¹⁷ Nevertheless, more recent
300 clinical trials evaluating 8-week grazoprevir/elbasvir have demonstrated high SVR in HCV genotype
301 1b or 4 infected patients with liver fibrosis levels at METAVIR F0-F2^{25,26} or in genotype 1b when
302 plasma HCV-RNA is low.²⁷ Given that MSM recently acquiring HCV are unlikely to have severe liver
303 fibrosis²⁸, 8-week duration of grazoprevir/elbasvir could be sufficient for treating acute HCV. The
304 recent DAHHS2 trial conducted in the Netherlands demonstrated that short-term duration of
305 grazoprevir/elbasvir is possible in acutely infected individuals, reporting SVR12 of 99% in 80 patients
306 with acute HCV genotype 1 (all with genotype 1a or 4 infection after 8-week grazoprevir/elbasvir).¹⁹
307 Our data confirm this finding with high SVR12 in both ITT and PP analysis. Furthermore, as reported
308 by Boerekamps *et al.*, SVR12 was also able to be achieved in all patients harbouring HCV genotype
309 1a (one of the most prevalent genotypes among European MSM³) and all but one genotype 1a-
310 infected patient with plasma HCV-RNA >800,000 IU/mL at baseline, which is also associated with
311 virologic failure during chronic HCV infection.²⁹

312
313 Of note, there were four patients with detectable plasma HCV-RNA at EOT, 3 of whom eventually
314 achieved SVR12. This phenomenon has been described in chronic HCV infected individuals enrolled in
315 registration trials of peg-IFN-free DAA containing regimens.³⁰ The other patient with detectable HCV-
316 RNA at EOT eventually developed virological failure with the emergence of 28V and 36M RAS, both of
317 which have been previously identified in HCV mono-infected patients with virological failure.³¹ This
318 patient might not have been consistently using study medication, as indicated by the undetectable
319 plasma grazoprevir/elbasvir and ARV concentrations at W4. Emergence of these substitutions would

likely be the result of selective pressures from poor adherence and not necessarily insufficient treatment duration.

Poor quality of life has been associated with suboptimal adherence and early treatment discontinuation^{32, 33}, thereby increasing the risk of virologic failure. In our study, we were able to observe improvements in both role physical and role emotional domains. Studies on patient-reported outcomes during and after grazoprevir/elbasvir treatment for chronic HCV have observed improvements in many domains, including mental health and role physical, but not with role emotional.³⁴ Differences between studies could be the result of included study populations (acute versus chronic HCV infection) and Quality of Life inventories (SF-36 versus SF-12). Yet considering the limited data on quality of life during DAA-treated acute infection, it is difficult to explain these discrepancies any further.

Despite some improvements in quality of life, no change in the mental health domain was observed. It is also noteworthy that, after EOT, one patient attempted suicide and another committed suicide. Illicit drug use within the 12 months prior to baseline was observed in 75% of patients with available data. Half of these individuals had consumed drugs that are known to be associated with high risk sexual behaviour and STI acquisition³⁵ and whose use could be potentially driven by internalizing mental health disorders.³⁶ These observations underscore the need to incorporate mental health and addiction services when treating HCV in this key population.

The proportion with a probable or possible drug-related adverse event was low and in line with previous studies using grazoprevir/elbasvir in HCV mono-infection or HIV/HCV co-infected populations.^{15,18,19} Besides the serious mental health problems identified during follow-up, no other serious adverse events and no serious laboratory abnormalities were observed. Other studies have included STIs as a part of adverse events, reported in roughly 24% of MSM twelve weeks after

346 grazoprevir/elbasvir treatment.¹⁹ We did not consider this as an adverse event; nevertheless, 27% of
347 included patients reported at least one bacterial STI within the year prior to treatment initiation and
348 among them, 88% reported having at least one adverse event during follow-up. Assuming that STIs
349 were only to occur in patients with previously detected infections, the proportion with adverse
350 events during treatment would not have changed.

351
352 In HIV-positive individuals, potential drug-drug interactions between grazoprevir/elbasvir and certain
353 antiretroviral agents are problematic and reconsideration of ART regimen might be required for a
354 number of patients planning to initiate grazoprevir/elbasvir.³⁷ ART regimens had to be modified in
355 only one of the twenty-eight HIV-positive individuals without concern and no rebounds in plasma
356 HIV-RNA or changes in CD4+ T cell count were observed in ART-experienced patients during
357 grazoprevir/elbasvir treatment.

358
359 There are certain limitations to our study. First, treatment efficacy was not assessed using a non-
360 inferior study design. Nevertheless, we argue that any non-inferiority limit assuming an SVR12 rate
361 below 90% would be uninformative given the efficacy of current DAA treatments. Second, we included
362 a limited number of acute HCV infected individuals, reducing the potential numbers of virological
363 failures and hence our ability to determine reasons for failure. There were also few HIV-negative
364 participants enrolled, which limits generalizability to this group. Finally, data on baseline RAS were
365 not collected and thus we cannot conclude on the efficacy of grazoprevir/elbasvir in the presence of
366 these substitutions. NS5A RAS appear to mostly affect virologic response in patients with high HCV
367 viral loads²⁷ or those infected with HCV genotype 1a and less so with genotype 1b or 4.^{15,38-40} Half of
368 included patients were infected with genotype 1a, which would make data on RAS particularly
369 helpful. Nevertheless, in the DAHHS2 study, where sequencing was performed at baseline for all
370 patients, no one harbouring a strain with RAS experienced virological failure.¹⁹

In conclusion, given the 96% SVR12 rate and the single patient with RAS, eight-week grazoprevir/elbasvir is an efficacious treatment option for recently acquired HCV infection. As adverse events were uncommon, particularly those related to the study drug, this treatment would appear well-tolerated. These data support the use of grazoprevir/elbasvir for rapid treatment against HCV, appropriate for reducing ongoing HCV transmission in the current epidemic of HCV infection among European MSM with genotypes 1 and 4. The suicide attempt and completed suicide coupled with frequent illicit drugs use observed in this study population stress the need for mental health services in MSM at risk for HCV infection.

ACKNOWLEDGEMENTS

We are grateful to the patients and clinical teams involved in the SAHIV Study.

The SAHIV clinical study team: Hôpital Saint-Antoine – Julie Bottero, Jessica Krause, Michel Pauchard, Julie Lamarque; Hôpital Tenon – Julie Chas, Anne Adda; Hôpital La Pitié-Salpêtrière – Marc-Antoine Valantin, Ludovic Lenclume; Hôpital Bichat – Yazdan Yazdanpanah, Malikhone Chansombat; CHU de Lyon – Patrick Mialhes, Valérie Galvan; CHU de Nice – Eric Rosenthal, Sophie Bréaud. Coordinating pharmacy: Hôpital Saint-Antoine – Anne Dagueneil-Nguyen. Data management team: Inserm U1136 – F. Chau, G. Mawuvi, M. Angot, F. Carrat.

Scientific advisory board: Karine Lacombe, Pierre-Marie Girard, Jessica Krause, Julie Chas, Marc-Antoine Valantin, Yasdan Yazdanpanah, Patrick Mialhes, Eric Rosenthal, Stéphane Chevaliez, Lionel Piroth, Amir Guidoum, Anders Boyd, Hayette Rougier.

Data Safety Monitoring Board: Pascal Trimoulet (CHU Bordeaux, Bordeaux), Philippe Flandre (Inserm, Paris), Patrick Ingiliz (Zentrum für Infektiologie, Berlin)

FUNDING

MSD provided an unrestricted grant for the study and was not involved in any part of the data collection, analysis, and writing.

TRANSPARENCY DECLARATIONS

KL reports receiving travel grants and advisory board fees from Gilead, Abbvie and MSD.

SC reports receiving advisory board fees from Abbott and Cepheid.

All other authors: none to declare

REFERENCES

1. European Association for the Study of the Liver. EASL Recommendations on Treatment of Hepatitis C 2018. *J Hepatol* 2018; **69**: 461-511.

2. World Health Organization. Global Hepatitis Programme. Global hepatitis report. <http://apps.who.int/iris/bitstream/10665/255016/1/9789241565455-eng.pdf?ua=1>.

3. van de Laar T, Pybus O, Bruisten S *et al*. Evidence of a large, international network of HCV transmission in HIV-positive men who have sex with men. *Gastroenterology* 2009; **136**: 1609-17.

- 422 4. van Santen DK, van der Helm JJ, Del Amo J *et al.* Lack of decline in hepatitis C virus incidence
423 among HIV-positive men who have sex with men during 1990-2014. *J Hepatol* 2017; **67**: 255-62.
- 424 5. Hoornenborg E, Achterbergh RCA, Schim van der Loeff MF *et al.* MSM starting preexposure
425 prophylaxis are at risk of hepatitis C virus infection. *AIDS* 2017; **31**: 1603-10.
- 426 6. Braun DL, Hampel B, Martin E *et al.* High Number of Potential Transmitters Revealed in a
427 Population-based Systematic Hepatitis C Virus RNA Screening Among Human Immunodeficiency
428 Virus-infected Men Who Have Sex With Men. *Clin Infect Dis* 2019; **68**: 561-8.
- 429 7. Bethea ED, Chen Q, Hur C *et al.* Should we treat acute hepatitis C? A decision and cost-
430 effectiveness analysis. *Hepatology* 2018; **67**: 837-46.
- 431 8. MacGregor L, Martin NK, Mukandavire C *et al.* Behavioural, not biological, factors drive the
432 HCV epidemic among HIV-positive MSM: HCV and HIV modelling analysis including HCV treatment-
433 as-prevention impact. *Int J Epidemiol* 2017; **46**: 1582-92.
- 434 9. Popping S, Hulleger SJ, Boerekamps A *et al.* Early treatment of acute hepatitis C infection is
435 cost-effective in HIV-infected men-who-have-sex-with-men. *PLoS One* 2019; **14**: e0210179.
- 436 10. Palaniswami PM, El Sayed A, Asriel B *et al.* Ledipasvir and Sofosbuvir in the Treatment of
437 Early Hepatitis C Virus Infection in HIV-Infected Men. *Open Forum Infect Dis* 2018; **5**: ofy238.
- 438 11. Rockstroh JK, Bhagani S, Hyland RH *et al.* Ledipasvir-sofosbuvir for 6 weeks to treat acute
439 hepatitis C virus genotype 1 or 4 infection in patients with HIV coinfection: an open-label, single-arm
440 trial. *Lancet Gastroenterol Hepatol* 2017; **2**: 347-53.
- 441 12. Naggie S, Marks KM, Hughes M *et al.* Sofosbuvir Plus Ribavirin Without Interferon for
442 Treatment of Acute Hepatitis C Virus Infection in HIV-1-Infected Individuals: SWIFT-C. *Clin Infect Dis*
443 2017; **64**: 1035-42.
- 444 13. Martinello M, Bhagani S, Gane E *et al.* Shortened therapy of eight weeks with
445 paritaprevir/ritonavir/ombitasvir and dasabuvir is highly effective in people with recent HCV
446 genotype 1 infection. *J Viral Hepat* 2018; **25**: 1180-8.

- 447 14. Lawitz E, Gane E, Pearlman B *et al.* Efficacy and safety of 12 weeks versus 18 weeks of
448 treatment with grazoprevir (MK-5172) and elbasvir (MK-8742) with or without ribavirin for hepatitis
449 C virus genotype 1 infection in previously untreated patients with cirrhosis and patients with
450 previous null response with or without cirrhosis (C-WORTHY): a randomised, open-label phase 2 trial.
451 Lancet 2015; **385**: 1075-86.
- 452 15. Zeuzem S, Ghalib R, Reddy KR *et al.* Grazoprevir-Elbasvir Combination Therapy for Treatment-
453 Naive Cirrhotic and Noncirrhotic Patients With Chronic Hepatitis C Virus Genotype 1, 4, or 6 Infection:
454 A Randomized Trial. Ann Intern Med 2015; **163**: 1-13.
- 455 16. Rockstroh JK, Nelson M, Katlama C *et al.* Efficacy and safety of grazoprevir (MK-5172) and
456 elbasvir (MK-8742) in patients with hepatitis C virus and HIV co-infection (C-EDGE CO-INFECTION): a
457 non-randomised, open-label trial. Lancet HIV 2015; **2**: e319-27.
- 458 17. Sulkowski M, Hezode C, Gerstoft J *et al.* Efficacy and safety of 8 weeks versus 12 weeks of
459 treatment with grazoprevir (MK-5172) and elbasvir (MK-8742) with or without ribavirin in patients
460 with hepatitis C virus genotype 1 mono-infection and HIV/hepatitis C virus co-infection (C-WORTHY):
461 a randomised, open-label phase 2 trial. Lancet 2015; **385**: 1087-97.
- 462 18. Braun DL, Hampel B, Kouyos R *et al.* High Cure Rates With Grazoprevir-Elbasvir With or
463 Without Ribavirin Guided by Genotypic Resistance Testing Among Human Immunodeficiency
464 Virus/Hepatitis C Virus-coinfected Men Who Have Sex With Men. Clin Infect Dis 2019; **68**: 569-76.
- 465 19. Boerekamps A, De Weggheleire A, van den Berk GE *et al.* Treatment of acute hepatitis C
466 genotypes 1 and 4 with 8 weeks of grazoprevir plus elbasvir (DAHHS2): an open-label, multicentre,
467 single-arm, phase 3b trial. Lancet Gastroenterol Hepatol 2019; **4**: 269-77.
- 468 20. Jung BH, Rezk NL, Bridges AS *et al.* Simultaneous determination of 17 antiretroviral drugs in
469 human plasma for quantitative analysis with liquid chromatography-tandem mass spectrometry.
470 Biomed Chromatogr 2007; **21**: 1095-104.
- 471 21. van Seyen M, de Graaff Teulen MJA, van Erp NP *et al.* Quantification of second generation
472 direct-acting antivirals daclatasvir, elbasvir, grazoprevir, ledipasvir, simeprevir, sofosbuvir and

473 velpatasvir in human plasma by UPLC-MS/MS. J Chromatogr B Analyt Technol Biomed Life Sci 2019;
474 **1110-1111**: 15-24.

475 22. Merck. ZEPATIER Product information.
476 www.merck.com/product/usa/pi_circulars/z/zepatier/zepatier_pi.pdf.

477 23. Hezode C, Chevaliez S, Scoazec G *et al*. Retreatment with sofosbuvir and simeprevir of
478 patients with hepatitis C virus genotype 1 or 4 who previously failed a daclatasvir-containing
479 regimen. Hepatology 2016; **63**: 1809-16.

480 24. Rodriguez C, Mercier-Darty M, Soulier A *et al*. Performance assessment of a fully automated
481 deep sequencing platform for HCV resistance testing. Antivir Ther 2019.

482 25. Abergel A, Hezode C, Asselah T *et al*. High efficacy and safety of grazoprevir and elbasvir for 8
483 weeks in treatment-naïve, non-severe fibrosis HCV GT1b-infected patients: Interim Results of the
484 STREAGER Study. J Hepatol 2018; **68**: S110.

485 26. Asselah T, Pol S, Hezode C *et al*. Efficacy and safety of elbasvir/grazoprevir for 8 or 12 weeks
486 in participants with hepatitis C virus genotype 4 infection. Hepatology 2018; **68**: 385A.

487 27. Huang CF, Hung CH, Cheng PN *et al*. An open-label, randomized, active control trial of 8
488 versus 12 weeks of elbasvir/grazoprevir for treatment-naïve chronic hepatitis C genotype 1b patients
489 with mild fibrosis (EGALITE): Impact of baseline viral loads and NS5A resistance-associated
490 substitution. J Infect Dis 2019.

491 28. Newsum AM, Kooij KW, Boyd A *et al*. Progression of liver fibrosis following acute hepatitis C
492 virus infection in HIV-positive MSM. AIDS 2019; **33**: 833-44.

493 29. Serfaty L, Jacobson I, Rockstroh J *et al*. The accuracy of baseline viral load for predicting the
494 efficacy of elbasvir/grazoprevir in participants with hepatitis C virus genotype 1a infection: An
495 integrated analysis. J Viral Hepat 2019; **26**: 329-36.

496 30. Harrington PR, Deming DJ, Komatsu TE *et al*. Hepatitis C Virus RNA Levels During Interferon-
497 Free Combination Direct-Acting Antiviral Treatment in Registrational Trials. Clin Infect Dis 2015; **61**:
498 666-7.

- 499 31. Lahser F, Galloway A, Hwang P *et al*. Interim analysis of a 3-year follow-up study of NS5A and
500 NS3 resistance-associated substitutions after treatment with grazoprevir-containing regimens in
501 participants with chronic HCV infection. *Antivir Ther* 2018; **23**: 593-603.
- 502 32. Bernstein D, Kleinman L, Barker CM *et al*. Relationship of health-related quality of life to
503 treatment adherence and sustained response in chronic hepatitis C patients. *Hepatology* 2002; **35**:
504 704-8.
- 505 33. Younossi ZM, Stepanova M, Nader F *et al*. The patient's journey with chronic hepatitis C from
506 interferon plus ribavirin to interferon- and ribavirin-free regimens: a study of health-related quality
507 of life. *Aliment Pharmacol Ther* 2015; **42**: 286-95.
- 508 34. Ng X, Nwankwo C, Arduino JM *et al*. Patient-reported outcomes in individuals with hepatitis C
509 virus infection treated with elbasvir/grazoprevir. *Patient Prefer Adherence* 2018; **12**: 2631-8.
- 510 35. Achterbergh RCA, de Vries HJC, Boyd A *et al*. Identification and characterization of latent
511 classes based on drug use among men who have sex with men at risk of sexually transmitted
512 infections in Amsterdam, the Netherlands. *Addiction* 2019.
- 513 36. King M, Semlyen J, Tai SS *et al*. A systematic review of mental disorder, suicide, and
514 deliberate self harm in lesbian, gay and bisexual people. *BMC Psychiatry* 2008; **8**: 70.
- 515 37. Martinello M, Dore GJ, Skurowski J *et al*. Antiretroviral Use in the CEASE Cohort Study and
516 Implications for Direct-Acting Antiviral Therapy in Human Immunodeficiency Virus/Hepatitis C Virus
517 Coinfection. *Open Forum Infect Dis* 2016; **3**: ofw105.
- 518 38. Asselah T, Reesink H, Gerstoft J *et al*. Efficacy of elbasvir and grazoprevir in participants with
519 hepatitis C virus genotype 4 infection: A pooled analysis. *Liver Int* 2018; **38**: 1583-91.
- 520 39. Kwo P, Gane EJ, Peng CY *et al*. Effectiveness of Elbasvir and Grazoprevir Combination, With or
521 Without Ribavirin, for Treatment-Experienced Patients With Chronic Hepatitis C Infection.
522 *Gastroenterology* 2017; **152**: 164-75 e4.
- 523 40. Zeuzem S, Serfaty L, Vierling J *et al*. The safety and efficacy of elbasvir and grazoprevir in
524 participants with hepatitis C virus genotype 1b infection. *J Gastroenterol* 2018; **53**: 679-88.

525

526

527

528

TABLES

Table 1. Description of the study population at treatment initiation

General characteristics	N=30
Male gender*	30 (100)
Birth country/region*	
France	22 (73)
Europe (outside of France)	3 (10)
Asia	3 (10)
Other	2 (7)
Age, years [†]	44 (34-52)
BMI, kg/m ^{2†}	22.4 (20.6-25.4)
Hypertension* [‡]	10 (33)
Diabetes* [§]	1 (3)
Co-infection with HIV*	28 (93)
HCV-related characteristics	N=30
Transmission route* [¶]	
Nasal drug-use	6 (20)
Injection drug-use	5 (17)
Sexual contact	23 (77)
Unknown	2 (7)
Involved first HCV reinfection*	9 (30)
Involved second HCV reinfection*	1 (10)
Time since last infection, years [†]	2.8 (1.2-3.8)

534 **Table 1 (con't).**

HIV-related characteristics	N=28
Known duration of HIV infection [†]	8.5 (4.5-14.0)
AIDS-defining illness*	3 (11)
Undergoing ART**μ	28 (100)
HIV RNA viral load >50 copies/mL*	2 (7)
HIV RNA viral loads, copies/mL**	(51, 90)
CD4+ T cell count, /mm ^{3†}	580 (467-706)
CD8+ T cell count, /mm ^{3†}	644 (508-964)
CD4:CD8 ratio [†]	0.84 (0.55-1.24)

535 *n (%) [†]Median (IQR)

536 [†]Systolic blood pressure >130 mm/Hg or diastolic blood pressure >80 mm/Hg.

537 [§]Declared by treating physician.

538 [¶]More than one transmission route can occur within individuals.

539 **For individuals with HIV RNA >50 copies/mL.

540 μ TDF/emtricitabine (FTC)/rilpivirine (RPV), n=9; abacavir/lamivudine/dolutegravir (DTG), n=9;
 541 TDF/FTC/DTG, n=6; TDF/FTC/raltegravir (RAL), n=3; or TDF/FTC dual therapy, n=1. Of note, one
 542 patient switched from TDF/FTC/cobicistat (COBI)-boosted darunavir to TDF/FTC/DTG at treatment
 543 initiation (following recommendations for concomitant antiretroviral agents).

544 **Table 2. Adverse and serious adverse events during follow-up**

All clinical adverse events, <i>n</i>	60
Patients with ≥ 1 clinical adverse event, <i>n</i> (%)	23 (77)
Serious clinical adverse events, <i>n</i>	3
Panic attack	1
Suicidal ideation (without suicide)	1
Suicide	1
Serious drug-related clinical adverse events, <i>n</i>	0
Discontinuation due to clinical adverse events, <i>n</i> (%)	0 (0)
Deaths, <i>n</i> (%)	1 (3)
Cause of death	
Suicide	1
All possible drug-related adverse events	8
Diarrhoea	2
Abdominal distension	1
Stomach spasms	1
Myalgia	1
Xerostomia	1
Hyperhidrosis	1
Osteoarthritis of knee and hands	1
All probable drug-related adverse events	8
Insomnia	2
Diarrhea	2
Fatigue	2
Binge eating	1

546 **Table 2 (con't).**

Dysgeusia	1
Patients with ≥1 grade 2 laboratory abnormality*, <i>n</i> (%)	12 (40)
Creatinine clearance 60-90 mL/min/1.73m ²	12
Patients with ≥1 grade 3 or 4 laboratory abnormality*, <i>n</i> (%)	0 (0)

547 *Specifically during treatment.

FIGURE LEGENDS

Figure 1. Patient flow during the study

Figure 2. Hepatitis C virus (HCV) and alanine transaminase levels during and after treatment with grazoprevir/elbasvir

Number of patients with undetectable HCV over total number of patients are given during treatment at week 4 (W4) and week 8 (W8) as well as post-treatment at week 4 (PTW4) and week 12 (PTW12) in PP and ITT analysis (A). Individual (grey lines) and median (black line) ALT levels are given at each study visit in (B).

Figure 3. Virological and biochemical evolution of the patient with treatment failure

Hepatitis C virus (HCV)-RNA (genotype 1a) and ALT levels during treatment at week 4 (W4) and week 8 (W8) as well as post-treatment at week 4 (PTW4) and week 12 (PTW12) for the only patient with treatment failure. Antiretroviral (ARV), elbasvir (EBR) and grazoprevir (GZR) plasma levels were quantified at W4 and W8 and RASs were determined at PTW12.