



HAL
open science

Two case reports of local envenoming by the Spotted grass snake, *Psammophylax rhombeatus* (Linnæus, 1758) (Serpentes, Psammophiidae)

Ivan Ineich, Francis Girard, Tyrone Ping, Jordy Reynes, Scott A Weinstein

► To cite this version:

Ivan Ineich, Francis Girard, Tyrone Ping, Jordy Reynes, Scott A Weinstein. Two case reports of local envenoming by the Spotted grass snake, *Psammophylax rhombeatus* (Linnæus, 1758) (Serpentes, Psammophiidae). *Toxicon*, 2021, 195, pp.24-28. 10.1016/j.toxicon.2021.02.013 . hal-03273641

HAL Id: hal-03273641

<https://hal.sorbonne-universite.fr/hal-03273641>

Submitted on 29 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Two Case Reports of Local Envenoming by the Spotted grass**
2 **snake, *Psammophylax rhombeatus* (Linnæus, 1758) (Serpentes,**
3 ***Psammophiidae*)**

4

5 Ivan Ineich^{a,*}, Francis Girard^b, Tyrone Ping^c, Jordy Reynes^d, Scott A.
6 Weinstein^e

7

8 ^a Institut de Systématique, Évolution, Biodiversité (ISYEB)
9 Muséum national d'Histoire naturelle, CNRS, Sorbonne Université,
10 EPHE, Université des Antilles - CP 30, 57 rue Cuvier, 75005 Paris,
11 France

12

13 ^b 167 boulevard Vincent Auriol, 75013 Paris, France

14

15 ^c 7 Oak Tree Avenue, Glen Anil, KwaZulu-Natal 4051, South Africa

16

17 ^d 31 rue des Mouleyres, 11120 Bize Minervois, France

18

19 ^e Department of Toxinology, Women's and Children's Hospital, 72
20 King William St., North Adelaide, South Australia, Australia 5000

21

22 * Corresponding author. E-mail: ivan.ineich@mnhn.fr

23

24

25 ABSTRACT

26 Two cases of bites by a South African psammophiid snake, *Psammophylax rhombeatus*, are described and
27 analyzed. These are the first detailed reports of local envenoming by a *Psammophylax* spp. While handling
28 a wild-collected 1 m *P. rhombeatus*, the snake inflicted a protracted bite proximal to the
29 metacarpophalangeal joint of digit #5, left hand of a 24-year-old male amateur herpetologist. Local edema
30 persisted for three days, but no pain or other signs or symptoms including non-specific autonomic effects
31 (e.g. headache, nausea) occurred. In a second case, a 28-year-old male herpetologist-photographer was
32 repositioning a 0.58 m female *P. rhombeatus* in order to photograph the snake and her egg clutch, when
33 the snake bit the metacarpophalangeal joint of digit #5, left hand, and briefly advanced its jaws. The bite
34 caused mild local pain, progressive edema of the left hand, and arthralgia; resolution required almost 1
35 week. Bites from non-front-fanged snakes such as these by *P. rhombeatus* are uncommonly reported in
36 comparison with those described for front-fanged snakes (e.g. Viperidae, Elapidae). Therefore,
37 documentation of bites even with minimal effects provides information essential for the construction of an
38 accurate medical risk profile for these less-known species.

39

40 Keywords:

41 Non-front fanged snake;

42 Colubroid;

43 Psammophiidae;

44 Envenoming;

45 Snake bites;

46 *Psammophylax rhombeatus*;

47 Spotted grass snake;

48 Rhombic Skaapsteker

49

50 **1. Introduction**

51

52 Grass Snakes or ‘Skaapstekers’ (genus *Psammophylax* Fitzinger,
53 1843) are terrestrial Central, South and East African non-front-fanged
54 colubroid snakes (NFFCs) with low-pressure venom glands (or,
55 ‘Duvernoy glands’) (Taub, 1967; Heymans, 1977; McKinstry, 1983).
56 Sub-equal maxillary teeth are followed after a diastema by a pair of
57 slightly enlarged, grooved posterior maxillary teeth. The venom of
58 studied species is reportedly quite viscous, and has been compared
59 with glycerin (FitzSimons, 1921).

60

61 Six species are recognized in the genus *Psammophylax* (Table
62 1). They feed on small vertebrates including fishes (Broadley, 1977;
63 Branch, 1988; Shine et al., 2006; Cottone & Bauer, 2010; Chippaux &
64 Jackson, 2019; Keates et al., 2019; Wilkey, 2019). They are unusual
65 among snakes by variably engaging in parental care (Shine et al.,
66 2006). Their common name ‘Skaapsteker’ has an Afrikaans origin

67 related to their presumed habit of biting sheep (literally: ‘sheep
68 stabber’), a belief that unfortunately leads some ranchers to kill these
69 snakes on sight (Tyrone Ping, personal observations). However, cape
70 cobras (*Naja nivea* (Linnæus, 1758), Elapidae) are probably
71 responsible for most of the stock losses that are blamed on relatively
72 harmless *Psammophylax* (FitzSimons, 1921; Alexander & Marais,
73 2007), although some authors have considered puff adders, *Bitis*
74 *arietans* Merrem, 1820 (Viperidae), more likely culprits (Elstob in
75 Chippaux & Jackson, 2019).

76

77 Early workers, most prominently, Frei (1910), Andrews (1912)
78 and FitzSimons (1912) contemplated whether *Psammophylax* spp.
79 were venomous. Based on experiments with fowl reportedly bitten by
80 *Trimerorhinus rhombeatus* (most likely *P. rhombeatus*; *Trimerorhinus*
81 A. Smith, 1847 is a junior synonym of *Psammophylax*.), FitzSimons
82 (1921: 488) considered the possibility that this species could be as
83 dangerous as the Boomslang (*Dispholidus typus* (A. Smith, 1828)) and
84 its venom more potent than that of elapid species of notable medical
85 importance (e.g., cobras, *Naja* spp. and mambas, *Dendroaspis* spp.).
86 However, Phisalix (1922) reported delayed lethal effects in a small
87 sampling of rodents and chicken bitten by specimens of two
88 *Psammophylax* species from South Africa.

89

90 Christensen (1955: 2) later commented that *Trimerorhinus*
91 *tritaeniatus* possess “potent venom”. Alexander and Marais (2007)

92 stated that the venom of *Psammophylax* spp. has no medical
93 relevance. Spawls et al. (2018) suggested that the venom of
94 *Psammophylax* spp. is “fairly toxic”, but little is injected when a bite
95 is delivered to a human, and thus no serious symptoms have been
96 recorded. Wilkey (2019) considered the two species of *Psammophylax*
97 present in Malawi and reported that their venom is mild and may
98 cause local swelling and some mild pain. However, Spawls and
99 Branch (2020) noted that *P. tritaeniatus* reportedly have “toxic
100 venom” with that of *P. rhombeatus* having “potent” neurotoxicity.
101 Nonetheless, they concluded that no medically significant effects from
102 a *Psammophylax* bite have so far been recorded. Likewise, FitzSimons
103 and Smith (1958) noted the absence of any reported serious case of
104 human envenomation from *Trimerorhinus* (now *Psammophylax*), and
105 opined that all published cases are dubious. Consistent with the latter
106 impression, Chapman (1968) described the effects observed in three
107 cases of bites by *Psammophylax* spp. in Natal, and indicated that these
108 consisted of “a slight local reaction of bruising and swelling, one with
109 a rigor”. Branch (1982) reported similar signs and symptoms
110 following a bite from an East Cape *P. rhombeatus*. However, these
111 reports are described second hand and it is not clear if formal medical
112 review was ever conducted within a reasonable timeframe after the
113 bites.

114

115 Warrell (1995: 460) noted that *P. tritaeniatus* bites were
116 associated with non-specific systemic symptoms such as nausea,

117 headache and rigors. Warrell (2010) later opined that *Psammophylax*
118 is probably capable of mild envenomations that cause only local pain,
119 mild swelling and lymphangitis, and emphasized the absence of any
120 reported serious bites. Kuch and Mebs (2002, 167-168) quoted
121 FitzSimons (1910) apparently referring to a case of human
122 envenomation from *P. rhombeatus* that reportedly included
123 “giddiness, lassitude, cold clammy skin, cold sweat on the forehead, a
124 little swelling at the site of the fang punctures, with discolouration of
125 the surrounding tissue”.

126

127 The previously reported cases were also reviewed by Minton
128 (1990), and later, Weinstein et al. (2011) evaluated them and assigned
129 these with a low-quality evidence rating (C/D). This rating indicated
130 limitations noted in these reports including lack of formal medical
131 review and/or detailed information/documentation about the species
132 assigned responsibility for the bite(s). Therefore, there are no detailed
133 descriptions of bites by *Psammophylax* spp., or first-hand reports that
134 have documented their medical effects.

135

136 While performing field work and photography, two of the
137 authors were bitten by specimens *P. rhombeatus* and developed
138 effects consistent with local envenoming. We report here on these two
139 first-hand cases of bites by *P. rhombeatus*, present the details of these
140 bites and their reported effects.

141

2. Case reports

Case 1

While performing a herpetological field survey on September 30 2004 near the Sterkfontein dam (Orange Free State) in South Africa (28°27'N, 29°01'E) in the morning around 1000-1100 hrs, one of us (JR; 24 yr old male, amateur herpetologist, with no significant medical history, no current medications or known allergies) was bitten by an approximately 1± m (total length) *P. rhombeatus* (gender undetermined) (Fig. 1). The victim was previously (2002) bitten by a Brazilian lancehead, *Bothrops moojeni* Hoge, 1966 (Viperidae, Crotalinae) and severely envenomed (progressive edema and consumptive coagulopathy); he was treated with six vials of Bothrofav[®] (Sanofi-Pasteur, Lyon, France; a monovalent F(ab')₂ antivenom against venom of the fer-de-lance or Martinique lancehead, *Bothrops lanceolatus* Bonnaterre, 1790; this antivenom has no clinically proven paraspecificity for *B. moojeni* venom).

The *P. rhombeatus* was discovered under a piece of pottery (pipe) in a humid habitat that had recently burned (2-3 months prior) and was beginning to reestablish floral growth. The snake was captured without difficulty, and was kept for several minutes in order to photograph the specimen. While handling the snake, the victim momentarily loosened his grip, and was promptly bitten proximal to

167 the metacarpophalangeal joint of the digit #5, left hand. The snake
168 maintained a firm grip for approximately 10-15 secs while it
169 implanted the enlarged posterior maxillary teeth into the victim's
170 hand. Once manually disengaged from the bite site, two symmetrical
171 puncture marks consistent with the enlarged posterior maxillary teeth,
172 as well as small blood drops produced by several of the anterior teeth
173 were observed (Fig. 2). The local, proportional bleeding stopped after
174 several minutes. There was no first aid applied, and the wound was
175 not disinfected. Several minutes post-bite, edema and erythema were
176 first noted around the bite site. Within about seven hours post-bite, the
177 edema involved the whole hand; moving the hand and fingers was not
178 painful, but there was a nearly complete limitation of flexion and
179 extension; manual dexterity was significantly affected (e.g., handling
180 silverware was very difficult). Subsequently, the local edema of the
181 bitten hand remained unchanged for three days and resolved on only
182 the fourth day post-bite. The pain was estimated as 1/10 (using the
183 verbal pain 0 to 10 with 10 being 'unbearable' pain) during the bite,
184 but there was no pain (0/10), even when moving the hand and fingers.
185 The victim described the most persistent effect as marked digital
186 stiffness noted especially with attempted flexion of the fingers.
187 Ecchymosis was absent and there were no other symptoms or signs
188 including non-specific complaints such as headache and/or nausea.
189 Complete resolution was observed within 4 days.

190

191 **Case 2**

192

193 A 28-yr-old male herpetologist/professional photographer with
194 no significant medical history or allergies was investigating the
195 herpetofauna at Glen Austen Gauteng, South Africa (25.975536°N,
196 28.169737°E) at 1120 hrs on October 18 2014 when he encountered
197 an approximately 600 mm (total length) female *P. rhombeatus*, that
198 was coiled around a recently deposited egg clutch (Fig. 3A). The
199 snake was gently moved in order to photograph the eggs, together
200 with the specimen. As the snake was handled, it inflicted a bite
201 delivered to the medial-dorsal surface of the thumb, left hand, that
202 initially consisted of contact with only the anterior maxillary and
203 mandibular teeth; however, the snake began to advance its jaws and
204 the victim sensed being punctured by both of the enlarged posterior
205 maxillary teeth (Fig. 3B). The snake was firmly attached and resisted
206 removal by the victim; it was gently coaxed to release by manually
207 manipulating its head and had remained attached for approximately 45
208 seconds. The victim noted that immediately after detachment of the
209 snake, the wounds immediately bled and continued bleeding for an
210 estimated 2-3 minutes; the victim opined that the wounds bled a bit
211 disproportionally in relation to the lacerations and punctures
212 comprising the wound site. There was no attempted first-aid or
213 interventions for the bite. For several hours post-bite, the victim only
214 noted a “slight burning sensation”, but approximately 7 hours later the
215 victim reported being awakened by his partner who stated that she felt
216 “heat radiating” from his hand. The victim noted local moderate

217 edema that involved the entire left hand; the edema mildly inhibited
218 digit flexion/extension and the skin was notably warm (“hot to
219 touch”). The victim reported a mild “throbbing-type pain” ranked 2/10
220 (“very mild, but pulsating, making it uncomfortable particularly when
221 trying to sleep”), and did not ingest any analgesics or any other
222 medications/substances. The edema was still present and moderately
223 inhibited manual dexterity 36 hours post-bite (Figs 4A, 4B), and
224 persisted for almost 1 week. Stiffness was present in the all of the
225 metacarpophalangeal joints and digits of the left hand, but this
226 completely subsided after approximately 4 days. Thereafter, the signs
227 and symptoms fully resolved without any sequelae. The victim
228 reported several previous asymptomatic and uneventful brief bites
229 from several species of non-front-fanged snakes including
230 *Psammophylax* (also, the psammophiids, short-snouted grass snake,
231 *Psammophis brevirostris* Peters, 1881; cross-marked grass snake,
232 *Psammophis crucifer* (Daudin, 1803), and the colubrine colubrid,
233 marbled tree snake, *Dipsadoboa aulica* (Günther, 1864)); there was no
234 history of any envenoming from any front-fanged species, and no
235 history of having received antivenom. The victim has many years of
236 experience photographing reptiles in the field often requiring close
237 contact with many species of reptiles.

238

239 **3. Discussion**

240

241 The victims in these cases received firm bites with some jaw
242 advancement from adult *P. rhombeatus*. Although the snakes were
243 attached for <1 min, both victims developed mild-moderate signs and
244 symptoms consisting of significant local edema with some local
245 progression and minimal pain at the bite site. These resolved without
246 sequelae in four to seven days (respectively, victims #1 and #2).
247 Notably, both victims experienced similar clinical evolution of edema
248 and local pain. The victims did not receive formal medical review and
249 this limits the evidence quality (Level C/D, following the evidence
250 rankings by Weinstein et al. (2011). However, careful observations
251 were recorded and accurately correlated with images taken in
252 chronological order, thus providing precise records of the effects of
253 the bites.

254

255 Observations of *Psammophylax* spp. have suggested that these
256 snakes rarely attempt to bite unless roughly handled (FitzSimons,
257 1921). Our own observations (FG, TP) suggest that some *P.*
258 *rhombeatus* specimens are quick to bite when handled, as was also
259 reported by Branch (1988). However, the effects of the bites described
260 here may indicate that an especially prolonged bite by a large adult
261 specimen might produce more significant local envenoming. Spawls
262 and Branch (2020) commented that *P. rhombeatus*, “has a potent
263 neurotoxic venom, but no adverse symptoms ever recorded from a
264 bite”, while Bates (1996) referring to *Psammophylax* venom and that
265 of several other NFFC stated, “...is comparatively weak and bites

266 usually result in only localized swelling and pain similar to that
267 caused by a bee sting”. Perceived venom neurotoxicity of
268 *Psammophylax* is probably based on the previously outlined
269 experiments by Phisalix (1922), FitzSimons (1921), and the comments
270 by Christensen (1955). However, there is so far no biomedical
271 evidence of neurotoxins in any *Psammophylax* venom, nor any
272 clinical evidence of neurotoxicity from their bites, although as noted
273 here, detailed reports about their venom and bites are absent in the
274 literature. Therefore, there is insufficient documentation to firmly
275 characterize the clinical syndrome that may be caused by prolonged
276 bites by *Psammophylax*. Additionally, 3-finger-fold neurotoxins are
277 common in many NFFCs including other psammophiids (Lumsden et
278 al., 2007; Jackson et al., 2019; Modahl and Mackessy, 2019), and
279 these may occur in *Psammophylax* venoms, but could have prey
280 specificity (e.g., lizards). We decline speculation about the venom
281 components that may have caused the effects that developed in the
282 victims described here. Several venom components probably
283 contribute to this e.g., snake venom metalloproteases; however, victim
284 hypersensitivity may also play a role in the acute effects of some
285 NFFC bites such as these (Weinstein et al., 2011). The minimal pain
286 associated with the rapidly progressive edema suggests the specific
287 investigation of this whenever possible e.g., laboratory evaluation of
288 inflammatory cellular subsets and immunoglobulins. For example,
289 Th2-related markers, including the interleukins (IL) and C-motif
290 chemokines (CCL): IL-5, IL-13, IL-10, IL-31, CCL13, CCL18, and

291 many others, are prominently expressed in lesions manifested in acute
292 atopic dermatitis (Malik et al., 2017).

293

294 Management of bites by *Psammophylax* spp. is governed by the
295 severity of the victim's distress; simple wounds and local effects can
296 be managed with meticulous wound care, while progressive edema,
297 persistent pain, bleeding and greater distress suggestive of systemic
298 effects should be promptly reviewed by a physician whenever
299 possible. There is no antivenom for bites by *Psammophylax* spp. and
300 no antivenom of any kind should be given because: it subjects the
301 patient to unnecessary risks; it would be ineffective, and in any case is
302 not clinically indicated. There is also no evidence supporting the
303 administration of parenteral steroids, or antihistamines for treatment
304 of bites by NFFC, nor for antibiotic prophylaxis unless there has been
305 interference with the wound (e.g., incorrect first-aid, application of
306 local remedies, etc.), environmental contamination, or shows signs of
307 early necrosis (Weinstein et al., 2011; Weinstein, 2017).

308

309 Further studies of the venoms of *Psammophylax* spp. are
310 desirable, and formal reports of any bites inflicted by these snakes can
311 further document their medical effects and clinical management.

312

313 **References**

314

315 Alexander, G., Marais, J. 2007. A guide to the Reptiles of Southern
316 Africa. Struik Publishers, Cape Town, South Africa. 408 pp.

317

318 Andrews, W.H. 1912. Experiments with snakes. Report Director Vet.
319 Res., Dept. Agric., Union of S. Africa 2, 406–483.

320

321 Bates, M.F. 1995. Poisonous snakes of the Free State. *Culna* 49, 12–
322 14.

323

324 Branch, B. 1988. Field Guide to the Snakes and Other Reptiles of
325 Southern Africa (First edition). New Holland, London. 328 pp. + 96
326 pls.

327

328 Branch, B. 2005. A photographic guide to snakes and other reptiles
329 and amphibians of East Africa. Struik Publishers, Cape Town, South
330 Africa. 144 pp.

331

332 Branch, W.R. 1982. Venomous snakes of southern Africa, 3.
333 Concluding part: Colubridae. *The Snake* 14, 1–17.

334

335 Branch, W.R., Baptista, N., Keates, C., Edwards, S. 2019.
336 Rediscovery, taxonomic status, and phylogenetic relationships of two
337 rare and endemic snakes (Serpentes: Psammophiinae) from the
338 southwestern Angolan plateau. *Zootaxa* 4590, 342–366.

339

340 Branch, W.R., Braak, H.H. 1987. Reptiles and amphibians of the
341 Addo Elephant National Park. *Koedoe* 30, 61–111.

342

343 Broadley, D.G. 1977. A revision of the African snakes of the genus
344 *Psammophylax* Fitzinger (Colubridae). *Occas. Pap. Natl. Mus. Rhod.*,
345 *Ser. B* 6(1), 1–44.

346

347 Chapman, D.S. 1968. The symptomatology, pathology, and treatment
348 of the bites of venomous snakes of central and southern Africa.

349 Bücherl, W., Buckley, E.E., Deulofeu, V., eds. *Venomous Animals*
350 *and Their Venoms. Vol. I: Venomous vertebrates.* Academic Press,
351 New York, 463–527.

352

353 Chippaux, J.-P., Jackson, K. 2019. *Snakes of Central and Western*
354 *Africa.* Johns Hopkins University Press, Baltimore. x + 429 pp.

355

356 Christensen, P.A. 1955. South African snake venoms and antivenoms.
357 *The South African Institute for Medical Res., Johannesburg, v,* 129
358 pp.

359

360 Cottone, A.M., Bauer, A.M. 2010. Sexual dimorphism, diet,
361 reproduction, and their geographic variation in sympatric
362 *Psammophiids, Psammophis crucifer and Psammophylax rhombeatus*
363 *rhombeatus,* from Southern Africa. *Copeia* 4, 578–590.

364

365 FitzSimons, F.W. 1910. The Snakes of South Africa; Their Venom
366 and the Treatment of Snake Bite. Walton, Port Elizabeth, South
367 Africa. 160 pp.

368

369 FitzSimons, F.W. 1912. The snakes of Southern Africa: Their Venom
370 and Treatment of Snake Bite. T.M. Miller, Cape Town, South Africa.
371 547 pp.

372

373 FitzSimons, F.W. 1921. The snakes of South Africa, their venom and
374 the treatment of snake bite. T. Maskew Miller, Cape Town. xvi + 550
375 pp.

376

377 FitzSimons, D.C., Smith, H.M. 1958. Another rear-fanged South
378 African snake lethal to humans. *Herpetologica* 14, 198–202.

379

380 Frei, W. 1910. Über einige Experimente mit Giften und
381 Speicheldrüsenextracten südafrikanischer Schlangen. *Zeitschr. Hyg.*
382 *Infektionskr.* 3, 211–217

383

384 Heymans, J.C. 1977. Variations des glandes parotidiennes chez les
385 Serpents Colubridae d'Afrique Centrale. *Rev. Zool. Afr.* 91, 893–905.

386

387 Jackson, T.N.W., Jouanne, H., Vidal, N. 2019. Snake Venom in
388 Context: Neglected Clades and Concepts. *Front. Ecol. Evol.* 7, 332.

389

390 Keates, C., Conradie, W., Greenbaum, E., Edwards, S. 2019. A snake
391 in the grass: Genetic structuring of the widespread African grass snake
392 (*Psammophylax* Fitzinger 1843), with the description of a new genus
393 and a new species. *J. Zool. Syst. Evol. Res.* 57, 1039–1066.

394

395 Kuch, U., Mebs, D. 2002. Envenomations by colubrid snakes in
396 Africa, Europe and the Middle East. *J. Toxicol. - Toxin Reviews* 21,
397 159–179.

398

399 Lumsden, N.G., Banerjee, Y., Kini, R.M., Kuruppu, S., Hodgson,
400 W.C. 2007. Isolation and characterization of rufoxin, a novel protein
401 exhibiting neurotoxicity from venom of the psammophiine,
402 *Rhamphiophis oxyrhynchus* (Rufous beaked snake).
403 *Neuropharmacology* 52, 1065–1070. doi:
404 10.1016/j.neuropharm.2006.11.002

405

406 Malik, K., Heitmiller, K.D., Czarnowicki, T. 2017. An update on the
407 pathophysiology of atopic dermatitis. *Dermatol. Clin.* 35(3), 317–326.

408

409 McKinstry, D.M. 1983. Morphologic evidence of toxic saliva in
410 colubrid snakes: a checklist of world genera. *Herpetological Review*
411 14, 12–15.

412

413 Minton, S.A., Jr. 1990. Venomous bites by nonvenomous snakes: an
414 annotated bibliography of colubrid envenomation. *J. Wild. Med.* 1(2),
415 119–127.

416

417 Modahl, C., Mackessy, S.P. 2019. Venoms of rear-fanged snakes:
418 New proteins and novel activities. *Front. Ecol. Evol.* 7 (doi:
419 10.3389/fevo.2019.00279).

420

421 Phisalix, M. 1922. Animaux venimeux et venins : la fonction
422 venimeuse chez tous les animaux ; les appareils venimeux, les venins
423 et leurs propriétés ; les fonctions et usages des venins ; l'envenimation
424 et son traitement. Tome 2. Masson & Cie Éditeurs - Libraires de
425 l'Académie de Médecine, Paris. xii + 864 pp. + [1 p., errata], 17 pls.

426

427 Shine, R., Branch, W.R., Webb, J.K., Harlow, P.S., Shine, T. 2006.
428 Sexual dimorphism, reproductive biology, and dietary habits of
429 Psammophiine snakes (Colubridae) from Southern Africa. *Copeia*
430 2006, 650–664.

431

432 Spawls, S., Branch, B. 2020. *The Dangerous Snakes of Africa*.
433 Bloomsbury, London (Oxford, New York, New Dehli, Sydney). 336
434 pp.

435

436 Spawls, S., Howell, K., Hinkel, H., Menegon, M. 2018. Field Guide to
437 East African Reptiles, 2nd edition. Bloomsbury Wildlife, London. 624
438 pp.

439

440 Taub, A.M. 1967. Comparative histological studies on Duvernoy's
441 gland of Colubrid snakes. Bull. Amer. Mus. Nat. Hist. 138, 1–50, 58
442 plates.

443

444 Warrell, D.A. 1995. Clinical toxicology of snakebite in Africa and the
445 Middle East / Arabian Peninsula. In Meier, J., and White, J. Handbook
446 of clinical toxicology of animal venoms and poisons. CRC Press,
447 Boca Raton, New York, London, Tokyo.

448

449 Warrell, D.A. 2010. Guidelines for the prevention and clinical
450 management of snakebite in Africa. World Health Organization,
451 Regional Office for Africa, Brazzaville. xiv + 129 pp.

452

453 Weinstein, S.A. 2017. Non-Front-Fanged Colubroid Snakes. In Brent,
454 J., Burkhart, K., Dargan, P., Hatten, B. et al. Textbook of Critical
455 Care Toxicology. Springer International, pp. 2453–2492.

456

457 Weinstein, S., Warrell, D., White, J., Keyler, D.E. 2011. “Venomous”
458 Bites from Non-Venomous Snakes. 1st Edition. A Critical Analysis of

459 Risk and Management of “Colubrid” Snake Bites. Elsevier, London.
460 xxvii + 336 pp.

461

462 Wilkey, R. 2019. Snakes of Malawi. A field guide to the snake species
463 of Malawi (2019 revised edition, 5th revision). Biddles Books. 329 pp.

464

465 **Table 1**466 A brief inventory of *Psammophylax* spp. and related species.

467

Valid name	Common names	Previous name(s); subspecies	Distribution	Reference(s)
<i>Psammophylax kellyi</i> Conradie, Keates & Edwards in Keates, Conradie, Greenbaum & Edwards, 2019	Tanzanian Grass Snake, Tanzanian Skaapsteker	<i>Psammophylax multisquamis</i> [pro parte]; [no recognized subspecies]	Mount Meru, N Tanzania	Keates et al., 2019
<i>Psammophylax multisquamis</i> (Loveridge, 1932)	Kenyan Grass Snake, Kenyan Striped Skaapsteker	<i>Trimerorhinus tritaeniatus multisquamis</i> , <i>Psammophylax tritaeniatus multisquamis</i> , <i>Psammophis variabilis multisquamis</i> ; [no recognized subspecies]	Ethiopia, Kenya, N Tanzania, N Rwanda	Branch, 2005; Spawls et al., 2018; Chippaux & Jackson, 2019
<i>Psammophylax ocellatus</i> (Bocage, 1873)	Angolan Grass Snake, Angolan Skaapsteker	<i>Psammophylax rhombeatus ocellatus</i> [pro parte]; [no recognized subspecies]	SW Angola, ?NW Namibia	Branch et al., 2019; Keates et al., 2019
<i>Psammophylax rhombeatus</i> (Linnæus, 1758)	Spotted Grass Snake, Spotted Skaapsteker, Rhombic Skaapsteker	<i>Coluber rhombeatus</i> , <i>Trimerorhinus rhombeatus</i> , <i>Psammophylax rhombeatus ocellatus</i> [pro parte]; [no recognized subspecies]	S Namibia, Rep. of South Africa, Lesotho, Swaziland, SW Angola	Phisalix, 1922; Branch, 1988; Alexander & Marais, 2007; Chippaux & Jackson, 2019; Spawls & Branch, 2020
<i>Psammophylax tritaeniatus</i> (Günther, 1868)	Striped Grass Snake, Striped Skaapsteker, Three-lined Grass Snake	<i>Trimerorhinus tritaeniatus</i> , <i>T. tritaeniatus</i> , <i>Rhagerhis tritaeniatus</i> , <i>Rhagerhis tritaeniata</i> ; [recognized subspecies: <i>P. t. tritaeniatus</i> , <i>P. t. subniger</i>]	NE Namibia, N Botswana, Zimbabwe, NE Rep. of South Africa, Angola, S Tanzania, Zambia, Malawi, S Dem. Rep. Congo, Zambia, Mozambique	Phisalix, 1922; Branch, 1988, 2005; Alexander & Marais, 2007; Spawls et al., 2018; Chippaux & Jackson, 2019; Wilkey, 2019; Spawls & Branch, 2020
<i>Psammophylax variabilis</i> Günther, 1893	Grey-bellied Grass Snake, Grey-bellied Skaapsteker	<i>Trimerorhinus tritaeniatus</i> [pro parte], <i>Trimerorhinus tritaeniatus variabilis</i> ; [recognized subspecies: <i>P.v.variabilis</i> , <i>P.v. vanoyei</i>]	N Botswana, Dem. Rep. Congo, Tanzania, Burundi, Rwanda, Uganda, Kenya, Ethiopia, Malawi, Zambia, Mozambique, Namibia	Branch, 1988; Alexander & Marais, 2007; Spawls et al., 2018; Chippaux & Jackson, 2019; Wilkey, 2019
<i>Kladirostratus acutus</i> (Günther, 1888)	Striped Beaked Snake, Beaked Skaapsteker	<i>Psammophis acutus</i> ; <i>Psammophylax acutus</i> ; <i>Rhamphiophis acutus</i> ; [recognized subspecies: <i>K. a. acutus</i> , <i>K. a. jappi</i>]	Angola, NW/W Zambia, S Dem. Rep. Congo, W Tanzania, N Malawi, N Rwanda	Keates et al., 2019
<i>Kladirostratus togoensis</i> (Matschie, 1893)	Northern Sharp-nosed Skaapsteker	<i>Psammophis togoensis</i> ; <i>Rhamphiophis togoensis</i> ; <i>Rhamphiophis acutus garambensis</i> ; <i>Psammophylax acutus togoensis</i> ; <i>Psammophylax togoensis</i> ; [no recognized subspecies]	Ghana, Togo, Nigeria, Cameroon, Central Afr. Rep., N Dem. Rep. Congo, Uganda	Keates et al., 2019

468

469

Figure legends

470

471 Figure 1. The Spotted Grass Snake or Rhombic Skaapsteker
472 (*Psammophylax rhombeatus*) that inflicted the bite (Case #1). The
473 snake was approximately 1 m total length; gender is unknown.

474



475

476

477

478 Figure 2. The left hand shortly after having been bitten by
479 *Psammophylax rhombeatus* (Case #1). The snake remained attached
480 for approximately 10-15 seconds, but did not advance its jaws. The
481 bite produced symmetrical punctures that corresponded with the
482 enlarged posterior maxillary teeth; only scant bleeding was noted.
483 Note the early edema proximal to the metacarpophalangeal joint.

484

485

486



487

488

489 Figures 3A, B. (A) The Spotted Grass Snake or Rhombic Skaapsteker
490 (*Psammophylax rhombeatus*) that inflicted the bite (Case #2). The
491 snake (female) was approximately 580 mm total length, and was
492 found coiled around her clutch of recently deposited eggs. The snake
493 inflicted the bite when she was re-positioned in order to take
494 photographs (image courtesy of Tyrone Ping). (B) The *Psammophylax*
495 *rhombeatus* specimen shown in Fig. 3A, inflicting the bite on the
496 medial-dorsal surface of the left thumb, victim #2. The wound
497 reportedly bled “freely”, but bleeding ceased within approximately 2
498 minutes. The victim reported that the subsequent local edema was
499 accompanied only by mild throbbing pain, but rendered sleep difficult
500 (image courtesy of Tyrone Ping).

501

502



503


504

505

506

507 Figures 4A, B. Persistent local effects, left hand 36 hrs after bite by
508 *Psammophylax rhombeatus* on medial-dorsal surface of left thumb
509 (Case #2). The snake remained attached for approximately 45 seconds
510 and briefly advanced its jaws. The bite caused progressive local
511 edema that eventually involved the entire hand and caused functional
512 limitations of digital flexion and extension; only mild local pain was
513 noted. Note the significant local edema of the thenar eminence in
514 Panel B (images courtesy of Tyrone Ping).

515



516

517