

Bilateral visual projections exist in non-teleost bony fish and predate the emergence of tetrapods

Robin J Vigouroux, Karine Duroure, Juliette Vougny, Shahad Albadri, Peter Kozulin, Eloisa Herrera, Kim Nguyen-Ba-Charvet, Ingo Braasch, Rodrigo Suárez, Filippo del Bene, et al.

▶ To cite this version:

Robin J Vigouroux, Karine Duroure, Juliette Vougny, Shahad Albadri, Peter Kozulin, et al.. Bilateral visual projections exist in non-teleost bony fish and predate the emergence of tetrapods. Science, 2021, 372 (6538), pp.150-156. 10.1126/science.abe7790. hal-03280289

HAL Id: hal-03280289 https://hal.sorbonne-universite.fr/hal-03280289

Submitted on 7 Jul2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

2 3 4	Bilateral visual projections exist in non-teleost bony fish and predate the emergence of tetrapods
5	One sentence summary: Bilateral vision preceded terrestrial life
6	
7 8 9 10	Robin J. Vigouroux ¹ , Karine Duroure ¹ , Juliette Vougny ² , Shahad Albadri ¹ , Peter Kozulin ³ , Eloisa Herrera ⁴ , Kim Nguyen-Ba-Charvet ¹ , Ingo Braasch ⁵ , Rodrigo Suárez ³ , Filippo Del Bene ¹ * and Alain Chédotal ¹ *
11	*Corresponding authors.
12	Email: filippo.delbene@inserm.fr and Email: alain.chedotal@inserm.fr
13	
14	Sorbonne Université, INSERM, CNRS, Institut de la Vision, 17 Rue Moreau, 75012 Paris,
15	France
16	² Institut Curie, PSL Research University, INSERM U934, CNRS UMR3215, Paris, France
17	³ Queensland Brain Institute, The University of Queensland, Building 79, St Lucia Campus,
18	Brisbane, QLD 4072, Australia
19	Instituto de Neurociencias Av. Ramón y Cajal s/n San Juan de Alicante 03550 Spain
20	³ Department of Integrative Biology and Program in Ecology, Evolution and Behavior,
21	Michigan State University, 288 Farm Lane, East Lansing MI 48824, USA
22	
23	
24	
25	
26	
27	
28	
29	

30 Abstract:

31 In most vertebrates, camera-style eyes contain retinal ganglion cell neurons projecting to 32 visual centers on both sides of the brain. However, in fish, ganglion cells are thought to 33 only innervate the contralateral side. This suggested that bilateral visual projections 34 appeared in tetrapods. Here, we show that bilateral visual projections exist in non-teleost 35 fishes and that the appearance of ipsilateral projections does not correlate with terrestrial 36 transition or predatory behavior. We also report that the developmental program 37 specifying visual system laterality differs between fishes and mammals as the Zic2 38 transcription factor which specifies ipsilateral retinal ganglion cells in tetrapods appears 39 absent from fish ganglion cells. However, overexpressing human ZIC2 induces ipsilateral 40 visual projections in zebrafish. Therefore, the existence of bilateral visual projections 41 likely preceded the emergence of binocular vision in tetrapods.

42

43

44

Eye position on the head is highly variable between species, but frontal eyes have long been considered critical for depth perception (stereopsis) by increasing the overlap of the right and left eye visual fields (1). In vertebrates, ganglion cell axons from each eye cross through each other at the optic chiasm and enter the brain on the contralateral side. In mammals, visual axons from each eye meet and interweave at the chiasm. However, optic nerve crossing modalities are more diverse in fish and in most species the two optic nerves remain fully separated and only overlap at the chiasm (2, 3).

52 Classic neuroanatomical studies showed that in mammals, eye projections are bilateral with a 53 variable fraction of retinal ganglion cell (referred to as ganglion cell thereafter) axons 54 continuing in the ipsilateral optic tract after crossing the chiasm. The proportion of ipsilateral 55 projections is low (2-3%) in rodents but reaches around 40% in primates (4, 5). The comparative 56 analysis of many vertebrate species conducted over several decades suggests that ipsilateral 57 visual axons exist in all mammals, anuran amphibians, some reptiles, and that they are 58 essentially absent or were secondarily lost in birds (5-8). Accordingly, developmental 59 transcriptional programs specifying ipsilateral ganglion cells described in mammals are 60 conserved in Xenopus but not in chick and zebrafish (5, 9). This textbook view implies that 61 visual axon bilaterality emerged in early tetrapods and might have provided a visual advantage, 62 in particular for nocturnal and predatory terrestrial species (10). However, a review of the 63 extensive literature on fish visual systems gives a more complex image with reports, sometimes 64 contradictory, of ipsilateral ganglion cell projections in some fish species (6, 11). Most of these 65 pioneering studies relied on imprecise histological staining methods such as the Nauta-Gygax 66 staining method or autoradiography (12). Here, we assessed the laterality of visual projections 67 in bony fishes (Fig. 1A) with the B fragment of the cholera toxin (12) coupled to fluorescent 68 dyes. Dye-coupled Cholera toxins have not been previously used in fish although they proved 69 to be highly reliable tracers for visual projections in rodents due to their efficient endocytosis 70 by neurons, slow elimination, high photostability and brightness (13). They are also compatible 71 with whole-brain clearing and thereby allow mapping visual pathways in intact brain using 3D 72 light sheet fluorescence microscopy (14, 15).

With more than 30,000 species, fishes account for at least half of the extant vertebrate species (16). We initially focused on ray-finned fishes (Actinopterygians; **Fig. 1A** and **fig. S1**) which separated from lobe-finned fish (Sarcopterygians, including tetrapods) around 450 Million years ago (Ma)(17). Within ray-finned fishes, we initially selected 6 species among the clupeocephalan lineage (**fig. S1**), the largest of the three lineages of teleost fishes which account for most (about 96%) of extant teleosts (18). Within clupeocephalans, three represent ostariophysians (Mexican tetra *Astyanax mexicanus*, redeye piranha *Serrasalmus rhombeus*, and zebrafish *Danio rerio*) and three are percomorphs (green-spotted pufferfish *Tetraodon nigroviridis*, Atlantic mudskipper *Periophtalmus barbarus*, and four-eyed fish *Anableps anableps*). With 9,000 and 16,000 species respectively, ostariophysians and percomorphs are the two largest clades of teleosts. They have diverse eye positions, feeding behaviors, and habitats and some were previously reported to have ipsilateral visual projections (*11*).

85

86 Totally crossed visual projections in teleosts

87 The visual system of the zebrafish has been extensively studied using lipophilic dye tracing or 88 genetic methods and shown to be exclusively contralateral (19). Accordingly, we found that 89 fluorescent (Cholera toxin labelled) axons were only present on the contralateral side of adult 90 zebrafish brains (Fig. 1B and movie S1; n=6). Previously identified retino-recipient visual 91 nuclei (20) could be detected with cholera toxin, thereby validating the use of this tracing 92 method in fish (Fig. 1C). Light-sheet microscopy imaging of cholera toxin injected fish, showed 93 that visual projections were only contralateral in Mexican tetra surface fish (Fig. 1, D and E 94 and movie S1; n=7), contradicting a previous study (21). Likewise, eyes in the redeye piranha 95 only projected contralaterally (Fig. 1, F and G and movie S1; n= 3), in disagreement with earlier 96 work in other piranha species (22, 23). This suggests that ostariophysians only have crossed 97 visual projections.

In the fresh water green-spotted pufferfish (24), a percomorph, the 2 optic nerves stay separated at the chiasm and visual projections were exclusively contralateral (**Fig. 1, H** and **I** and **fig. S2** and **movie S1**; n=4) as previously described in another pufferfish (25). We next studied the four-eyed fish, a percomorph surface dweller fish whose large protruding eyes with duplicated corneas and pupils allows seeing under and above the water (26). Again, one optic nerve passed over the other at the chiasm (**fig. S2**) and cholera toxin tracing showed that in this species, visual projections were also completely crossed (**Fig. 1, J** and **K**; n=3). Similar results were obtained in the mudskipper, a percomorph with amphibious lifestyle (Fig. 1, L and M and fig. S2 and
movie S1; n= 3). Together, these results show that in percomorph eyes, ganglion cells also
likely only project to visual nuclei on the opposite side of the brain (fig. S1).

108 Osteoglossomorphs (bonytongues), a teleost sister groups of clupeocephalans (Fig. 1A and 109 fig. S1), are considered a group of basal (i.e the group which gave rise to later forms) teleosts 110 constituted of about 200 living species (16). We chose to trace visual projections in the African 111 butterflyfish (Pantodon buchholzi), a predator living close to the surface of freshwater system, 112 and found a small contingent of retinal axons (Fig. 2, A and B and movie S1, 2.33±0.23 % 113 ipsilateral projections in the optic tectum; n=3) project to the ipsilateral side, corroborating an 114 earlier report (27). The main portion of ipsilateral axons targeted the tectum and some others 115 targeted pretectal nuclei. Ipsilateral visual axons have also been described in a mormyrid 116 electric fish (Gnathonemus petersii) (28), another osteoglossomorph with a more nocturnal 117 predatory behavior that can orient by active electrolocation (29). These results show that 118 bilateral visual projections exist in osteoglossomorph teleosts regardless of their predatory 119 strategy and lifestyle history. Therefore, within teleosts, ipsilateral projections could have been 120 secondarily lost in clupeocephalans or independently acquired in osteoglossomorphs. In 121 mammals, binocular inputs to visual targets/areas are either segregated (thalamus, colliculus) 122 or intermingled (suprachiasmatic nucleus). As both eyes were injected with 2 distinct Alexa-123 conjugated-cholera toxins, we also studied the relative distribution of ipsilateral and 124 contralateral ganglion cell axons in butterflyfish brain areas innervated by both eyes. This 125 showed that in African butterflyfish (Fig. 2C) retinal inputs from both sides segregated, as in 126 the thalamus and superior colliculus of mammals.

127

128 Bilateral visual projections exist in basal ray-finned fishes

129 These results on teleosts led us to study retinal projections in non-teleost ray-finned fish 130 lineages (Holosteans, Acipenseriforms, and Polypteriforms, Fig. 1A) which split from teleosts 131 before the teleost whole genome duplication event (TGD, Fig. 1A) that occurred around 320 132 Ma in the ancestor of extant teleosts (reviewed in (30)). Holosteans and Acipenseriforms are 133 considered to have evolved slowly since they branched from other vertebrates 350 Ma (31). 134 Bilateral visual projections (movie S2, n=5/5) were observed in the spotted gar (Lepisosteus 135 oculatus), one of the seven extant species of garfish, and a representative of the holosteans. The 136 spotted gar is a unique vertebrate model system as its genome is thought to provide a "bridge" 137 between tetrapods and teleosts (32). Bilateral cholera toxin injections revealed an ipsilateral 138 projection in the rostral optic tectum of the spotted gar, with visual axons targeting several 139 pretectal nuclei. There was no overlap of the contralateral and ipsilateral axons (Fig. 2, D to F 140 and movie S2, n=5/5) which represented 4.78±0.46 % of visual inputs in the tectum, a ratio 141 comparable to rodents. Next, we traced visual projections in the acipenseriform sterlet sturgeon 142 (Acipenser ruthenus, n=2/2). Cholera toxin tracing demonstrated the existence of a binocular 143 domain in the tectum of the sterlet sturgeon as well as in several pretectal nuclei (Fig. 2, G to 144 J and movie S2; 9.77±1.28 % ipsilateral projections in the tectum, n=2). No re-crossing of 145 visual inputs after entering the brain were detected contrary to previous observations in the 146 Russian sturgeon Acipenser güldenstädtii (33). We then studied the armored bichir (Polypterus 147 *delhezi*; n=2/2), a carnivorous nocturnal fish representing the most basally diverging lineage of 148 extant ray-finned fishes, the Polypteriforms. In the bichir, the two optic nerves meet at the 149 chiasm and ganglion cells axons interweave during decussation (fig. S2). Ipsilateral axons 150 projected to numerous pretectal nuclei such as the nucleus opticus dorsolateralis anterior 151 thalami, the area optica ventrolateralis thalami, and the nucleus commissurae posterior par 152 magnocellularis (Fig. 2, K to N and movie S2). This corroborates previous studies in gray bichir 153 *Polypterus senegalus (34).* These results, together with similar observations in the holosteans 154 longnose gar (Lepisosteus osseus) and the bowfin (Amia calva) (35, 36) indicate that bilateral 155 visual projections likely are ancestral among actinopterygians and arose before their 156 diversification and that the ipsilateral component likely was subsequently lost in 157 clupeocephalans. To further test this hypothesis, we analyzed visual projections in the 158 Australian lungfish (Neoceratodus forsteri) a basal member of the lobe-finned fishes 159 (Sarcopterygians) the monophyletic group that includes tetrapods. Lobe-finned fishes diverged 160 from ray-finned fishes about 450 Ma and lungfish are now considered the closest living fish 161 relative of tetrapods (17). In all injected animals (n=6/6) a small ipsilateral projection was found 162 innervating the optic tectum (Fig. 2, O to S and movie S3, n=6). In contrast to other fish species 163 analyzed here, ipsilateral projections intermingled with contralateral ones (Fig. 2, R and S; 164 n=6). This was consistent with an earlier analysis of a single specimen of Australian lungfish 165 (37) and supports the existence of bilateral visual projections in the bony vertebrate ancestor of 166 actinopterygians and sarcopterygians (fig.S1). This intermingling of ipsilateral and contralateral 167 axons could have some functional implications as it suggests that some tectal neurons might 168 receive and integrate inputs from both eyes. Alternatively, it could represent an immature stage 169 of visual system development that could be resolved in adult animals in an activity-dependent 170 manner as it is the case in mammals. Together, these results indicate that the bilateral 171 organization of the visual system likely did not appear in amniotes but that it is an ancestral 172 vertebrate feature that emerged much earlier in evolution, before water-to-land transition and 173 aerial vision adaptation in tetrapods.

174

175 Zic2 expression in the ipsilateral human embryonic retina

Our results raised questions about the evolution and conservation of the genetic mechanisms underlying visual system binocularity. Are they conserved in ray-finned fish with bilateral visual projections? The zinc-finger transcription factor Zic2 specifies the ipsilateral identity of 179 ganglion cells in developing mice, ferrets and Xenopus (5). In Xenopus, Zic2 is absent from 180 the neural retina of pre-metamorphic tadpoles that have only crossed visual projections, but 181 Zic2 is expressed in ipsilaterally projecting ganglion cells after metamorphosis (5). To further 182 evaluate and support the implication of Zic2 in the control of mammalian ganglion cell 183 laterality, we analyzed the expression of ZIC2 in the human eye and compared it to mice by 184 performing immunohistochemistry in whole-mount retinas. In human embryos, ganglion cell 185 axons reach the brain by the 7th post-conception week (pcw7) and the optic nerve is well formed 186 at pcw10 (38). Using the EyeDISCO clearing protocol (15) in mice, we observed that at 187 embryonic day 16 (E16), the peak of Zic2 expression in mice (5), the retinal domain positive 188 for Zic2 represented about 5.34±0.36 % of the total retinal surface (Fig. 3, A to C and fig. S3A; 189 n=6 eyes). Post-mitotic ganglion cells, the first neurons generated in the retina(39), migrate to 190 the basal side to accumulate at the inner surface of the retina and express the transcription 191 factors Islet1 (40) and RNA-binding protein with multiple splicing (RBPMS)(41) (Fig. 3). At 192 pcw9, RBPMS+ and ISLET1+ ganglion cells were present all over the retina (Fig. 3, D to I). 193 Flat-mounted and sections of human retinas from pcw9 embryos, an age equivalent to E16 in 194 mice (40), showed that ZIC2+ cells were restricted to the temporal quadrant of the retina 195 (representing about 18.95±0.98 % of the retina surface), which contains ipsilaterally projecting 196 ganglion cells in primates (Fig. 3, D to J and movie S4; n=3 eyes). ZIC2+ were ganglion cells 197 as they co-expressed ISLET1 and RBPMS (Fig. 3, F to H). In the temporal retina, the density 198 of ZIC2+/ISLET1+ ganglion cells was higher close to the ciliary marginal zone, at the edge of 199 the retina (Fig. 3G), than in more medial regions where it was absent from the most superficial 200 ISLET1+ (Fig. 3H and fig. S3B) and RBPMS+ ganglion cells (Fig. 3F and fig. S3). ZIC2 was 201 not detectable in the nasal retina (Fig.3, D, E and I). Unlike in the mouse (42), ZIC2 was not 202 present in the neuroblastic layer which contains SOX2 (sex determining region Y-box 2) + 203 progenitors (Fig. 3, J to L and fig. S3C). At pcw14, at the end of ganglion cell neurogenesis (39), ZIC2 was still only present in the temporal retina (Fig.3, K and L). Although we could not access later stages of development, the absence of ZIC2 from the most superficial ganglion cells in the inner retina suggest that human ZIC2 is expressed in recently differentiated ipsilateral ganglion cells in the temporal retina and might be down-regulated as they mature, as is the case in mice.

209

210 Zic2 is not expressed in differentiating ganglion cells in fish with bilateral visual

211 projections

212 The pivotal role for Zic2 in the specification of an ipsilateral axonal growth program in 213 mammals was also well correlated with the absence of transcripts of the two *zic2* co-orthologs 214 (zic2a and zic2b, generated in the TGD) in zebrafish ganglion cells (43) (fig. S4, A to C and E 215 to G). Double fluorescent in situ hybridization for zic2b and atoh7 (a committed precursor 216 marker) from 24 to 48 hours post fertilization confirmed that *zic2b* did not colocalize with 217 differentiating *atoh7*+ ganglion cells. By contrast, and as reported for zic2 in the mouse (44), 218 *zic2b* was detected in the ciliary marginal zone, (fig. S4M, n=3) which contains dividing 219 progenitors and stem cells producing all retinal cell types, even in adult teleosts (45, 46) (fig. 220 **S4, H to** L; n=5).

221 The presence of an ipsilateral visual projection in the spotted gar, as extensive as that of mice, 222 together with its well characterized genome and the accessibility of gar embryos (32, 47), led 223 us to evaluate the expression of zic2 in the developing gar retina. We first analyzed the 224 development of the gar visual system using whole-mount immunolabelling, iDISCO+ clearing 225 and light-sheet microscopy (Table S1) (15). In the spotted gar, a few Islet1- immunoreactive 226 ganglion cells were detected at 2-3 days post fertilization (dpf) (Fig. 4, A and B and fig.S5C 227 and movie S5; n=5). At 6-7dpf, optic nerves could be observed and had reached the optic 228 chiasm (Fig. 4, C and D and movie S5; n=5). Development is slower in gar than in zebrafish 229 and it is temperature-dependent (48). The next 10 days of development are only characterized 230 by changes in fin opercular and gill formation but not in eye morphogenesis (48). By 17-18dpf 231 the retina contained many ganglion cells and the optic tract was well developed (Fig. 4, E to G 232 and **movie S5**; n=5). The highly proliferative ciliary marginal zone could be identified by the 233 presence of cells expressing the S-phase marker, proliferating cell nuclear antigen marker and 234 overlapping with zic2 expression, which was absent from neighboring post-mitotic Islet1+ 235 ganglion cells (Fig. 4, H to L). At 2-3dpf and 6-7 dpf, zic2 mRNA was detected in proliferating 236 cells of the developing neuroretina and progressively became restricted to the ciliary marginal 237 zone (fig. S5, A to D). Its paralogs, *zic1* and *zic5*, also enriched in ipsilateral ganglion cell in 238 mice (49), were absent from the embryonic gar retina (fig. S5, E to L). These results show that 239 neither Zic2, Zic1 nor Zic5, specify ipsilateral ganglion cells in the spotted gar, suggesting that 240 *zic* genes might be dispensable for gar ipsilateral projections. By contrast, the presence of Zic2 241 in the ciliary marginal zone of fish and mammals suggests that Zic2 might have a function in 242 retinal precursors that is evolutionarily conserved.

In mammals, Zic2 acts in part by activating the expression of the receptor tyrosine kinase EphB1 in ipsilateral axons (8, 50), whose ligand ephrinB2, localized at the chiasm, prevents crossing (7). According to the lack of Zic2 in the gar ganglion cells, we did not detect *ephB1* mRNA in the developing retina and *ephrinB2* was absent from the chiasm (**fig. S5, M** to **W**).

247

248 Zic2 overexpression induces ipsilateral projections in zebrafish

In mice, *Zic2* overexpression in the retina, outside the ipsilateral domain, increases the proportion of ipsilaterally projecting ganglion cell axons (*50*). Therefore, we tested the hypothesis that despite its absence in fish ganglion cells, the forced expression of Zic2 in zebrafish ganglion cells could affect their axonal targeting. We used a human *ZIC2-T2A-GFP* overexpression construct to express ZIC2 and GFP in the zebrafish eye under the control of the 254 atoh7 promoter. Vertebrate Zic2 proteins are highly conserved (51), with 81.1% identity 255 between Human ZIC2 and zebrafish Zic2a and more than 93% similarity in the zinc finger 256 domain (fig. S6). To visualize the projections coming from ZIC2 overexpressing ganglion cells 257 derived from a single eye, we removed one eye at 2 dpf. As previously reported, under normal 258 conditions retinal fibers from the remaining eye projected exclusively to the contralateral 259 tectum (Fig. 4, M and N; n=10) (52, 53). In contrast, ZIC2-expressing ganglion cells generated 260 ipsilateral retinotectal afferent fibers representing a mean of 19.6±8.5% of the GFP+ axons 261 (Fig. 4, O and P; n=10/13). Expression of ZIC2 did not seem to bias the targeting of ganglion 262 cell axons to their topographic position as previously reported in mouse (54). These results 263 show that *zic2*, although not normally expressed in zebrafish ganglion cells, can still specify an 264 ipsilateral program. In mice, the receptor tyrosine kinase EphB1 is expressed by ipsilateral 265 ganglion cell axons and the ectopic expression of Zic2 in the contralateral retina induces EphB1 266 and reduce midline crossing at the chiasm (8, 50). However, we could not detect EphB1 protein 267 or mRNA (fig. S7) in zebrafish ganglion cells, neither in controls injected with GFP (n=13) nor 268 in ZIC2-overexpressing fish (n=11) suggesting that these may be guided by alternative cues.

269

270 Discussion

271 It has been proposed that the evolution of terrestrial vertebrates followed an increase of eye size 272 in aquatic vertebrates able to see through air, in a process that has occurred also in modern 273 crocodiles or fish species similar to the four-eye fish and mudskippers (55). The existence of 274 ipsilateral projections in the most basally branching groups of both actinopterygians and 275 sarcopterygians indicates that ipsilateral connections were likely already present in the common 276 ancestor of bony vertebrates, a bony fish, thus preceding aerial vision adaptation of tetrapods. 277 This example highlights how the comparative study of a variety of species outside the list of 278 the classical model species allows drawing evolutionary conclusions that may otherwise remain

obscured (56). Moreover, all the teleosts species analyzed in our study can be described as diurnal predators that heavily rely on visual cues to detect and consume their preys. These preys may vary in size from large vertebrates (redeye piranha) to small invertebrates (Mexican tetra, zebrafish, green-spotted puffer, Atlantic mudskipper and the four-eyed fish). In all cases our data show that ipsilateral projections in teleosts are not required for a visually mediated predatory behavior as it is usually assumed in mammals. Recent studies have confirmed this also in larval zebrafish where possible alternative neuronal circuits have been described (*53*).

286 On the other hand, lungfish and some basally branching actinopterygians, where ipsilateral 287 projections are present, show reduced visual system development, as they are bottom dwellers 288 that show nocturnal predatory behavior (lungfish and bichir) or feed on benthic organisms 289 (sturgeon). Overall, our data show that the presence of ipsilateral projections in the visual 290 system of fishes appears to correlate with phylogeny and not with life style or predatory 291 behavior. Along these lines, it is therefore unlikely that ipsilateral retinal projections serve a 292 function similar to what is commonly considered in mammals. On the contrary, visual system 293 bilaterality might have been used as the neural substrate to compute stereopsis following the 294 acquisition in diurnal mammals of visual-based predatory abilities after the Cretaceous-295 Paleogene (K–Pg) extinction event. Supporting this view, the number of ipsilateral projections 296 in reptiles (chelonians and squamates) correlates neither with eye position nor with the degree 297 of binocular field (11, 57). It has been hypothesized that ipsilateral ganglion cells facilitate 298 motor coordination by providing a direct visual feedback to the limb steering brain centers (6). 299 However, the function of ipsilateral ganglion cells in fish remains elusive and behavioral studies 300 in non-canonical model species such as the gar will be required to address this question.

The conservation of the main families of axon guidance cues and receptors in Bilateria suggested that the mechanisms underlying the development of neuronal connectivity are evolutionarily conserved (58, 59). However, the loss of the gene encoding Deleted in Colorectal 304 Cancer receptor in some bird species (60) and the uniqueness of the Roundabout3 receptor in 305 mammals (61) have challenged this view. Here we show that the guidance program specifying 306 visual axon ipsilaterality does not appear to be entirely conserved as we failed to detect 307 expression of a *zic2* and other *zic* genes or *ephB1* in spotted gar ganglion cells. Hence, the 308 textbook model of Zic2 and EphB1 in orchestrating retinal ganglion cell laterality does not 309 simply translate to fish. In rodents, the contralateral identity of ganglion cells is specified by 310 Islet2 (62) and SoxC (63) transcription factors but whether they influence the development of 311 visual axons in fish is unknown. Therefore, further experiments are needed to address the 312 molecular mechanisms underlying visual bilaterality in bony fish species. Are there other 313 pathways, apart from Zic2, that could direct ipsilateral projections or which could block the 314 contralateral fate that occurs normally? A recent study (64) shows that in mice, contralateral 315 ganglion cells activate a non-canonical Wnt signaling pathway to cross the midline. In 316 ipsilateral ganglion cells, Zic2 prevents midline crossing by inducing a genetic module that 317 changes the expression of a set of genes to jointly inhibit this non-canonical Wnt pathway.

318 In fact, the ectopic expression of human ZIC2 in the developing zebrafish retina still induces 319 the formation of ipsilateral visual axons without causing other targeting defects. This finding 320 suggests a possible conservation in fishes of downstream components of the genetic program 321 specifying ipsilateral axons in mammals although the factor initiating its expression and its 322 relationship to Zic2 remains unclear in the gar and in other non-teleost fish. By parsimony 323 principle, given our data, we propose that the presence of ipsilaterality in the bony vertebrate 324 ancestor is the most likely explanation. In particular, the Australian lungfish data show that 325 ipsilateral projections were likely present in the sarcopterygian fish ancestor of tetrapods. 326 Lungfish also functions as outgroup to the actinoptery gians and thereby making an independent 327 origin of ipsilateral visual axons within actinopterygians a less likely hypothesis. An alternative,

- 328 yet less parsimonious explanation, could be that ipsilaterality has evolved independently
- 329 multiple times among bony vertebrates, using different genetic mechanisms.

330

- 331 **References**
- 332

333	1.	S. Ramón y Cajal, <i>Histologie du système nerveux de l'homme & des vertébrés</i>
334	2	(Maloine, Paris, 1909). $C = 1$
335 336	2.	G. L. Walls, <i>The vertebrate eye and its daptative radiation</i> (Hafner Publishing company, Bloomfield Hills Michigan, 1942).
337	3.	K. Mogi <i>et al.</i> , Optic chiasm in the species of order Clupeiformes, family Clupeidae:
338		Optic chiasm of Spratelloides gracilis shows an opposite laterality to that of Etrumeus
339		teres. Laterality. 14, 495–514 (2009).
340	4.	G. Jeffery, L. Erskine, Variations in the architecture and development of the vertebrate
341		optic chiasm. Prog. Retin. Eye Res. 24, 721–753 (2005).
342	5.	E. Herrera et al., Zic2 Patterns Binocular Vision by Specifying the Uncrossed Retinal
343		Projection. Cell. 114, 545–557 (2003).
344	6.	M. L. Larsson, Binocular vision, the optic chiasm, and their associations with
345		vertebrate motor behavior. Front. Ecol. Evol. 3, 89 (2015).
346	7.	S. Nakagawa et al., Ephrin-B regulates the Ipsilateral routing of retinal axons at the
347		optic chiasm. Neuron. 25, 599-610 (2000).
348	8.	S. E. Williams et al., Ephrin-B2 and EphB1 Mediate Retinal Axon Divergence at the
349		Optic Chiasm. <i>Neuron</i> . 39 , 919–935 (2003).
350	9.	A. Seth et al., Belladona/(lhx2) is required for neural patterning and midline axon
351		guidance in the zebrafish forebrain. Development. 133, 725–735 (2006).
352	10.	C. P. Heesy, Seeing in stereo: The ecology and evolution of primate binocular vision
353		and stereopsis. Evol. Anthropol. 18, 21–35 (2009).
354	11.	R. Ward, J. Repérant, S. Hergueta, D. Miceli, M. Lemire, Ipsilateral visual projections
355		in non-eutherian species: random variation in the central nervous system? Brain Res.
356		<i>Rev.</i> 20 , 155–170 (1995).
357	12.	C. Saleeba, B. Dempsey, S. Le, A. Goodchild, S. McMullan, A Student's Guide to
358		Neural Circuit Tracing. Front. Neurosci. 13, 897 (2019).
359	13.	W. L. Conte, H. Kamishina, R. L. Reep, The efficacy of the fluorescent conjugates of
360		cholera toxin subunit B for multiple retrograde tract tracing in the central nervous
361		system. Brain Struct. Funct. 213, 367–373 (2009).
362	14.	X. Luo et al., Three-dimensional evaluation of retinal ganglion cell axon regeneration
363		and pathfinding in whole mouse tissue after injury. Exp. Neurol. 247, 653–662 (2013).
364	15.	R. J. Vigouroux et al., Revisiting the role of Dcc in visual system development with a
365		novel eye clearing method. Elife. 9, e51275 (2020).
366	16.	L. C. Hughes et al., Comprehensive phylogeny of ray-finned fishes (Actinopterygii)
367		based on transcriptomic and genomic data. Proc. Natl. Acad. Sci. U. S. A. 115, 6249-
368		6254 (2018).
369	17.	C. T. Amemiya et al., The African coelacanth genome provides insights into tetrapod
370		evolution. Nature. 496, 311-316 (2013).
371	18.	R. Betancur-R et al., Phylogenetic classification of bony fishes. BMC Evol. Biol. 17,
372		162 (2017).
		14

- R. O. Karlstrom *et al.*, Zebrafish mutations affecting retinotectal axon pathfinding. *Development*. **123**, 427–38 (1996).
- 20. E. Robles, E. Laurell, H. Baier, The Retinal Projectome Reveals Brain-Area-Specific
 Visual Representations Generated by Ganglion Cell Diversity. *Curr. Biol.* 24, 2085–
 2096 (2014).
- T. J. Voneida, C. M. Sligar, A comparative neuroanatomic study of retinal projections
 in two fishes: Astyanax hubbsi (the blind cave fish), and Astyanax mexicanus. J. *Comp. Neurol.* 165, 89–105 (1976).
- 22. E. Fiebig, S. O. E. Ebbesson, D. L. Meyer, Afferent connections of the optic tectum in
 the piranha (Serrasalmus natteren). *Cell Tissue Res.* 231, 55–72 (1983).
- 383 23. S. E. Ebbesson, H. Ito, B. R. Projections, Bilateral Retinal Projections in the Black
 384 Piranah (Serrasalmus niger). *Cell Tissue Res.* 213, 483–495 (1980).
- 385 24. O. Jaillon *et al.*, Genome duplication in the teleost fish Tetraodon nigroviridis reveals
 386 the early vertebrate proto-karyotype. *Nature*. 431, 946–957 (2004).
- 25. D. L. Meyer, E. Fiebig, S. O. E. Ebbesson, A note on the reciprocal connections
 between the retina and the brain in the puffer fish Tetraodon fluviatilis. *Neurosci. Lett.*23, 111–115 (1981).
- L. N. Perez *et al.*, Eye development in the four-eyed fish Anableps anableps: Cranial
 and retinal adaptations to simultaneous aerial and aquatic vision. *Proc. R. Soc. B Biol. Sci.* 284 (2017), doi:10.1098/rspb.2017.0157.
- A. B. Butler, W. M. Saidel, Retinal Projections in the Freshwater Butterfly Fish,
 Pantodon buchholzi (Osteoglossoidei). *Brain. Behav. Evol.* 38, 127–140 (1991).
- 395 28. G. Lázár, S. Libouban, T. Szabo, The mormyrid mesencephalon. III. Retinal
 396 projections in a weakly electric fish, Gnathonemus petersii. J. Comp. Neurol. 230, 1–12
 397 (1984).
- 398 29. G. von der Emde *et al.*, Active electrolocation in Gnathonemus petersii: Behaviour,
 399 sensory performance, and receptor systems. *J. Physiol. Paris.* 102, 279–290 (2008).
- 400 30. I. Braasch, J. H. Postlethwait, in *Polyploidy and Genome Evolution*, P. Soltis, D. Soltis,
 401 Eds. (Springer Berlin Heidelberg, 2012), pp. 341–383.
- 402 31. K. Du *et al.*, The sterlet sturgeon genome sequence and the mechanisms of segmental
 403 rediploidization. *Nat. Ecol. Evol.* 4, 841–852 (2020).
- 40432.I. Braasch *et al.*, The spotted gar genome illuminates vertebrate evolution and405facilitates human-teleost comparisons. *Nat. Genet.* 48, 427–437 (2016).
- 406 33. J. Repérant *et al.*, The Retinofugal Pathways in a primitive actinopterygian, the
 407 chondrostean Acipenser güldenstädti. An experimental study using degeneration,
 408 radioautographic and HRP methods. *Brain Res.* 251, 1–23 (1982).
- 409 34. J. Repérant, J. P. Rio, D. Miceli, M. Amouzou, J. Peyrichoux, The retinofugal
 410 pathways in the primitive African bony fish Polypterus senegalus (Cuvier, 1829). *Brain*411 *Res.* 217, 225–43 (1981).
- 412 35. R. G. Northcutt, A. B. Butler, Retinofugal pathways in the longnose gar Lepisosteus osseus (linnaeus). J. Comp. Neurol. 166, 1–15 (1976).
- 414 36. A. B. Butler, R. G. Northcutt, Retinal Projections in the Bowfin, Amia calva:
- 415 Cytoarchitectonic and Experimental Analysis. *Brain. Behav. Evol.* **39**, 169–194 (1992).
- 416 37. R. Glenn Northcutt, Retinal projections in the australian lungfish. *Brain Res.* 185, 85–
 417 90 (1980).
- 418 38. J. M. Provis, D. Van Driel, F. A. Billson, P. Russell, Human fetal optic nerve:
 419 Overproduction and elimination of retinal axons during development. *J. Comp. Neurol.*420 238, 92–100 (1985).
- 421 39. Y. Lu *et al.*, Single-Cell Analysis of Human Retina Identifies Evolutionarily Conserved
 422 and Species-Specific Mechanisms Controlling Development. *Dev. Cell.* 53, 473-491.e9

423		(2020).
424	40.	A. Hoshino et al., Molecular Anatomy of the Developing Human Retina. Dev. Cell. 43,
425		763-779.e4 (2017).
426	41.	J. M. K. Kwong, J. Caprioli, N. Piri, RNA Binding Protein with Multiple Splicing: A
427		New Marker for Retinal Ganglion Cells. Investig. Opthalmology Vis. Sci. 51, 1052
428		(2010).
429	42.	O. Wang, F. Marcucci, I. Cerullo, C. Mason, <i>eNeuro</i> 3 , ENEURO.0169-16.2016
430		(2016).
431	43.	R. Toyama, D. M. Gomez, M. D. Mana, I. B. Dawid, Sequence relationships and
432		expression patterns of zebrafish zic2 and zic5 genes. Gene Expr. Patterns. 4, 345–350
433		(2004).
434	44.	F. Marcucci <i>et al.</i> , The Ciliary Margin Zone of the Mammalian Retina Generates
435		Retinal Ganglion Cells. Cell Rep. 17 3153–3164 (2016).
436	45.	D. C. Otteson, A. R. D'Costa, P. F. Hitchcock, Putative stem cells and the lineage of
437	101	rod photoreceptors in the mature reting of the goldfish. <i>Dev. Biol.</i> 232 62–76 (2001).
438	46	M Fernández-Nogales V Murcia-Belmonte H Y Chen E Herrera The peripheral
439	101	eve: A neurogenic area with potential to treat retinal pathologies? <i>Prog. Retin. Eve Res.</i>
440		68 (2019) pp 110–123
441	47	I Braasch <i>et al</i> A new model army. Emerging fish models to study the genomics of
442	• • •	vertebrate Evo-Devo I Exp Zool B Mol Dev Evol 324 316–41 (2015)
443	48	W L Long W W Ballard Normal embryonic stages of the longnose gar Lepisosteus
443	40.	RMC Dev Riol 1 1–8 (2001)
445	49	Ω L α Giudice <i>et al.</i> Single-cell transcriptional logic of cell-fate specification and axon
446	12.	guidance in early-born retinal neurons <i>Development</i> 146 dev178103 (2019)
AA7	50	C García-Frigola M I Carreres C Vegar C A Mason E Herrera Zic2 promotes
/ 448	50.	avonal divergence at the optic chiasm midline by EnbB1_dependent and _independent
449		mechanisms Development 135 1833–1841 (2008)
450	51	I Aruga M Hatayama Comparative genomics of the Zic family genes Adv Ern
450	51.	Med Biol 1046 3_26 (2018)
452	52	A I Dell F Fried-Cassoria H Xu I A Raper cAMP-Induced Expression of
452	52.	Neuropilin Promotes Retinal Axon Crossing in the Zebrafish Ontic Chiasm I
454		Neurosci 33 11076–11088 (2013)
455	53	C Gebhardt <i>et al</i> An interhemispheric neural circuit allowing binocular integration in
456	55.	the optic tectum Nat Commun 10 5471 (2019)
457	54	C García-Frigola E Herrera Zic2 regulates the expression of Sert to modulate eve-
458	57.	specific refinement at the visual targets <i>FMBO I</i> 29 3170–3183 (2010)
450	55	M A MacIver I Schmitz II Mugan T D Murphey C D Mobley Massive
457	55.	increase in visual range preceded the origin of terrestrial vertebrates <i>Proc</i> Natl Acad
461		Sci 114 F2375_F2384 (2017)
462	56	G Laurent On the value of model diversity in neuroscience Nat Rev Neurosci 21
463	50.	$395_{396} (2020)$
464	57	S Hergueta <i>et al</i> Overlapping visual fields and insilateral retinal projections in turtles
465	57.	Brain Res Rull 29 427_433 (1992)
466	58	C S Goodman The likeness of being: Phylogenetically conserved molecular
467	50.	mechanisms of growth cone guidance <i>Cell</i> 78 353_356 (1994)
468	59	P Heger W Zheng A Rottmann K A Panfilio T Wiehe The genetic factors of
469	57.	hilaterian evolution <i>Flife</i> 9 (2020) doi:10.7554/eI ife 45530
470	60	E Friocourt et al Recurrent DCC gene losses during bird evolution Sci Ren 7 37560
<u>4</u> 70 <u>4</u> 71	00.	(2017)
472	61	P Zelina et al Signaling Switch of the Axon Guidance Recentor Robo3 during
т/ 🚄	U1 .	i · Zenna er ar, orgnannig ownen of the rivon Outdance Receptor Robos during

473		Vertebrate Evolution, <i>Neuron</i> , 84 , 1258–1272 (2014).	
474	62.	W. Pak, R. Hindges, YS. S. Lim, S. L. Pfaff, D. D. M. M. O'leary, Magnitude of	
475		Binocular Vision Controlled by Islet-2 Repression of a Genetic Program that Specifies	
476		Laterality of Retinal Axon Pathfinding. Cell. 119, 567–578 (2004).	
477	63.	T. Kuwajima, C. A. Soares, A. A. Sitko, V. Lefebvre, C. Mason, SoxC Transcription	
478		Factors Promote Contralateral Retinal Ganglion Cell Differentiation and Axon	
479		Guidance in the Mouse Visual System. Neuron. 93, 1110-1125.e5 (2017).	
480	64.	C. Morenilla-Palao et al., A Zic2-regulated switch in a noncanonical Wnt/Bcatenin	
481		pathway is essential for the formation of bilateral circuits. Sci. Adv. 6 (2020),	
482		doi:10.1126/sciadv.aaz8797.	
483	65.	I. Braasch et al., Connectivity of vertebrate genomes: Paired-related homeobox (Prrx)	
484		genes in spotted gar, basal teleosts, and tetrapods. Comp. Biochem. Physiol. Part C	
485		<i>Toxicol. Pharmacol.</i> 163 , 24–36 (2014).	
486	66.	V. Marillat et al., Spatiotemporal expression patterns of slit and robo genes in the rat	
487		brain. J. Comp. Neurol. 442, 130–155 (2002).	
488	67.	C. Thisse, B. Thisse, High-resolution in situ hybridization to whole-mount zebrafish	
489		embryos. <i>Nat. Protoc.</i> 3 , 59–69 (2008).	
490 491	68.	V. Bercier et al., Dynactin1 depletion leads to neuromuscular synapse instability and functional abnormalities. <i>Mol. Neurodegener.</i> 14 , 27 (2019).	
492	69.	R. C. Edgar, MUSCLE: Multiple sequence alignment with high accuracy and high	
493		throughput. Nucleic Acids Res. 32, 1792–1797 (2004).	
494	70.	F. Oteri, F. Nadalin, R. Champeimont, A. Carbone, BIS2Analyzer: A server for co-	
495		evolution analysis of conserved protein families. <i>Nucleic Acids Res.</i> 45 , 307–314	
496	- 1	(2017).	
497	71.	D. Santos, S. M. Monteiro, A. Luzio, in <i>Methods in Molecular Biology</i> 1/97, 365–371	
498		(2018).	
499 500			
500			
502	Figur	a lagands	
502	rigui	e regenus	
505	D1 1	• 7• • • • • • • • • • • • • • • • • • •	
504	F1g.1	. Visual projections are only contralateral in clupeocephalan teleosts.	
505	(A) S	implified phylogenetic tree of the major groups of vertebrates. Divergence of each major	
506	group	is displayed in million years. Asterisks indicate whole genome duplication events in the	
507	teleos	teleost (TGD) and sturgeon (AGD) ancestors. (B) Whole brain visualization of a juvenile	
508	zebrafish injected with an AlexaFluor-555-conjugated CTb (left eye) and AlexaFluor-647-		
509	conju	gated CTb (right eye) highlighting complete contralateral projections. (C) High	
510	magn	ification showing pre-tectal nuclei. (D to M) 3D rendering of visual projections labelled	
511	by in	jecting AlexaFluor-555-conjugated CTb (left eye) and AlexaFluor-647-conjugated CTb	
512	(right	eye) followed by iDISCO whole-brain clearing and 3D imaging using light-sheet	

fluorescence microscopy. (D and E) Mexican tetra. (F and G) Redeye piranha. (H and I) Greenspotted pufferfish. (J and K) Four-eyed fish. (L and M) Atlantic mudskipper. In all species,
visual axons only project to the brain on the contralateral side. Abbreviations: A, anterior; Cb,
Cerebellum; Ctb, Cholera toxin B; D, dorsal; OB, Olfactory bulb; ON, Optic nerve; P, posterior;
V, ventral; Sil, Silurian. Scale bars are 500 µm in (B) and (D) to (M) and 200 µm in (C).

518

519 Fig. 2. Bilateral visual projections in basal ray-finned fishes and lobe-finned fishes

520 (A to S) 3D light-sheet fluorescence microscopy images of iDISCO+ cleared brains from fish 521 injected into the eyes with 2 cholera toxins. The left panels show only one channel. (C, F, J, N 522 and S) are optical sections through the brain region receiving bilateral visual inputs. (I and M 523 and **R**) high magnification from whole-mount brains. (A to C) Butterflyfish. (D to F) Spotted 524 gar. (G to J) Sterlet sturgeon. (K to N) Armored bichir. (O to S) Australian lungfish. (C, F, J 525 and N) In all fishes, contralateral and ipsilateral projections segregate in two different optic 526 tectum (OT) layers except in the lungfish (S) where they are intermingled in the OT. 527 Arrowheads indicate ipsilateral projections. Abbreviations: Cb, Cerebellum; OB, Olfactory 528 bulb; OE, Olfactory epithelium; ON, Optic nerve; OC, Optic chiasm; pT, pretectal nuclei; T, 529 Optic tract. Scale bars are 500 µm in (A, B, D, E, G, H, K, L, P and Q) and 80 µm in (C, G, K, 530 N and S) and 100 μ m in (M, O) and 150 μ m in (I, R).

531

532 Fig. 3. Zic2 is expressed in the temporal retina of mammal embryos

(A to C) Whole-mount immunohistochemistry of an E16 mouse eye labeled with the pan ganglion cell marker RBPMS and the ipsilateral ganglion cell marker Zic2. (A) frontal view and (B) top view. (C) Optical section at the level indicated by the dashed line in (C). (D to F), 3D light-sheet fluorescence microscopy of pcw9 human embryonic eye cleared using EyeDISCO and labeled for RBPMS and ZIC2. (D) frontal view and (E) top view. (F) Optical 538 section at the level indicated by the dashed line in (C). G, H, I indicate the approximate positions 539 of images in panels G-I. (G to I) retinal cryosections of a pcw9 human embryo eye labeled with 540 ZIC2 and ISLET1 at 3 different levels: temporal and close to retinal outer limit (G), temporo-541 medial (H) and nasal (I). ZIC2 cells are in the ganglion cell layer (GCL) and co-express 542 ISLET1. They are absent from the neuroblastic layer (NBL). (J) retinal cryosection of a pcw9 543 human embryo eye labeled with ZIC2 and SOX2. ZIC2 is absent from the NBL which contains 544 SOX2+ progenitors. (K and L) retinal cryosections of a pcw14 human embryo eye labeled with 545 ZIC2, SOX2 and ISLET1. ZIC2 is only present in ISLET1+ ganglion cells in the temporal 546 retina and absent from SOX2+ progenitors (K). Abbreviations: D, dorsal; V, ventral; N, nasal; 547 T, temporal ; ON, optic nerve. Scale bars are 70 µm in (A and B) and 20 µm in (C) and 300 µm 548 in (D and E) and 50 µm in (F to L).

549

550 Fig. 4. Zic2 is not expressed by ganglion cells in spotted gar and zebrafish

551 (A to F) Development of the visual system in the spotted gar. All images are 3D light-sheet 552 fluorescence microscopy images of EyeDISCO-cleared spotted gar embryos labeled with Islet1 553 and acetylated Tubulin. (A and C and E) Top (dorsal) views of spotted gars at 2-3dpf (A), 6-554 7dpf (C), and 17-18dpf (E). (B and D and F) frontal views of whole spotted gar eyes at 2-3dpf 555 (B), 6-7dpf (D), and 17-18dpf (F). The optic nerve (ON and asterisk) starts to form by 6-7dpf 556 and is well developed by 17-18dpf. The optic chiasm (OC) is formed by 6-7dpf. (G) Coronal 557 cryosection of spotted gar embryos at 17-18dpf labeled for ßIII-tubulin and Islet1. (H to K) 558 cryosection from 17-18dpf spotted gar eyes hybridized with zic2 riboprobe (H and J) and 559 labeled for proliferating cell nuclear antigen (PCNA) and Islet1 (I, K and L). (J to L) higher 560 magnification of the ciliary margin zone (area framed in I). (M to P) 3D rendering of whole-561 brain viewed from the top of zebrafish injected with Tg(atoh7:Gal4,14UASubc:T2A-eGFP-pA) 562 (M and N) or Tg(atoh7:Gal4,14UASubc:ZIC2-T2A-eGFP-pA (O and P). N and P show 563 segmented ganglion cell projections. (P) A large ipsilateral projection (arrowhead) is seen in 564 the Tg(atoh7:Gal4,14UASubc:ZIC2-T2A-eGFP-pA)-injected fish. Abbreviations: ON, Optic 565 nerve; OC, Optic chiasm; GCL, Ganglion cell layer; INL, Inner nuclear layer; ONL, Outer 566 nuclear layer. Scale bars are 50 μ m in (A, D, F, J, K, L, M to P) and 15 μ m in (B) and 80 μ m 567 in (C, H and I) and 150 μ m in (E) and 200 μ m in (G).

568

569 Acknowledgements

570 The authors wish to thank Anais Favre and Ali Abjaghou for assistance with light-sheet 571 microscopy, Fabio Cortesi for help with lungfish experiments, Sylvie Dufour and all members 572 of Alain Chédotal and Del Bene teams for helpful discussions and input during the project. We 573 thank Sylvie Rétaux (Paris Saclay University) for help with Mexican tetra work and helpful 574 comments on the manuscript. We thank Allyse Ferrara and Solomon David (Nicholls State 575 University) and members of the Braasch Lab for help with gar work. Funding: Programme 576 Investissements d'Avenir IHU FOReSIGHT (ANR-18-IAHU-01) (AC, FDB). INSERM cross-577 cutting program HuDeCA 2018 (AC). NIH R010D011116 (IB). UQ Amplify Fellowship (RS). 578 Authors contribution: AC, FDB and RV designed the experiments. FDB, KD, KNBC, JV, 579 SA, RV, PK, RS conducted the experiments. AC, KNBC and FDB obtained funding for the 580 project. IB provided Lepisosteus samples and EH the human ZIC2 cDNA. AC, FDB and RV 581 designed the figures and wrote the manuscript. All authors discussed the results and commented 582 on the manuscript. Competing interests: The authors declare no competing interests. Data and 583 **materials availability:** All data are available in the manuscript or the supplementary materials. 584

585 List of supplementary materials

586	Materials and Methods
587	Figs. S1 to S7
588	Table S1

589 Movies S1 to S5











Supplementary Materials for

Bilateral visual projections exist in non-teleost bony fish and predate the emergence of tetrapods

Robin J. Vigouroux, Karine Duroure, Juliette Vougny, Shahad Albadri, Peter Kozulin, Eloisa Herrera, Kim Nguyen-Ba-Charvet, Ingo Braasch, Rodrigo Suarez, Filippo Del Bene* and Alain Chédotal*

Correspondence to: filippo.del-bene@inserm.fr and alain.chedotal@inserm.fr

This PDF file includes:

Materials and Methods Figs. S1 to S7 Table S1 Captions for Movies S1 to S5

Other Supplementary Materials for this manuscript include the following:

Movies S1 to S5

Material and Methods

<u>Animals</u>

Juvenile Mexican tetra (San Solomon Spring, Balmorhea State Park, Texas, USA) were maintained at 26°C (surface fish) on a 12:12 h light:dark cycle. Juvenile zebrafish and embryos were maintained at 28.5°C on a 14 h light/10 h dark cycle. Juvenile Australian lungfish (10.2-13.5 cm body length; Jardini Pty Ltd, Brisbane, Australia) were on freshwater at 26°C on a 12:12 h light:dark cycle. Juvenile armored bichir, sterlet sturgeon, African butterflyfish, redeye piranha, atlantic mudskipper, green puffer fish and four-eyed fish, were acquired from commercial vendors. Spotted gar embryos were spawned at Nicholls State University in Louisiana and then raised and maintained at Michigan State University as previously described (65). Embryos were raised at 18°C which leads to a comparatively slow progression through the Long & Ballard stages of gar development (*48*). Sizes of each specimen were recorded for future analysis. Juvenile specimens of either sex were used. All animal procedures were performed under the in accordance with protocols approved by Sorbonne Université and Institute (#QBI/041/20/France) and Michigan State University (#10/16-179-00).

Human eye samples

Human fetal eyes from terminated pregnancies were obtained from the INSERM-funded Human Developmental Cell Atlas collection (HuDeCA, https://hudeca.genouest.org/). All tissues were collected with appropriate maternal consent and approval from the French National Biomedicine agency (N° PFS19-012).

In Situ Hybridization

Spotted gar sections were hybridized with digoxigenin-labeled riboprobes as described in (66). Briefly, tissue sections were postfixed for 10 min in 4% paraformaldehyde (PFA) before being treated with Proteinase K (10 µg/ml; Invitrogen, #03115852001) for 2 min and subsequently postfixed for 5 min in 4% PFA. Sections were then acetylated and permeabilized in PBS, 1% Triton X-100. Sections were first homogenized with hybridization buffer (50% formamide (VWR #24311.291), 5× SSC (Euromedex, #EU0300-A), 1× Denhardt's, 250 µg/ml yeast tRNA, and 500 µg/ml herring sperm DNA, pH 7.4) for 2 h at RT and then hybridized overnight at 72°C with riboprobes (1/200), see Table S1 for probe sequences. The next day, sections were rinsed for 2 h in 2× SSC at 72°C, and blocked in 0.1 M Tris, pH 7.5, 0.15 M NaCl (B1) containing 10% normal goat serum (NGS) for 1 h at RT. After blocking, slides were incubated o/n at 4°C with anti-DIG antibody conjugated with the alkaline phosphatase (1/5000, Roche Diagnostics) or anti-DIG antibody conjugated with peroxidase in B1 containing 1% NGS. After washing in B1 buffer, the alkaline phosphatase activity was detected by using nitroblue tetrazolium chloride (337.5 µg/ml) and 5-bromo-4-chloro-3-indolyl phosphate (175 µg/ml) (Roche Diagnostics). The peroxidase activity was detected by using Tyramide Signal Amplification (TSA) (PerkinElmer, #NEL741001KT) and incubated with Fluorescein fluorophore Tyramide diluted at 1:50 in TSA. Sections were mounted in Mowiol (Calbiochem/Merck, Carlstadt, Germany).

Whole-mount in situ hybridization were carried out on zebrafish as previously described (67). Embryos were then embedded in gelatin/albumin with 4% of glutaraldehyde and sectioned (20 μ m) on a VT1000 S vibrating blade microtome (Leica). Slides were scanned with either a Nanozoomer (Hamamatsu) or laser scanning confocal microscope (Olympus, FV1000).

Fluorescent in Situ Hybridization

To generate anti-sense probes, DNA fragments were obtained by PCR using Phusion™ High-Fidelity DNA polymerase (Thermo Scientific, #F530L) with the primers listed in Table S1. Total cDNA from 1 to 5 dpf zebrafish were used as a template. PCR fragments were cloned into the pCRII-TOPO vector (Invitrogen, #K280040) according to manufacturer's instructions. All plasmids used were sequenced for confirmation. Anti-sense DIG or fluorescein-labeled riboprobes were in vitro transcribed using the RNA labeling kit (Roche, #11685619910 or #11277073910) according to manufacturer's instructions. De-chorionated embryos at the appropriate developmental stages were fixed in fresh 4% PFA in 1X PBS (pH7.4) containing 0.1% Tween20 (PBSTw) for 4 h at RT and stored o/n in 100% methanol. Embryos were rehydrated by immersing them in subsequent baths of 50% methanol/PBSTw (Sigma, #34860) and then twice in PBSTw, baths followed by a 10 min incubation in a 3% H₂O₂/0.5%KOH (Sigma, #P5958) solution. Embryos were then rinsed in 50% methanol and post-fixed in 100% methanol at -20°C for 2 h. Embryos were then re-hydrated in methanol/PBSTw (75%/50%/25%) followed by treatment in 10 µg/ml proteinase K at RT (1 dpf = 5 min, 2 dpf = 15 min, 3 dpf = 20 min), and post-fixed for 20 min in 4% PFA in PBSTw. Embryos were pre-hybridized at 68°C, and hybridized with either a fluorescein-labelled probe or DIGlabelled probe or both probes for dFISH assays o/n at 68°C with gentle shaking. Embryos were then rinsed at 68°C in 50% formamide/2XSSC/0.1%Tween-20 twice, 2XSSC/0.1%Tween-20, 0.2XSSC/0.1% Tween-20 twice and finally in TNT buffer (0.1 M Tris pH7.5, 0.15 M NaCl, 0.1% Tween-20). Blocking was done in TNB buffer (2% DIG block (Roche, #11096176001) in TNT) for 2 h at RT and incubated o/n with anti-Fluo-Fab-POD (Roche, #11426346910) diluted at 1:50 in TNB buffer at 4°C. All steps were performed in the dark. Embryos were then washed several times in TNT, rinsed using $100 \,\mu$ l Tyramide Signal Amplification (TSA) (PerkinElmer, #NEL741001KT) and incubated with Fluorescein fluorophore Tyramide diluted at 1:50 in TSA.

The reaction was stopped by 5 rapid washes of TNT. For dFISH assays, the DIG-labelled probe was then revealed by carrying out a 20 min incubation in 1%H₂O₂/TNT (Sigma, #18312-1L), then washed several times in TNT. A second blocking step was carried out for 1 h in TNB buffer prior to incubating embryos in anti-DIG-POD (Roche, #11207733910) diluted at 1:100 in TNB buffer o/n at 4°C. Revelation was done with Cy3 Fluorophore Tyramide solution (PerkinElmer, NEL#744001KT), washed with TNT and processed for imaging upon DAPI staining.

Molecular cloning

14xUAS:ubc-ZIC2-T2A-GFP-pA or *14xUAS:ubc-T2A-GFP-pA* were obtained via Gibson assembly using the *pT1UciMP Tol1* (Addgene, #62215) destination vector described by (68). *Tol1*mRNA was synthesized from the plasmid (Addgene, #61388) digested by NotI (NEB, #R3189S) and retro-transcribed with SP6 RNA polymerase (Roche, #10810274001). Human ZIC2 (*hZIC2*), *GFP*, and *T2A* were amplified via PCR from pCAG-hZIC2 and *pUAS:Cas9T2AGFP;U6:sgRNA1;U6sgRNA2* (Addgene, #74009) respectively using the NEBuilder HiFi DNA Assembly Cloning kit (NEB, #E5520). Appropriate sequences were inserted after the *UBC* intron of the *pT1UciMP Tol1* destination vector opened by restriction digest with NcoI-HF (NEB).

Alignment between the amino acid sequences of the Zic2 proteins zing finger domains

A multiple sequence alignment was performed for the region covering the ZIC2 zinc finger domains of NCBI Reference Sequence proteins of mouse (NP_033600), human (NP_009060) and zebrafish (ZIC2a NP_571633, ZIC2b NP_001001820), as well as for genome-predicted ZIC2 proteins the spotted gar (XP_006638968). The UniProtKB/Swiss-Prot curated zinc finger sequences of human ZIC2 (O95409) were used to delineate the domain positions within the

alignment. Protein sequence alignment was performed using MUSCLE version 3.8.31 (69), the amino acid conservation at each aligned position visualised using BIS2Analyzer (70).

Eye enucleation

The transgenic line Tg(atoh7:gal4-vp16) (RRID: ZFIN_ZDB-GENO-130306-1) was used. Prior to eye enucleation, fish were selected for the atoh7 expression in green. At 2 dpf, eye enucleation was performed. The embryos were anesthetized in 0.004% tricaine MS222 in a 2% agarose gel solution (Life technologies, #16520050). One eye was surgically removed using a pulled capillary and mouth pipetting. Embryos were then transferred into fish medium (egg medium with penicillin/streptomycin (Life Technologies, #15140122) and 0.003% 1-phenyl-2thiouera (Sigma, #189235) until 5 dpf, for whole-mount immunohistochemistry.

Immunohistochemistry

Cryosections

Spotted gar embryos were fixed by immersion in 4% PFA in 0.12 M phosphate buffer (VWR, 28028.298 and 28015.294), pH 7.4 (PFA) o/n at 4°C. Following three washes in 1XPBS, the samples were incubated in 10% sucrose (VWR, 27478.296) in 0.12 M phosphate buffer o/n at 4°C. The next day, samples were transferred to a 30% sucrose solution in 0.12 M phosphate buffer o/n at 4°C. Samples were then embedded in 0.12 M phosphate containing 7.5% gelatin (Sigma, 62500) and 10% sucrose, frozen in isopentane at -40°C and then cut at 16 µm with a cryostat (Leica, CM3050S). Sections were blocked in PBS containing 0.2% gelatin (VWR) and 0.25% Triton-X100 (PBS-GT) for 1 h at RT. Following the blocking, sections were incubated with primary antibodies (see Table S1) diluted in a PBS-GT solution o/n at RT. Following three washes in PBST (0.05% Trinton-X100) secondary antibodies coupled to the appropriate fluorophore (see Table S1) were diluted in PBS-GT and incubated for 2 h at RT. Sections were

counterstained with Hoechst (Sigma, B2883, 1:1000) or DAPI (Life Technologies, D3571, 1/500). For PCNA staining, an antigen retrieval step was performed by boiling sections in a 1X Sodium Citrate solution pH 6.0 for 5 min using a microwave. This step was skipped when the samples were first used for an *in-situ* hybridization assay. Slides were scanned with either a Nanozoomer (Hamamatsu) or laser scanning confocal microscope (Olympus, FV1000).

Whole-mount Immunohistochemistry

Zebrafish whole-mount immunohistochemistry was adapted from (*61*). Briefly, embryos were fixed in 4% PFA diluted in PBS containing 0.1% Tween-20 (VWR, #0777-1L) (PBSTw) for 4 h at RT and stored o/n in 100% methanol. After re-hydration, embryos were incubated for 20 min at -20°C in already pre-chilled acetone (Sigma, #650501). The embryos were rinsed several times with PBSTw and blocked for 2 h in blocking solution (10% bovine serum albumin (BSA) (Euromedex, #04-100-812-C), 10% normal goat serum (LifeTechnologies, #1000C), 1% DMSO (Sigma Aldrich, #D8418) in PBSTw). The primary antibodies were incubated o/n at 4°C in 1% BSA, 1% normal goat serum, 0.1% DMSO in PBSTw according to the dilutions in Table S1. After several washes in PBSTw, the secondary antibodies were incubated o/n at 4°C. The next day, embryos were rinsed in PBSTw and processed for imaging.

Whole-mount immunostaining on spotted gar embryos was carried out as previously described (*15*). Briefly, embryos were depigmented in a solution of 11% H₂O₂ (VWR, 216763) at 70 rpm exposed to an 11W warm white Light-Emitting Diode (LED) (3000° Kelvin) for 1-3 days. Samples were then blocked and permeabilized before being incubated with the primary antibodies for 7 days at RT (see Table S1) in a solution containing: 0.5% Tirton-X100, 5% donkey normal serum, 20% Dimethyl Sulfoxide, 1XPBS, 0.1 g/L thimerosal. The samples were further labeled with secondary antibodies (see Table S1) for 2 days at RT under agitation.

Retinal flat-mounts

For retinal flat mounts, human eyes were harvested and fixed in 4% PFA, followed by three

washes in 1XPBS. Eyes were then de-pigmented using the EyeDISCO protocol as previously described (*15*). For immunohistochemistry, retinas were permeabilized and blocked in a solution containing 0.5% Triton-X100, 5% donkey normal serum, 1XPBS, 0.1 g/L thimerosal for 1 day at RT under agitation. Primary antibodies (see Table S1) were diluted in a solution containing 0.5% Triton-X100, 5% donkey normal serum, 20% Dimethyl Sulfoxide, 1XPBS, 0.1 g/L thimerosal for 3 days at RT under agitation. The retinas were then washed for 1 day in PBST (1XPBS, 0.5% Triton-X100). The secondary antibodies (see Table S1) were diluted in the same solution as primary antibodies and left for 2 days. After washing retinas for 1 day, they were mounted on slides and imaged using a scanning confocal microscope (Olympus, FV1000).

Tracing of visual projections

All fish were anesthetized with 0,04% MS222, tricaine-methanesulfonate (Sigma, #E10521) diluted in fish water. Australian lungfish were anesthetized with 0.05% clove oil in fresh water. Injection of cholera toxin β subunit was carried out as described in (*15*). Briefly, using a capillary approximately 1µl of 2 µg/µl of Alexa Fluor-conjugated cholera toxin β subunit (Thermo Fischer, Alexa Fluor555-CTb C22843 and Alexa Fluor647-CTb C34778) was injected intravitreally. 72-96 h following CTb injection, specimens were transcardially perfused with 4%PFA and the heads and/or brains were dissected for tissue clearing.

Tissue clearing and imaging

<u>Clearing</u>

Prior to clearing, spotted gar *embryos* were embedded in 1.5% agarose (Roth) in 1X TAE (Life Technologies). Clearing was carried out as previously described (*15*). Briefly, samples were gently de-hydrated in ascending baths of methanol (1.5 h). Samples were further treated with a

solution containing 2/3 Dichloromethane (DCM, Sigma) 1/3 methanol o/n. The next day, samples were placed in DCM for 30 min prior to being immersed in Di-benzyl Ether (DBE, Sigma).

Imaging

Acquisitions were performed by using an UltraMicroscope I (Miltenyi Biotec, Germany) or UltraMicroscope Blaze (Miltenyi Biotec, Germany) with the ImspectorPro software (Miltenyi Biotec, Germany, 5.1.328 version). The light sheet was generated by a laser (wavelength 488, 561, 647 Coherent Sapphire Laser, LaVision BioTec, Miltenyi Biotec, Germany) or a secondgeneration laser beam combiner (wavelengths 488 nm, 561 nm and 647 nm; LaVision BioTec, Miltenyi Biotec, Germany). All light sheets were matched within their Rayleigh lengths for optimal illumination at the sample site. Either a binocular stereomicroscope (Olympus, MXV10) with a 2x objective (Olympus, MVPLAPO) was used Or a MI Plan 1.1x (NA = 0.1), a MI Plan 4x (NA = 0.35), and a MI Plan 12x (NA = 0.53) objectives were used (Miltenyi Biotec, Germany). Samples were placed in an imaging reservoir made of 100% quartz (LaVision BioTec, Miltenyi Biotec) filled with DBE and illuminated from the side by the laser light. A Zyla sCMOS camera (Andor, Oxford Instrument; 2,048 × 2,048, 6.5 x 6.5 μ m, peak QE 82%) was used to acquire images. The step size between each image was fixed at 1 or 2 μ m (NA = 0.5, 150 ms time exposure). All tiff images are generated in 16-bit.

Confocal microscopy

Whole-mount 5 dpf zebrafish larvae were mounted in a labtex plates (LabTex) in 2.5% agarose or in 1% low-melting agarose on FluoroDish Cell Culture dish (FD3510-100, World Precision Instruments). For imaging, a scanning inverted confocal microscope (FV1200, Olympus) was used with a 30x objective (Olympus, UPLSAPO30XS, NA = 1.05, WD = 0.8 mm) as well as the LSM780 and LSM880 scanning inverted confocal microscopes (Zeiss) for high resolution microscopy. 40x water immersion objective for whole mount dFISH stained zebrafish embryos

and 63x oil objective for zebrafish retinal cryosections were used and a 10x air objective was used to image the spotted gar cryosections.

Image Processing

3D rendering of light sheet and confocal stacks were converted to an Imaris file (.ims) using ImarisFileConverter (Bitplane, 9.5.1 version) and then visualized using the Imaris x64 software (Bitplane, 9.5.1). To quantify ipsilateral territories, entire tectum volume and ipsilateral projections were automatically segmented with a surface detail of $5.00 \,\mu\text{m}$, automatic threshold. Volumes were extracted from the surface. Movies were generated using the animation tool on Imaris x64 software (Bitplane, version 9.1.2) and movie reconstruction with .tiff series were done using ImageJ (1.50e, Java 1.8.0_60, 64-bit). All movie editing (text and transitions) was performed using iMovie (Apple Inc., version 10.1.1).

To quantify the ipsilateral projections in the hZIC2 overexpression experiments, a fixed region of interest was identified for each zebrafish (corresponding to the ipsilateral and contralateral optic tecta). Retinal projections were segmented with a surface detail of $0.5 \,\mu$ m using an automatic threshold. Ipsilateral and contralateral volumes were extracted and summed to constitute the "total visual projections" using Imaris x64 software (Bitplane, version 9.1.2). The volume of ipsilateral projections was isolated as a ratio of ipsilateral projections:total projections.

Statistical analyses

All data are described are listed as biological replicates (n) and all experiments (N) were carried out at least in triplicates unless indicated otherwise). An observer blinded to the experimental conditions realized all the quantifications. No data were excluded from the statistical analyses. All data are represented as mean values ± SEM. Statistical significance was estimated using two-tailed unpaired tests for non-parametric tendencies (Kruskall-Wallis or Mann-Whitney), two-way ANOVA and Bonferroni's multiple comparison test. * = p < 0.05; ** = p < 0.01; *** = p < 0.001, *** = p < 0.0001. All statistical measurements were carried out using GraphPad Prism 7.

Supplementary Figures



Fig. S1. Simplified chart of fish taxonomy indicating the species analyzed in this study.

Fish with bilateral/ipsilateral visual projections appear in magenta and fish with only contralateral visual projections appear in grey.


Fig. S2. Two types of optic nerve crossing modalities in ray-finned fishes.

Ventral views (A to D) and coronal optical sections (E to H) at the level of the optic chiasm of iDISCOcleared brains and optic nerves. A surface rendering with normal shading (Imaris) was applied to generate the ventral view images. The arrowheads (A to D) indicate the level of the chiasm optical section in (E to H). In all fishes, one eye was injected with Alexa Fluor-555-conjugated CTb and the other one with Alexa Fluor-647-conjugated CTb. The right (R) and left (L) optic nerves were pseudocolored in magenta and green respectively. In Pufferfish (A and E), Four-eyed (B and F) and Muskipper (C and G), the two optic nerves pass over and overlap at the chiasm but remain separated up to the brain. By contrast, in the Armored bichir (D and H), the right and left nerves meet at the chiasm and retinal ganglion cell axons from both eyes interweave during crossing (asterisk). Abbreviations: Ant, anterior; Post, Posterior. Scale bars are: 2 mm in (B), 1 mm in (A, C, D, F), 800 μ m in (G), 600 μ m in (E, H).



Fig. S3. ZIC2 expression pattern in human embryo retina.

(A) Box and whiskers representation of the ZIC2-positive surface in E16 mouse and pcw9 human retinas. (**B** and **C**) flat-mount pcw9 human retina labeled for ZIC2 and RBPMS (B) or SOX2 (C). (B) In the most superficial (basal side) regions of the temporal retina, ganglion cells expressing low levels of ZIC2 and RBPMS (arrowheads) are seen but ZIC2 and RBPMS are mostly exclusive. (C) image at the level of the interface between the neuroblastic layer showing that ZIC2+ cells are not SOX2+. Scale bars are: 50 μ m in (B and C).



Fig. S4. Zic2 ortholog expression in zebrafish is restricted to the ciliary margin zone. (A to H) Whole-mount *in situ* hybridization of zebrafish embryos for *zic2a* at 1 day post fertilization (1 dpf; A), 2 dpf (B) and 3 dpf (C) as well as *zic2b* at 1 dpf (E), 2 dpf (F) and 3 dpf (G). Zic2b is expressed in the ciliary marginal zone (CMZ, arrowheads in F and G) and in the dorsal diencephalon (Di). (D) Schematic drawing of the zebrafish CMZ in the developing retina showing spatial distribution of stem cells, cycling progenitors, committed progenitors and differentiated neurons. (H), Lateral view of whole-mount double fluorescent in situ hybridization for zic2b and atoh7 on 3 dpf zebrafish embryos with DAPI counterstaining. (I to L) Confocal sections through the central retina of wild-type embryos hybridized with antisense RNA probes for *zic2b* and *atoh7*. At 24 hpf, *zic2b* is expressed in the entire proliferative neuroepithelium and later from a central to peripheral wave-like manner (arrowheads) in complementarity to the neurogenic transient expression of *atoh7* (asterisks) as shown here for 36, 48 and 72 hpf. (L to M) Confocal sections through the central retina of 72 hpf wild-type zebrafish embryos hybridized with antisense RNA probes for *zic2b* and *retinal homeo- box transcription factor2 (rx2, contexperimentation)* a marker or dividing progenitors and stem cells in the CMZ). Zic2b expression overlaps with the expression of the rx2 (arrowheads). All retinae were counterstained with the nuclear marker DAPI. Scale bars are 50 μ m (A to C and E to H) and 40 μ m (I to M).



Fig. S5. Mammalian ipsilateral markers are not expressed in the spotted gar visual system.

(A) 3D light-sheet fluorescence microscopy images of iDISCO-cleared 17-18 dpf spotted gar indicating with dotted lines the anatomical levels of the cryosections. (**B** to **D**) *In situ* hybridization for *zic2* on retinal cryosections of the developing spotted gar at 2-3 dpf (B), 6-7 dpf (C, left panel), 17-18 dpf (D). Only proliferating cells in the neuroblastic layer (NBL) express *zic2*. The right panel in (C) is an immunostaining for PCNA and Islet1. The arrowheads in (C) indicate the region where the first ganglion cells (Islet1+) are present at this stage in the retina. *zic2* is also found in the diencephalon (Di). (**E** to **T**) Rostral-to-caudal coronal cryosections from 17-18 dpf spotted gar. *zic1* (E to H) and *zic5* (I to L) are only expressed in the ciliary marginal zone (CMZ; arrow). (**M** to **P**) *ephB1* is absent from the retina and weakly expressed in the CMZ. (**Q** to **T**) *ephrinB2* is expressed in the dorsal retina (arrow). (**U** to **W**) Cryosections of the diencephalon of a 17-18dpf spotted gar hybridized for *ephrinB2*. *ephrinB2* is absent from the optic chiasm (asterisk). Immuno-reactive regions are highlighted (arrowhead). Abbreviations: NBL, Neuroblastic layer; ON, Optic nerve, OC, Optic chiasm; GCL, Ganglion cell layer; INL, Inner nuclear layer; ONL, Outer nuclear layer. Scale bars: A, 200 µm; B to D, 50 µm; C,U to W, 100 µm; D to T, 250 µm.



Fig. S6. Alignment between the amino acid sequences of the Zic2 protein zing finger domains of fish and mammals.

Alignment of Zic proteins across the zinc finger domains shows the high level of conservation between ray-finned fish and mammals. Amino acids are color coded according to the physiochemical class they belong to. Abbreviations: HU, human; MM, mouse; SG, spotted gar; ZE, zebrafish.



Tg(atoh7:gal4), p14UASubc:T2A-eGFP

Fig.S7. Ectopic *ZIC2* expression in *atoh7* retinal progenitor cells does not induce EphB1 expression in retinal ganglion cells.

Confocal images of cryostat (A-F) or optical (G-L) sections of 3 dpf retinae from Tg(atoh7:gal4) embryos injected at 1-cell stage with either a *p14UASubc:T2A-eGFP* (A to C and G to I) control construct or a *p14UASubc:ZIC2-T2A-eGFP* construct (D to F and J to L). No signal for EphB1 in GFP-positive cells is detectable in all injected retinae either double stained with anti-EphB1 antibody (A to F) or hybridized with an *ephB1* antisense riboprobe (G to L). All retinae were counterstained with the nuclear marker DAPI. Scale bars are 50 µm.

			In situ probes						
Name	Sequence			RRID	Dilution	In situ hybridization			
L-zic1 fwd	ACCTCCAGACATCACTCAAC			n/a	1:200	Cryosections			
L-zic1 rev	GGAACAC	GGAACACTCTTCCCAGAAAC			1:200	Cryosections			
L-zic2 fwd	AAACTTAACCACGACCTCTCTC			n/a	1:200	Cryosections			
L-zic2 rev	CTCGTGCATTGTGCTGAAAG			n/a	1:200	Cryosections			
L-zic5 fwd	CTTTGAGCAAGAGGAATCCGGC			n/a	1:200	Cryosections			
L-zic5 rev	CCTGCCG	CGATGTTCACA	TTTA	n/a	1:200	Cryosections			
L-efnb2 fwd	TCCCCATI	TATGAGAAGGT	GAGCGG	n/a	1:200	Cryosections			
L-efnb2 rev	ACAGGCTA	ACCACTTCAGA	AGGCAG	n/a	1:200	Cryosections			
L-ephb1 fwd	AGAACCTO	GAACACAATCO	CGCAC	n/a	1:200	Cryosections			
L-ephb1 rev	ACAGTITA	ATGGGCACGT	CCAC	n/a	1:200	Cryosections			
zf-zic2a fwd	ACAACAA	ICTGTCGCCTT	CCTC	n/a	1:200	whole-mount			
zf-zic2a rev	ACAAATGCCCCTGTTTAGCCC			n/a	1:200	whole-mount			
zf-zic2b fwd	TCTTCCGCTACATGCGACAAC			n/a	1:200	whole-mount			
zf-zic2b rev	GCAACACCGACATGCTGAGAAC			n/a	1:200	whole-mount			
zf-ephb1 fwd	CGCGTGTGGATGGATTACGG			n/a	1:200	whole-mount			
zf-ephb1 rev	CATCCCCACCAGCTGGATCA			n/a	1:200	whole-mount			
zf-atoh7 fwd	GGAGAAG	GGAGAAGTTTGAGAGTGCTATGCGG			1:200	whole-mount			
zf-atoh7 rev	CGACTITIC	CGACTTTGAGCTGAGCACACACC			1:200	whole-mount			
zf-rx2 fwd	GATACCAT	IGAACATGGTG	GACGATGG	n/a	1:200	whole-mount			
zf-rx2 fwd	CCATCGAC	CTGAATGTGCT	CCTTGG	n/a	1:200	whole-mount			
			Primary antibodi	es					
Antigen	Species	Catalog #	Company	RRID	Dilution	Immunohistochemistry			
Islet1	Rabbit	GTX128201	GeneTex	Ab_2868422	1:300	Cryosections/whole-mount			
Acetylated-tubulin	Mouse	T6793	Sigma	Ab_477585	1:300	Cryosections/whole-mount			
PCNA	Mouse	P8825	Sigma	Ab_477413	1:500	Cryosections			
Islet1+2	Mouse	39.4D5	DSHB	Ab_2314683	1:50	Cryosections			
GFP	Chicken	GTX13970	GeneTex	Ab_371416	1:5000	whole-mount			
Rbpms	Guinea Pig	ABN1376	Millipore	Ab_2687403	1:400	Cryosections/flat-mount/ whole-			
Zic2	Rabbit	Ab150404	Abcam	Ab_2868423	1:300	Cryosections/flat-mount/ whole-			
Sox2	Goat	Sc17320	Santa-Cruz	Ab_2286684	1:300	flat-mount			
EphB1	Mouse	MAb EfB1-3	DSBH	Ab_2314357	1:5	Cryosection			
			Secondary antibod	lies					
Anti-Rabbit cy3	Donkey	711-165-152	Jackson	Ab 2307443	1:500	cryosections/whole-mount			
Anti Dabbit Alava Eluar	Dankay	711 605 152	ImmunoResearch			amagastions/Elst mount/whole			
647	Donkey	711-005-152	ImmunoResearch	Ab_2492288	1:500	mount			
Anti-Goat Alexa Fluor 488	Donkey	A11055	Life Technologies	Ab 2534102	1.500	cryosections/Flat-mount/whole-			
	-			A0_2334102	1.500	mount			
Anti-Goat Alexa Fluor 555	Donkey	A21432	Life Technologies	Ab_2535853	1:500	cryosections/Flat-mount/whole-			
Anti-Goat Alexa Fluor 647	Bovine	805-605-180	Jackson			cryosections			
			ImmunoResearch	AB_2340885	1:600	5			
Anti-Goat cy3	Donkey	705-165-147	Jackson	Ab 2307351	1:500	cryosections/Flat-mount/whole-			
Anti mouse Alexa Eluor	Donkey	A21202	ImmunoResearch	_		mount			
488	Donkey	A21202	Life recimologies	Ab_141607	1:500	mount			
Anti-Guinea-Pig Alexa cy3	Donkey	706-165-148	Jackson	Ab 2340460	1.500	cryosections/Flat-mount/whole-			
			ImmunoResearch	A0_2340400	1.500	mount			
Anti-mouse Alexa Fluor	Donkey	715-605-150	Jackson Jackson	Ab_2340862	1:500	cryosections/Flat-mount/whole-			
Anti-Mouse, Alexa Fluor	Goat	A31574	Life Technologies			cryosections			
635	-		8	Ab_2536184	1:500	5			
Anti-Rabbit, Alexa Fluor	Goat	A11036	Life Technologies	Ab 10563566	1:500	cryosections			
568 Anti Mouso, Alava Eluor	Cont	411004	Life Technologies	-		orrespections			
568	Goal	AI 1004	Life fectiliologies	Ab_2534072	1:500	cryosections			
Alexa Fluor 488 anti-	Goat	A11039	Life Technologies	Ab 142024	1.500	cryosections			
chicken			-	A0_142924	1:500				
Tracers									
Cholera toxin subunit B-	n/a	C22843	Life technologies	n/a	$2 \ \mu g/\mu l$	Whole-mount			
Cholera toxin subunit R-			-						
AlexaFluor647	n/a	C34778	Life technologies	n/a	$2 \ \mu g/\mu l$	Whole-mount			

Table S1. Comprehensive table summarizing the antibodies and probes sequences.

Movie S1.

Visual projections in teleosts.

Whole brain rendering of visual projections in 5 teleosts, the zebrafish, Mexican tetra, greenspotted pufferfish, mudskipper and butterflyfish. All species shows a complete decussation of retinal projections except the butterflyfish. All fish had bilateral eye injections of CTb coupled to either an Alexa Fluor-555 or and Alexa Fluor-647.

Movie S2.

Bilateral visual projections in non teleosts.

Whole brain rendering of visual projections in spotted gar, sterlet and armored bichir. Ipsilateral projections are seen in all species observed. All fish had bilateral eye injections of CTb coupled to either an Alexa Fluor-555 or and Alexa Fluor-647.

Movie S3.

The Australian lungfish possesses non-segregated ipsilateral projections.

Whole brain rendering of visual projections in the Australian lungfish, a sarcopterygian, injected with either an Alexa Fluor-555 or an Alexa Fluor-647. Many ipsilateral projections are observed, with a major component in the optic tectum. Ipsilateral projections are intermingled with contralateral projections in the optic tectum.

Movie S4.

ZIC2 expression is evolutionarily conserved in Humans.

Whole-mount immunohistochemistry of pcw9 human eyes using EyeDISCO clearing and labeled for the ipsilateral transcription factor ZIC2 (magenta) and the pan-retinal ganglion cell marker RBPMS (green). A large ZIC2-positive region can be seen in the temporal retina.

Movie S5.

Development of the Lepisosteus oculatus visual system.

3D rendering of 2-3 dpf, 6-7 dpf, and 17-18 dpf spotted gar embryos using EyeDISCO clearing and light-sheet fluorescence microscopy. Spotted gar embryos were labeled with the panneuronal marker acetylated tubulin (a-tubulin, green) and the LIM/homeodomain family of transcription factor Islet1, which is critical for the proper specification of retinal ganglion cells and motor neurons (magenta).

Supplementary Materials for

Bilateral visual projections exist in non-teleost bony fish and predate the emergence of tetrapods

Robin J. Vigouroux, Karine Duroure, Juliette Vougny, Shahad Albadri, Peter Kozulin, Eloisa Herrera, Kim Nguyen-Ba-Charvet, Ingo Braasch, Rodrigo Suarez, Filippo Del Bene* and Alain Chédotal*

Correspondence to: filippo.del-bene@inserm.fr and alain.chedotal@inserm.fr

This PDF file includes:

Materials and Methods Figs. S1 to S7 Table S1 Captions for Movies S1 to S5

Other Supplementary Materials for this manuscript include the following:

Movies S1 to S5

Material and Methods

<u>Animals</u>

Juvenile Mexican tetra (San Solomon Spring, Balmorhea State Park, Texas, USA) were maintained at 26°C (surface fish) on a 12:12 h light:dark cycle. Juvenile zebrafish and embryos were maintained at 28.5°C on a 14 h light/10 h dark cycle. Juvenile Australian lungfish (10.2-13.5 cm body length; Jardini Pty Ltd, Brisbane, Australia) were on freshwater at 26°C on a 12:12 h light:dark cycle. Juvenile armored bichir, sterlet sturgeon, African butterflyfish, redeye piranha, atlantic mudskipper, green puffer fish and four-eyed fish, were acquired from commercial vendors. Spotted gar embryos were spawned at Nicholls State University in Louisiana and then raised and maintained at Michigan State University as previously described (65). Embryos were raised at 18°C which leads to a comparatively slow progression through the Long & Ballard stages of gar development (*48*). Sizes of each specimen were recorded for future analysis. Juvenile specimens of either sex were used. All animal procedures were performed under the in accordance with protocols approved by Sorbonne Université and Institute (#QBI/041/20/France) and Michigan State University (#10/16-179-00).

Human eye samples

Human fetal eyes from terminated pregnancies were obtained from the INSERM-funded Human Developmental Cell Atlas collection (HuDeCA, https://hudeca.genouest.org/). All tissues were collected with appropriate maternal consent and approval from the French National Biomedicine agency (N° PFS19-012).

In Situ Hybridization

Spotted gar sections were hybridized with digoxigenin-labeled riboprobes as described in (66). Briefly, tissue sections were postfixed for 10 min in 4% paraformaldehyde (PFA) before being treated with Proteinase K (10 µg/ml; Invitrogen, #03115852001) for 2 min and subsequently postfixed for 5 min in 4% PFA. Sections were then acetylated and permeabilized in PBS, 1% Triton X-100. Sections were first homogenized with hybridization buffer (50% formamide (VWR #24311.291), 5× SSC (Euromedex, #EU0300-A), 1× Denhardt's, 250 µg/ml yeast tRNA, and 500 µg/ml herring sperm DNA, pH 7.4) for 2 h at RT and then hybridized overnight at 72°C with riboprobes (1/200), see Table S1 for probe sequences. The next day, sections were rinsed for 2 h in 2× SSC at 72°C, and blocked in 0.1 M Tris, pH 7.5, 0.15 M NaCl (B1) containing 10% normal goat serum (NGS) for 1 h at RT. After blocking, slides were incubated o/n at 4°C with anti-DIG antibody conjugated with the alkaline phosphatase (1/5000, Roche Diagnostics) or anti-DIG antibody conjugated with peroxidase in B1 containing 1% NGS. After washing in B1 buffer, the alkaline phosphatase activity was detected by using nitroblue tetrazolium chloride (337.5 µg/ml) and 5-bromo-4-chloro-3-indolyl phosphate (175 µg/ml) (Roche Diagnostics). The peroxidase activity was detected by using Tyramide Signal Amplification (TSA) (PerkinElmer, #NEL741001KT) and incubated with Fluorescein fluorophore Tyramide diluted at 1:50 in TSA. Sections were mounted in Mowiol (Calbiochem/Merck, Carlstadt, Germany).

Whole-mount in situ hybridization were carried out on zebrafish as previously described (67). Embryos were then embedded in gelatin/albumin with 4% of glutaraldehyde and sectioned (20 μ m) on a VT1000 S vibrating blade microtome (Leica). Slides were scanned with either a Nanozoomer (Hamamatsu) or laser scanning confocal microscope (Olympus, FV1000).

Fluorescent in Situ Hybridization

To generate anti-sense probes, DNA fragments were obtained by PCR using Phusion™ High-Fidelity DNA polymerase (Thermo Scientific, #F530L) with the primers listed in Table S1. Total cDNA from 1 to 5 dpf zebrafish were used as a template. PCR fragments were cloned into the pCRII-TOPO vector (Invitrogen, #K280040) according to manufacturer's instructions. All plasmids used were sequenced for confirmation. Anti-sense DIG or fluorescein-labeled riboprobes were in vitro transcribed using the RNA labeling kit (Roche, #11685619910 or #11277073910) according to manufacturer's instructions. De-chorionated embryos at the appropriate developmental stages were fixed in fresh 4% PFA in 1X PBS (pH7.4) containing 0.1% Tween20 (PBSTw) for 4 h at RT and stored o/n in 100% methanol. Embryos were rehydrated by immersing them in subsequent baths of 50% methanol/PBSTw (Sigma, #34860) and then twice in PBSTw, baths followed by a 10 min incubation in a 3% H₂O₂/0.5%KOH (Sigma, #P5958) solution. Embryos were then rinsed in 50% methanol and post-fixed in 100% methanol at -20°C for 2 h. Embryos were then re-hydrated in methanol/PBSTw (75%/50%/25%) followed by treatment in 10 µg/ml proteinase K at RT (1 dpf = 5 min, 2 dpf = 15 min, 3 dpf = 20 min), and post-fixed for 20 min in 4% PFA in PBSTw. Embryos were pre-hybridized at 68°C, and hybridized with either a fluorescein-labelled probe or DIGlabelled probe or both probes for dFISH assays o/n at 68°C with gentle shaking. Embryos were then rinsed at 68°C in 50% formamide/2XSSC/0.1%Tween-20 twice, 2XSSC/0.1%Tween-20, 0.2XSSC/0.1% Tween-20 twice and finally in TNT buffer (0.1 M Tris pH7.5, 0.15 M NaCl, 0.1% Tween-20). Blocking was done in TNB buffer (2% DIG block (Roche, #11096176001) in TNT) for 2 h at RT and incubated o/n with anti-Fluo-Fab-POD (Roche, #11426346910) diluted at 1:50 in TNB buffer at 4°C. All steps were performed in the dark. Embryos were then washed several times in TNT, rinsed using $100 \,\mu$ l Tyramide Signal Amplification (TSA) (PerkinElmer, #NEL741001KT) and incubated with Fluorescein fluorophore Tyramide diluted at 1:50 in TSA.

The reaction was stopped by 5 rapid washes of TNT. For dFISH assays, the DIG-labelled probe was then revealed by carrying out a 20 min incubation in 1%H₂O₂/TNT (Sigma, #18312-1L), then washed several times in TNT. A second blocking step was carried out for 1 h in TNB buffer prior to incubating embryos in anti-DIG-POD (Roche, #11207733910) diluted at 1:100 in TNB buffer o/n at 4°C. Revelation was done with Cy3 Fluorophore Tyramide solution (PerkinElmer, NEL#744001KT), washed with TNT and processed for imaging upon DAPI staining.

Molecular cloning

14xUAS:ubc-ZIC2-T2A-GFP-pA or *14xUAS:ubc-T2A-GFP-pA* were obtained via Gibson assembly using the *pT1UciMP Tol1* (Addgene, #62215) destination vector described by (68). *Tol1*mRNA was synthesized from the plasmid (Addgene, #61388) digested by NotI (NEB, #R3189S) and retro-transcribed with SP6 RNA polymerase (Roche, #10810274001). Human ZIC2 (*hZIC2*), *GFP*, and *T2A* were amplified via PCR from pCAG-hZIC2 and *pUAS:Cas9T2AGFP;U6:sgRNA1;U6sgRNA2* (Addgene, #74009) respectively using the NEBuilder HiFi DNA Assembly Cloning kit (NEB, #E5520). Appropriate sequences were inserted after the *UBC* intron of the *pT1UciMP Tol1* destination vector opened by restriction digest with NcoI-HF (NEB).

Alignment between the amino acid sequences of the Zic2 proteins zing finger domains

A multiple sequence alignment was performed for the region covering the ZIC2 zinc finger domains of NCBI Reference Sequence proteins of mouse (NP_033600), human (NP_009060) and zebrafish (ZIC2a NP_571633, ZIC2b NP_001001820), as well as for genome-predicted ZIC2 proteins the spotted gar (XP_006638968). The UniProtKB/Swiss-Prot curated zinc finger sequences of human ZIC2 (O95409) were used to delineate the domain positions within the

alignment. Protein sequence alignment was performed using MUSCLE version 3.8.31 (69), the amino acid conservation at each aligned position visualised using BIS2Analyzer (70).

Eye enucleation

The transgenic line Tg(atoh7:gal4-vp16) (RRID: ZFIN_ZDB-GENO-130306-1) was used. Prior to eye enucleation, fish were selected for the atoh7 expression in green. At 2 dpf, eye enucleation was performed. The embryos were anesthetized in 0.004% tricaine MS222 in a 2% agarose gel solution (Life technologies, #16520050). One eye was surgically removed using a pulled capillary and mouth pipetting. Embryos were then transferred into fish medium (egg medium with penicillin/streptomycin (Life Technologies, #15140122) and 0.003% 1-phenyl-2thiouera (Sigma, #189235) until 5 dpf, for whole-mount immunohistochemistry.

Immunohistochemistry

Cryosections

Spotted gar embryos were fixed by immersion in 4% PFA in 0.12 M phosphate buffer (VWR, 28028.298 and 28015.294), pH 7.4 (PFA) o/n at 4°C. Following three washes in 1XPBS, the samples were incubated in 10% sucrose (VWR, 27478.296) in 0.12 M phosphate buffer o/n at 4°C. The next day, samples were transferred to a 30% sucrose solution in 0.12 M phosphate buffer o/n at 4°C. Samples were then embedded in 0.12 M phosphate containing 7.5% gelatin (Sigma, 62500) and 10% sucrose, frozen in isopentane at -40°C and then cut at 16 µm with a cryostat (Leica, CM3050S). Sections were blocked in PBS containing 0.2% gelatin (VWR) and 0.25% Triton-X100 (PBS-GT) for 1 h at RT. Following the blocking, sections were incubated with primary antibodies (see Table S1) diluted in a PBS-GT solution o/n at RT. Following three washes in PBST (0.05% Trinton-X100) secondary antibodies coupled to the appropriate fluorophore (see Table S1) were diluted in PBS-GT and incubated for 2 h at RT. Sections were

counterstained with Hoechst (Sigma, B2883, 1:1000) or DAPI (Life Technologies, D3571, 1/500). For PCNA staining, an antigen retrieval step was performed by boiling sections in a 1X Sodium Citrate solution pH 6.0 for 5 min using a microwave. This step was skipped when the samples were first used for an *in-situ* hybridization assay. Slides were scanned with either a Nanozoomer (Hamamatsu) or laser scanning confocal microscope (Olympus, FV1000).

Whole-mount Immunohistochemistry

Zebrafish whole-mount immunohistochemistry was adapted from (*61*). Briefly, embryos were fixed in 4% PFA diluted in PBS containing 0.1% Tween-20 (VWR, #0777-1L) (PBSTw) for 4 h at RT and stored o/n in 100% methanol. After re-hydration, embryos were incubated for 20 min at -20°C in already pre-chilled acetone (Sigma, #650501). The embryos were rinsed several times with PBSTw and blocked for 2 h in blocking solution (10% bovine serum albumin (BSA) (Euromedex, #04-100-812-C), 10% normal goat serum (LifeTechnologies, #1000C), 1% DMSO (Sigma Aldrich, #D8418) in PBSTw). The primary antibodies were incubated o/n at 4°C in 1% BSA, 1% normal goat serum, 0.1% DMSO in PBSTw according to the dilutions in Table S1. After several washes in PBSTw, the secondary antibodies were incubated o/n at 4°C. The next day, embryos were rinsed in PBSTw and processed for imaging.

Whole-mount immunostaining on spotted gar embryos was carried out as previously described (*15*). Briefly, embryos were depigmented in a solution of 11% H₂O₂ (VWR, 216763) at 70 rpm exposed to an 11W warm white Light-Emitting Diode (LED) (3000° Kelvin) for 1-3 days. Samples were then blocked and permeabilized before being incubated with the primary antibodies for 7 days at RT (see Table S1) in a solution containing: 0.5% Tirton-X100, 5% donkey normal serum, 20% Dimethyl Sulfoxide, 1XPBS, 0.1 g/L thimerosal. The samples were further labeled with secondary antibodies (see Table S1) for 2 days at RT under agitation.

Retinal flat-mounts

For retinal flat mounts, human eyes were harvested and fixed in 4% PFA, followed by three

washes in 1XPBS. Eyes were then de-pigmented using the EyeDISCO protocol as previously described (*15*). For immunohistochemistry, retinas were permeabilized and blocked in a solution containing 0.5% Triton-X100, 5% donkey normal serum, 1XPBS, 0.1 g/L thimerosal for 1 day at RT under agitation. Primary antibodies (see Table S1) were diluted in a solution containing 0.5% Triton-X100, 5% donkey normal serum, 20% Dimethyl Sulfoxide, 1XPBS, 0.1 g/L thimerosal for 3 days at RT under agitation. The retinas were then washed for 1 day in PBST (1XPBS, 0.5% Triton-X100). The secondary antibodies (see Table S1) were diluted in the same solution as primary antibodies and left for 2 days. After washing retinas for 1 day, they were mounted on slides and imaged using a scanning confocal microscope (Olympus, FV1000).

Tracing of visual projections

All fish were anesthetized with 0,04% MS222, tricaine-methanesulfonate (Sigma, #E10521) diluted in fish water. Australian lungfish were anesthetized with 0.05% clove oil in fresh water. Injection of cholera toxin β subunit was carried out as described in (*15*). Briefly, using a capillary approximately 1µl of 2 µg/µl of Alexa Fluor-conjugated cholera toxin β subunit (Thermo Fischer, Alexa Fluor555-CTb C22843 and Alexa Fluor647-CTb C34778) was injected intravitreally. 72-96 h following CTb injection, specimens were transcardially perfused with 4%PFA and the heads and/or brains were dissected for tissue clearing.

Tissue clearing and imaging

<u>Clearing</u>

Prior to clearing, spotted gar *embryos* were embedded in 1.5% agarose (Roth) in 1X TAE (Life Technologies). Clearing was carried out as previously described (*15*). Briefly, samples were gently de-hydrated in ascending baths of methanol (1.5 h). Samples were further treated with a

solution containing 2/3 Dichloromethane (DCM, Sigma) 1/3 methanol o/n. The next day, samples were placed in DCM for 30 min prior to being immersed in Di-benzyl Ether (DBE, Sigma).

Imaging

Acquisitions were performed by using an UltraMicroscope I (Miltenyi Biotec, Germany) or UltraMicroscope Blaze (Miltenyi Biotec, Germany) with the ImspectorPro software (Miltenyi Biotec, Germany, 5.1.328 version). The light sheet was generated by a laser (wavelength 488, 561, 647 Coherent Sapphire Laser, LaVision BioTec, Miltenyi Biotec, Germany) or a secondgeneration laser beam combiner (wavelengths 488 nm, 561 nm and 647 nm; LaVision BioTec, Miltenyi Biotec, Germany). All light sheets were matched within their Rayleigh lengths for optimal illumination at the sample site. Either a binocular stereomicroscope (Olympus, MXV10) with a 2x objective (Olympus, MVPLAPO) was used Or a MI Plan 1.1x (NA = 0.1), a MI Plan 4x (NA = 0.35), and a MI Plan 12x (NA = 0.53) objectives were used (Miltenyi Biotec, Germany). Samples were placed in an imaging reservoir made of 100% quartz (LaVision BioTec, Miltenyi Biotec) filled with DBE and illuminated from the side by the laser light. A Zyla sCMOS camera (Andor, Oxford Instrument; 2,048 × 2,048, 6.5 x 6.5 μ m, peak QE 82%) was used to acquire images. The step size between each image was fixed at 1 or 2 μ m (NA = 0.5, 150 ms time exposure). All tiff images are generated in 16-bit.

Confocal microscopy

Whole-mount 5 dpf zebrafish larvae were mounted in a labtex plates (LabTex) in 2.5% agarose or in 1% low-melting agarose on FluoroDish Cell Culture dish (FD3510-100, World Precision Instruments). For imaging, a scanning inverted confocal microscope (FV1200, Olympus) was used with a 30x objective (Olympus, UPLSAPO30XS, NA = 1.05, WD = 0.8 mm) as well as the LSM780 and LSM880 scanning inverted confocal microscopes (Zeiss) for high resolution microscopy. 40x water immersion objective for whole mount dFISH stained zebrafish embryos

and 63x oil objective for zebrafish retinal cryosections were used and a 10x air objective was used to image the spotted gar cryosections.

Image Processing

3D rendering of light sheet and confocal stacks were converted to an Imaris file (.ims) using ImarisFileConverter (Bitplane, 9.5.1 version) and then visualized using the Imaris x64 software (Bitplane, 9.5.1). To quantify ipsilateral territories, entire tectum volume and ipsilateral projections were automatically segmented with a surface detail of $5.00 \,\mu\text{m}$, automatic threshold. Volumes were extracted from the surface. Movies were generated using the animation tool on Imaris x64 software (Bitplane, version 9.1.2) and movie reconstruction with .tiff series were done using ImageJ (1.50e, Java 1.8.0_60, 64-bit). All movie editing (text and transitions) was performed using iMovie (Apple Inc., version 10.1.1).

To quantify the ipsilateral projections in the hZIC2 overexpression experiments, a fixed region of interest was identified for each zebrafish (corresponding to the ipsilateral and contralateral optic tecta). Retinal projections were segmented with a surface detail of $0.5 \,\mu$ m using an automatic threshold. Ipsilateral and contralateral volumes were extracted and summed to constitute the "total visual projections" using Imaris x64 software (Bitplane, version 9.1.2). The volume of ipsilateral projections was isolated as a ratio of ipsilateral projections:total projections.

Statistical analyses

All data are described are listed as biological replicates (n) and all experiments (N) were carried out at least in triplicates unless indicated otherwise). An observer blinded to the experimental conditions realized all the quantifications. No data were excluded from the statistical analyses. All data are represented as mean values ± SEM. Statistical significance was estimated using two-tailed unpaired tests for non-parametric tendencies (Kruskall-Wallis or Mann-Whitney), two-way ANOVA and Bonferroni's multiple comparison test. * = p < 0.05; ** = p < 0.01; *** = p < 0.001, *** = p < 0.0001. All statistical measurements were carried out using GraphPad Prism 7.

Supplementary Figures



Fig. S1. Simplified chart of fish taxonomy indicating the species analyzed in this study.

Fish with bilateral/ipsilateral visual projections appear in magenta and fish with only contralateral visual projections appear in grey.



Fig. S2. Two types of optic nerve crossing modalities in ray-finned fishes.

Ventral views (A to D) and coronal optical sections (E to H) at the level of the optic chiasm of iDISCOcleared brains and optic nerves. A surface rendering with normal shading (Imaris) was applied to generate the ventral view images. The arrowheads (A to D) indicate the level of the chiasm optical section in (E to H). In all fishes, one eye was injected with Alexa Fluor-555-conjugated CTb and the other one with Alexa Fluor-647-conjugated CTb. The right (R) and left (L) optic nerves were pseudocolored in magenta and green respectively. In Pufferfish (A and E), Four-eyed (B and F) and Muskipper (C and G), the two optic nerves pass over and overlap at the chiasm but remain separated up to the brain. By contrast, in the Armored bichir (D and H), the right and left nerves meet at the chiasm and retinal ganglion cell axons from both eyes interweave during crossing (asterisk). Abbreviations: Ant, anterior; Post, Posterior. Scale bars are: 2 mm in (B), 1 mm in (A, C, D, F), 800 μ m in (G), 600 μ m in (E, H).



Fig. S3. ZIC2 expression pattern in human embryo retina.

(A) Box and whiskers representation of the ZIC2-positive surface in E16 mouse and pcw9 human retinas. (**B** and **C**) flat-mount pcw9 human retina labeled for ZIC2 and RBPMS (B) or SOX2 (C). (B) In the most superficial (basal side) regions of the temporal retina, ganglion cells expressing low levels of ZIC2 and RBPMS (arrowheads) are seen but ZIC2 and RBPMS are mostly exclusive. (C) image at the level of the interface between the neuroblastic layer showing that ZIC2+ cells are not SOX2+. Scale bars are: 50 μ m in (B and C).



Fig. S4. Zic2 ortholog expression in zebrafish is restricted to the ciliary margin zone. (A to H) Whole-mount *in situ* hybridization of zebrafish embryos for *zic2a* at 1 day post fertilization (1 dpf; A), 2 dpf (B) and 3 dpf (C) as well as *zic2b* at 1 dpf (E), 2 dpf (F) and 3 dpf (G). Zic2b is expressed in the ciliary marginal zone (CMZ, arrowheads in F and G) and in the dorsal diencephalon (Di). (D) Schematic drawing of the zebrafish CMZ in the developing retina showing spatial distribution of stem cells, cycling progenitors, committed progenitors and differentiated neurons. (H), Lateral view of whole-mount double fluorescent in situ hybridization for zic2b and atoh7 on 3 dpf zebrafish embryos with DAPI counterstaining. (I to L) Confocal sections through the central retina of wild-type embryos hybridized with antisense RNA probes for *zic2b* and *atoh7*. At 24 hpf, *zic2b* is expressed in the entire proliferative neuroepithelium and later from a central to peripheral wave-like manner (arrowheads) in complementarity to the neurogenic transient expression of *atoh7* (asterisks) as shown here for 36, 48 and 72 hpf. (L to M) Confocal sections through the central retina of 72 hpf wild-type zebrafish embryos hybridized with antisense RNA probes for *zic2b* and *retinal homeo- box transcription factor2 (rx2, contexperimentation)* a marker or dividing progenitors and stem cells in the CMZ). Zic2b expression overlaps with the expression of the rx2 (arrowheads). All retinae were counterstained with the nuclear marker DAPI. Scale bars are 50 μ m (A to C and E to H) and 40 μ m (I to M).



Fig. S5. Mammalian ipsilateral markers are not expressed in the spotted gar visual system.

(A) 3D light-sheet fluorescence microscopy images of iDISCO-cleared 17-18 dpf spotted gar indicating with dotted lines the anatomical levels of the cryosections. (**B** to **D**) *In situ* hybridization for *zic2* on retinal cryosections of the developing spotted gar at 2-3 dpf (B), 6-7 dpf (C, left panel), 17-18 dpf (D). Only proliferating cells in the neuroblastic layer (NBL) express *zic2*. The right panel in (C) is an immunostaining for PCNA and Islet1. The arrowheads in (C) indicate the region where the first ganglion cells (Islet1+) are present at this stage in the retina. *zic2* is also found in the diencephalon (Di). (**E** to **T**) Rostral-to-caudal coronal cryosections from 17-18 dpf spotted gar. *zic1* (E to H) and *zic5* (I to L) are only expressed in the ciliary marginal zone (CMZ; arrow). (**M** to **P**) *ephB1* is absent from the retina and weakly expressed in the CMZ. (**Q** to **T**) *ephrinB2* is expressed in the dorsal retina (arrow). (**U** to **W**) Cryosections of the diencephalon of a 17-18dpf spotted gar hybridized for *ephrinB2*. *ephrinB2* is absent from the optic chiasm (asterisk). Immuno-reactive regions are highlighted (arrowhead). Abbreviations: NBL, Neuroblastic layer; ON, Optic nerve, OC, Optic chiasm; GCL, Ganglion cell layer; INL, Inner nuclear layer; ONL, Outer nuclear layer. Scale bars: A, 200 µm; B to D, 50 µm; C,U to W, 100 µm; D to T, 250 µm.



Fig. S6. Alignment between the amino acid sequences of the Zic2 protein zing finger domains of fish and mammals.

Alignment of Zic proteins across the zinc finger domains shows the high level of conservation between ray-finned fish and mammals. Amino acids are color coded according to the physiochemical class they belong to. Abbreviations: HU, human; MM, mouse; SG, spotted gar; ZE, zebrafish.



Tg(atoh7:gal4), p14UASubc:T2A-eGFP

Fig.S7. Ectopic *ZIC2* expression in *atoh7* retinal progenitor cells does not induce EphB1 expression in retinal ganglion cells.

Confocal images of cryostat (A-F) or optical (G-L) sections of 3 dpf retinae from Tg(atoh7:gal4) embryos injected at 1-cell stage with either a *p14UASubc:T2A-eGFP* (A to C and G to I) control construct or a *p14UASubc:ZIC2-T2A-eGFP* construct (D to F and J to L). No signal for EphB1 in GFP-positive cells is detectable in all injected retinae either double stained with anti-EphB1 antibody (A to F) or hybridized with an *ephB1* antisense riboprobe (G to L). All retinae were counterstained with the nuclear marker DAPI. Scale bars are 50 µm.

			In situ probes						
Name	Sequence			RRID	Dilution	In situ hybridization			
L-zic1 fwd	ACCTCCAGACATCACTCAAC			n/a	1:200	Cryosections			
L-zic1 rev	GGAACAC	GGAACACTCTTCCCAGAAAC			1:200	Cryosections			
L-zic2 fwd	AAACTTAACCACGACCTCTCTC			n/a	1:200	Cryosections			
L-zic2 rev	CTCGTGCATTGTGCTGAAAG			n/a	1:200	Cryosections			
L-zic5 fwd	CTTTGAGCAAGAGGAATCCGGC			n/a	1:200	Cryosections			
L-zic5 rev	CCTGCCG	CGATGTTCACA	TTTA	n/a	1:200	Cryosections			
L-efnb2 fwd	TCCCCATI	TATGAGAAGGT	GAGCGG	n/a	1:200	Cryosections			
L-efnb2 rev	ACAGGCTA	ACCACTTCAGA	AGGCAG	n/a	1:200	Cryosections			
L-ephb1 fwd	AGAACCTO	GAACACAATCO	CGCAC	n/a	1:200	Cryosections			
L-ephb1 rev	ACAGTITA	ATGGGCACGT	CCAC	n/a	1:200	Cryosections			
zf-zic2a fwd	ACAACAA	ICTGTCGCCTT	CCTC	n/a	1:200	whole-mount			
zf-zic2a rev	ACAAATGCCCCTGTTTAGCCC			n/a	1:200	whole-mount			
zf-zic2b fwd	TCTTCCGCTACATGCGACAAC			n/a	1:200	whole-mount			
zf-zic2b rev	GCAACACCGACATGCTGAGAAC			n/a	1:200	whole-mount			
zf-ephb1 fwd	CGCGTGTGGATGGATTACGG			n/a	1:200	whole-mount			
zf-ephb1 rev	CATCCCCACCAGCTGGATCA			n/a	1:200	whole-mount			
zf-atoh7 fwd	GGAGAAG	GGAGAAGTTTGAGAGTGCTATGCGG			1:200	whole-mount			
zf-atoh7 rev	CGACTITIC	CGACTTTGAGCTGAGCACACACC			1:200	whole-mount			
zf-rx2 fwd	GATACCAT	IGAACATGGTG	GACGATGG	n/a	1:200	whole-mount			
zf-rx2 fwd	CCATCGAC	CTGAATGTGCT	CCTTGG	n/a	1:200	whole-mount			
			Primary antibodi	es					
Antigen	Species	Catalog #	Company	RRID	Dilution	Immunohistochemistry			
Islet1	Rabbit	GTX128201	GeneTex	Ab_2868422	1:300	Cryosections/whole-mount			
Acetylated-tubulin	Mouse	T6793	Sigma	Ab_477585	1:300	Cryosections/whole-mount			
PCNA	Mouse	P8825	Sigma	Ab_477413	1:500	Cryosections			
Islet1+2	Mouse	39.4D5	DSHB	Ab_2314683	1:50	Cryosections			
GFP	Chicken	GTX13970	GeneTex	Ab_371416	1:5000	whole-mount			
Rbpms	Guinea Pig	ABN1376	Millipore	Ab_2687403	1:400	Cryosections/flat-mount/ whole-			
Zic2	Rabbit	Ab150404	Abcam	Ab_2868423	1:300	Cryosections/flat-mount/ whole-			
Sox2	Goat	Sc17320	Santa-Cruz	Ab_2286684	1:300	flat-mount			
EphB1	Mouse	MAb EfB1-3	DSBH	Ab_2314357	1:5	Cryosection			
			Secondary antibod	lies					
Anti-Rabbit cy3	Donkey	711-165-152	Jackson	Ab 2307443	1:500	cryosections/whole-mount			
Anti Dabbit Alava Eluar	Dankay	711 605 152	ImmunoResearch			amagastions/Elst mount/whole			
647	Donkey	711-005-152	ImmunoResearch	Ab_2492288	1:500	mount			
Anti-Goat Alexa Fluor 488	Donkey	A11055	Life Technologies	Ab 2534102	1.500	cryosections/Flat-mount/whole-			
	-			A0_2334102	1.500	mount			
Anti-Goat Alexa Fluor 555	Donkey	A21432	Life Technologies	Ab_2535853	1:500	cryosections/Flat-mount/whole-			
Anti-Goat Alexa Fluor 647	Bovine	805-605-180	Jackson			cryosections			
			ImmunoResearch	AB_2340885	1:600	5			
Anti-Goat cy3	Donkey	705-165-147	Jackson	Ab 2307351	1:500	cryosections/Flat-mount/whole-			
Anti mouse Alexa Eluor	Donkey	A21202	ImmunoResearch	_		mount			
488	Donkey	A21202	Life recimologies	Ab_141607	1:500	mount			
Anti-Guinea-Pig Alexa cy3	Donkey	706-165-148	Jackson	Ab 2340460	1.500	cryosections/Flat-mount/whole-			
			ImmunoResearch	A0_2340400	1.500	mount			
Anti-mouse Alexa Fluor	Donkey	715-605-150	Jackson Jackson	Ab_2340862	1:500	cryosections/Flat-mount/whole-			
Anti-Mouse, Alexa Fluor	Goat	A31574	Life Technologies			cryosections			
635	-		8	Ab_2536184	1:500	5			
Anti-Rabbit, Alexa Fluor	Goat	A11036	Life Technologies	Ab 10563566	1:500	cryosections			
568 Anti Mouso, Alava Eluor	Cont	411004	Life Technologies	-		orrespections			
568	Goal	AI 1004	Life fectiliologies	Ab_2534072	1:500	cryosections			
Alexa Fluor 488 anti-	Goat	A11039	Life Technologies	Ab 142024	1.500	cryosections			
chicken			-	A0_142924	1:500				
Tracers									
Cholera toxin subunit B-	n/a	C22843	Life technologies	n/a	$2 \ \mu g/\mu l$	Whole-mount			
Cholera toxin subunit R-			-						
AlexaFluor647	n/a	C34778	Life technologies	n/a	$2 \ \mu g/\mu l$	Whole-mount			

Table S1. Comprehensive table summarizing the antibodies and probes sequences.

Movie S1.

Visual projections in teleosts.

Whole brain rendering of visual projections in 5 teleosts, the zebrafish, Mexican tetra, greenspotted pufferfish, mudskipper and butterflyfish. All species shows a complete decussation of retinal projections except the butterflyfish. All fish had bilateral eye injections of CTb coupled to either an Alexa Fluor-555 or and Alexa Fluor-647.

Movie S2.

Bilateral visual projections in non teleosts.

Whole brain rendering of visual projections in spotted gar, sterlet and armored bichir. Ipsilateral projections are seen in all species observed. All fish had bilateral eye injections of CTb coupled to either an Alexa Fluor-555 or and Alexa Fluor-647.

Movie S3.

The Australian lungfish possesses non-segregated ipsilateral projections.

Whole brain rendering of visual projections in the Australian lungfish, a sarcopterygian, injected with either an Alexa Fluor-555 or an Alexa Fluor-647. Many ipsilateral projections are observed, with a major component in the optic tectum. Ipsilateral projections are intermingled with contralateral projections in the optic tectum.

Movie S4.

ZIC2 expression is evolutionarily conserved in Humans.

Whole-mount immunohistochemistry of pcw9 human eyes using EyeDISCO clearing and labeled for the ipsilateral transcription factor ZIC2 (magenta) and the pan-retinal ganglion cell marker RBPMS (green). A large ZIC2-positive region can be seen in the temporal retina.

Movie S5.

Development of the Lepisosteus oculatus visual system.

3D rendering of 2-3 dpf, 6-7 dpf, and 17-18 dpf spotted gar embryos using EyeDISCO clearing and light-sheet fluorescence microscopy. Spotted gar embryos were labeled with the panneuronal marker acetylated tubulin (a-tubulin, green) and the LIM/homeodomain family of transcription factor Islet1, which is critical for the proper specification of retinal ganglion cells and motor neurons (magenta).








			In situ probes			
Name	Sequence			RRID	Dilution	In situ hybridization
L-zic1 fwd	ACCTCCAC	ACCTCCAGACATCACTCAAC		n/a	1:200	Cryosections
L-zic1 rev	GGAACACTCTTCCCAGAAAC		n/a	1:200	Cryosections	
L-zic2 fwd	AAACTTAACCACGACCTCTCTC			n/a	1:200	Cryosections
L-zic2 rev	CTCGTGCATTGTGCTGAAAG			n/a	1:200	Cryosections
L-zic5 fwd	CTTTGAGCAAGAGGAATCCGGC			n/a	1:200	Cryosections
L-zic5 rev	CCTGCCGCGATGTTCACATTTA			n/a	1:200	Cryosections
L-efnb2 fwd	TCCCCATTATGAGAAGGTGAGCGG			n/a	1:200	Cryosections
L-efnb2 rev	ACAGGCTACCACTTCAGAAGGCAG			n/a	1:200	Cryosections
L-ephb1 fwd	AGAACCTGAACACAATCCGCAC			n/a	1:200	Cryosections
L-ephb1 rev	ACAGTTTAATGGGCACGTCCAC			n/a	1:200	Cryosections
zf-zic2a fwd	ACAACAATCTGTCGCCTTCCTC			n/a	1:200	whole-mount
zf-zic2a rev	ACAAATGCCCCTGTTTAGCCC			n/a	1:200	whole-mount
zf-zic2b fwd	TCTTCCGCTACATGCGACAAC			n/a	1:200	whole-mount
zf-zic2b rev	GCAACACCGACATGCTGAGAAC			n/a	1:200	whole-mount
zf-ephb1 fwd	CGCGTGTGGATGGATTACGG			n/a	1:200	whole-mount
zf-ephb1 rev	CATCCCCACCAGCTGGATCA			n/a	1:200	whole-mount
zf-atoh7 fwd	GGAGAAGTTTGAGAGTGCTATGCGG			n/a	1:200	whole-mount
zf-atoh7 rev	CGACTTTGAGCTGAGCACACACC		n/a	1:200	whole-mount	
zf-rx2 fwd	GATACCATGAACATGGTGGACGATGG			n/a	1:200	whole-mount
zf-rx2 fwd	CCATCGACTGAATGTGCTCCTTGG			n/a	1:200	whole-mount
Primary antibodies						
Antigen	Species	Catalog #	Company	RRID	Dilution	Immunohistochemistry
Islet1	Rabbit	GTX128201	GeneTex	Ab 2868422	1.300	Cryosections/whole-mount
Acetylated_tubulin	Mouse	T6793	Sigma	Ab 477585	1.300	Cryosections/whole-mount
PCNA	Mouse	P8825	Sigma	Ab_477413	1.500	Cryosections
I CINA	Mausa	20.4D5	DSUB	Ab_2214682	1.500	Cryosections
CED	Chicken	GTV12070	ConoTox	Ab 371416	1.5000	whole mount
Rhome	Guinea Pig	ABN1376	Millipore	Ab_371410	1:400	Cryosections/flat mount/ whole
Zio2	Duffica Fig	Ab150404	Ahaam	Ab_2687403	1.400	Cryosections/flat_mount/ whole-
S2	Cast	R-17200	Santa Cruz	Ab_2808423	1.300	fit menut
5082	Goal	Sc17320	Santa-Cruz	Ab_2286684	1:500	
Ерпы	Mouse	WIAU EIDI-3	Eccondom ontihod	A0_2314337	1.5	cryosection
Anti Pabhit av2	Donkay	711 165 152	Jackson	lles	1	any associans/whole mount
Anti-Kabbit Cy5	Donkey	/11-105-152	ImmunoResearch	Ab_2307443	1:500	cryosections/ whole-mount
Anti-Rabbit Alexa Fluor	Donkey	711-605-152	Jackson	AL 2402288	1,500	cryosections/Flat-mount/whole-
647			ImmunoResearch	AD_2492288	1:300	mount
Anti-Goat Alexa Fluor 488	Donkey	A11055	Life Technologies	Ab_2534102	1:500	cryosections/Flat-mount/whole-
Anti-Goat Alexa Fluor 555	Donkey	A21432	Life Technologies			cryosections/Flat-mount/whole-
	,		8	Ab_2535853	1:500	mount
Anti-Goat Alexa Fluor 647	Bovine	805-605-180	Jackson	AB 2340885	1.600	cryosections
	D 1	705 165 147	ImmunoResearch	TID_23 10003	1.000	
Anti-Goat cy3	Donkey	/05-165-14/	Jackson ImmunoResearch	Ab_2307351	1:500	cryosections/Flat-mount/whole-
Anti-mouse Alexa Fluor	Donkey	A21202	Life Technologies			cryosections/Flat-mount/whole-
488	,		U	Ab_141607	1:500	mount
Anti-Guinea-Pig Alexa cy3	Donkey	706-165-148	Jackson	Ab 2340460	1:500	cryosections/Flat-mount/whole-
Anti-mener Alene Eleren	Devilence	715 605 150	ImmunoResearch			mount
647	Donkey	/13-003-130	ImmunoResearch	Ab_2340862	1:500	mount
Anti-Mouse, Alexa Fluor	Goat	A31574	Life Technologies	AL 052(104	1.500	cryosections
635				Ab_2536184	1:500	-
Anti-Rabbit, Alexa Fluor	Goat	A11036	Life Technologies	Ab 10563566	1:500	cryosections
568 Anti Mausa Alava Eluan	Cont	411004	Life Technologies	_		amogaatiana
568	Goal	AI 1004	Life Technologies	Ab_2534072	1:500	cryosections
Alexa Fluor 488 anti-	Goat	A11039	Life Technologies	Ab 140004	1,500	cryosections
chicken			Ű	Ab_142924	1:500	
			Tracers			·
Cholera toxin subunit B-	n/a	C22843	Life technologies	n/a	2 μg/ul	Whole-mount
AlexaFluor555					1.9. 64	
AlexaFluor647	n/a	C34778	Life technologies	n/a	$2 \ \mu g/\mu l$	Whole-mount

Table S1. Comprehensive table summarizing the antibodies and probes sequences.