



**HAL**  
open science

## Spinal Epidural AV Fistula Embolization with EVOH Copolymer Using the Scepter Mini Dual Lumen Balloon

Mehdi Drir, Frédéric Clarençon, Eimad Shotar, Arnaud Pouvelle, Kevin Premat, Stéphanie Lenck, Gonzague Guillaumet, Elisabeth Maillart, Nader-Antoine Sourour

### ► To cite this version:

Mehdi Drir, Frédéric Clarençon, Eimad Shotar, Arnaud Pouvelle, Kevin Premat, et al.. Spinal Epidural AV Fistula Embolization with EVOH Copolymer Using the Scepter Mini Dual Lumen Balloon. *Journal of Neurointerventional Surgery*, 2021, 13 (5), pp.492-492. 10.1136/neurintsurg-2020-016395 . hal-03283852

**HAL Id: hal-03283852**

**<https://hal.sorbonne-universite.fr/hal-03283852>**

Submitted on 12 Jul 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Spinal Epidural AV Fistula Embolization with EVOH Copolymer Using the Scepter  
Mini Dual Lumen Balloon**

Frédéric Clarençon <sup>1,2</sup>, M.D., Ph.D, Eimad Shotar <sup>1</sup>, M.D., Arnaud Pouvelle <sup>1</sup>, M.D., Kevin Premat <sup>1,2</sup>, M.D., Stéphanie Lenck <sup>1</sup>, M.D., Mehdi Drir <sup>2,3</sup>, M.D., Gonzague Guillaumet <sup>4</sup>, M.D., Elisabeth Maillart <sup>5</sup>, M.D., Nader-Antoine Sourour <sup>1</sup>, M.D.

<sup>1</sup> Department Neuroradiology, Pitié-Salpêtrière Hospital, 47 Boulevard de l'Hôpital 75013 Paris, France.

<sup>2</sup> Sorbonne Université, Paris, France.

<sup>3</sup> Department of Neurosurgical Anesthesiology and Critical Care, Pitié-Salpêtrière Hospital, Paris, France.

<sup>4</sup> Department of Neurosurgery, Pitié-Salpêtrière Hospital, Paris, France.

<sup>5</sup> Department of Neurology, Pitié-Salpêtrière Hospital, Paris, France.

*Corresponding author:*

Pr Frédéric Clarençon (MD, PhD)

Department of Neuroradiology. Pitié-Salpêtrière Hospital.

47, Bd de l'Hôpital. 75013. Paris. FRANCE

Tel: + 33 1 42 16 35 99

Fax: +33 1 42 16 35 95

e-mail: fredclare5@gmail.com

Word count: 150

Number of figures: 0

Number of videos: 1

Number of tables: 0

**Acknowledgment:**

The authors thank Mme Valérie Klodawski and Dr Mathilde Aubertin for their kind help in performing the movie of the balloon preparation.

**Funding:** none

**Disclosures:**

Pr F. Clarençon reports conflict of interest with Medtronic, Guerbet, Balt Extrusion, Penumbra (payment for readings; non-related to the study), Codman Neurovascular and Microvention (core lab; non-related to the study).

Dr Nader-Antoine Sourour is consultant for Medtronic, Balt Extrusion, Microvention. The other authors report no conflict of interest concerning the materials or methods used in this study or the findings specified in this paper. The manuscript is not supported by industry.

## **Abstract**

Left unattended, spinal epidural arteriovenous fistulas (EAVF) have a potentially severe clinical course. Embolization using ethylene vinyl alcohol (EVOH) copolymers through regular dual lumen balloons has emerged as a potential option for the treatment of spinal arteriovenous fistulas<sup>1, 2, 3</sup>; the main issue of this technique being the navigability of these balloons. The Scepter mini is a low-profile dual lumen balloon, which may be helpful for EVOH embolization of spinal AV fistulas; it may help to overcome the navigation drawbacks. In this technical video, we present a case of EVOH embolization of a right T6 spinal EAVF through a Scepter mini balloon. Of note, great attention should be paid to radiculomedullary arteries arising at the same level or at adjacent levels to avoid severe neurologic complication related to uncontrolled migration of the liquid embolic agent. Moreover, excessive use of embolic material should be avoided to prevent spinal cord compression.

## **References**

1. Trivelato FP, Rezende MTS, Ulhoa AC, Nakiri GS, Abud DG. Dual-lumen balloon to increase onyx venous penetration in the treatment of spinal dural arteriovenous fistulas. *J Neuroradiol.* 2018;45(2):142-146. doi: 10.1016/j.neurad.2017.09.006.
2. Arslan M, Cınar C, Oran I. Dual-Lumen Balloon Catheter Technique for Onyx Embolization of Spinal Dural Arteriovenous Fistula. *J Vasc Interv Neurol.* 2019;10(3):34-37.
3. Nakae R, Nagaishi M, Hyodo A, Suzuki K. Embolization of a spinal dural arteriovenous fistula with ethylene-vinyl alcohol copolymer (Onyx) using a dual-lumen microballoon catheter and buddy wire technique. *Surg Neurol Int.* 2017;8:166. doi: 10.4103/sni.sni\_193\_17.

**Contributorship statement:**

Frédéric Clarençon: Manuscript redaction, video preparation

Eimad Shotar: Manuscript preparation, critical review of the manuscript

Arnaud Pouvelle: Data collection, critical review of the manuscript

Kevin Premat: Critical review of the manuscript

Stéphanie Lenck: Critical review of the manuscript

Mehdi Drir: Critical review of the manuscript

Gonzague Guillaumet: Data collection

Elisabeth Maillart: Critical review of the manuscript

Nader-Antoine Sourour: Data collection, Critical review of the manuscript