



HAL
open science

Editorial: Clinical Cardiopulmonary Exercise Testing

Denis E O'Donnell, Pierantonio Laveneziana, J. Alberto Neder

► **To cite this version:**

Denis E O'Donnell, Pierantonio Laveneziana, J. Alberto Neder. Editorial: Clinical Cardiopulmonary Exercise Testing. *Frontiers in Physiology*, 2021, 12, pp.711505. 10.3389/fphys.2021.711505 . hal-03289940

HAL Id: hal-03289940

<https://hal.sorbonne-universite.fr/hal-03289940v1>

Submitted on 19 Jul 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Editorial: Clinical Cardiopulmonary Exercise Testing

Denis E. O'Donnell^{1*}, Pierantonio Laveneziana² and J. Alberto Neder¹

¹ Respiratory Investigation Unit and the Laboratory of Clinical Exercise Physiology, Queen's University and Kingston Health Sciences Centre, Kingston, ON, Canada, ² Sorbonne Université, Faculté de Médecine Pierre et Marie Curie & APHP and Service d'Explorations Fonctionnelles de la Respiration, de l'Exercice et de la Dyspnée, Hôpital Universitaire Pitié-Salpêtrière, Tenon et Saint Antoine, Paris, France

Keywords: cardiopulmonary exercise testing, inspiratory neural drive, dynamic respiratory mechanics, exertional dyspnea, physiological characterization

Editorial on the Research Topic

Clinical Cardiopulmonary Exercise Testing

The value of cardiopulmonary exercise testing (CPET) is well-established as an adjunct to clinical evaluation. Worldwide, exercise “stress” testing is vastly under-utilized, especially for individuals with pulmonary disorders. This is lamentable because modern refinements in the measurements of physiological impairment during CPET have improved our ability to evaluate such patients more comprehensively (Domnik et al.; Phillips et al.; James et al.). Thus, the overarching objective of the current Research Topic series was to review some of these recent advancements in CPET methods and interpretation and to discuss their relevance, in both clinical and research settings, in patients with various respiratory conditions.

Uniquely, CPET can identify specific abnormalities of the integrated metabolic, cardiopulmonary, locomotor muscle, and neurosensory systems, under graded physiologic stress. Based on foundational physiological principles, CPET interpretation has traditionally focused on measuring peak oxygen uptake (VO_2)—a key metric of exercise performance—and an independent predictor of mortality in various cardio-pulmonary disorders (Killian and Jones, 1984; Jones, 1997; Wasserman et al., 1999, 2012; Jones and Killian, 2000). Traditional interpretative algorithms incorporate quantitative estimates of cardiac and ventilatory reserves, as well as aerobic capacity, to help explain reduced peak VO_2 in the individual. However, it must be remembered that low peak VO_2 often reflects intolerable exertional symptoms (e.g., dyspnea and leg discomfort), well before physiological maxima are actually attained (Neder et al.). Thus, measurement of exertional symptoms, using validated scales, is integral to any assessment of exercise capacity (Laviolette and Laveneziana, 2014). Moreover, it is now clear that traditional estimates of breathing and cardiac reserves, based on ratios of peak ventilation and heart rate, to their respective maxima, can sometimes be misleading. Thus, apparent preservation of breathing and cardiac reserves can often obscure important respiratory mechanical constraints and severe dyspnea at relatively low exercise intensities which, in turn, explain low peak VO_2 (Neder et al.).

Recent experimental work during CPET examined the simultaneous associations between inspiratory neural drive (IND) to the respiratory muscles, dynamic respiratory mechanics and subjective breathing discomfort and has shed light on fundamental mechanisms of exercise limitation in health and various respiratory conditions (Domnik et al.) (Laveneziana et al., 2009, 2011, 2013a,b, 2014, 2015; Laviolette and Laveneziana, 2014; O'Donnell et al., 2017, 2019; Neder et al., 2019; Boucly et al., 2020). In healthy participants, several acute physiological adaptations to the increasing metabolic demand of exercise, ensure admirable preservation of harmonious coupling between increasing IND and the ventilatory response of the dynamic respiratory system,

OPEN ACCESS

Edited and reviewed by:

Johannes Van Lieshout,
University of Amsterdam, Netherlands

*Correspondence:

Denis E. O'Donnell
odonnell@queensu.ca

Specialty section:

This article was submitted to
Clinical and Translational Physiology,
a section of the journal
Frontiers in Physiology

Received: 18 May 2021

Accepted: 03 June 2021

Published: 28 June 2021

Citation:

O'Donnell DE, Laveneziana P and
Neder JA (2021) Editorial: Clinical
Cardiopulmonary Exercise Testing.
Front. Physiol. 12:711505.
doi: 10.3389/fphys.2021.711505

such that arterial blood gas and acid-base homeostasis are achieved, and dyspnea is minimized. However, these remarkable, integrated cardio-respiratory adjustments to exercise become variably undermined by the presence of pulmonary diseases, even when resting pulmonary function appears normal. Thus, disease-related disruption of normal cardio-respiratory adaptations to exercise requires large compensatory increases in IND to maintain adequate alveolar ventilation. This, in conjunction with a widening disparity between increasing IND and the abnormal mechanical response of the impaired respiratory system (neuro-mechanical dissociation, NMD) has serious deleterious sensory consequences at higher exercise intensities, leading to premature termination of exercise (Domnik et al.).

Most chronic respiratory conditions are characterized by excessive ventilation for a given metabolic load during exercise, when compared with healthy controls. This ultimately points to increased IND due to multiple physiological perturbations, including: the effect of high physiological dead space, increased respiratory muscle loading, early lactic acidosis, and critical hypoxemia. Additionally, increased sympathetic nervous system activation and altered afferent activity from receptors in the respiratory and locomotor muscles, baroreceptors and chemoreceptors are undoubtedly contributory, but difficult to measure. High ventilatory equivalent for CO₂ (V_E/V_{CO_2}) is linked to increased IND and dyspnea intensity and is common to several respiratory conditions, including chronic obstructive pulmonary disease (COPD) (Phillips et al.), interstitial lung disease (ILD), (Molgat-Seon et al.) and pulmonary arterial hypertension (PAH) (Laveneziana and Weatherald). It is no surprise, therefore, that V_E/V_{CO_2} is increasingly represented in CPET interpretative algorithms (Phillips et al.). High V_E/V_{CO_2} nadir (e.g., > 34) generally indicates presence of poor ventilatory efficiency, due to high physiological dead space (and compromised lung CO₂ elimination), alveolar hyperventilation (enhanced chemo-sensitivity) or both. Measurement of arterial partial pressure of CO₂ (PaCO₂), and not end-tidal CO₂, (as it may grossly underestimate PaCO₂ when physiological dead space is increased), is needed to determine the relative contribution of each to high V_E/V_{CO_2} . As repeatedly highlighted in this series, V_E/V_{CO_2} must be considered in the context of the prevailing mechanical constraints, which can blunt the normal ventilatory response and compound interpretation. For simplicity, measurement of V_E/V_{CO_2} nadir is the preferred parameter: it generally occurs at relatively low exercise intensities, when arterial O₂ saturation is normal and before the onset of both metabolic acidosis (with ventilatory compensation) and significant restrictive mechanical constraints on tidal volume (V_T) expansion (O'Donnell et al., 2017, 2019; Neder et al., 2019).

Recent refinements in the “non-invasive” assessment of dynamic respiratory mechanics during exercise have increased our ability to ascertain—with reasonable precision—the nature and severity of the physiological impairment in the symptomatic individual (Milne et al.). The combination of operating lung volumes [measured by serial inspiratory capacity (IC)

maneuvers] and breathing pattern can help detect important inspiratory mechanical constraints, which are relevant to dyspnea and exercise limitation (Laveneziana et al., 2009, 2011, 2013a,b, 2014, 2015; Laviolette and Laveneziana, 2014; O'Donnell et al., 2017, 2019; Neder et al., 2019; Boucly et al., 2020). This approach is arguably more sensitive than traditional assessments of breathing reserve (V_E /derived maximal ventilatory capacity), especially in milder forms of obstructive and restrictive disorders or other cardio-respiratory conditions (Laveneziana et al., 2009, 2011, 2013a,b, 2014, 2015; Boucly et al., 2020). Additionally, examination of tidal vs. maximal flow-volume loops throughout exercise provides qualitative assessments of inspiratory and expiratory flow reserves. For practical purposes, the IC, and not the vital capacity, represents the true operating limits for V_T expansion in older individuals with common respiratory conditions, especially if expiratory flow limitation is present. Resting IC is variably reduced in both obstructive and restrictive lung disorders (James et al.; Molgat-Seon et al.). The lower the resting IC, the closer V_T is positioned to total lung capacity (TLC) and to the upper extremity of the respiratory system's sigmoidal pressure-volume relation. Thus, as V_T expands to occupy ~70% of IC and inspiratory reserve volume (IRV) reaches its minimal value, higher IND and breathing effort is needed to achieve a given V_T . This reflects the increased elastance of the respiratory system, when breathing close to TLC. Appearance of the V_T inflection or plateau (minimal IRV), as a function of increasing V_E during exercise, is an important mechanical event and heralds the onset of NMD and an abrupt rise in dyspnea. As resting IC diminishes over time in obstructive and restrictive disorders, critical inspiratory mechanical constraints and onset of dyspnea will occur at progressively lower V_E during exercise (O'Donnell et al., 2017, 2019; Neder et al., 2019).

This series has provided a comprehensive physiological characterization of patients with COPD, ILD, and PAH, (James et al.; Molgat-Seon et al.; Laveneziana and Weatherald) and taught us that exercise response pattern abnormalities are surprisingly similar, despite vastly different pathological origins. For patients referred with unexplained exertional dyspnea, who do not manifest any of the described abnormal patterns during CPET, alternative diagnoses such as dysfunctional breathing of psychogenic origin should be considered (Ionescu et al.). We hope that the pragmatic approach to CPET proposed here will help the clinician to individualize management plans by providing clear answers to a few key questions: *how severe is dyspnea and exercise intolerance in my patient?; is ventilatory demand increased and if so, why?; and are dynamic mechanical abnormalities present that contribute to poor exercise performance?*

AUTHOR CONTRIBUTIONS

DEO wrote the first draft of the manuscript. All authors were responsible for critically reviewing the manuscript for intellectual content and approved the final version of the manuscript submitted for publication.

REFERENCES

- Boucly, A., Morélot-Panzini, C., Garcia, G., Weatherald, J., Jaïs, X., Savale, L., et al. (2020). Intensity and quality of exertional dyspnoea in patients with stable pulmonary hypertension. *Eur. Respir. J.* 55:1802108. doi: 10.1183/13993003.02108-2018
- Jones, N. L. (1997). *Clinical Exercise Testing*. Philadelphia, PA: WB Saunders.
- Jones, N. L., and Killian, K. J. (2000). Exercise limitations in health and disease. *N. Engl. J. Med.* 343, 632–549. doi: 10.1056/NEJM200008313430907
- Killian, K. J., and Jones, N. L. (1984). The use of exercise testing and other methods in the investigation of dyspnea. *Clin. Chest Med.* 5, 99–108. doi: 10.1016/S0272-5231(21)00235-5
- Laveneziana, P., Bruni, G. I., Presi, I., Stendardi, L., Duranti, R., and Scano, G. (2013a). Tidal volume inflection and its sensory consequences during exercise in patients with stable asthma. *Respir. Physiol. Neurobiol.* 185, 374–379. doi: 10.1016/j.resp.2012.08.026
- Laveneziana, P., Garcia, G., Joureau, B., Nicolas-Jilwan, F., Brahimi, T., Lavolette, L., et al. (2013b). Dynamic respiratory mechanics and exertional dyspnoea in pulmonary arterial hypertension. *Eur. Respir. J.* 41, 578–587. doi: 10.1183/09031936.00223611
- Laveneziana, P., Humbert, M., Godinas, L., Joureau, B., Malrin, R., Straus, C., et al. (2015). Inspiratory muscle function, dynamic hyperinflation and exertional dyspnoea in pulmonary arterial hypertension. *Eur. Respir. J.* 45, 1495–1498. doi: 10.1183/09031936.00153214
- Laveneziana, P., Montani, D., Dorfmueller, P., Girerd, B., Sitbon, O., Jaïs, X., et al. (2014). Mechanisms of exertional dyspnoea in pulmonary veno-occlusive disease with EIF2AK4 mutations. *Eur. Respir. J.* 44, 1069–1072. doi: 10.1183/09031936.00088914
- Laveneziana, P., O'Donnell, D. E., Ofir, D., Agostoni, P., Padeletti, L., Ricciardi, G., et al. (2009). Effect of biventricular pacing on ventilatory and perceptual responses to exercise in patients with stable chronic heart failure. *J. Appl. Physiol.* 106, 1574–1583. doi: 10.1152/jappphysiol.90744.2008
- Laveneziana, P., Webb, K. A., Ora, J., Wadell, K., and O'Donnell, D. E. (2011). Evolution of dyspnea during exercise in COPD: Impact of critical volume constraints. *Am. J. Respir. Crit. Care Med.* 184, 1367–1373. doi: 10.1164/rccm.201106-1128OC
- Lavolette, L., and Laveneziana, P. (2014). Dyspnoea: a multidimensional and multidisciplinary approach. *Eur. Respir. J.* 43, 1750–1762. doi: 10.1183/09031936.00092613
- Neder, J. A., Berton, D. C., Marillier, M., Bernard, A. C., and O'Donnell, D. E. (2019). Inspiratory constraints and ventilatory inefficiency are superior to breathing reserve in the assessment of exertional dyspnea in COPD. *COPD* 16, 174–181. doi: 10.1080/15412555.2019.1631776
- O'Donnell, D. E., Elbehairy, A. F., Berton, D. C., Domnik, N. J., and Neder, J. A. (2017). Advances in the evaluation of respiratory pathophysiology during exercise in chronic lung diseases. *Front. Physiol.* 8:82. doi: 10.3389/fphys.2017.00082
- O'Donnell, D. E., Milne, K. M., Vincent, S. G., and Neder, J. A. (2019). Unraveling the causes of unexplained dyspnea: the value of exercise testing. *Clin. Chest Med.* 40, 471–499. doi: 10.1016/j.ccm.2019.02.014
- Wasserman, K., Hansen, J. E., Sue, D. Y., and Stringer, W. W. (2012). *Principles of Exercise Testing and Interpretation: Including Pathophysiology and Clinical Applications*. Philadelphia, PA: Lippincott Williams and Wilkins.
- Wasserman, K., Hansen, J. E., Sue, D. Y., Stringer, W. W., Sietsema, K. E., Sun, X. G., et al. (1999). *Principles of Exercise Testing and Interpretation*. Baltimore, MD: Lippincott Williams and Wilkins.

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 O'Donnell, Laveneziana and Neder. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.