



HAL
open science

Updated values of solar gravitational moments J_{2n} using HMI helioseismic inference of internal rotation

R. Mecheri, Mustapha Meftah

► To cite this version:

R. Mecheri, Mustapha Meftah. Updated values of solar gravitational moments J_{2n} using HMI helioseismic inference of internal rotation. Monthly Notices of the Royal Astronomical Society, 2021, 506 (2), pp.2671-2676. 10.1093/mnras/stab1827 . hal-03298817

HAL Id: hal-03298817

<https://hal.sorbonne-universite.fr/hal-03298817>

Submitted on 2 Aug 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Updated values of solar gravitational moments J_{2n} using HMI helioseismic inference of internal rotation

R. Mecheri¹★ and M. Meftah²★

¹Centre de Recherche en Astronomie, Astrophysique et Géophysique, CRAAG, BP 63, 16340 Bouzaréah, Algiers, Algeria

²Laboratoire Atmosphères, Milieux, Observations Spatiales (CNRS-LATMOS), 11 Boulevard d'Alembert, F-78280 Guyancourt, France

Accepted 2021 June 24. Received 2021 June 24; in original form 2021 May 12

ABSTRACT

The solar gravitational moments J_{2n} are important astronomical quantities whose precise determination is relevant for solar physics, gravitational theory and high precision astrometry, and celestial mechanics. Accordingly, we propose in the present work to calculate new values of J_{2n} (for $n = 1, 2, 3, 4$, and 5) using recent two-dimensional rotation rates inferred from the high-resolution Solar Dynamics Observatory / Helioseismic and Magnetic Imager helioseismic data spanning the whole solar activity cycle 24. To this aim, a general integral equation relating J_{2n} to the solar internal density and rotation is derived from the structure equations governing the equilibrium of slowly rotating stars. For comparison purpose, the calculations are also performed using rotation rates obtained from a recently improved analysis of Solar and Heliospheric Observatory / Michelson Doppler Imager helioseismic data for solar cycle 23. In agreement with earlier findings, the results confirmed the sensitivity of high-order moments ($n > 1$) to the radial and latitudinal distribution of rotation in the convective zone. The computed value of the quadrupole moment J_2 ($n = 1$) is in accordance with recent measurements of the precession of Mercury's perihelion deduced from high precision ranging data of the *MESSENGER* spacecraft. The theoretical estimate of the related solar oblateness Δ_{\odot} is consistent with the most accurate space-based determinations, particularly the one from *Reuven Ramathy High-Energy Solar Spectroscopic Imager*/Solar Aspect Sensor.

Key words: Sun: helioseismology – Sun: interior – Sun: rotation.

1 INTRODUCTION

Solar gravitational moments J_{2n} are coefficients that describe the rotation-induced deviation of the Sun's outer gravitational potential ϕ_{out} from a spherical configuration. Assuming an axial symmetry around the rotation axis, they intervene in the expression of ϕ_{out} as projection coefficients on the basis of Legendre polynomials:

$$\phi_{\text{out}}(r, u) = -\frac{GM_{\odot}}{r} \left[1 - \sum_{n=1}^{\infty} \left(\frac{R_{\odot}}{r} \right)^{2n} J_{2n} P_{2n}(u) \right]. \quad (1)$$

The odd terms have been omitted from the series in equation (1) because of equatorial symmetry. The quantities G , M_{\odot} , r , R_{\odot} , P_{2n} and $u = \cos \theta$, are respectively the gravitational constant, the solar mass, the distance from the centre of the Sun, the mean solar radius, the Legendre polynomials of degree $2n$, and the cosine of the colatitude of the Sun θ (angle to the rotation axis). The accurate determination of J_{2n} is of interest not only in solar physics but also in many other astrophysical applications. The most famous one is undoubtedly the test of general relativity (GR) resulting from the combination of the value of the quadrupole moment J_2 with the measurements of the anomalous precession of Mercury's orbit (Dicke 1964; Shapiro et al. 1972; Campbell et al. 1983; Lydon & Sofia 1996; Chapman 2008; Gough 2013). In the same way, J_2 can be used to constraint the Eddington-Robertson parameters in

the parametrized-post-newtonian theory of gravity, an alternative gravitation theory to GR (Pireaux & Rozelot 2003; Iorio 2005). In astrometry, an estimate of J_2 makes possible to study its effect on the astrometric (Kislik 1983; Bursa 1986) and celestial mechanics (Xu et al. 2011, 2017; Vaishwar, Kushvah & Mishra 2018) determination of planetary orbits and also on the dynamics of the earth–moon system (Bois & Girard 1999). For detailed reviews on the implication of J_{2n} in alternative theories of gravitation, high precision astrometry and celestial mechanics, readers are referred to the two articles by Rozelot, Damiani & Pireaux (2009) and Rozelot & Fazel (2013). In solar physics, J_{2n} indicate non-uniform mass and angular velocity distribution inside the Sun and their accurate knowledge would provide a good constraint on internal structure and rotation (Dicke & Goldenberg 1967; Ulrich & Hawkins 1981a, b; Paterno, Sofia & di Mauro 1996; Godier & Rozelot 1999; Armstrong & Kuhn 1999; Mecheri et al. 2004), and on solar cycle models through the study of their temporal evolution (Antia, Chitre & Gough 2008), complementing thus the constraints imposed by helioseismology.

Several observational and theoretical works have been undertaken to determine solar gravitational moments J_{2n} (mainly J_2). In general, the observational determinations are either from oblateness estimates based on the profile of the Sun's limb [Dicke & Goldenberg (1967), Dicke, Kuhn & Libbrecht (1986) using the Solar Distortion telescope, Hill & Stebbins (1975) using the *SCLERA telescope*, Lydon & Sofia (1996) using the Solar Disc Sextant (SDS) instrument, Rösch et al. (1996), Rozelot & Roesch (1997) using the Pic du Midi heliometer, Fivian et al. (2008) using the Solar Aspect Sensor (SAS) onboard of the *Reuven Ramathy High-Energy Solar Spectroscopic Imager*

* E-mail: redouane.mecheri@craag.edu.dz (RM);
Mustapha.Meftah@latmos.ipsl.fr (MM)

(*RHESSI*) satellite], or from astrometric observations of planetary orbit of Mercury and other minor planets such as Icarus (Lieske & Null 1969; Anderson et al. 1978; Afanaseva et al. 1990; Landgraf 1992; Pitjeva 2005) or form lunar laser ranging (LLR) data (Rozelot & Bois 1998). Theoretical expressions relating the solar gravitational moments J_{2n} to the inner structure and dynamics of a star can be determined using the theory of slowly rotating stars (Schwarzschild 1947; Sweet 1950). Early application of this theory to the Sun was done by Roxburgh (1964), Goldreich & Schubert (1967), and Gough (1981) in the context of analysing internal rotation. It was used for the determination of J_{2n} by Ulrich & Hawkins (1981a, b) using a simple quadratic rotation law. Several theoretical determinations followed Ulrich & Hawkins work, using two-dimensional helioseismically inferred rotation rates either in a parametric form (Paterno et al. 1996; Godier & Rozelot 1999; Roxburgh 2001; Mecheri et al. 2004) or through direct inversion of rotational frequency splitting (Gough 1982; Campbell et al. 1983; Duvall et al. 1984; Brown et al. 1989; Pijpers 1998; Armstrong & Kuhn 1999; Antia, Chitre & Thompson 2000; Antia et al. 2008). All these contributions computed values of J_{2n} either from a differential or an integral equation that was derived explicitly for the special case of $n = 1$ or $n = 2$. Exception is made to works by Armstrong & Kuhn (1999), Roxburgh (2001), and particularly Mecheri et al. (2004) who derived a convenient general form of the Poisson equation whose solution at the surface gives J_{2n} for any value of n .

In this work, we take over the above-mentioned equation [see Mecheri et al. 2004, equation (4)] and perform further algebraic calculations to derive a general integral equation relating J_{2n} to the internal rotation following the Green's functions method described by Pijpers (1998). This integral equation is then used to compute values of J_{2n} for $n = 1, 2, 3, 4$, and 5 taking into account new constraints on internal rotation provided by the high-resolution HMI (Helioseismic and Magnetic Imager) aboard of SDO (Solar Dynamics Observatory) helioseismic data covering the whole solar cycle 24. Our main equations are presented in Section 2. The results of our computations of J_{2n} are presented and discussed in Section 3. Finally, we give our principal conclusions in Section 4.

2 GENERAL INTEGRAL EQUATION FOR J_{2n}

Theoretical expressions relating the distortions of a star to the internal mass, density, and rotation can be obtained under the assumption of a slow rotation (i.e. centrifugal acceleration small compared to the gravitational acceleration) where all stellar structure quantities are described in terms of perturbations (with subscript 1) of the spherically symmetric non-rotating star (with subscript 0). The perturbations are thereby expanded on the basis of Legendre polynomials giving a gravitational potential inside the Sun ϕ_{int} as follows:

$$\phi_{\text{int}}(r, u) = \phi_0(r) + \phi_1(r, u) = \phi_0(r) + \sum_{n=1}^{\infty} \phi_{12n}(r) P_{2n}(u), \quad (2)$$

where ϕ_0 is the gravitational potential of a spherical Sun and ϕ_{12n} represent the projections of the perturbed gravitational potential ϕ_1 on the Legendre polynomials basis. The gravitational moments J_{2n} are given assuming the continuity of the gravitational potential at the solar surface, i.e. $\phi_{\text{int}}(R_{\odot}, u) = \phi_{\text{out}}(R_{\odot}, u)$, as follows:

$$J_{2n} = \frac{R_{\odot}}{GM_{\odot}} \phi_{12n}(R_{\odot}). \quad (3)$$

Applying this perturbation technique to stellar structure equations, Mecheri et al. (2004) derived a convenient form of the Poisson

equation for a general n that is given as follows:

$$\begin{aligned} \frac{d^2 \phi_{12n}}{dr^2} + \frac{2}{r} \frac{d\phi_{12n}}{dr} - (2n(2n+1) + UV) \frac{\phi_{12n}}{r^2} \\ = U \left((V+2) A_{2n} + r \frac{dA_{2n}}{dr} + B_{2n} \right) \end{aligned} \quad (4)$$

which was obtained by combining linearized equations governing the equilibrium of rotating star in which only first-order terms have been retained (Goldreich & Schubert 1968; Ulrich & Hawkins 1981a, b). The quantities $U = 4\pi \rho_0 r^3 / M_r$ and $V = d \ln \rho_0 / d \ln r$, which refer to a spherical non-rotating Sun, are obtained from solar models through the density ρ_0 and the mass M_r contained in a sphere of radius r inside the Sun. For a solar angular velocity $\Omega(r, u)$, the quantities A_{2n} and B_{2n} are given by

$$\begin{aligned} A_{2n}(r) &= \int_{-1}^1 a_{2n}(u) \Omega(r, u)^2 du \\ &= -\frac{1}{2n!} \frac{4n+1}{2^{2n+1}} \int_{-1}^1 u \Omega(r, u)^2 \frac{d^{2n-1}}{du^{2n-1}} (u^2 - 1)^{2n} du, \\ B_{2n}(r) &= \int_{-1}^1 b_{2n}(u) \Omega(r, u)^2 du \\ &= \frac{4n+1}{2} \int_{-1}^1 (1-u^2) P_{2n}(u) \Omega(r, u)^2 du. \end{aligned} \quad (5)$$

Following closely the treatment of Pijpers (1998) using the Green's functions method, it is possible to derive from the above general differential equation (4), a general integral equation giving ϕ_{12n} at the surface of the Sun:

$$\begin{aligned} \phi_{12n}(R_{\odot}) &= -\frac{R_{\odot}^{-2n}}{GM_{\odot}} \left[\frac{r^{2n}}{(2n+1)\psi_{2n} + r\psi'_{2n}} \right]_{r=R_{\odot}} \\ &\times \int_0^{R_{\odot}} r^2 U \left((V+2) A_{2n} + r \frac{dA_{2n}}{dr} + B_{2n} \right) \psi_{2n} dr, \end{aligned} \quad (6)$$

where $\psi_{2n}(r)$ is a regular solution at the origin (i.e. $\psi_{2n}(r) \propto r^{2n}$ as $r \rightarrow 0$) of equation (4) with a right-hand side identical to zero and $\psi'_{2n}(r)$ is its derivative with respect to r . Finally, using equation (3) and dimensionless variables $x = r/R_{\odot}$, $\omega^2 = \Omega^2(R_{\odot}^3/GM_{\odot})$, J_{2n} is given by

$$\begin{aligned} J_{2n} &= - \left[\frac{x^{2n}}{(2n+1)\psi_{2n} + x\psi'_{2n}} \right]_{x=1} \\ &\times \int_0^1 ((x^2(U-4)U\psi_{2n} - x^3U\psi'_{2n})A_{2n} + x^2U\psi_{2n}B_{2n}) dx \\ &= \int_0^1 \int_{-1}^1 F_{2n}(x, u) \omega(x, u)^2 du dx \end{aligned} \quad (7)$$

The normalized integration kernel $F_{2n}(x, u)$ is therefore given by

$$\begin{aligned} F_{2n}(x, u) &= - \left[\frac{x^{2n}}{(2n+1)\psi_{2n} + x\psi'_{2n}} \right]_{x=1} \\ &\times ((x^2(U-4)U\psi - x^3U\psi'_{2n})a_{2n} + x^2U\psi_{2n}b_{2n}) \end{aligned} \quad (8)$$

Note that for $n = 1$, equation (7) reduces to equation (23) of Pijpers (1998) in the case of general angular rotation $\omega(x, u)$ and to equation (12) of Gough (1981) for a radially dependent angular rotation $\omega(x)$.

3 RESULTS AND DISCUSSION

The calculated values of J_{2n} for $n = 1, 2, 3, 4$, and 5 together with previously published results also obtained using a helioseismic

Table 1. Values of solar gravitational moments J_{2n} ($n = 1, 2, 3, 4,$ and 5) computed using solar models from CESAM and ASTEC stellar evolution codes and rotation rates obtained from HMI and MDI helioseismic data, together with values from other authors also computed using helioseismic estimates of internal rotation.

Authors	Rotation data	$J_2 (\times 10^{-7})$	$J_4 (\times 10^{-9})$	$J_6 (\times 10^{-10})$	$J_8 (\times 10^{-11})$	$J_{10} (\times 10^{-12})$
Present work	SDO/HMI (CESAM)	2.211	-4.252	-1.282	5.897	-4.372
	SDO/HMI (ASTEC)	2.216	-4.256	-1.283	5.901	-4.375
	SoHO/MDI (CESAM)	2.204	-4.064	-1.136	5.404	-3.993
	SoHO/MDI (ASTEC)	2.208	-4.069	-1.137	5.408	-3.996
Antia et al. (2008)	GONG	2.22	-3.97	-0.8	1.1	7.4
	SoHO/MDI	2.18	-4.70	-2.4	-0.8	7.1
Mecheri et al. (2004)	SoHO/MDI	2.205	-4.455			
Roxburgh (2001)	SoHO/MDI (ISM)	2.208	-4.46	-2.80	1.49	
	SoHO/MDI (CSM)	2.206	-4.44	-2.79	1.48	
Antia et al. (2000)	GONG + SoHO/MDI	2.18	-4.64			
Armstrong & Kuhn (1999)	SoHO/MDI	2.22	-3.84			
Godier & Rozelot (1999)	SoHO/MDI	1.6				
Pijpers (1998)	GONG + SoHO/MDI	2.18				
Paterno et al. (1996)	IRIS + BISON + LOWL	2.22				
Brown et al. (1989)	SPO/Fourier Tachometer	1.7				
Duvall et al. (1984)	KPNO/McMath telescope	1.7				

estimates of internal rotation are given in Table 1, where a difference in sign convention has been taken into account concerning the results of Armstrong & Kuhn (1999) and Antia et al. (2000). They have been computed using equation (7), in which the function ψ_{2n} and the kernel F_{2n} are evaluated using the quantities U and V from two solar models obtained from CESAM (Morel & Lebreton 2008) and ASTEC (Christensen-Dalsgaard 2008) stellar evolution codes. For ω , we use time-averaged two-dimensional rotation rates obtained from SDO/HMI helioseismic data of full-disc (fd_V) dopplergrams available in the SDO HMI-AIA Joint Science Operations Center (JSOC) data base covering the period between 2010 April and 2020 July. For comparison purpose, we also compute J_{2n} using rotation rates provided by the Michelson Doppler Imager (MDI) onboard of the Solar and Heliospheric Observatory (SoHO), available in the same data base for the period between 1996 May and 2008 March. This comparison is all the more interesting as, unlike previous contributions of Table 1, it uses rotation rates obtained from an improved recent analysis of fd_V MDI helioseismic data (Larson & Schou 2015, 2018), which corrects for several geometric effects during spherical harmonic decomposition as well as some other physical effects such as the distortion of eigenfunctions by the differential rotation and the horizontal displacement at the solar surface. The HMI fd_V data, which require less geometric corrections, have been processed exactly in the same manner as the MDI fd_V data. The rotation rates for both data sets, have been calculated using two-dimensional regularized least-squares inversions (Schou et al. 1998) of odd rotational splitting coefficients of f-mode and p-mode frequencies. Fig. 1 shows superimposed time-averaged radial profiles at different latitudes of HMI (solid lines) and MDI (dashed lines) rotation. The two rotation profiles are very similar with only small differences at high latitude in the convective zone. However, a more pronounced difference can be noticed in deeper region inside the Sun below approximately $0.4R_{\odot}$. It should be noted that these two locations are regions in the Sun where rotation estimates are considered unreliable, but nevertheless, we use them in our calculations in the absence of other alternatives. Table 1 shows that, for the same solar model, the calculated values of J_{2n} from HMI and MDI rotation data have the same order of magnitude with however a slightly larger absolute values for HMI results. The difference is approximately of the order of 0.3 per cent




Figure 1. Time-averaged radial profiles of HMI (solid lines) and MDI (dashed lines) rotation, obtained from helioseismic data of full disc (fd_V) dopplergrams, given each 15° from equator (top) to pole (bottom).

for J_2 and increases for higher multipole moments to 4 per cent for J_4 , 11 per cent for J_6 , 8 per cent for J_8 , and 9 per cent for J_{10} , presumably due to the difference in the rotation deep inside the Sun for J_2 and in the outer layers for higher multipole moments. Indeed, as already emphasized by Antia et al. (2008), high-order multipole moments are predominantly determined from the contributions of the outer layers of the Sun where their integration kernels are principally concentrated as shown in Figs 2 and 3 (for $n = 2, 3, 4,$ and 5), exhibiting substantial variation with latitude, with local minima and maxima positioned approximately at radial distances between $0.8R_{\odot}$ and $0.9R_{\odot}$. On the other hand, the major contribution to J_2 comes from deeper regions where the corresponding integration kernel (see Figs 2 and 3, for $n = 1$) exhibits its greatest value also at $r \approx 0.77R_{\odot}$ principally at low latitudes around 34° . Note that the sensitivity of high-order multipole moments to the differential rotation in the outer layers of the Sun has been evidenced for J_4 by Mecheri et al. (2004), particularly the effect due to the presence of a subsurface radial gradient. More pronounced differences in the values of J_{2n} have been found by Antia et al. (2008) using GONG and MDI rotation rates (Table 1) which, according to the authors, are the




Figure 2. Three-dimensional plots (top panels) of the normalized kernel F_{2n} as a function of $x = r/R_{\odot}$ and latitude for $n = 1, 2, 3, 4,$ and 5 and their corresponding contour plots (bottom panels).




Figure 3. Plots of latitudinal (top panels) and radial (bottom panels) cuts of the normalized kernel F_{2n} for $n = 1, 2, 3, 4,$ and $5,$ respectively for different values of $x = r/R_{\odot} = 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8,$ and 0.9 and colatitude θ ($^{\circ}$) = $10^{\circ}, 20^{\circ}, 30^{\circ}, 40^{\circ}, 50^{\circ}, 60^{\circ}, 70^{\circ},$ and $80^{\circ}.$

direct consequence of the differences between the measured splitting coefficients. For J_2 , our results are in close agreement with most of the evaluations reported in Table 1, except for those of Godier & Rozelot (1999), Brown et al. (1989), and Duvall et al. (1984), which are considerably smaller. For Duvall et al. (1984) and Brown et al. (1989), this difference is principally due to the very early helioseismic data used in the inference of internal rotation, restricted to regions close to the equator for the former. Surprisingly, Godier & Rozelot's value of J_2 is also largely inferior to the ones obtained by Mecheri et al. (2004) and Roxburgh (2001) despite of using exactly the same rotation law. Higher order multipole moments $J_6, J_8,$ and J_{10} have the same order of magnitude as those of Roxburgh (2001) and Antia et al. (2008), with however sensitively different exact values. It is worth mentioning that Roxburgh's results have been obtained using a rotation model in a parametric form which roughly approximate the internal rotation inferred from helioseismology. Note from Table 1, that for the same rotation data, our results from the two solar models

are in very good agreement with insignificant differences inferior to 0.2 per cent. Similar compatibility was found by Roxburgh (2001) for $J_2, J_4, J_6,$ and J_8 computed using inverted (ISM) and calculated (CSM) solar models (see Table 1). This compatibility is also verified when comparing the values of J_2 and J_4 obtained respectively by Roxburgh (2001) and Mecheri et al. (2004) using distinct solar models but the same model of rotation of Kosovichev (1996). Both authors pointed out that the differential rotation in the convective zone introduces only a diminution of 0.5 per cent of the value of J_2 with comparison to the one obtained for a Sun rotating uniformly at the rotation rate of the radiative interior. This indicates that the quadrupole moment J_2 is basically determined by a spherically averaged rotation whose departure from interior rotation is relatively small (Roxburgh 2001).

On the other hand, the sensitivity of high-order multipole moments to the differential rotation in the convective zone makes them responsive to the observed temporal variation of the latitudinal

component of the angular rotation (Howe 2009) exhibiting changes either correlated or anti-correlated with magnetic activity (Antia et al. 2008), whereas by contrast, J_2 , which is more sensitive to the radiative zone rotation, do not present significant variation basically because the angular rotation in deeper layers inside the Sun do not show reliable temporal fluctuations. However, observational temporal changes of J_2 have been recently evidenced by Rozelot & Eren (2020) from the analysis of the perihelion precession measurements of several planets taken at different periods. Rozelot & Eren reported a mean weighted value of $J_2 = (2.17 \pm 0.06) \times 10^{-7}$, which is very compatible with our results. We mention also the good compatibility of our results with the value $J_2 = (2.25 \pm 0.09) \times 10^{-7}$ deduced from the measurements of the precession of Mercury's perihelion obtained from ranging data of the *MESSENGER* (MErcury Surface, Space ENvironment, GEochemistry, and Ranging) spacecraft (Park et al. 2017). They are however not compatible with the earlier values of $J_2 = (1.8 \pm 5.1) \times 10^{-7}$ and $J_4 = (9.8 \pm 4.6) \times 10^{-7}$ found by Lydon & Sofia (1996) from the SDS balloon-borne experiment.

The calculated quadrupole moment J_2 gives an approximate estimate of the theoretical solar oblateness Δ_\odot via the formula $\Delta_\odot \approx (3/2)J_2 + (\delta r/R_\odot)$, where $\delta r/R_\odot = 8.1 \times 10^{-6}$ (Dicke 1970), yielding $\Delta_\odot \approx 8.43 \times 10^{-6}$. This value is in fair agreement with most of the observational oblateness estimates from the analysis of space-based solar limb shape measurements, namely by SoHO/MDI (Emilio et al. 2007), SODISM (solar diameter imager and surface mapper) onboard of PICARD spacecraft (Irbah et al. 2014; Meftah et al. 2015), and SDO/HMI (Meftah et al. 2016; Irbah et al. 2019). It is worth to note also its excellent agreement with the most accurate oblateness measurement to date ($8.35 \pm 0.15) \times 10^{-6}$ obtained from RHESSI/SAS limb data (Fivian et al. 2008).

Finally, the calculation of J_{2n} and resulting Δ_\odot for all MDI and HMI rotation data available for an entire period of two solar cycles can make possible to explore their temporal variation and possible relation to magnetic activity and therefore allow for a direct comparison with optical limb shape inference of solar oblateness. The study of the dynamic evolution of these quantities from model calculations is an ongoing work that will be the subject of a future publication.

4 CONCLUSIONS

The precise theoretical estimate of solar gravitational moment J_{2n} is very important in many astrophysical applications. In this work, we have used new HMI solar rotation rates to calculate updated values of J_{2n} (for $n = 1, 2, 3, 4$, and 5) by mean of a general integral equation derived in the framework of the theory of slowly rotating stars. The results revealed a good agreement with most of the earlier helioseismic estimates particularly for J_2 and J_4 , whereas J_6 , J_8 , and J_{10} agree as an order of magnitude but however differ in their exact values. On the other hand, the comparison with the calculation results obtained using MDI rotation rates yielded a difference of the order of ≈ 0.3 per cent for the quadrupole moment J_2 . This difference increases by one order of magnitude for higher order multipole moments indicating their greater sensitivity, as compared to J_2 , to the differences between HMI and MDI rotation rates, particularly in the outer layers of the Sun. The calculated value of $J_2 \approx 2.21 \times 10^{-7}$ is in agreement with the observational value $J_2 = 2.25 \times 10^{-7}$ provided by the high precision measurements of the precession of Mercury's perihelion obtained from ranging data of the *MESSENGER* spacecraft. The resulting theoretical value of the solar oblateness Δ_\odot was found to be approximately equal to 8.43×10^{-6}

that is in perfect accordance with the most accurate space-based observational estimate of 8.35×10^{-6} obtained by RHESSI/SAS. The dynamic evolution of J_{2n} and Δ_\odot and its eventual correlation with magnetic activity during solar cycles 23 and 24 is an ongoing work for a planned subsequent contribution.

ACKNOWLEDGEMENTS

This work has been performed with support of the Centre National d'Etudes Spatiales (CNES), Stanford University, and the University of Hawai'i System. The HMI data used are courtesy of NASA/SDO and the HMI science teams. This work also utilizes data from the SoHO/MDI instrument. SoHO is a project of international cooperation between ESA and NASA. The authors thank the referee for the constructive remarks and suggestions that helped improving the quality of the paper.

DATA AVAILABILITY

All MDI and HMI rotation data used in this study are available online from the global helioseismology pipeline on the website of the JSOC at http://jsoc.stanford.edu/MDI/Global_products.html for MDI and likewise at http://jsoc.stanford.edu/HMI/Global_products.html for HMI, as cited in Larson & Schou (2015, 2018).

REFERENCES

- Afanaseva T. I., Kislik M. D., Kolyuka Y. F., Tikhonov V. F., 1990, *SvA*, 34, 670
- Anderson J. D., Keesey M. S. W., Lau E. L., Standish E. M. J., Newhall X. X., 1978, *Acta Astronaut.*, 5, 43
- Antia H. M., Chitre S. M., Thompson M. J., 2000, *A&A*, 360, 335
- Antia H. M., Chitre S. M., Gough D. O., 2008, *A&A*, 477, 657
- Armstrong J., Kuhn J. R., 1999, *ApJ*, 525, 533
- Bois E., Girard J. F., 1999, *Celest. Mech. Dyn. Astron.*, 73, 329
- Brown T. M., Christensen-Dalsgaard J., Dziembowski W. A., Goode P., Gough D. O., Morrow C. A., 1989, *ApJ*, 343, 526
- Bursa M., 1986, *Bull. Astron. Inst. Czech.*, 37, 312
- Campbell L., McDow J. C., Moffat J. W., Vincent D., 1983, *Nature*, 305, 508
- Chapman G. A., 2008, *Science*, 322, 535
- Christensen-Dalsgaard J., 2008, *Ap&SS*, 316, 13
- Dicke R. H., 1964, *Nature*, 202, 432
- Dicke R. H., 1970, *ApJ*, 159, 1
- Dicke R. H., Goldenberg H. M., 1967, *Phys. Rev. Lett.*, 18, 313
- Dicke R. H., Kuhn J. R., Libbrecht K. G., 1986, *ApJ*, 311, 1025
- Duvall T. L. J., Dziembowski W. A., Goode P. R., Gough D. O., Harvey J. W., Leibacher J. W., 1984, *Nature*, 310, 22
- Emilio M., Bush R. I., Kuhn J., Scherrer P., 2007, *ApJ*, 660, L161
- Fivian M. D., Hudson H. S., Lin R. P., Zahid H. J., 2008, *Science*, 322, 560
- Godier S., Rozelot J.-P., 1999, *A&A*, 350, 310
- Goldreich P., Schubert G., 1967, *ApJ*, 150, 571
- Goldreich P., Schubert G., 1968, *ApJ*, 154, 1005
- Gough D. O., 1981, *MNRAS*, 196, 731
- Gough D. O., 1982, *Nature*, 298, 334
- Gough D., 2013, *Sol. Phys.*, 287, 9
- Hill H. A., Stebbins R. T., 1975, *ApJ*, 200, 471
- Howe R., 2009, *Living Reviews in Sol. Phys.*, 6, 1
- Iorio L., 2005, *A&A*, 433, 385
- Irbah A., Meftah M., Hauchecorne A., Djafer D., Corbard T., Bocquier M., Momar Cisse E., 2014, *ApJ*, 785, 89
- Irbah A., Mecheri R., Damé L., Djafer D., 2019, *ApJ*, 875, L26
- Kislik M. D., 1983, *SvA Lett.*, 9, 296
- Kosovichev A. G., 1996, *ApJ*, 469, L61
- Landgraf W., 1992, *Sol. Phys.*, 142, 403

- Larson T. P., Schou J., 2015, *Sol. Phys.*, 290, 3221
Larson T. P., Schou J., 2018, *Sol. Phys.*, 293, 29
Lieske J. H., Null G. W., 1969, *AJ*, 74, 297
Lydon T. J., Sofia S., 1996, *Phys. Rev. Lett.*, 76, 177
Mecheri R., Abdelatif T., Irbah A., Provost J., Berthomieu G., 2004, *Sol. Phys.*, 222, 191
Meftah M. et al., 2015, *Sol. Phys.*, 290, 673
Meftah M., Hauchecorne A., Bush R. I., Irbah A., 2016, *Adv. Space Res.*, 58, 1425
Morel P., Lebreton Y., 2008, *Ap&SS*, 316, 61
Park R. S., Folkner W. M., Konopliv A. S., Williams J. G., Smith D. E., Zuber M. T., 2017, *AJ*, 153, 121
Paterno L., Sofia S., di Mauro M. P., 1996, *A&A*, 314, 940
Pijpers F. P., 1998, *MNRAS*, 297, L76
Pireaux S., Rozelot J. P., 2003, *Ap&SS*, 284, 1159
Pitjeva E. V., 2005, *Astron. Lett.*, 31, 340
Rösch J., Rozelot J. P., Deslandes H., Desnoux V., 1996, *Sol. Phys.*, 165, 1
Roxburgh I. W., 1964, *Icarus*, 3, 92
Roxburgh I. W., 2001, *A&A*, 377, 688
Rozelot J., Bois E., 1998, in Balasubramaniam K. S., Harvey J., Rabin D., eds, ASP Conf. Ser. Vol. 140, Synoptic Solar Physics. Astron. Soc. Pac., San Francisco, p. 75
Rozelot J. P., Eren S., 2020, *Adv. Space Res.*, 65, 2821
Rozelot J. P., Fazel Z., 2013, *Sol. Phys.*, 287, 161
Rozelot J. P., Roesch J., 1997, *Sol. Phys.*, 172, 11
Rozelot J. P., Damiani C., Pireaux S., 2009, *ApJ*, 703, 1791
Schou J. et al., 1998, *ApJ*, 505, 390
Schwarzschild M., 1947, *ApJ*, 106, 427
Shapiro I. I., Pettengill G. H., Ash M. E., Ingalls R. P., Campbell D. B., Dyce R. B., 1972, *Phys. Rev. Lett.*, 28, 1594
Sweet P. A., 1950, *MNRAS*, 110, 548
Ulrich R. K., Hawkins G. W., 1981a, *ApJ*, 246, 985
Ulrich R. K., Hawkins G. W., 1981b, *ApJ*, 249, 831
Vaishwar A., Kushvah B. S., Mishra D. P., 2018, *Few-Body Syst.*, 59, 4
Xu Y., Yang Y., Zhang Q., Xu G., 2011, *MNRAS*, 415, 3335
Xu Y., Shen Y., Xu G., Shan X., Rozelot J.-P., 2017, *MNRAS*, 472, 2686

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.