



HAL
open science

Stratified Radiative Transfer for Multidimensional Fluids

François Golse, Olivier Pironneau

► **To cite this version:**

François Golse, Olivier Pironneau. Stratified Radiative Transfer for Multidimensional Fluids. 2021.
hal-03306461v1

HAL Id: hal-03306461

<https://hal.sorbonne-universite.fr/hal-03306461v1>

Preprint submitted on 29 Jul 2021 (v1), last revised 7 Nov 2023 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Stratified Radiative Transfer for Multidimensional Fluids¹

François Golse², Olivier Pironneau³

Abstract

New mathematical and numerical results are given for the coupling of the temperature equation of a fluid with Radiative Transfer: existence and uniqueness and a convergent monotone numerical scheme. The technique is shown to be feasible for studying the temperature of lake Lemman heated by the sun and for the earth atmosphere to study the effects of greenhouse gases.

Keywords: Radiative Transfer, Navier-Stokes equations, Integral equations, Numerical Method, Convergence, Climate.

1. Introduction

Fifty years ago, the second author was admitted to the prestigious Dept of Applied Math. & Theoretical Physics at Cambridge, UK, headed then by Sir James Lighthill. Two `IBM` card punchers connected to the computing center – also one of the best in the world in those days – had been relegated to the basement; to use them was frowned upon as a threat to the speciality of the lab: clever analytic approximations and other multiple scales expansions of special cases of the Navier-Stokes equations.

It took a decade to prove that computer simulations for fluids were not only possible, but also useful to industry. A colleague from the wind tunnels in Modane told us then that an airplane could never be designed and validated by a numerical simulation. True to this wrong prediction however, many ad-hoc turbulence models had to be devised: it was only by a combined theoretical, experimental and computational (TEC) effort that the world's first complete airplane could be simulated at Dassault Aviation in 1979 and that airplanes have since be flown safely without the difficult certification stamps of wind tunnels.

It was also a success of the top-down approach to CFD. The “JLL” (Lions) school of applied mathematics had the luck of being taken seriously by a few French high-tech industry labs. This was not the case in the USA where the head of a national research funding agency had ruled out variational methods (leading to finite volumes and finite elements for fluids) as “incomprehensible by aeronautical engineers”, thereafter forcing all numerical schemes to be in the class of body fitted structured meshes, an impossible task for airplanes.

The top-down approach to a problem could be defined by saying that the mathematical model is defined first, then shown to be well posed and then approximated numerically by

¹Will be submitted for publication in *Compte-Rendus de Mécanique*

²francois.golse@polytechnique.edu, CMLS, Ecole polytechnique, 91128 Palaiseau Cedex, France

³olivier.pironneau@sorbonne-universite.fr, LJLL, Sorbonne Université, Paris, France.

convergent algorithms. The bottom-up approach is when the problem is made of several modules, studied independently, and patched together at the algorithmic level.

The downside of the top-down approach - from functional analysis to numerical methods - is that it may discard important faster algorithms for which convergence are not known. This was the case for compressible flows in the nineties for which the bottom-up approach pragmatically patched different turbulence and/or numerical models in different zones with the drawback that it was difficult to assert that the computed solution was one of the original problem.

In the numerical simulations which fill the supercomputing centers today, CFD is often only one part of a multi-physics model. Such are the combustion and climate computations. Both need, at least, radiative transfer and chemistry modules.

While the top-down approach is successful in computational chemistry [CDK⁺03], mathematical analysis of climate models is still in progress. The three dimensional Primitive Equations with hydrostatic and geostrophic approximations have been shown to be well posed (see [LTW94],[AG01],[CCT20] and the bibliography therein) and so are the multi-layered Shallow Water equations for the oceans [CLGP13]; but even if the coupled ocean-atmosphere is mathematically well-posed, it is very far from the complete model used in climatology. No doubt when a new numerical climate project is proposed, such as [DDT⁺15], a top-down approach is made [EDK19], but soon overwhelmed by the complexity of the task when more modules are added.

Radiative transfer – one such module that needs to be added – is essential in astrophysics [Cha50] to derive the composition of stars, in nuclear engineering to predict plasma [DL00], in combustion for engines [ACP⁺09], and many other fields like solar panels [ZPS⁺21] and even T-shirts [ZPS⁺21]!

In the eighties, at CEA, R. Dautray [DL00] headed a team of applied mathematicians who used the top-down approach in nuclear engineering. The first author was in close contact with them. But turning his expertise on radiative transfer to climate modeling is not straightforward.

Books on radiative transfer for the atmosphere are numerous, such as [GY61], [Boh06] and [ZT03]; but to speed-up codes, the documentation manual of climate models reveal that many approximations are made. For instance LMDZ refers to a model proposed by Fouquart [Fou88][Mor91] which suggests that empirical formulas are used in addition to simplified numerical schemes to speed-up the computations. The formulas for the absorption, scattering and albedo coefficients are complex and adapted to reproduce the experimental data. In other words the gap is wide between practice and fundamentals as seen by Fowler [Fow11] and Chandrasekhar [Cha50], for instance.

Coupling radiative transfer to the Navier-Stokes system using the top-down approach is the topic of this article. The problem is shown well posed in the context of a stratified atmosphere and a numerical method – derived from the mathematical proof of well posedness – is proposed. It is accurate in the sense that there are no singular functions or integrals to approximate. It is fast compared to the fluid solver to which it is coupled but of course not as fast as empirical formulas.

2. Radiative transfer and the temperature equation

Let us begin with a simple problem: the effect of sunlight on a lake Ω . Let $I_\nu(\mathbf{x}, \omega, t)$ be the light intensity of frequency ν at $\mathbf{x} \in \Omega$, in the direction $\omega \in \mathbb{S}^2$, the unit sphere, at time $t \in (0, T)$. Let T, ρ, \mathbf{u} be the temperature, density and velocity in the lake. Energy, momentum and mass conservations (see [Pom73],[Fow11]) yields (1),(2),(3):

2.1. The fundamental equations

Given I_ν, T at time zero, find I_ν, T for all $\{\mathbf{x}, \omega, t, \nu\} \in \Omega \times \mathbb{S}^2 \times (0, T) \times \mathbb{R}^+$ such that

$$\frac{1}{c} \partial_t I_\nu + \boldsymbol{\omega} \cdot \nabla I_\nu + \rho \bar{\kappa}_\nu a_\nu \left[I_\nu - \frac{1}{4\pi} \int_{\mathbb{S}^2} p(\boldsymbol{\omega}, \boldsymbol{\omega}') I_\nu(\boldsymbol{\omega}') d\boldsymbol{\omega}' \right] = \rho \bar{\kappa}_\nu (1 - a_\nu) [B_\nu(T) - I_\nu],$$

$$\partial_t T + \mathbf{u} \cdot \nabla T - \kappa_T \Delta T = -\nabla \cdot \int_0^\infty \int_{\mathbb{S}^2} I_\nu(\boldsymbol{\omega}') \boldsymbol{\omega} d\boldsymbol{\omega} d\nu. \quad (2)$$

$$\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} - \frac{\mu_F}{\rho} \Delta \mathbf{u} + \frac{1}{\rho} \nabla p = \mathbf{g}, \quad \nabla \cdot \mathbf{u} = 0, \quad \partial_t \rho + \nabla \cdot (\rho \mathbf{u}) = 0, \quad (3)$$

where ∇, Δ are with respect to \mathbf{x} , $B_\nu(T) = \frac{2\hbar\nu^3}{c^2 [e^{\frac{\hbar\nu}{kT}} - 1]}$, is the Planck function, \hbar is the

Planck constant, c is the speed of light in the medium and k is the Boltzmann constant. The absorption coefficient $\kappa_\nu := \rho \bar{\kappa}_\nu$ is the percentage of light absorbed per unit length, $a_\nu \in (0, 1)$ is the scattering albedo, $\frac{1}{4\pi} p(\boldsymbol{\omega}, \boldsymbol{\omega}')$ is the probability that a ray in the direction $\boldsymbol{\omega}'$ scatters in the direction $\boldsymbol{\omega}$. The constants κ_T and μ_F are the thermal and molecular diffusions; \mathbf{g} is the gravity.

Existence of solution for (3) has been established by P-L. Lions [Lio96].

As $c \gg 1$, in a regime where $\frac{1}{c} \partial_t I_\nu \ll 1$, integrating (1) in $\boldsymbol{\omega}$ leads to an alternative form for (2):

$$\partial_t T + \mathbf{u} \cdot \nabla T - \kappa_T \Delta T = - \int_0^\infty \rho \bar{\kappa}_\nu (1 - a_\nu) \left(4\pi B_\nu(T) - \int_{\mathbb{S}^2} I_\nu(\boldsymbol{\omega}) d\boldsymbol{\omega} \right) d\nu. \quad (4)$$

As usual, boundary conditions must be given. Dirichlet or Neumann conditions may be prescribed for \mathbf{u} and T on $\partial\Omega$. For the light intensity equation, I_ν should be given at all times on $\{(\mathbf{x}, \boldsymbol{\omega}) \in \partial\Omega \times \mathbb{S}^2 : \mathbf{n}(\mathbf{x}) \cdot \boldsymbol{\omega} < 0\}$, where \mathbf{n} is the outer unit normal of $\partial\Omega$. Finally ρ should be specified on $\partial\Omega$ when $\mathbf{u} \cdot \mathbf{n} < 0$.

2.2. Grey Medium

When κ_ν and a_ν are independent of ν - a so-called *grey medium* (cf. [Fow11], p. 70)- the problem can be written in terms of $I = \int_0^\infty I_\nu d\nu$:

$$\boldsymbol{\omega} \cdot \nabla I + \kappa a \left[I - \frac{1}{4\pi} \int_{\mathbb{S}^2} p(\boldsymbol{\omega}, \boldsymbol{\omega}') I(\boldsymbol{\omega}') d\boldsymbol{\omega}' \right] = \kappa (1 - a) (B_0 T^4 - I), \quad (5)$$

$$\partial_t T + \mathbf{u} \cdot \nabla T - \kappa_T \Delta T = -\kappa (1 - a) 4\pi \left(B_0 T^4 - \frac{1}{4\pi} \int_{\mathbb{S}^2} I(\boldsymbol{\omega}) d\boldsymbol{\omega} \right), \quad (6)$$

where B_0 comes from the Boltzmann-Stefan law:

$$\int_0^\infty \frac{2\hbar\nu^3}{c^2 [e^{\frac{\hbar\nu}{kT}} - 1]} d\nu = \left(\frac{\hbar}{kT} \right)^{-4} \frac{2\hbar}{c^2} \int_0^\infty \frac{\left(\frac{\hbar\nu}{kT} \right)^3}{e^{\frac{\hbar\nu}{kT}} - 1} d\frac{\hbar\nu}{kT} = B_0 T^4 \text{ with } B_0 := \frac{2k^4 \pi^4}{\hbar^3 c^2 15}.$$

2.3. Vertically stratified cases: spatial invariance

Let (x, y, z) be a cartesian frame with z the altitude/depth. The sun being very far, the light source on the lake is independent of x and y . Then, assuming that T' varies slowly with x and y , in the sense that

$$(H) \quad \partial_z I_\nu \gg \partial_x I_\nu, \quad \partial_z I_\nu \gg \partial_y I_\nu, \quad (7)$$

then (1),(2) become [ZT03]

$$\mu \partial_z I_\nu + \kappa_\nu I_\nu = \kappa_\nu (1 - a_\nu) B_\nu(T) + \frac{\kappa_\nu a_\nu}{2} \int_{-1}^1 p(\mu, \mu') I_\nu(z, \mu') d\mu' \quad (8)$$

$$I_\nu(z_M, \mu)|_{\mu < 0} = Q^-(\mu) B_\nu(\bar{T}_S), \quad I(z_m, \mu)|_{\mu > 0} = 0, \quad (9)$$

$$\partial_t T + \mathbf{u} \cdot \nabla T - \kappa_T \Delta T = -4\pi \int_0^\infty \kappa_\nu (1 - a_\nu) \left(B_\nu(T) - \frac{1}{2} \int_{-1}^1 I_\nu d\mu \right) d\nu, \quad \partial_n T|_{\partial\Omega} = 0. \quad (10)$$

where $z_M(x, y)$ and $z_m(x, y)$ are max and min of z such that $(x, y, z) \in \Omega$, μ is the cosine of the angle ω to the vertical axis, $Q^-(\mu) = -\mu Q' \cos \theta$ is the sunlight intensity when θ is the latitude, and \bar{T}_S is the temperature of the sun; we have assumed that the sun is a black body and that no light comes back from the bottom of the lake. Here \mathbf{u} is given, solenoidal and regular enough for (10) to make sense.

Remarks 1.

- Hypothesis (H) will hold if T varies slowly with x, y . It will be so if \mathbf{u} is almost horizontal and the vertical cross sections of Ω depend slowly on x, y . Turbulent flows do not satisfy this criteria.
- According to our definition of top-down analysis, the problem investigated is (8),(9),(10), not (1),(2),(3), justifying the restriction “stratified” in the title.
- All terms of (10) must be kept, except maybe, $\kappa_T \partial_{xx} T$ and $\kappa_T \partial_{yy} T$, but neglecting them renders the boundary conditions mathematically difficult.
- We shall ignore the mathematical difficulty induced by the boundary condition $\partial_n T|_{\partial\Omega} = 0$ when the intersection of the side of the lake with the water surface is not at right angle.

2.4. The vertically stratified grey problem

For a grey medium (8),(10) become

$$(P^1) \begin{cases} \mu \partial_z I + \kappa I = \kappa(1 - a) B_0 T^4 + \frac{\kappa a}{2} \int_{-1}^1 p I d\mu', \quad I|_{z_M, \mu < 0} = -\mu Q B_0 \bar{T}_S^4, \quad I|_{z_m, \mu > 0} = 0, \\ \partial_t T + \mathbf{u} \cdot \nabla T - \kappa_T \Delta T = -4\pi \kappa(1 - a) \left(B_0 T^4 - \frac{1}{2} \int_{-1}^1 I d\mu \right), \quad \partial_n T|_{\partial\Omega} = 0. \end{cases} \quad (11)$$

2.5. Elimination of I when the scattering is isotropic

Denote the exponential integral and the mean light intensity respectively by

$$E_m(x) := \int_0^1 \mu^{m-2} e^{-\frac{x}{\mu}} d\mu, \quad J(z) := \frac{1}{2} \int_{-1}^1 I(z, \mu) d\mu.$$

Then the method of characteristics applied to (11) gives

$$(P^2) \begin{cases} J(z) = \frac{1}{2} Q B_0 \bar{T}_S^4 E_3(\kappa(z_M - z)) + \frac{1}{2} \int_{z_m}^{z_M} \kappa E_1(\kappa|s - z|) ((1-a)B_0 T_s^4 + aJ(s)) ds, \\ \partial_t T + \mathbf{u} \cdot \nabla T - \kappa_T \Delta T = -4\pi\kappa(1-a) (B_0 T^4(z) - J(z)) \end{cases} \quad (12)$$

Note that to improve readability, we write indifferently $T(z)$ or T_z .

2.6. No scattering

Let $T_e(z) = (\frac{1}{2} Q E_3(\kappa|z_M - z|))^{\frac{1}{4}} \bar{T}_S$ and assume that $a = 0$, then

$$(P^3) \begin{cases} (4\pi\kappa B_0)^{-1} (\partial_t T + \mathbf{u} \cdot \nabla T - \kappa_T \Delta T) + T^4 = T_e^4 + \frac{1}{2} \int_{z_m}^{z_M} \kappa E_1(\kappa|s - z|) T_s^4 ds, & \partial_n T|_{\partial\Omega} = 0. \end{cases} \quad (13)$$

2.7. Algorithm for (P^3) in the stationary static case

Assume T stationary and $\mathbf{u} = 0$. Let $\bar{\kappa}_T = (4\pi\kappa B_0)^{-1} \kappa_T$.

Generate $\{T^n\}_{n \geq 0}$ from $T^0 = 0$ by,

$$\begin{cases} (T^{n+\frac{1}{2}})^4 := T_e^4 + \frac{1}{2} \int_{z_m}^{z_M} \kappa E_1(\kappa|s - z|) T_s^{n4} ds, & T^{n+\frac{1}{2}} \geq 0 \\ -\bar{\kappa}_T \Delta T^{n+1} + (T_+^{n+1})^4 = (T^{n+\frac{1}{2}})^4, & \partial_n T^{n+1}|_{\partial\Omega} = 0. \end{cases} \quad (14)$$

where $T_+ = \max(T, 0)$. Note that $T \mapsto -\bar{\kappa}_T \Delta T + T_+^4$ is a monotone operator for which Newton or fixed point iterations can be applied to solve the PDE. To prove monotone convergence, the following result is needed.

Lemma 1. $C_1(\kappa) := \frac{1}{2} \max_z \int_0^Z \kappa E_1(\kappa|s - z|) ds < 1$.

Proof

$$\begin{aligned} \int_0^X E_1(x) dx &= \int_1^\infty \int_0^X \frac{e^{-xt}}{t} dx dt = \int_1^\infty \frac{1 - e^{-Xt}}{t^2} dt < \int_1^\infty \frac{1}{t^2} dt = 1. \\ \Rightarrow \kappa \int_0^Z E_1(\kappa|\tau - t|) dt &= \int_0^{\kappa Z} E_1(|s - \kappa\tau|) ds = \int_0^{\kappa\tau} E_1(\kappa\tau - s) ds + \int_{\kappa\tau}^{\kappa Z} E_1(s - \kappa\tau) ds \\ &= \int_0^{\kappa\tau} E_1(\theta) d\theta + \int_0^{\kappa(Z-\tau)} E_1(\theta) d\theta < 2. \end{aligned} \quad (15)$$

□

Theorem 1.

$\{T^n\}_{n \geq 0}$ generated by Algorithm (14) converges to a solution of (13) and the convergence is monotone: $T^{n+1}(\mathbf{x}) > T^n(\mathbf{x})$ for all \mathbf{x} and all n .

Proof: From (14)

$$(T^{n+\frac{1}{2}})^4 \leq |T_e^4|_\infty + C_1(\kappa)|T^n|_\infty^4.$$

By the maximum principle for the PDE in (14), $T^{n+1} \geq 0$ and $|T^{n+1}|_\infty \leq |T^{n+\frac{1}{2}}|_\infty$, therefore

$$|T^{n+1}|^4 \leq |T_e^4|_\infty + C_1(\kappa)|T^n|_\infty^4.$$

Hence $|T^{n+1}|_\infty$ is bounded. Assume that $T^n \geq T^{n-1}$. The convergence is monotone because

$$(T^{n+\frac{1}{2}})^4 - (T^{n-\frac{1}{2}})^4 = \frac{1}{2} \int_{z_m}^{z_M} \kappa E_1(\kappa|s-z|) [(T_s^n)^4 - (T_s^{n-1})^4] \geq 0,$$

and as

$$-\bar{\kappa}_T \Delta(T^{n+1} - T^n) + b(T^{n+1} - T^n) = (T^{n+\frac{1}{2}})^4 - (T^{n-\frac{1}{2}})^4 \quad (16)$$

with $b = ((T^{n+1})^2 + (T^n)^2)(T^{n+1} + T^n) \geq 0$, the maximum principle implies that $T^{n+1} - T^n \geq 0$. □

Remark 1. Generalization of the above result to (P^3) is straightforward because the maximum principle holds also for the temperature equation with convection. Consequently it seems doable to extend the above to the system (2),(3). When the density variations with the temperature are small the Boussinesq approximation can be used in conjunction with (13):

$$(P^4) \left\{ \begin{array}{l} (4\pi\kappa B_0)^{-1}(\partial_t T + \mathbf{u} \cdot \nabla T - \kappa_T \Delta T) + T^4 = T_e^4 + \frac{1}{2} \int_{z_m}^{z_M} \kappa E_1(\kappa|t-z|) T_s^4 ds, \\ \partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} - \nu_F \Delta \mathbf{u} + \nabla p = -b(T - T_0)\mathbf{g}, \quad \nabla \cdot \mathbf{u} = 0, \end{array} \right. \quad (17)$$

with \mathbf{u}, T given at $t = 0$ and \mathbf{u} or $\partial_n \mathbf{u}$ or $p\mathbf{n} + \nu_T \partial_n \mathbf{u}$ and $\partial_n T = 0$ or T given on $\partial\Omega$. The kinematic viscosity $\nu_F = \mu_F/\rho$ is taken constant; b is a measure of $\partial_T \rho$ and T_0 is the average temperature. See [Att09], for instance, for the mathematical analysis of the Boussinesq-Stefan problem (similar to (P^4) without the T^4 terms).

3. Numerical tests

Earth sees the sun as a black body at temperature $\bar{T}_S = 5800\text{K}$ radiating with an intensity $Q' = 1370\text{W}/\text{m}^2$ of which 70% reach the ground, giving at noon in Milano $Q = 1370 \times 0.7 \cos \frac{\pi}{4} = 678$.

For water $\rho = 1000\text{kg}/\text{m}^3$; light absorption is $\kappa = 0.1$ for one meter and thermal diffusivity of water is $\kappa_T = 1.5 \times 10^{-7}\text{m}^2/\text{s}$ giving $\bar{\kappa}_T = 0.66 \times 10^{11}$.

To avoid those large numbers we scale T by 10^{-3} . Then $\bar{T}_S = 5.8$, $(\frac{Q}{2})^{\frac{1}{4}} \bar{T}_S = 24.9$, $\tilde{\kappa}_T = 10^{-9} \bar{\kappa}_T = 66$.

Table 1: *The physical constants.*

c	\hbar	k	B_0
2.998×10^8	6.6261×10^{-34}	1.381×10^{-23}	1.806657×10^{-19}




Figure 1: Convergence of T^n solution of triangulation is also shown, adapted from T computed at iteration 5. (18)




Figure 2: Color map of $T(x, z)$ at iteration 10. The triangulation is also shown, adapted from T computed at iteration 5.

3.1. A one dimensional test

If $\Omega = (0, 10)$, we need to solve with Algorithm (14) the integro-differential equation in z :

$$\begin{aligned}
 -66T'' + T^4 &= 12.5E_3(0.1|10 - z|) + 0.05 \int_0^{10} E_1(0.1|s - z|)T^4(s)ds, \\
 T(0) &= (12.5E_3(0))^{1/4}, \quad T'(10) = 0.
 \end{aligned} \tag{18}$$

To solve $-66T'' + T^4 = f$, 3 iterations of a fixed point loop are used: $-830T''^{m+1} + T^{m3}T^{m+1} = f$.

The results are shown on Figure 1. The convergence is monotone as expected, even though Theorem 1 hasn't been proved when a Dirichlet condition is applied to T on part of $\partial\Omega$. Notice that in absence of sunlight the temperature would be $T(0)$ everywhere.

3.2. A two dimensional test for a lake

Now Ω is half of the vertical cross section of a symmetric lake. The lower right quarter side of the unit circle is stretched by $x, z \mapsto 30x, 10z$. The bottom boundary has an equation named $z = z_m(x)$. The same problem is solve in 2D:

$$\begin{aligned}
 -66\Delta T + T^4 &= 12.5E_3(0.1|z_m(x) - z|) + 0.05 \int_{z_m(x)}^{10} E_1(0.1|s - z|)T^4(s)ds, \\
 T(x, z_m(x)) &= (12.5E_3(0.1z_m(x)))^{1/4}, \quad \partial_z T(10) = 0.
 \end{aligned} \tag{19}$$

The same 3×10 double iteration loop is used ; the results are shown on Figure 2.

3.3. A 3D case with convection in Lake Lemman

Lake Lemman is discretized into 33810 tetrahedra. The surface has 1287 triangles. The Finite Element method of degree 1 is used. This is too coarse for a Navier-Stokes simulation but appropriate for a potential flow. Pressure is imposed on the left and right tips to simulate the debit of the Rhône. The pressure p solves $-\Delta p = 0$ with $\partial_n p = 0$ on the remaining boundaries; the velocity is $\mathbf{u} = \nabla p$. The top plot in Figure 3 shows p and \mathbf{u} . The full temperature equation of Problem (P^4) is solved with the same physical constant as above. The temperature is set at T_e initially and on the bottom and side boundaries of the lake. The time step is $t = 0.1$; the method is fully implicit for the temperature. At each time step 3 iterations are needed to handle the T^4 term. Figure 3 shows the temperature after 15 time steps; it appears to have reached a steady state. The top right view of Figure 3 shows a region in red where the water at the surface is the hottest.




Figure 3: Top left: velocity vectors and pressure isolines at the surface of the lake. Top right: iso lines of the surface temperature. Bottom left: perspective view of a 3D color map of the temperature on the side of the lake past a middle vertical plane. Bottom right: perspective view showing some temperature level surfaces inside the lake.

This computation is merely a feasibility study to prove that the implementation of the RT module in a standard CFD code is easy and fast. Computing time on an intel core i9 takes less than a minute.

3.4. Comments on the programming tools

In fifty years the research problems have become increasingly complex and without the joint development of computers and programming tools it would not be possible for a single individual to contribute or even test his ideas. The second author is part of the team which developed the PDE solver FreeFem++ [Hec12]

(see <https://fr.wikipedia.org/wiki/FreeFem%2B%2B>).

The algorithms discussed here have been implemented with this tool in a very short time. The discretization of lake Lemman is part of the examples in [Hec12], written by F. Hecht.

4. The general case, κ_ν, a_ν non constant

Photons interact with the atomic structure of the medium which implies that κ_ν depends strongly on ν but also on the temperature and pressure. For the earth atmosphere the pressure and the temperature are approximately decaying exponentially with altitude.

Assume that variations with altitude are known: $\rho\bar{\kappa}_\nu = \varphi(z)\kappa_\nu$ with $z = z_m = 0$ on the ground. Let $\tau = \int_0^z \varphi(s)ds$; for instance $\tau = 1 - e^{-z}$ when $\varphi(z) = e^{-z}$. Now (8),(10) hold with $0 < \tau < Z := 1 - e^{-z_M}$ instead of $z_m < z < z_M$).

Consider two types of scattering kernels: a Rayleigh scattering kernel $p^r(\mu, \mu') = \frac{3}{8}[3 - \mu^2 + 3(\mu^2 - 1)\mu'^2]$ and an isotropic scattering kernel $p = 1$. Let a_ν^r and $a_\nu^i := a_\nu - a_\nu^r$ be the scattering coefficients for both. The problem is

$$\begin{aligned} \mu\partial_\tau I_\nu + \kappa_\nu I_\nu &= \kappa_\nu(1 - a_\nu)B_\nu(T) + \frac{1}{2}\kappa_\nu \int_{-1}^1 (a_\nu^r p^r + a_\nu^i) I_\nu d\mu' \\ I(0, \mu)|_{\mu>0} &= \alpha I(0, -\mu) + Q_\nu^+(\mu), \quad I(Z, \mu)|_{\mu<0} = Q_\nu^-(\mu) \end{aligned} \quad (20)$$

The boundary condition at $\tau = 0$ is a simplified Lambert condition which says that a portion α of the incoming light is reflected back (Earth albedo) and adds to the prescribed upgoing light Q_ν^+ . Sun light is prescribed at high altitude, Z , to be $Q_\nu^-(\mu)$.

Let

$$J_\nu(\tau) = \frac{1}{2} \int_{-1}^1 I_\nu(\tau, \mu) d\mu, \quad K_\nu(\tau) = \frac{1}{2} \int_{-1}^1 \mu^2 I_\nu(\tau, \mu) d\mu.$$

An integral formulation can be derived from (20) as in [Cha50], section 11.2:

$$\begin{aligned} (\mu\partial_\tau + \kappa_\nu)I_\nu &= H_\nu(\tau, \mu) \\ &:= \kappa_\nu \left((1 - a_\nu)B_\nu(T_\tau) + [a_\nu^i + \frac{3}{8}a_\nu^r(3 - \mu^2)]J_\nu(\tau) + \frac{9}{8}a_\nu^r(\mu^2 - 1)K_\nu(\tau) \right) \end{aligned} \quad (21)$$

$$\begin{aligned} \Rightarrow I(\tau, \mu) &= \mathbf{1}_{\mu>0} \left[R_\nu^+(\mu) e^{-\kappa_\nu \frac{\tau}{\mu}} + \int_0^\tau \frac{e^{\kappa_\nu \frac{t-\tau}{\mu}}}{\mu} \kappa_\nu H_\nu(t, \mu) dt \right] \\ &+ \mathbf{1}_{\mu<0} \left[Q_\nu^-(\mu) e^{\kappa_\nu \frac{Z-\tau}{\mu}} - \int_\tau^Z \frac{e^{\kappa_\nu \frac{t-\tau}{\mu}}}{\mu} \kappa_\nu H_\nu(t, \mu) dt \right], \end{aligned} \quad (22)$$

where $R^+(\mu) = Q_\nu^+(\mu) + \alpha I(0, -\mu)$, i.e.

$$R_\nu^+(\mu)|_{\mu>0} = Q_\nu^+(\mu) + \alpha \left[Q_\nu^-(-\mu) e^{-\kappa_\nu \frac{Z}{\mu}} + \int_0^Z \frac{e^{-\kappa_\nu \frac{t}{\mu}}}{\mu} \kappa_\nu H_\nu(t, -\mu) dt \right]. \quad (23)$$

From (22), since $H_\nu = H_\nu^0 + \mu^2 H_\nu^2$, with H_ν^0, H_ν^2 independent of μ , linear functions of J_ν and K_ν :

$$\begin{aligned} H_\nu^0(\tau) &= \kappa_\nu(1 - a_\nu)B_\nu(T) + \kappa_\nu \left((a_\nu^i + \frac{9a_\nu^r}{8})J_\nu - \frac{9a_\nu^r}{8}K_\nu \right), \\ H_\nu^2(\tau) &= -\kappa_\nu \frac{3a_\nu^r}{8} [J_\nu - 3K_\nu], \\ J_\nu(\tau) &= \frac{1}{2} \int_0^1 \left(e^{-\kappa_\nu \frac{\tau}{\mu}} Q_\nu^+(\mu) + \left[e^{-\kappa_\nu \frac{(Z-\tau)}{\mu}} + \alpha e^{-\kappa_\nu \frac{(Z+\tau)}{\mu}} \right] Q_\nu^-(-\mu) \right) d\mu \\ &+ \frac{1}{2} \int_0^Z \left([E_1(\kappa_\nu|\tau - t|) + \alpha E_1(\kappa_\nu(\tau + t))] H_\nu^0(\tau) + [E_3(\kappa_\nu|\tau - t|) + \alpha E_3(\kappa_\nu(\tau + t))] H_\nu^2(\tau) \right) dt \end{aligned} \quad (24)$$

(25)

$$\begin{aligned}
K_\nu(\tau) &= \frac{1}{2} \int_0^1 \mu^2 \left(e^{-\kappa_\nu \frac{\tau}{\mu}} Q_\nu^+(\mu) + \left[e^{-\kappa_\nu \frac{(Z-\tau)}{\mu}} + \alpha e^{-\kappa_\nu \frac{(Z+\tau)}{\mu}} \right] Q_\nu^-(-\mu) \right) d\mu \\
&+ \frac{1}{2} \int_0^Z ([E_3(\kappa_\nu|\tau-t|) + \alpha E_3(\kappa_\nu(\tau+t))] H_\nu^0(\tau) + [E_5(\kappa_\nu|\tau-t|) + \alpha E_5(\kappa_\nu(\tau+t))] H_\nu^2(\tau)) dt,
\end{aligned} \tag{26}$$

The system is coupled to

$$\partial_t T + \mathbf{u} \cdot \nabla T - \kappa_T \Delta_{x,y,z} T + 4\pi \int_0^\infty \kappa_\nu (1 - a_\nu) B_\nu(T_\tau) d\nu = 4\pi \int_0^\infty \kappa_\nu (1 - a_\nu) J_\nu(\tau) d\nu, \tag{27}$$

4.1. Iterative method for the general case

In the spirit of (14), consider

4.2. Algorithm 2

1. Starting from $T^0 = 0$, $J_\nu^0 = 0$, $K_\nu^0 = 0$.
2. Compute $J_\nu^{n+1}(\tau)$, $K_\nu^{n+1}(\tau)$ by (25)(26) with T^n , J^n, K^n . in place of T, J, K .
3. Compute T^{n+1} by solving (27) with $J_\nu^{n+1}(\tau)$ in the r.h.s.

Note that for isotropic scattering K_ν is not needed. Then the following convergence results hold when thermal diffusion is neglected.

Theorem 2. *Assume $\alpha = 0$, $\mathbf{u} = 0$, $\kappa_T = 0$, $\partial_t T = 0$. Assume κ_ν strictly positive and uniformly bounded, and $0 \leq a_\nu < 1$ for all $\nu > 0$. Let $Q_\nu^\pm \geq 0$ satisfy, for some T_M and some Q*

$$0 \leq Q_\nu^\pm(\mu) \leq Q B_\nu(T_M) \quad \forall \mu, \nu \in (-1, 1) \times \mathbb{R}^+. \tag{28}$$

Then Algorithm 4.2 defines a sequence of radiative intensities I_ν^n and temperatures T^n converging pointwise to I_ν and T respectively, which is a solution of (20),(27) and the convergence is uniformly increasing.

Remarks 2.

1. Starting with $T^0 = 0$ is a sure way to initialise the recurrence and have $T^1 > T^0$.
2. Most likely, monotone convergence holds also in the general case $\alpha > 0$, \mathbf{u} , κ_T and $\partial_t T$ non-zero because, just like $T \mapsto T^4$, the function $T \mapsto \int_0^\infty \kappa_\nu (1 - a_\nu) B_\nu(T) d\nu$ is monotone increasing (its derivative is strictly positive).
3. In the special case $a_\nu^r = 0$, and $Q_\nu^\pm(\mu) = |\mu| Q_\nu^\pm$ the problem is

$$\begin{aligned}
(\mu \partial_\tau + \kappa_\nu) I_\nu(\tau, \mu) &= \kappa_\nu a_\nu J_\nu(\tau) + \kappa_\nu (1 - a_\nu) B_\nu(T_\tau), & J_\nu(\tau) &= \frac{1}{2} \int_{-1}^1 I_\nu(\tau, \mu) d\mu, \\
I_\nu(0, \mu) &= Q_\nu^+ \mu, & I_\nu(Z, -\mu) &= Q_\nu^- \mu, & 0 < \mu < 1, \\
\int_0^\infty \kappa_\nu (1 - a_\nu) B_\nu(T_\tau) d\nu &= \int_0^\infty \kappa_\nu (1 - a_\nu) J_\nu(\tau) d\nu.
\end{aligned} \tag{29}$$

The iterative process is then to start with $T^0 = 0$, and compute T^{n+1} from T^n by

$$J_\nu^{n+1}(\tau) = \frac{1}{2}Q_\nu^+ E_3(\kappa_\nu \tau) + \frac{1}{2}Q_\nu^- E_3(\kappa_\nu(Z - \tau)) + \kappa_\nu \int_0^Z E_1(\kappa_\nu|\tau - t|) (a_\nu J_\nu^n(t) + (1 - a_\nu)B_\nu(T_t^n)) dt, \quad (30)$$

$$\int_0^\infty \kappa_\nu(1 - a_\nu)B_\nu(T_\tau^{n+1})d\nu = \int_0^\infty \kappa_\nu(1 - a_\nu)J_\nu^{n+1}(\tau)d\nu. \quad (31)$$

4. Note that $T \mapsto \int_0^\infty \kappa_\nu(1 - a_\nu)B_\nu(T)d\nu$ is continuous, strictly increasing, hence invertible. Thus (31) defines T_τ^{n+1} uniquely.
5. One may recover the light intensity by

$$I_\nu^{n+1}(\tau, \mu) = e^{-\kappa_\nu \frac{\tau}{\mu}} Q_\nu^+(\mu) \mathbf{1}_{\mu > 0} + e^{-\kappa_\nu \frac{(Z-\tau)}{|\mu|}} Q_\nu^-(\mu) \mathbf{1}_{\mu < 0} + \mathbf{1}_{\mu > 0} \int_0^\tau e^{-\kappa_\nu \frac{(\tau-t)}{\mu}} \frac{\kappa_\nu}{\mu} (a_\nu J_\nu^n(t) + (1 - a_\nu)B_\nu(T_t^n)) dt + \mathbf{1}_{\mu < 0} \int_\tau^Z e^{-\kappa_\nu \frac{(t-\tau)}{\mu}} \frac{\kappa_\nu}{\mu} (a_\nu J_\nu^n(t) + (1 - a_\nu)B_\nu(T_t^n)) dt. \quad (32)$$

but numerically these are singular integrals while (30),(31) are not. Indeed $e^{-\frac{x}{\mu}}/\mu$ tends to infinity when x and μ tend to 0.

6. Theorem 2 extends a result given in [Pir21] which had unnecessary restrictions on κ_ν .

Proof

The complete proof will appear in [BP21]. Here, for simplicity, we consider the case $a_\nu = 0$. Let $S(\tau) := \int_0^\infty \frac{\kappa_\nu}{2} \int_0^1 \left(e^{-\kappa_\nu \frac{\tau}{\mu}} Q_\nu^+(\mu) + e^{-\kappa_\nu \frac{Z-\tau}{\mu}} Q_\nu^-(-\mu) \right) d\mu d\nu$. By (30)

$$\begin{aligned} \int_0^\infty \kappa_\nu B_\nu(T_\tau^{n+1})d\nu &= \int_0^\infty \kappa_\nu J_\nu^{n+1}(\tau)d\nu = S(\tau) + \frac{1}{2} \int_0^\infty \int_0^Z \kappa_\nu^2 E_1(\kappa_\nu|\tau - t|) B_\nu(T_t^n) dt d\nu \\ &\leq S(\tau) + \frac{1}{2} \max_\kappa \int_0^Z \kappa E_1(\kappa|\tau - t|) dt \sup_{t \in (0, Z)} \int_0^\infty \kappa_\nu B_\nu(T_t^n) d\nu \\ &\leq C_2 + C_1(\kappa_M) \sup_{t \in (0, Z)} \int_0^\infty \kappa_\nu B_\nu(T_t^n) d\nu, \end{aligned}$$

with $C_2 = \sup_{t \in (0, Z)} S(t)$ and $\kappa_M = \sup_\nu \kappa_\nu$, because $\kappa \mapsto C_1(\kappa)$ is monotone increasing. As $C_1(\kappa_M) < 1$ it implies that $B_\nu^n(\tau) := B_\nu(T_\tau^n)$ is bounded for all τ .

Now assume that $T_\tau^n > T_\tau^{n-1}$ for all $\tau > 0$. Then $T \mapsto B_\nu(T)$ being increasing, $B_\nu(T_\tau^n) > B_\nu(T_\tau^{n-1})$, $\forall \tau, \nu$, and so for all τ :

$$\begin{aligned} \int_0^\infty \kappa_\nu (B_\nu(T_\tau^{n+1}) - B_\nu(T_\tau^n)) d\nu &= \int_0^\infty \kappa_\nu (J_\nu^{n+1}(\tau) - J_\nu^n(\tau)) d\nu \\ &= \int_0^\infty \frac{\kappa_\nu^2}{2} \int_0^Z E_1(\kappa_\nu|\tau - t|) (B_\nu(T_t^n) - B_\nu(T_t^{n-1})) dt d\nu > 0. \end{aligned} \quad (33)$$

As $T \mapsto B_\nu(T)$ is continuous, it implies that $T_\tau^{n+1} > T_\tau^n$, $\forall \tau$. Hence for some $T^*(\tau)$, possibly $+\infty$, $T^n \rightarrow T^*$. By continuity $B_\nu(T_t^n) \rightarrow B_\nu(T_t^*)$, but it has been show above that $B_\nu(T_t^n) = B_\nu^n \rightarrow B_\nu^*$, so $B_\nu(T_t^*)$ is finite and so is T_t^* . Recall that a bounded increasing sequence converges, so $B_\nu(T_t^n) \rightarrow B_\nu(T_t^*)$ for all t and ν and the convergence of $E_1(\kappa_\nu|\tau - t|)B_\nu(T_t^n) \rightarrow E_1(\kappa_\nu|\tau - t|)B_\nu(T_t^*)$ being monotone, the integral converges to the integral of the limit (Beppo Levi's lemma). This shows that T_τ^* is the solution of the problem. \square

5. Uniqueness, Maximum Principle

This section follows computations in [Gol87] (in the case $Z = +\infty$ and with $a_\nu = 0$) and in [Mer87].

Theorem 3. *Assume $0 < \kappa_\nu \leq \kappa_M$, $0 \leq a_\nu < 1$ for all $\nu > 0$. Let $Q^\pm, R^\pm \in L^1((0, 1) \times \mathbb{R}^+)$ satisfy*

$$0 \leq Q_\nu^\pm(\mu) \leq R_\nu^\pm(\mu) \quad \text{for a.e. } (\mu, \nu) \in (0, 1) \times (0, \infty).$$

Then, the solutions (I_ν, T) and (I'_ν, T') of (37) with $Q_\nu^\pm(\mu)$ and $R_\nu^\pm(\mu)$ respectively, satisfy

$$I_\nu(\tau, \mu) \leq I'_\nu(\tau, \mu) \quad \text{and} \quad T_\tau \leq T'_\tau \quad \text{for a.e. } (\tau, \mu) \in (-1, 1) \times (0, \infty).$$

In particular, $Q_\nu^\pm(\mu) = R_\nu^\pm(\mu)$ for a.e. $(\mu, \nu) \in (0, 1) \times (0, \infty)$ implies

$$I_\nu(\tau, \mu) = I'_\nu(\tau, \mu) \quad \text{and} \quad T_\tau = T'_\tau \quad \text{for a.e. } (\tau, \mu) \in (-1, 1) \times (0, \infty).$$

One has also the following form of a Maximum Principle.

Corollary 1. *Let the hypotheses of Theorem 2 hold. Let $Q_\nu^\pm(\mu) \leq B_\nu(T_M)$ (resp. $Q_\nu^\pm(\mu) \geq B_\nu(T_m)$) for a.e. $(\mu, \nu) \in (0, 1) \times (\mathbb{R}^+)$. Then a.e. $(\tau, \mu) \in (-1, 1) \times (0, \infty)$,*

$$I_\nu(\tau, \mu) \leq B_\nu(T_M) \quad \text{and} \quad T_\tau \leq T_M \quad \text{resp.} \quad I_\nu(\tau, \mu) \geq B_\nu(T_m) \quad \text{and} \quad T_\tau \geq T_m$$

The proof relies partially on a difficult argument due to [Mer87]. It will be published in [GP21].

6. An application to the temperature in the Earth atmosphere

A numerical test is reported on Figures 4 and 5. It is an attempt at the simulation of the effect of an increase of CO_2 in the atmosphere. Our purpose is only to assess that the numerical method can detect such a small change of κ_ν .

Equation (31) is solved by a few steps of dichotomy followed by a few New steps. When κ_ν is larger than 4 some instabilities occur, probably in the exponential integrals. This point will be investigated in the future.

The physical and numerical parameters are

- Atmosphere thickness: 12km
- Scaled sunlight power hitting the top of the atmosphere: 3.042×10^{-5}

- Percentage of sunlight reaching the ground unaffected: 0.99
- Percentage reemitted (Earth albedo): 10%.
- Percentage of sunlight being a source at high altitude (Q^-): 0.1%
- Cloud (isotropic) scattering: 20%. Cloud position : between 6 and 9km
- Rayleigh scattering: 20% above 9km
- average absorption coefficient $\kappa_0 = 1.225$
- density drop versus altitude : $\rho_0 \exp(-z)$
- Discretization: 60 altitude stations, 300 frequencies (unevenly distributed)
- Number of iterations 22. Computing time 30" per cases.

The results are very sensitive to the value of Q^- and th the earth albedo. The values for κ_ν are taken from Russian measurements posted on wikipedia
https://commons.wikimedia.org/wiki/File:Atmosfaerisk_spredning-ru.svg.




Figure 4: Transmittance t_e versus wave-length digitized from commons.wikimedia.org/wiki/File:Atmosfaerisk_spredning-ru.svg The window around 3 is blocked by CO₂. The absorption is related to the transmittance t_e by $\kappa_\nu = -\log t_e$.

Figure transmittance https://commons.wikimedia.org/wiki/File:Atmosfaerisk_spredning-ru.svg

7. Conclusion

Results obtained here are in continuation of [Gol87],[Mer87],[BGPS88], recently reviewed for possible applications to climatology in [BP21] and [Pir21]. Existence and uniqueness for the radiative transfer equations had remained open in the context of nuclear engineering. For incompressible fluids it is not unrealistic to assume that the dependence of the absorption coefficient κ_ν upon the temperature can be replaced by an explicit dependence on altitude. This is the key simplification by which existence, uniqueness and




Figure 5: Scaled temperatures (left) : 3 curves $z \rightarrow T(z)$ are plotted. One computed with $\kappa_0 = 1.225$ which corresponds to a grey atmosphere. One with κ_ν shown on the right in pink color which corresponds to Figure 4. The third one is with κ_ν shown in green on the right where the transparent window around frequency 1 has been blocked. On the right the mean light intensity at altitude Z are shown (mostly outgoing waves). Filling the transparent window results in an elevation of temperature.

monotone fast and accurate numerical schemes could be found. Hence, adding RT to a Navier-Stokes solver is easy and fast when radiations come from one direction only.

As a final remark note that it seems doable to extend the method to the general case where κ_ν depends on τ and T . Indeed if the dependency $\tau \mapsto \kappa_\nu(\tau)$ is guessed only approximately, then knowing $\kappa_\nu^M > \kappa_\nu(\tau)$ independent of τ is enough to apply the method with κ_M on the left of the equation for I_ν with a correction on the right equal to $(\kappa_\nu^M - \kappa_\nu(\tau))I_\nu$; this correction seems compatible with the monotone convergence of the temperature. Then the method could also be extended to the case κ function of T by an additional algorithmic m-loop using $\kappa(T^m)$ instead of $\kappa(T)$ and then updating T^m to the T just computed.

In this article the numerical computations are only given for showing the potential of the method. Real life applications, coupling RT to the full Navier-Stokes equations requires supercomputing power and will be done later.

References

- [ACP⁺09] J. Amaya, O. Cabrit, D. Poitou, B. Cuenot, and M. ElHafi. Unsteady coupling of navier-stokes and radiative heat transfer solvers applied to an anisothermal multicomponent turbulent channel flow. In P.J. Coelho et al, editor, *Computational Thermal Radiation in Participating Media III 15–17 April 2009, Lisbon*. FCT report, 2009.
- [AG01] P. Azérad and F. Guillén. Mathematical justification of the hydrostatic approximation in the primitive equations of geophysical fluid dynamics. *SIAM J. Math. Anal.*, 33:847–859, 2001.
- [Att09] A. Attaoui. Existence of solutions for a nonlinear boussinesq-stefan system. *Adv. Differential Equations*, 14(9):985–1018, 2009.

- [BGPS88] C. Bardos, F. Golse, B. Perthame, and R. Sentis. The nonaccretive radiative transfer equations: existence of solutions and rosseland approximation. *J. Functional Analysis*, 77:434–460, 1988.
- [Boh06] Craig F. Bohren. *Fundamentals of Atmospheric Radiation*. Cambridge U Press, 2006.
- [BP21] C. Bardos and O Pironneau. Radiative Transfer for the Greenhouse Effect. submitted to SeMA J. Springer, 2021.
- [BSS84] C. Bardos, R. Santos, and R. Sentis. Diffusion approximation and computation of the critical size. *Transaction of the American Math Society*, 284(2):617–649, 1984.
- [CCT20] Q. Lin C. Cao and E. S. Titi. On the well-posedness of reduced 3d primitive geostrophic adjustment model with weak dissipation. *J. Math. Fluid Mech*, 22(32), 2020.
- [CDK⁺03] Eric Cancès, Mireille Defranceschi, Werner Kutzelnigg, Claude Le Bris, and Yvon Maday. Computational quantum chemistry: A primer. In *Special Volume, Computational Chemistry*, volume 10 of *Handbook of Numerical Analysis*, pages 3–270. Elsevier, 2003.
- [Cha50] S. Chandrasekhar. *Radiative Transfer*. Clarendon Press, Oxford, 1950.
- [CLGP13] M.J. Castro Díaz, J.A. López-García, and Carlos Parés. High order exactly well-balanced numerical methods for shallow water systems. *Journal of Computational Physics*, 246:242–264, 2013.
- [Cor15] C. Cornet. *Transfert radiatif dans une atmosphère tridimensionnelle*. PhD thesis, Université de Lille, Labo d’optique atmosphérique, 2015.
- [DDT⁺15] T. Dubos, S. Dubey, M. Tort, R. Mittal, Y. Meurdesoif, and F. Hourdin. Dynamico-1.0, an icosahedral hydrostatic dynamical core designed for consistency and versatility. *Geosci. Model Dev*, 8:3131–3150, 2015.
- [DECG20] J. Dufresne, V. Eymet, C. Crevoisier, and J. Grandpeix. Greenhouse effect: The relative contributions of emission height and total absorption. *Journal of Climate, American Meteorological Society*, 33(9):3827–3844, 2020.
- [DL00] R. Dautray and J. Lions. *Mathematical Analysis and Numerical Methods for Science and Technology*, volume 3. Springer Verlag, NY, 2000.
- [dmdL14] L’équipe de modélisation du LMD. *LMDZ5: a documentation*. Laboratoire de Météo Dynamique, <https://lmdz.lmd.jussieu.fr/>, 2014.
- [EDK19] C. Eldred, T. Dubos, and E. Kritsikis. A quasi-hamiltonian discretization of the thermal shallow water equations. *Journal of Computational Physics*, 379:1–31, 2019.

- [Fou88] Y. Fouquart. Radiative transfer in climate models. In M.E. Schlesinger ed., editor, *NATO ASI Series.*, 1988.
- [Fow11] A. Fowler. *Mathematical Geoscience*. Springer Verlag, New York, 2011.
- [GHM20] Mohamed Ghattassi, Xiaokai Huo, and Nader Masmoudi. On the diffusive limits of radiative heat transfer system i: well prepared initial and boundary conditions, 2020.
- [Gol87] F. Golse. The milne problem for the radiative transfer equations (with frequency dependence). *Trans. Amer. Math. Soc.*, 303:125–143, 1987.
- [GP21] F. Golse and O. Pironneau. Existence and uniqueness for the radiative transfer equations for the atmosphere. *volume in honor of Prof. Pedro Diaz*, to appear, 2021.
- [GY61] R.M. Goody and Y.L. Yung. *Atmospheric Radiation*. Oxford U Press, 1961.
- [Hec12] F Hecht. New developments in freefem++. *J. Numer. Math.*, 20:251–265, 2012.
- [KMRW00] G. Kanschat, E. Meinköhn, R. Rannacher, and R. Wehrse, editors. *Numerical Methods in Multidimensional Radiative Transfer*. Springer Verlag, 2000.
- [Lie01] Loss M. Lieb, E. *Analysis*. Amer. Math. Soc., Providence, 2 edition, 2001.
- [Lio96] Pierre-Louis Lions. *Mathematical Topics in Fluid Dynamics, Vol. 1 Incompressible Models.*, Oxford U Press, 1996.
- [LTW94] J.-L. Lions, R. Temam, and S. Wang. Geostrophic asymptotics of the primitive equations of the atmosphere. *Topological Methods in Nonlinear Analysis*, 4(2):253 – 287, 1994.
- [Mer87] B. Mercier. Application of accretive operators theory to the radiative transfer equations. *SIAM J. Math. Anal.*, 18(2):393–408, 1987.
- [Mor91] J-J. Morette. Radiation and cloud radiative properties in the european centre for medium range weather forecasts forecasting system. *J. Geophys. Research: Atmospheres*, 91(D5):9121–9132, 1991.
- [Pir21] O. Pironneau. A fast and accurate numerical method for radiative transfer in the atmosphere. October 2021.
- [Pom73] G. Pomraning. *The equations of Radiation Hydrodynamics*. Pergamon Press, NY, 1973.
- [XSW19] Yu Xie, Manajit Sengupta, and Chenxi Wang. A fast all-sky radiation model for solar applications with narrowband irradiances on tilted surfaces. *Solar Energy, Part II*, 2019.

[ZPS⁺21] Shaoning Zeng, Sijie Pian, Minyu Su, Zhuning Wang, Maoqi Wu, Xinhang Liu, Mingyue Chen, Yuanzhuo Xiang, Jiawei Wu, Manni Zhang, Qingqing Cen, Yuwei Tang, Xianheng Zhou, Zhiheng Huang, Rui Wang, Alitenai Tunuhe, Xiyu Sun, Zhigang Xia, Mingwei Tian, Min Chen, Xiao Ma, Lvyun Yang, Jun Zhou, Huamin Zhou, Qing Yang, Xin Li, Yaoguang Ma, and Guangming Tao. Hierarchical-morphology metafabric for scalable passive daytime radiative cooling. *Science*, 2021.

[ZT03] Wilford Zdunkowski and Thomas Trautmann. *Radiation in the Atmosphere*. Cambridge U Press, 2003.

8. APPENDIX not part of this Comptes-Rendus de Mécanique: Proofs of the results quoted above

Consider the problem

$$\begin{aligned} (\mu\partial_\tau + \kappa_\nu)I_\nu(\tau, \mu) &= \kappa_\nu a_\nu J_\nu(\tau) + \kappa_\nu(1 - a_\nu)B_\nu(T(\tau)), \\ I_\nu(0, \mu) &= Q_\nu^+(\mu), \quad I_\nu(Z, -\mu) = Q_\nu^-(\mu), \quad 0 < \mu < 1, \\ \partial_\tau \int_0^\infty \int_{-1}^1 \mu I_\nu(\tau, \mu) d\mu d\nu &= 0, \end{aligned} \tag{34}$$

with the notation

$$J_\nu(\tau) := \frac{1}{2} \int_{-1}^1 I_\nu(\tau, \mu) d\mu.$$

The last equality in (34) implies that

$$\int_0^\infty \kappa_\nu(1 - a_\nu)J_\nu(\tau) d\nu = \int_0^\infty \kappa_\nu(1 - a_\nu)B_\nu(T(\tau)) d\nu \tag{35}$$

and, assuming that $0 < \kappa_\nu \leq \kappa_M$ while $0 \leq a_\nu < 1$ for all $\nu > 0$, the r.h.s. of (35) defines T as a functional of J , henceforth denoted $T[J]$. Thus (34) can be recast as

$$\begin{aligned} (\mu\partial_\tau + \kappa_\nu)I_\nu(\tau, \mu) &= \kappa_\nu a_\nu J_\nu(\tau) + \kappa_\nu(1 - a_\nu)B_\nu(T[J](\tau)), \\ I_\nu(0, \mu) &= Q_\nu^+(\mu), \quad I_\nu(Z, -\mu) = Q_\nu^-(\mu), \quad 0 < \mu < 1, \end{aligned} \tag{36}$$

In order to solve numerically (34), one uses the method of iteration on the sources. Starting from some appropriate (I_ν^0, T^0) , one constructs a sequence (I_ν^n, T^n) by the following prescription

$$\begin{aligned} (\mu\partial_\tau + \kappa_\nu)I_\nu^{n+1}(\tau, \mu) &= \kappa_\nu a_\nu J_\nu^n(\tau) + \kappa_\nu(1 - a_\nu)B_\nu(T^n(\tau)), \quad T^n = T[J_\nu^n] \\ I_\nu^{n+1}(0, \mu) &= Q_\nu^+(\mu), \quad I_\nu^{n+1}(Z, -\mu) = Q_\nu^-(\mu), \quad 0 < \mu < 1, \end{aligned} \tag{37}$$

Applying the method of characteristics shows that

$$\begin{aligned} I_\nu^{n+1}(\tau, \mu) &= e^{-\frac{\kappa_\nu \tau}{\mu}} Q_\nu^+(\mu) \mathbf{1}_{\mu > 0} + e^{-\frac{\kappa_\nu(Z-\tau)}{|\mu|}} Q_\nu^-(|\mu|) \mathbf{1}_{\mu < 0} \\ &+ \mathbf{1}_{\mu > 0} \int_0^\tau e^{-\frac{\kappa_\nu(\tau-t)}{\mu}} \frac{\kappa_\nu}{\mu} (a_\nu J_\nu^n(t) + (1 - a_\nu)B_\nu(T^n(t))) dt \\ &+ \mathbf{1}_{\mu < 0} \int_\tau^Z e^{-\frac{\kappa_\nu(t-\tau)}{|\mu|}} \frac{\kappa_\nu}{|\mu|} (a_\nu J_\nu^n(t) + (1 - a_\nu)B_\nu(T^n(t))) dt. \end{aligned} \tag{38}$$

Since $B_\nu \geq 0$, this formula shows, by a straightforward induction argument, that

$$I_\nu^0 \geq 0, T^0 \geq 0, Q_\nu^\pm \geq 0 \implies I_\nu^n \geq 0.$$

Moreover

$$\begin{aligned} & I_\nu^{n+1}(\tau, \mu) - I_\nu^n(\tau, \mu) = \\ & + \mathbf{1}_{\mu > 0} \int_0^\tau e^{-\frac{\kappa_\nu(\tau-t)}{\mu}} \frac{\kappa_\nu}{\mu} a_\nu (J_\nu^n(t) - J_\nu^{n-1}(t)) dt \\ & + \mathbf{1}_{\mu > 0} \int_0^\tau e^{-\frac{\kappa_\nu(\tau-t)}{\mu}} \frac{\kappa_\nu}{\mu} (1 - a_\nu) (B_\nu(T^n(t)) - B_\nu(T^{n-1}(t))) dt \\ & + \mathbf{1}_{\mu < 0} \int_\tau^Z e^{-\frac{\kappa_\nu(t-\tau)}{|\mu|}} \frac{\kappa_\nu}{|\mu|} a_\nu (J_\nu^n(t) - J_\nu^{n-1}(t)) dt \\ & + \mathbf{1}_{\mu < 0} \int_\tau^Z e^{-\frac{\kappa_\nu(t-\tau)}{|\mu|}} \frac{\kappa_\nu}{|\mu|} (1 - a_\nu) (B_\nu(T^n(t)) - B_\nu(T^{n-1}(t))) dt. \end{aligned}$$

Since B_ν is nondecreasing for each $\nu > 0$, formula (35) shows that

$$J_\nu^n \geq J_\nu^{n-1} \implies T^n \geq T^{n-1}$$

we conclude from the equality above that

$$I_\nu^0 = 0, T^0 = 0, Q_\nu^\pm \geq 0 \implies \begin{cases} 0 \leq I_\nu^1 \leq I_\nu^2 \leq \dots \leq I_\nu^n \leq \dots \\ 0 \leq T^1 \leq T^2 \leq \dots \leq T^n \leq \dots \end{cases}$$

Since the term

$$(a_\nu J_\nu^n(t) + (1 - a_\nu) B_\nu(T^n(t)))$$

in both integrals on the r.h.s. is independent of μ , one has

$$\begin{aligned} J_\nu^{n+1}(\tau) &= \frac{1}{2} \int_0^1 \left(e^{-\frac{\kappa_\nu \tau}{\mu}} Q_\nu^+(\mu) + e^{-\frac{\kappa_\nu(Z-\tau)}{\mu}} Q_\nu^-(\mu) \right) d\mu \\ &+ \int_0^\tau \left(\int_0^1 e^{-\frac{\kappa_\nu(\tau-t)}{\mu}} \frac{d\mu}{2\mu} \right) \kappa_\nu (a_\nu J_\nu^n(t) + (1 - a_\nu) B_\nu(T^n(t))) dt \\ &+ \int_\tau^Z \left(\int_0^1 e^{-\frac{\kappa_\nu(t-\tau)}{\mu}} \frac{d\mu}{2\mu} \right) \kappa_\nu (a_\nu J_\nu^n(t) + (1 - a_\nu) B_\nu(T^n(t))) dt. \end{aligned}$$

One changes variables in the inner integral, so that

$$\int_0^1 e^{-\frac{x}{\mu}} \frac{d\mu}{\mu} = \int_1^\infty \frac{e^{-Xy}}{y} dy = \int_X^\infty \frac{e^{-z}}{z} dz =: E_1(X).$$

Thus

$$\begin{aligned} J_\nu^{n+1}(\tau) &= \frac{1}{2} \int_0^1 \left(e^{-\frac{\kappa_\nu \tau}{\mu}} Q_\nu^+(\mu) + e^{-\frac{\kappa_\nu(Z-\tau)}{\mu}} Q_\nu^-(\mu) \right) d\mu \\ &+ \frac{1}{2} \int_0^Z E_1(\kappa_\nu |\tau - t|) \kappa_\nu (a_\nu J_\nu^n(t) + (1 - a_\nu) B_\nu(T^n(t))) dt. \end{aligned}$$

Integrating over $[0, Z]$ in τ implies that

$$\begin{aligned}
\int_0^Z J_\nu^{n+1}(\tau) d\tau &= \frac{1}{2} \int_0^Z \int_0^1 \left(e^{-\frac{\kappa_\nu \tau}{\mu}} Q_\nu^+(\mu) + e^{-\frac{\kappa_\nu(Z-\tau)}{\mu}} Q_\nu^-(\mu) \right) d\mu d\tau \\
&+ \frac{1}{2} \int_0^Z \left(\int_0^Z E_1(\kappa_\nu |\tau - t|) \kappa_\nu d\tau \right) (a_\nu J_\nu^n(t) + (1 - a_\nu) B_\nu(T^n(t))) dt \\
&\leq \frac{1}{2} \int_0^Z \int_0^1 \left(e^{-\frac{\kappa_\nu \tau}{\mu}} Q_\nu^+(\mu) + e^{-\frac{\kappa_\nu(Z-\tau)}{\mu}} Q_\nu^-(\mu) \right) d\mu d\tau \\
&+ \frac{1}{2} \left(\sup_{0 \leq t \leq Z} \int_0^Z E_1(\kappa_\nu |\tau - t|) \kappa_\nu d\tau \right) \int_0^Z (a_\nu J_\nu^n(t) + (1 - a_\nu) B_\nu(T^n(t))) dt.
\end{aligned}$$

Let us estimate the quantity

$$\sup_{0 \leq t \leq Z} \int_0^Z E_1(\kappa_\nu |\tau - t|) \kappa_\nu d\tau = \sup_{0 \leq s \leq \kappa_\nu Z} \int_0^{\kappa_\nu Z} E_1(|\sigma - s|) d\sigma.$$

Observe that

$$\begin{aligned}
\int_0^{\kappa_\nu Z} E_1(|\sigma - s|) d\sigma &= \int_{\mathbf{R}} E_1(|\sigma - s|) 1_{[0, \kappa_\nu Z]}(\sigma) d\sigma \\
&= \int_{\mathbf{R}} E_1(|\theta|) 1_{[-s, \kappa_\nu Z - s]}(\theta) d\theta \\
&\leq \int_{\mathbf{R}} E_1(|\theta|) 1_{[-\kappa_\nu Z/2, \kappa_\nu Z/2]}(\theta) d\theta \\
&= 2 \int_0^{\kappa_\nu Z/2} E_1(\theta) d\theta \\
&\leq 2 \int_0^{Z \kappa_M/2} E_1(\theta) d\theta =: 2C_1.
\end{aligned}$$

The first inequality is the elementary rearrangement inequality (Theorem 3.4 in [Lie01]); the last one is based on the assumption $0 < \kappa_\nu \leq \kappa_M$. Thus

$$\begin{aligned}
\int_0^Z J_\nu^{n+1}(\tau) d\tau &\leq \frac{1}{2} \int_0^Z \int_0^1 \left(e^{-\frac{\kappa_\nu \tau}{\mu}} Q_\nu^+(\mu) + e^{-\frac{\kappa_\nu(Z-\tau)}{\mu}} Q_\nu^-(\mu) \right) d\mu d\tau \\
&+ C_1 \int_0^Z (a_\nu J_\nu^n(t) + (1 - a_\nu) B_\nu(T^n(t))) dt.
\end{aligned}$$

Multiply both sides of this inequality by κ_ν and integrate in ν : one finds that

$$\begin{aligned}
&\int_0^\infty \int_0^Z \kappa_\nu J_\nu^{n+1}(\tau) d\tau d\nu \\
&\leq \frac{1}{2} \int_0^\infty \int_0^Z \int_0^1 \kappa_\nu \left(e^{-\frac{\kappa_\nu \tau}{\mu}} Q_\nu^+(\mu) + e^{-\frac{\kappa_\nu(Z-\tau)}{\mu}} Q_\nu^-(\mu) \right) d\mu d\tau d\nu \\
&\quad + C_1 \int_0^\infty \int_0^Z \kappa_\nu (a_\nu J_\nu^n(t) + (1 - a_\nu) B_\nu(T^n(t))) dt d\nu.
\end{aligned}$$

At this point, we recall that $T^n = T[J_\nu^n]$, so that

$$\int_0^\infty \kappa_\nu (1 - a_\nu) B_\nu(T^n(t)) d\nu = \int_0^\infty \kappa_\nu (1 - a_\nu) J_\nu^n(t) d\nu,$$

and hence

$$\begin{aligned} & \int_0^\infty \int_0^Z \kappa_\nu J_\nu^{n+1}(\tau) d\tau d\nu \leq C_1 \int_0^\infty \int_0^Z \kappa_\nu J_\nu^n(t) dt d\nu \\ & + \frac{1}{2} \int_0^\infty \int_0^Z \int_0^1 \kappa_\nu \left(e^{-\frac{\kappa_\nu \tau}{\mu}} Q_\nu^+(\mu) + e^{-\frac{\kappa_\nu(Z-\tau)}{\mu}} Q_\nu^-(\mu) \right) d\mu d\tau d\nu. \end{aligned}$$

The expression of the source term can be slightly reduced, by integrating out the τ variable:

$$\int_0^Z \kappa_\nu e^{-\frac{\kappa_\nu \tau}{\mu}} d\tau = \int_0^Z \kappa_\nu e^{-\frac{\kappa_\nu(Z-\tau)}{\mu}} d\tau = \mu \left(1 - e^{-\frac{\kappa_\nu Z}{\mu}} \right),$$

so that

$$\begin{aligned} 0 & \leq \frac{1}{2} \int_0^\infty \int_0^Z \int_0^1 \kappa_\nu \left(e^{-\frac{\kappa_\nu \tau}{\mu}} Q_\nu^+(\mu) + e^{-\frac{\kappa_\nu(Z-\tau)}{\mu}} Q_\nu^-(\mu) \right) d\mu d\tau d\nu \\ & \leq \frac{1}{2} \int_0^\infty \int_0^1 (Q_\nu^+(\mu) + Q_\nu^-(\mu)) \mu d\mu =: \mathcal{Q}. \end{aligned}$$

Thus

$$\int_0^\infty \int_0^Z \kappa_\nu J_\nu^{n+1}(\tau) d\tau d\nu \leq C_1 \int_0^\infty \int_0^Z \kappa_\nu J_\nu^n(t) dt d\nu + \mathcal{Q}.$$

Initializing the sequence I_ν^n with $I_\nu^0 = 0$ and $T^0 = T[J_\nu^0] = 0$, one finds that

$$\begin{aligned} & \int_0^\infty \int_0^Z \kappa_\nu J_\nu^1(\tau) d\tau d\nu \leq \mathcal{Q}, \\ & \int_0^\infty \int_0^Z \kappa_\nu J_\nu^2(\tau) d\tau d\nu \leq C_1 \mathcal{Q} + \mathcal{Q} \\ & \int_0^\infty \int_0^Z \kappa_\nu J_\nu^3(\tau) d\tau d\nu \leq C_1^2 \mathcal{Q} + C_1 \mathcal{Q} + \mathcal{Q} \end{aligned}$$

and by induction

$$\int_0^\infty \int_0^Z \kappa_\nu J_\nu^{n+1}(\tau) d\tau d\nu \leq \mathcal{Q} \sum_{j=0}^n C_1^j.$$

Since

$$C_1 = \int_0^{Z\kappa_M/2} E_1(\theta) d\theta < \int_0^\infty E_1(\theta) d\theta = \int_0^\infty \left(\int_1^\infty \frac{e^{-\theta y}}{y} dy \right) dy = \int_1^\infty \frac{dy}{y^2} = 1,$$

the series above converges and one has the uniform bound

$$\int_0^\infty \int_0^Z \kappa_\nu J_\nu^{n+1}(\tau) d\tau d\nu \leq \frac{\mathcal{Q}}{1 - C_1}.$$

Since

$$0 \leq I_\nu^1 \leq I_\nu^2 \leq \dots \leq I_\nu^n \leq I_\nu^{n+1} \leq \dots$$

the bound above and the Monotone Convergence Theorem implies that the sequence $I_\nu^{n+1}(\tau, \mu)$ converges for a.e. $(\tau, \mu, \nu) \in (0, Z) \times (-1, 1) \times (0, +\infty)$ to a limit denoted $I_\nu(\tau, \mu)$ as $n \rightarrow \infty$. Since

$$0 \leq T^1 \leq T^2 \leq \dots \leq T^n \leq T^{n+1} \leq \dots$$

we conclude from (35) and the Monotone Convergence Theorem that $T^{n+1}(\tau)$ converges for a.e. $\tau \in (0, Z)$ to a limit denoted $T(\tau)$ as $n \rightarrow \infty$.

Then we can pass to the limit in (38) as $n \rightarrow \infty$ by monotone convergence, to find that

$$\begin{aligned} I_\nu(\tau, \mu) &= e^{-\frac{\kappa_\nu \tau}{\mu}} Q_\nu^+(\mu) \mathbf{1}_{\mu > 0} + e^{-\frac{\kappa_\nu (Z-\tau)}{|\mu|}} Q_\nu^- (|\mu|) \mathbf{1}_{\mu < 0} \\ &\quad + \mathbf{1}_{\mu > 0} \int_0^\tau e^{-\frac{\kappa_\nu (\tau-t)}{\mu}} \frac{\kappa_\nu}{\mu} (a_\nu J_\nu(t) + (1-a_\nu) B_\nu(T(t))) dt \\ &\quad + \mathbf{1}_{\mu < 0} \int_\tau^Z e^{-\frac{\kappa_\nu (t-\tau)}{|\mu|}} \frac{\kappa_\nu}{|\mu|} (a_\nu J_\nu(t) + (1-a_\nu) B_\nu(T(t))) dt \end{aligned}$$

for a.e. $(\tau, \mu, \nu) \in (0, Z) \times (-1, 1) \times (0, +\infty)$. One recognizes in this equality the integral formulation of (34) or (36).

Besides, since we have seen that

$$\begin{aligned} 0 &= I_\nu^0 \leq I_\nu^1 \leq I_\nu^2 \leq \dots \leq I_\nu^n \leq I_\nu^{n+1} \leq \dots \leq I_\nu \\ 0 &= T^0 \leq T^1 \leq T^2 \leq \dots \leq T^n \leq T^{n+1} \leq \dots \leq T \end{aligned}$$

so that

$$\begin{aligned} 0 &\leq \int_0^Z (J_\nu^{n+1} - J_\nu^n)(\tau) d\tau \\ &= \frac{1}{2} \int_0^Z \left(\int_0^Z E_1(\kappa_\nu |\tau - t|) \kappa_\nu d\tau \right) a_\nu (J_\nu^n - J_\nu^{n-1})(t) dt \\ &\quad + \frac{1}{2} \int_0^Z \left(\int_0^Z E_1(\kappa_\nu |\tau - t|) \kappa_\nu d\tau \right) (1-a_\nu) (B_\nu(T^n(t)) - B_\nu(T^{n-1}(t))) dt \\ &\leq C_1 \int_0^Z (a_\nu (J_\nu^n - J_\nu^{n-1})(t) + (1-a_\nu) (B_\nu(T^n(t)) - B_\nu(T^{n-1}(t)))) dt. \end{aligned}$$

Using again the equality

$$\int_0^\infty \kappa_\nu (1-a_\nu) B_\nu(T^n(t)) d\nu = \int_0^\infty \kappa_\nu (1-a_\nu) J_\nu^n(t) d\nu,$$

we conclude that

$$0 \leq \int_0^Z \int_0^\infty \kappa_\nu (J_\nu^{n+1} - J_\nu^n)(\tau) d\nu d\tau \leq C_1 \int_0^Z \int_0^\infty \kappa_\nu (J_\nu^n - J_\nu^{n-1})(t) dt.$$

Hence

$$0 \leq \int_0^Z \int_0^\infty \kappa_\nu (J_\nu^{n+1} - J_\nu^n)(\tau) d\nu d\tau \leq C_1^n \int_0^\infty \kappa_\nu J_\nu^1(\tau) d\nu d\tau \leq C_1^n \mathcal{Q},$$

so that

$$0 \leq \int_0^Z \int_0^\infty \kappa_\nu (J_\nu - J_\nu^n)(\tau) d\nu d\tau \leq C_1^n \int_0^\infty \kappa_\nu J_\nu^1(\tau) d\nu d\tau \leq \frac{C_1^n \mathcal{Q}}{1 - C_1}.$$

Summarizing, we have proved the following result.

Theorem 4. Assume that $0 < \kappa_\nu \leq \kappa_M$, while $0 \leq a_\nu < 1$ for all $\nu > 0$. Let $Q_\nu^\pm(\mu)$ satisfy

$$\int_0^\infty \int_0^1 \mu Q_\nu^\pm(\mu) d\mu d\nu \leq \mathcal{Q}.$$

Choose $I_\nu^0 = 0$ and $T^0 = 0$, and let I_ν^n and $T^n = T[J_\nu^n]$ be the solution of (37). Then

$$I_\nu^n(\tau, \mu) \rightarrow I_\nu(\tau, \mu) \quad \text{and} \quad T^n(\tau) \rightarrow T(\tau)$$

for $(\tau, \mu, \nu) \in (0, Z) \times (-1, 1) \times (0, +\infty)$ as $n \rightarrow \infty$, where (I_ν, T) is a solution of (34) or (36). This method converges exponentially fast, in the sense that

$$0 \leq \int_0^Z \int_0^\infty \kappa_\nu (J_\nu - J_\nu^n)(\tau) d\nu d\tau \leq \frac{C_1^n \mathcal{Q}}{1 - C_1},$$

and, if $0 \leq a_\nu \leq a_M < 1$ while $0 < \kappa_m \leq \kappa_\nu$, one has

$$0 \leq \int_0^Z \alpha (T(t)^4 - T^n(t)^4) dt \leq \frac{C_1^n \mathcal{Q}}{\kappa_m (1 - a_M) (1 - C_1)}.$$

The last bound comes from the defining equality for the temperature in terms of the radiative intensity

$$\begin{aligned} \kappa_m (1 - a_M) \alpha (T^4 - (T^n)^4) &= \kappa_m (1 - a_M) \int_0^\infty (B_\nu(T) - B_\nu(T^n)) d\nu \\ &\leq \int_0^\infty \kappa_\nu (1 - a_\nu) (B_\nu(T) - B_\nu(T^n)) d\nu \\ &= \int_0^\infty \kappa_\nu (1 - a_\nu) (J_\nu - J_\nu^n) d\nu \leq \int_0^\infty \kappa_\nu (J_\nu - J_\nu^n) d\nu. \end{aligned}$$

9. Uniqueness, Maximum Principle

This section follows computations in [Gol87] (in the case $Z = +\infty$ and with $a_\nu = 0$) and in [Mer87].

The rather subtle monotonicity structure of the radiative transfer equations is a striking result, discovered by Mercier in [Mer87]. In view of the complexity of the computations in [Mer87], it may be useful to keep in mind the following simple remarks, which should be viewed as a motivation.

Consider the steady radiative transfer equation (36) without scattering ($a_\nu = 0$) in the whole space with a source term $0 \leq S_\nu \in L^1(\mathbf{R} \times (-1, 1) \times (0, \infty))$:

$$\lambda I_\nu(\tau, \mu) + \mu \partial_\tau I_\nu(\tau, \mu) + \kappa_\nu I_\nu(\tau, \mu) = \kappa_\nu B_\nu(T[I]) + \lambda S_\nu(\tau, \mu), \quad \tau \in \mathbf{R}, \quad |\mu| < 1,$$

where $\lambda > 0$. By definition of $T[I]$, one easily checks that

$$\int_{-\infty}^\infty \int_{-1}^1 \int_0^\infty I_\nu(\tau, \mu) d\nu d\mu d\tau = \int_{-\infty}^\infty \int_{-1}^1 \int_0^\infty S_\nu(\tau, \mu) d\nu d\mu d\tau.$$

The radiative intensity is given in terms of the temperature $T[I]$ and the source S_ν by the explicit formula

$$I_\nu(\tau, \mu) = \mathbf{1}_{\mu > 0} \int_{-\infty}^{\tau} e^{-\frac{(\lambda + \kappa_\nu)(\tau - t)}{\mu}} \frac{\kappa_\nu B_\nu(T[I](t)) + \lambda S_\nu(t, \mu)}{\mu} dt \\ + \mathbf{1}_{\mu < 0} \int_{\tau}^{\infty} e^{-\frac{(\lambda + \kappa_\nu)(t - \tau)}{|\mu|}} \frac{\kappa_\nu B_\nu(T[I](t)) + \lambda S_\nu(t, \mu)}{|\mu|} dt.$$

Now, if one replaces the source of radiation S_ν in the right hand side of this equation with a larger source $S'_\nu \geq S_\nu$, it is natural to expect that the resulting radiation intensity I'_ν will be such that the associated temperature $T[I'] \geq T[I]$. Observe now that the function $T \mapsto B_\nu(T)$ is increasing on $(0, +\infty)$ for each $\nu > 0$; the explicit formula for I_ν in terms of S_ν and $T[I]$ shows that $I'_\nu(\tau, \mu) \geq I_\nu(\tau, \mu)$.

Of course, this argument is by no means rigorous, since it rests on the assumption that $S'_\nu \geq S_\nu \implies T[I'] \geq T[I]$, which, although physically plausible, has not been proved yet. (Notice however that

$$I'_\nu \geq I_\nu \implies T[I'] \geq T[I]$$

by (35), since the Planck function B_ν is increasing for each $\nu > 0$.) Thus, the map $S_\nu \mapsto I_\nu$ preserves both the integral and the order between radiation intensities. Now there is a clever characterization of order preserving maps on L^1 leaving the integral invariant, which is due to Crandall and Tartar [?]. Roughly speaking, a map from L^1 to itself that preserves the integral is order preserving iff it is nonexpansive in L^1 . This brings in the notion of L^1 -accretivity, which is at the heart⁴ of Mercier's remarkable discovery.

Indeed, the monotonicity argument above, together with Proposition 1 of [?] (with $C = L^1(\mathbf{R} \times (-1, 1) \times (0, \infty))^+$, which is the set of a.e. positive elements of $L^1(\mathbf{R} \times (-1, 1) \times (0, \infty))$) strongly suggest that it might be a good idea⁵ to study

$$\int_{-\infty}^{\infty} \int_{-1}^1 \int_0^{\infty} (I_\nu^2 - I_\nu^1)_+(\tau, \mu) d\nu d\mu d\tau$$

in terms of

$$\int_{-\infty}^{\infty} \int_{-1}^1 \int_0^{\infty} (S_\nu^2 - S_\nu^1)_+(\tau, \mu) d\nu d\mu d\tau$$

where $S_\nu^1, S_\nu^2 \in C$ and I_ν^1, I_ν^2 are the solutions of the steady radiative transfer equation above with source terms S_ν^1 and S_ν^2 respectively. (Mercier's original argument is even more complex, because he assumes that the opacity κ_ν depends on the temperature T , and is a decreasing function of T for each $\nu > 0$ while $T \mapsto \kappa_\nu(T)B_\nu(T)$ is nondecreasing; the reader can easily verify that the intuitive argument above still applies, provided of

⁴The Crandall-Tartar lemma appeared a few years before Mercier's paper [Mer87] on the radiative transfer equation. I learned of both results in 1984, during discussions in Mercier's lab, either from Mercier himself, or from Tartar. At the time of this writing, I cannot remember whether the Crandall-Tartar lemma was mentioned to me in connection with Mercier's result, or for some other reason.

⁵This may be a reconstruction of a discussion with Mercier in the early 1980's, unless he found the L^1 -accretivity structure of the radiative transfer equations by some other argument which I fail to remember.

course that our physically natural assumption that $S'_\nu \geq S_\nu \implies T[I'] \geq T[I]$ remains valid in this case as well.)

Define $s_+(z) = 1_{z \geq 0}$, and $z_+ = \max(z, 0)$ while $z_- = \max(-z, 0)$. Thus

$$z = z_+ - z_-, \quad |z| = z_+ + z_-, \quad z_+ = z s_+(z).$$

In accordance with the discussion above, we multiply both sides of the radiative transfer equation for two solutions I_ν and I'_ν by $s_+(I_\nu - I'_\nu)$ and integrate in all variables. This is precisely Mercier's computation (simpler because κ_ν is independent of the temperature). Denote

$$\langle \Phi \rangle := \int_0^\infty \int_{-1}^1 \Phi(\mu, \nu) d\mu d\nu$$

With $T = T[I]$ and $T' = T[I']$ defined by (35), let us compute

$$\begin{aligned} D &:= \langle \kappa_\nu((I_\nu - I'_\nu) - a_\nu(J_\nu - J'_\nu) - (1 - a_\nu)(B_\nu(T) - B_\nu(T')))s_+(I_\nu - I'_\nu) \rangle \\ &= \langle \kappa_\nu(1 - a_\nu)((I_\nu - I'_\nu) - (B_\nu(T) - B_\nu(T')))s_+(I_\nu - I'_\nu) \rangle \\ &\quad + \langle \kappa_\nu a_\nu((I_\nu - I'_\nu) - (J_\nu - J'_\nu))s_+(I_\nu - I'_\nu) \rangle =: D_1 + D_2 \end{aligned}$$

Observe that

$$\begin{aligned} (J_\nu - J'_\nu)s_+(I_\nu(\mu) - I'_\nu(\mu)) &= \frac{1}{2} \int_{-1}^1 (I_\nu - I'_\nu)(\mu') s_+(I_\nu - I'_\nu)(\mu) d\mu' \\ &\leq \frac{1}{2} \int_{-1}^1 (I_\nu - I'_\nu)_+(\mu') d\mu', \end{aligned}$$

so that $D_2 \geq 0$.

Next

$$D_1 = \langle \kappa_\nu(1 - a_\nu)((I_\nu - I'_\nu) - (B_\nu(T) - B_\nu(T')))(s_+(I_\nu - I'_\nu) - s_+(T - T')) \rangle$$

because

$$T = T[I] \text{ and } T' = T[I'] \implies \langle \kappa_\nu(1 - a_\nu)((I_\nu - I'_\nu) - (B_\nu(T) - B_\nu(T'))) \rangle = 0.$$

Since B_ν is increasing for each $\nu > 0$, one has

$$s_+(T - T') = s_+(B_\nu(T) - B_\nu(T'))$$

so that

$$D_1 = \langle \kappa_\nu(1 - a_\nu)((I_\nu - I'_\nu) - (B_\nu(T) - B_\nu(T')))(s_+(I_\nu - I'_\nu) - s_+(B_\nu(T) - B_\nu(T'))) \rangle$$

and

$$s_+ \text{ nondecreasing} \implies D_1 \geq 0.$$

Let I_ν and J_ν be two solutions of (36) with boundary data

$$\begin{aligned} I_\nu(0, \mu) &= Q_\nu^+(\mu), & I_\nu(Z, -\mu) &= Q_\nu^-(\mu), & 0 < \mu < 1, \\ J_\nu(0, \mu) &= R_\nu^+(\mu), & J_\nu(Z, -\mu) &= R_\nu^-(\mu), & 0 < \mu < 1. \end{aligned}$$

Assume that

$$Q_\nu^\pm(\mu) \leq R_\nu^\pm(\mu) \quad \text{for a.e. } (\mu, \nu) \in (0, 1) \times (0, \infty).$$

Then

$$\begin{aligned} \partial_\tau \langle \mu(I_\nu - J_\nu)_+ \rangle &\leq -\langle \kappa_\nu(1 - a_\nu)((I_\nu - J_\nu) - (B_\nu(T[I]) - B_\nu(T[J])))s_+(I_\nu - J_\nu) \rangle \\ &\quad -\langle \kappa_\nu a_\nu((I_\nu - J_\nu) - (J_\nu - \tilde{J}_\nu))s_+(I_\nu - J_\nu) \rangle \leq 0, \end{aligned}$$

so that $\tau \mapsto \langle \mu(I_\nu - J_\nu)_+ \rangle(\tau)$ is nonincreasing. Since

$$\begin{aligned} Q_\nu^- \leq R_\nu^- &\implies \langle \mu(I_\nu - J_\nu)_+ \rangle(Z) = \langle \mu_+(I_\nu - J_\nu)_+ \rangle(Z) \geq 0, \\ Q_\nu^+ \leq R_\nu^+ &\implies \langle \mu(I_\nu - J_\nu)_+ \rangle(0) = -\langle \mu_-(I_\nu - J_\nu)_+ \rangle(0) \leq 0, \end{aligned}$$

one has

$$\begin{aligned} &\text{for a.e. } \tau \in (0, Z) \quad 0 = \langle \mu(I_\nu - J_\nu)_+ \rangle \\ &= \langle \kappa_\nu a_\nu((I_\nu - J_\nu) - (J_\nu - \tilde{J}_\nu))s_+(I_\nu - J_\nu) \rangle \\ &= \langle \kappa_\nu(1 - a_\nu)((I_\nu - J_\nu) - (B_\nu(T[I]) - B_\nu(T[J])))s_+(I_\nu - J_\nu) \rangle, \end{aligned}$$

and

$$(I_\nu - J_\nu)_+(0, -\mu) = (I_\nu - J_\nu)_+(Z, \mu) = 0 \quad \text{for a.e. } \mu \in (0, 1).$$

Besides, since $\kappa_\nu(1 - a_\nu) > 0$ for all $\nu > 0$

$$\begin{aligned} 0 &= \langle \kappa_\nu(1 - a_\nu)((I_\nu - J_\nu) - (B_\nu(T[I]) - B_\nu(T[J])))s_+(I_\nu - J_\nu) \rangle \\ &= \langle \kappa_\nu(1 - a_\nu)((I_\nu - J_\nu) - (B_\nu(T[I]) - B_\nu(T[J])))s_+(I_\nu - J_\nu) - s_+(T[I] - T[J]) \rangle \\ &\implies s_+(I_\nu(\tau, \mu) - J_\nu(\tau, \mu)) = s_+(T[I] - T[J]) \text{ for a.e. } (\tau, \mu, \nu). \end{aligned}$$

At this point, we must appeal to an additional idea, which is not present in Mercier's paper [Mer87]. Since we are dealing with solutions of the radiative transfer equation having the slab symmetry, it is natural idea to use the K -invariant (in the terminology of section 10 in chapter I of Chandrasekhar [Cha50]). This idea⁶ is at the heart of the exponential decay estimate for the Milne problem obtained in [Gol87], and will be used here for a different purpose.

We compute

$$\begin{aligned} \partial_\tau \left\langle \frac{\mu^2}{\kappa_\nu}(I_\nu - J_\nu)_+ \right\rangle &= -\langle a_\nu \mu((I_\nu - J_\nu) - (J_\nu - \tilde{J}_\nu))s_+(T[I] - T[J]) \rangle \\ &\quad -\langle (1 - a_\nu)\mu((I_\nu - J_\nu) - (B_\nu(T[I]) - B_\nu(T[J])))s_+(T[I] - T[J]) \rangle \\ &= -\langle a_\nu \mu(I_\nu - J_\nu)s_+(T[I] - T[J]) \rangle - \langle (1 - a_\nu)\mu(I_\nu - J_\nu)s_+(T[I] - T[J]) \rangle \\ &= -\langle \mu(I_\nu - J_\nu)s_+(T[I] - T[J]) \rangle = -\langle \mu(I_\nu - J_\nu)_+ \rangle = 0, \end{aligned}$$

since

$$\int_{-1}^1 \mu(J_\nu(\tau) - \tilde{J}_\nu(\tau))d\mu = \int_{-1}^1 \mu(B_\nu(T[I]) - B_\nu(T[J]))d\mu = 0.$$

⁶A somewhat similar idea, unfortunately unpublished, had been used by R. Sentis to simplify the uniqueness proof for the linear Milne problem studied in [BSS84].

Next we integrate in $\tau \in (0, Z)$, and observe that

$$\begin{aligned} (I_\nu - J_\nu)_+(0, -\mu) = 0 \text{ and } Q_\nu^+(\mu) \leq R_\nu^+(\mu) \quad \text{for a.e. } \mu \in (0, 1) \\ \implies \left\langle \frac{\mu^2}{\kappa_\nu} (I_\nu - J_\nu)_+ \right\rangle (\tau) = \left\langle \frac{\mu^2}{\kappa_\nu} (I_\nu - J_\nu)_+ \right\rangle (0) = 0. \end{aligned}$$

Summarizing, we have proved the following result.

Theorem 5. *Assume that $0 < \kappa_\nu \leq \kappa_M$, while $0 \leq a_\nu < 1$ for all $\nu > 0$. Let $Q^\pm, R^\pm \in L^1((0, 1) \times (0, \infty))$ satisfy*

$$0 \leq Q_\nu^\pm(\mu) \leq R_\nu^\pm(\mu) \quad \text{for a.e. } (\mu, \nu) \in (0, 1) \times (0, \infty).$$

Then, the solutions $(I_\nu, T[I])$ of (36), and $(J_\nu, T[J])$ of (36) with boundary data $Q_\nu^\pm(\mu)$ replaced with $R_\nu^\pm(\mu)$ satisfy

$$I_\nu(\tau, \mu) \leq J_\nu(\tau, \mu) \text{ and } T[I](\tau) \leq T[J](\tau) \quad \text{for a.e. } (\tau, \mu) \in (-1, 1) \times (0, \infty).$$

In particular,

$$\begin{aligned} Q_\nu^\pm(\mu) = R_\nu^\pm(\mu) \text{ for a.e. } (\mu, \nu) \in (0, 1) \times (0, \infty) \\ \implies I_\nu(\tau, \mu) = J_\nu(\tau, \mu) \text{ and } T[I](\tau) = T[J](\tau) \quad \text{for a.e. } (\tau, \mu) \in (-1, 1) \times (0, \infty). \end{aligned}$$

One has also the following form of Maximum Principle.

Corollary 2. *Assume that $0 < \kappa_\nu \leq \kappa_M$, while $0 \leq a_\nu < 1$ for all $\nu > 0$. Let $Q_\nu^\pm(\mu) \leq B_\nu(T_M)$ (resp. $Q_\nu^\pm(\mu) \geq B_\nu(T_m)$) for a.e. $(\mu, \nu) \in (0, 1) \times (0, \infty)$. Then*

$$\begin{aligned} I_\nu(\tau, \mu) \leq B_\nu(T_M) \text{ and } T[I](\tau) \leq T_M \\ \text{resp. } I_\nu(\tau, \mu) \geq B_\nu(T_m) \text{ and } T[I](\tau) \geq T_m \\ \text{for a.e. } (\tau, \mu) \in (-1, 1) \times (0, \infty). \end{aligned}$$

Proof Indeed, $J_\nu = B_\nu(T_M)$ and $T[J] = T_M$ (resp. $J_\nu = B_\nu(T_m)$ and $T[J] = T_m$) is the solution of (36) with boundary data $R_\nu^\pm = B_\nu(T_M)$ (resp. $R_\nu^\pm = B_\nu(T_m)$). \square

In Theorem 4, if one has the stronger condition

$$0 \leq Q_\nu^\pm(\mu) \leq B_\nu(T_M) \quad \text{for a.e. } (\mu, \nu) \in (0, 1) \times (0, \infty)$$

one obtains the following bound for the numerical and theoretical solutions

$$0 \leq I_\nu^1 \leq I_\nu^2 \leq \dots \leq I_\nu^n \leq \dots \leq I_\nu \leq B_\nu(T_M)$$

while

$$0 \leq T^1 \leq T_2 \leq \dots \leq T^n \leq \dots \leq T \leq T_M.$$

10. Radiative Transfer with Rayleigh Scattering in a Slab

In this section, we discuss the same problem as in the previous section, with the isotropic scattering replaced by the Rayleigh phase function. In the case of slab symmetry, the Rayleigh phase function is

$$p(\mu, \mu') = \frac{3}{16}(3 - \mu^2) + \frac{3}{16}(3\mu'^2 - 1)\mu'^2$$

(see section 11.2 in chapter I of [Cha50]). Observe that

$$p(\mu, \mu') = \frac{3}{16}(3 + 3\mu^2\mu'^2 - \mu^2 - \mu'^2) \geq \frac{3}{16} > 0, \quad (39)$$

while

$$\frac{1}{2} \int_{-1}^1 p(\mu, \mu') d\mu = \frac{3}{16}(6 + 3 \cdot \frac{2}{3}\mu'^2 - \frac{2}{3} - 2\mu'^2) = 1. \quad (40)$$

Keeping (35) as the defining equation for $T[I]$, the problem (36) becomes

$$\begin{aligned} (\mu\partial_\tau + \kappa_\nu)I_\nu(\tau, \mu) &= \frac{3}{8}\kappa_\nu a_\nu((3 - \mu^2)J_\nu(\tau) + (3\mu^2 - 1)K_\nu(\tau)) \\ &\quad + \kappa_\nu(1 - a_\nu)B_\nu(T[J](\tau)), \\ I_\nu(0, \mu) &= Q_\nu^+(\mu), \quad I_\nu(Z, -\mu) = Q_\nu^-(\mu), \quad 0 < \mu < 1, \end{aligned} \quad (41)$$

with $K = \frac{1}{2} \int_{-1}^1 \mu^2 Id\mu$, and one easily checks that (35) and (40) imply that

$$\partial_\tau \int_0^\infty \int_{-1}^1 \mu I_\nu(\tau, \mu) d\mu d\nu = 0.$$

Starting from $I_\nu^0(\tau, \mu) = 0$ and $T^0(\tau) = 0$, one solves for I^{n+1}

$$\begin{aligned} (\mu\partial_\tau + \kappa_\nu)I_\nu^{n+1}(\tau, \mu) &= \frac{3}{8}\kappa_\nu a_\nu((3 - \mu^2)J_\nu^n(\tau) + (3\mu^2 - 1)K_\nu^n(\tau)) \\ &\quad + \kappa_\nu(1 - a_\nu)B_\nu(T^n(\tau)), \quad T^n := T[J^n] \\ I_\nu^{n+1}(0, \mu) &= Q_\nu^+(\mu), \quad I_\nu^{n+1}(Z, -\mu) = Q_\nu^-(\mu), \quad 0 < \mu < 1. \end{aligned} \quad (42)$$

Since B_ν is nondecreasing for each $\nu > 0$, one easily checks with (39) that

$$\begin{aligned} 0 &= I_\nu^0 \leq I_\nu^1 \leq I_\nu^2 \leq \dots \leq I_\nu^n \leq I_\nu^{n+1} \leq \dots \\ 0 &= T^0 \leq T^1 \leq T^2 \leq \dots \leq T^n \leq T^{n+1} \leq \dots \end{aligned}$$

Explicitly

$$\begin{aligned} I_\nu^{n+1}(\tau, \mu) &= e^{-\frac{\kappa_\nu \tau}{\mu}} Q_\nu^+(\mu) \mathbf{1}_{\mu > 0} + e^{-\frac{\kappa_\nu (Z - \tau)}{|\mu|}} Q_\nu^- (|\mu|) \mathbf{1}_{\mu < 0} \\ &\quad + \mathbf{1}_{\mu > 0} \int_0^\tau e^{-\frac{\kappa_\nu (\tau - t)}{\mu}} \frac{\kappa_\nu}{\mu} \frac{3}{8} a_\nu((3 - \mu^2)J_\nu^n(t) + (3\mu^2 - 1)K_\nu^n(t)) dt \\ &\quad + \mathbf{1}_{\mu > 0} \int_0^\tau e^{-\frac{\kappa_\nu (\tau - t)}{\mu}} \frac{\kappa_\nu}{\mu} (1 - a_\nu) B_\nu(T^n(t)) dt \\ &\quad + \mathbf{1}_{\mu < 0} \int_\tau^Z e^{-\frac{\kappa_\nu (t - \tau)}{|\mu|}} \frac{\kappa_\nu}{|\mu|} \frac{3}{8} a_\nu((3 - \mu^2)J_\nu^n(t) + (3\mu^2 - 1)K_\nu^n(t)) dt \\ &\quad + \mathbf{1}_{\mu < 0} \int_\tau^Z e^{-\frac{\kappa_\nu (t - \tau)}{|\mu|}} \frac{\kappa_\nu}{|\mu|} (1 - a_\nu) B_\nu(T^n(t)) dt. \end{aligned} \quad (43)$$

This scheme can be reduced to the following iteration to compute first J_ν and $\tilde{\mu}^2 I_\nu$ as follows:

$$\begin{aligned}
J_\nu^{n+1}(\tau) &= \frac{1}{2} \int_0^1 \left(e^{-\frac{\kappa_\nu \tau}{\mu}} Q_\nu^+(\mu) \mathbf{1}_{\mu>0} + e^{-\frac{\kappa_\nu(Z-\tau)}{|\mu|}} Q_\nu^- (|\mu|) \mathbf{1}_{\mu<0} \right) d\mu \\
&\quad + \frac{3}{16} \int_0^Z E_1(\kappa_\nu|\tau-t|) \kappa_\nu a_\nu (3J_\nu^n(t) - K_\nu^n(t)) dt \\
&\quad + \frac{3}{16} \int_0^Z E_3(\kappa_\nu|\tau-t|) \kappa_\nu a_\nu (3K_\nu^n(t) - J_\nu^n(t)) dt \\
&\quad + \frac{1}{2} \int_0^Z E_1(\kappa_\nu|\tau-t|) \kappa_\nu (1-a_\nu) B_\nu(T^n(t)) dt,
\end{aligned}$$

while

$$\begin{aligned}
K_\nu^{n+1}(\tau) &= \frac{1}{2} \int_0^1 \left(e^{-\frac{\kappa_\nu \tau}{\mu}} Q_\nu^+(\mu) \mathbf{1}_{\mu>0} + e^{-\frac{\kappa_\nu(Z-\tau)}{|\mu|}} Q_\nu^- (|\mu|) \mathbf{1}_{\mu<0} \right) \mu^2 d\mu \\
&\quad + \frac{3}{16} \int_0^Z E_3(\kappa_\nu|\tau-t|) \kappa_\nu a_\nu (3J_\nu^n(t) - K_\nu^n(t)) dt \\
&\quad + \frac{3}{16} \int_0^Z E_5(\kappa_\nu|\tau-t|) \kappa_\nu a_\nu (3K_\nu^n(t) - J_\nu^n(t)) dt \\
&\quad + \frac{1}{2} \int_0^Z E_3(\kappa_\nu|\tau-t|) \kappa_\nu (1-a_\nu) B_\nu(T^n(t)) dt,
\end{aligned}$$

where we have denoted

$$E_n(X) := \int_1^\infty \frac{e^{-Xy}}{y^n} dy = \int_X^\infty \frac{e^{-z}}{z^n} dz = \int_0^1 e^{-X/\mu} \mu^{n-2} d\mu.$$

Once J_ν and K_ν are known, the right hand side of (41) is known, and I_ν is obtained by a simple quadrature formula.

Returning to (43), assume that

$$0 \leq Q_\nu^\pm \leq B_\nu(T_M), \quad 0 \leq I_\nu^n \leq B_\nu(T_M) \quad \text{and} \quad 0 \leq T^n \leq T_M.$$

Then

$$\begin{aligned}
I_\nu^{n+1}(\tau, \mu) &\leq \left(e^{-\frac{\kappa_\nu \tau}{\mu}} \mathbf{1}_{\mu>0} + e^{-\frac{\kappa_\nu(Z-\tau)}{|\mu|}} \mathbf{1}_{\mu<0} \right) B_\nu(T_M) \\
&\quad + \mathbf{1}_{\mu>0} \int_0^\tau e^{-\frac{\kappa_\nu(\tau-t)}{\mu}} \frac{\kappa_\nu}{\mu} \frac{3}{8} a_\nu ((3 - \mu^2) B_\nu(T_M) + (\mu^2 - \frac{1}{3}) B_\nu(T_M)) dt \\
&\quad + \mathbf{1}_{\mu>0} \int_0^\tau e^{-\frac{\kappa_\nu(\tau-t)}{\mu}} \frac{\kappa_\nu}{\mu} (1 - a_\nu) B_\nu(T_M) dt \\
&\quad + \mathbf{1}_{\mu<0} \int_\tau^Z e^{-\frac{\kappa_\nu(t-\tau)}{|\mu|}} \frac{\kappa_\nu}{|\mu|} \frac{3}{8} a_\nu ((3 - \mu^2) B_\nu(T_M) + (\mu^2 - \frac{1}{3}) B_\nu(T_M)) dt \\
&\quad + \mathbf{1}_{\mu<0} \int_\tau^Z e^{-\frac{\kappa_\nu(t-\tau)}{|\mu|}} \frac{\kappa_\nu}{|\mu|} (1 - a_\nu) B_\nu(T_M) dt \\
&= \left(e^{-\frac{\kappa_\nu \tau}{\mu}} \mathbf{1}_{\mu>0} + e^{-\frac{\kappa_\nu(Z-\tau)}{|\mu|}} \mathbf{1}_{\mu<0} \right) B_\nu(T_M) \\
&\quad + \mathbf{1}_{\mu>0} \int_0^\tau e^{-\frac{\kappa_\nu(\tau-t)}{\mu}} \frac{\kappa_\nu}{\mu} \left(\frac{3}{8} a_\nu (3 - \frac{1}{3}) + (1 - a_\nu) \right) B_\nu(T_M) dt \\
&\quad + \mathbf{1}_{\mu<0} \int_\tau^Z e^{-\frac{\kappa_\nu(t-\tau)}{|\mu|}} \frac{\kappa_\nu}{|\mu|} \left(\frac{3}{8} a_\nu (3 - \frac{1}{3}) + (1 - a_\nu) \right) B_\nu(T_M) dt \\
&= \left(e^{-\frac{\kappa_\nu \tau}{\mu}} \mathbf{1}_{\mu>0} + e^{-\frac{\kappa_\nu(Z-\tau)}{|\mu|}} \mathbf{1}_{\mu<0} \right) B_\nu(T_M) \\
&\quad + B_\nu(T_M) \left(\mathbf{1}_{\mu>0} \int_0^\tau e^{-\frac{\kappa_\nu(\tau-t)}{\mu}} \frac{\kappa_\nu}{\mu} dt + \mathbf{1}_{\mu<0} \int_\tau^Z e^{-\frac{\kappa_\nu(t-\tau)}{|\mu|}} \frac{\kappa_\nu}{|\mu|} dt \right) \\
&= \left(e^{-\frac{\kappa_\nu \tau}{\mu}} \mathbf{1}_{\mu>0} + e^{-\frac{\kappa_\nu(Z-\tau)}{|\mu|}} \mathbf{1}_{\mu<0} \right) B_\nu(T_M) \\
&\quad + B_\nu(T_M) \left(\mathbf{1}_{\mu>0} \left(1 - e^{-\frac{\kappa_\nu \tau}{\mu}} \right) + \mathbf{1}_{\mu<0} \left(1 - e^{-\frac{\kappa_\nu(Z-\tau)}{|\mu|}} \right) \right) \\
&= B_\nu(T_M).
\end{aligned}$$

Besides

$$T^{n+1} = T[I^{n+1}] \leq T[B_\nu(T_M)] = T_M$$

(using again that $T \mapsto B_\nu(T)$ is increasing for each $\nu > 0$ while $\kappa_\nu(1 - a_\nu) > 0$ for all $\nu > 0$).

Summarizing, we have proved the following result.

Theorem 6. *Assume that $\kappa_\nu > 0$ while $0 \leq a_\nu < 1$ for all $\nu > 0$. Let Q^\pm satisfy*

$$0 \leq Q_\nu^\pm(\mu) \leq B_\nu(T_M) \quad \text{for all } \mu \in (-1, 1) \text{ and } \nu > 0.$$

The iteration method (42) starting from $I_\nu^0 = 0$ and $T^0 = 0$ defines a sequence of radiative intensities I_ν^n and temperatures T^n converging pointwise to I_ν and $T = T[I]$ respectively, which is a solution of (41).

The argument above is based on the monotonicity of the sequences I_ν^n and T^n , and does not give any information on the convergence rate.

Finally, Theorem 5 holds verbatim for the problem (41). Here are the (slight) modifications to the proof due to the Rayleigh phase function.

First, we slightly modify the argument concerning the term D_2 as follows. In the case of the Rayleigh phase function

$$D_2 = \frac{1}{2} \int_0^\infty \kappa_\nu a_\nu \int_{-1}^1 (I_\nu - I'_\nu)_+(\mu) d\mu d\nu \\ - \frac{1}{2} \int_0^\infty \kappa_\nu a_\nu \int_{-1}^1 \int_{-1}^1 p(\mu, \mu') (I_\nu - J'_\nu)(\mu') s_+(I_\nu - I'_\nu)(\mu) d\mu' d\mu d\nu.$$

Since $p \geq 0$, one has

$$p(\mu, \mu') (I_\nu - J'_\nu)(\mu') s_+(I_\nu - I'_\nu)(\mu) \leq p(\mu, \mu') (I_\nu - J'_\nu)_+(\mu')$$

so that

$$D_2 \geq \frac{1}{2} \int_0^\infty \kappa_\nu a_\nu \int_{-1}^1 (I_\nu - I'_\nu)_+(\mu) d\mu d\nu \\ - \frac{1}{2} \int_0^\infty \kappa_\nu a_\nu \int_{-1}^1 \int_{-1}^1 p(\mu, \mu') (I_\nu - J'_\nu)_+(\mu') d\mu' d\mu d\nu = 0$$

since

$$\frac{1}{2} \int_{-1}^1 p(\mu, \mu') d\mu = 1.$$

Therefore, following the proof of Theorem 5, we obtain in the same manner the following conclusions

$$\langle \mu (I_\nu - J_\nu)_+ \rangle(\tau) = 0 \text{ for a.e. } \tau \in (0, Z),$$

and

$$s_+(I_\nu(\tau, \mu) - J_\nu(\tau, \mu)) = s_+(T[I](\tau) - T[J](\tau)) \\ \text{for a.e. } (\tau, \mu, \nu) \in (0, Z) \times (-1, 1) \times (0, \infty),$$

while

$$(I_\nu - J_\nu)_+(0, -\mu) = (I_\nu - J_\nu)_+(Z, \mu) = 0 \quad \text{for a.e. } \mu \in (0, 1).$$

Next we compute

$$\partial_\tau \left\langle \frac{\mu^2}{\kappa_\nu} (I_\nu - J_\nu)_+ \right\rangle = -\frac{1}{2} \int_0^\infty a_\nu \int_{-1}^1 \mu (I_\nu - J_\nu)_+(\tau, \mu) d\mu d\nu \\ + \frac{1}{2} \int_0^\infty a_\nu \int_{-1}^1 \mu \int_{-1}^1 p(\mu, \mu') (I_\nu - J_\nu)_+(\tau, \mu') d\mu' d\nu s_+(T[I](\tau) - T[J](\tau)) \\ - \langle (1 - a_\nu) \mu ((I_\nu - J_\nu) - (B_\nu(T[I]) - B_\nu(T[J]))) s_+(T[I] - T[J]) \rangle \\ = -\langle a_\nu \mu (I_\nu - J_\nu) s_+(T[I] - T[J]) \rangle - \langle (1 - a_\nu) \mu (I_\nu - J_\nu) s_+(T[I] - T[J]) \rangle \\ = -\langle \mu (I_\nu - J_\nu) s_+(T[I] - T[J]) \rangle = -\langle \mu (I_\nu - J_\nu)_+ \rangle = 0,$$

since

$$\int_{-1}^1 \mu p(\mu, \mu') d\mu = \int_{-1}^1 \mu (B_\nu(T[I]) - B_\nu(T[J])) d\mu = 0.$$

Finally we integrate in $\tau \in (0, Z)$, and conclude as in the previous section that

$$(I_\nu - J_\nu)_+(0, -\mu) = 0 \text{ and } Q_\nu^+(\mu) \leq R_\nu^+(\mu) \quad \text{for a.e. } \mu \in (0, 1) \\ \implies \left\langle \frac{\mu^2}{\kappa_\nu} (I_\nu - J_\nu)_+ \right\rangle(\tau) = \left\langle \frac{\mu^2}{\kappa_\nu} (I_\nu - J_\nu)_+ \right\rangle(0) = 0.$$