



HAL
open science

On optimal cloaking-by-mapping transformations

Yves Capdeboscq, Michael S Vogelius

► **To cite this version:**

Yves Capdeboscq, Michael S Vogelius. On optimal cloaking-by-mapping transformations. ESAIM: Mathematical Modelling and Numerical Analysis, 2022, 56 (1), pp.303 - 316. 10.1051/m2an/2022004 . hal-03335152v1

HAL Id: hal-03335152

<https://hal.sorbonne-universite.fr/hal-03335152v1>

Submitted on 6 Sep 2021 (v1), last revised 10 Feb 2022 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

ON OPTIMAL CLOAKING-BY-MAPPING TRANSFORMATIONS

YVES CAPDEBOSCQ AND MICHAEL S. VOGELIUS

ABSTRACT. A central ingredient of cloaking-by-mapping is the diffeomorphism which transforms an annulus with a small hole into an annulus with a finite size hole, while being the identity on the outer boundary of the annulus. The resulting meta-material is anisotropic, which makes it difficult to manufacture. The problem of minimizing anisotropy among radial transformations has been studied in [4]. In this work, as in [4], we formulate the problem of minimizing anisotropy as an energy minimization problem. Our main goal is to provide strong evidence for the conjecture that for cloaks with circular boundaries, non-radial transformations do not lead to lower degree of anisotropy. In the final section, we consider cloaks with non-circular boundaries and show that in this case, non-radial cloaks may be advantageous, when it comes to minimizing anisotropy.

1. INTRODUCTION

A central ingredient in the construction of (approximate) cloaks by the passive cloaking technique, known as “cloaking by mapping”, is the diffeomorphism, which transforms an annulus with a small hole into an annulus with a finite size hole, and which is the identity on the outer boundary of the annulus. The push-forward of the background coefficient (say, the identity matrix) with the diffeomorphism represents the meta-material needed for the cloak, and the finite size hole is the area that may be used as a “hiding place” [6]. The fact that the diffeomorphism is the identity on the outer boundary ensures that the perturbation in the “far field” is that corresponding to a small inhomogeneity. The corresponding “lack of cloaking”/visibility can be estimated by the volume of the small inhomogeneity. The required meta-material is anisotropic, which presents a problem when it comes to actual manufacture of the cloak. Typically a radial affine transformation has been used [2, 3, 6, 7, 8], however, a very natural question arises, namely : “are there transformations that lead to lower degree of anisotropy than the radial affine transformation?” In [4] it was shown that there are indeed better radial transformations than the affine, when it comes to minimizing anisotropy. In that paper the meta-material obtained by “optimal radial transformation” is also shown to be quite related to meta-materials obtained by other cloak enhancement strategies, employing additional layers [1, 5]. The focus of this note is to produce very strong evidence for the conjecture that when the cloak takes the shape of a classical annulus, non-radial transformations do not help in reducing the degree of anisotropy. Like in [4], we formulate the problem of minimizing anisotropy as a variational problem (minimization of an appropriate energy). Corollary 7 summarizes our main results. Broadly speaking, we show that

- There exists a radial transformation, which is a stationary point for the energy.

- This radial transformation has smaller energy than all other transformations with “directional field” $\frac{x}{|x|}$.
- If the amplitude is kept fixed and radial, then any change in the “directional field” away from $\frac{x}{|x|}$ will increase energy.

In the final section of this note we consider the case when the outer (and inner) boundary of the cloak are not circles, and we illustrate how the optimal radial transformation for the circular case translates into a non-radial (optimal) transformation for a non-circular cloak.

2. PRELIMINARIES

For $r > 0$ we set

$$B_r = \{x \in \mathbb{R}^2 : |x| < r\} \quad , \quad \text{and} \quad C_r = \{x \in \mathbb{R}^2 : |x| = r\} \quad .$$

Given $\epsilon > 0$, we shall use the notation Φ for a bijective diffeomorphism $\overline{B_1} \setminus B_\epsilon \rightarrow \overline{B_1} \setminus B_{\frac{1}{2}}$ with $\Phi \in C^1(\overline{B_1} \setminus B_\epsilon; \overline{B_1} \setminus B_{\frac{1}{2}})$, and $\Phi^{-1} \in C^1(\overline{B_1} \setminus B_{\frac{1}{2}}; \overline{B_1} \setminus B_\epsilon)$. We furthermore impose that

$$\Phi|_{C_1} = Id \quad , \quad \text{and} \quad \Phi(C_\epsilon) = C_{\frac{1}{2}} \quad .$$

One such transformation is the radial affine transformation, given by

$$x \rightarrow \left(\frac{|x| - 1}{2(1 - \epsilon)} + 1 \right) \frac{x}{|x|} \quad .$$

The push-forward of the identity matrix with the diffeomorphism Φ is given by

$$\Phi_* [I] (\Phi(x)) = \frac{D\Phi D\Phi^T}{|\det D\Phi|} (x) \quad .$$

This is a positive definite matrix, and since we are in two dimensions, with determinant 1. Let $0 < \lambda_1(x) \leq 1 \leq \lambda_2(x)$ denote the eigenvalues of $\Phi_* [I] (\Phi(x))$. A natural measure of the degree of anisotropy of $\Phi_* [I]$ at the point $\Phi(x)$ is

$$\begin{aligned} |\lambda_1(x) - 1| + |\lambda_2(x) - 1| = \lambda_2(x) - \lambda_1(x) &= \sqrt{(\lambda_2(x) - \lambda_1(x))^2} \\ &= \sqrt{(\lambda_1(x) + \lambda_2(x))^2 - 4} \quad . \end{aligned}$$

To minimize this we must minimize trace $\Phi_* [I] (\Phi(x))$. As a way of minimizing the aggregate anisotropy we shall seek to minimize¹

$$I_p(\Phi) = \int_{B_1 \setminus B_\epsilon} (\text{trace } \Phi_* [I])^p (\Phi(x)) \, dx$$

for a fixed choice of $1 \leq p < \infty$, and

$$I_\infty(\Phi) = \max_{x \in \overline{B_1} \setminus B_\epsilon} \text{trace } \Phi_* [I] (\Phi(x)) = \max_{y \in \overline{B_1} \setminus B_{\frac{1}{2}}} \text{trace } \Phi_* [I] (y) \quad ,$$

corresponding to $p = \infty$. Note that λ is an eigenvalue for $\Phi_* [I] (\Phi(x))$, with eigenvector v , if and only if λ is an eigenvalue for

$$\frac{D\Phi^T D\Phi}{|\det D\Phi|} (x) \quad ,$$

¹In a slight deviation from [4], the domain of integration of the energy functional is $B_1 \setminus B_\epsilon$, not the transformed domain $B_1 \setminus B_{\frac{1}{2}}$.

with eigenvector $D\Phi^T(x)v$, and thus

$$\text{trace } \Phi_* [I] (\Phi(x)) = \text{trace} \left[\frac{D\Phi^T D\Phi}{|\det D\Phi|} \right] (x) .$$

Proposition 1. *Let Φ be represented in terms of its polar decomposition*

$$\Phi = \exp(\psi)\phi ,$$

where the directional field ϕ is in $C^1(\overline{B_1} \setminus B_\epsilon; \mathcal{S}^1)$ and logarithmic amplitude ψ is in $C^1(\overline{B_1} \setminus B_\epsilon; \mathbb{R})$. Then

$$\text{trace} (D\Phi^T D\Phi) = |\Phi|^2 \left(|D\phi|^2 + |D\psi|^2 \right) .$$

Proof. Differentiating we find

$$D\Phi = \exp(\psi)\phi D\psi^T + \exp(\psi)D\phi .$$

Since $\phi^T \phi = 1$, we have

$$\phi^T D\phi = 0 , \text{ and } D\phi^T \phi = 0 ,$$

and therefore

$$\begin{aligned} D\Phi^T D\Phi &= \exp(2\psi) (D\psi\phi^T + D\phi^T) (\phi D\psi^T + D\phi) \\ &= |\Phi|^2 (D\phi^T D\phi + D\psi D\psi^T) . \end{aligned}$$

By taking the trace we arrive at the desired conclusion. \square

It is well known that ϕ , being in $C^1(\overline{B_1} \setminus B_\epsilon; \mathcal{S}^1)$, admits a canonical lift $\theta = \arg(\phi) \in C^1(\overline{B_1} \setminus B_\epsilon; \mathbb{R}/2\pi\mathbb{Z})$ ² such that

$$\phi = (\cos \theta, \sin \theta)^T .$$

We write

$$J = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, \quad e_r = \frac{x}{|x|}, \quad \text{and} \quad e_\theta = J \frac{x}{|x|} .$$

Proposition 2. *The matrix $D\phi$ has rank one; furthermore*

$$\text{Range}(D\phi) = \text{Span}(\phi)^\perp , \quad \text{and} \quad \text{Ker}(D\phi) = \text{Span}(D\theta)^\perp .$$

We denote by $\widehat{D\psi}, \widehat{D\theta}$ the angle defined by

$$\begin{aligned} \cos(\widehat{D\psi}, \widehat{D\theta}) &= \frac{1}{|D\psi| |D\theta|} D\psi \cdot D\theta , \quad \text{and} \\ \sin(\widehat{D\psi}, \widehat{D\theta}) &= \frac{1}{|D\psi| |D\theta|} \det(D\psi, D\theta) . \end{aligned}$$

Then

$$\text{trace } \Phi_* [I] (\Phi(x)) = \frac{1}{\left| \sin(\widehat{D\psi}, \widehat{D\theta}) \right|} \left(\frac{|D\theta|}{|D\psi|} + \frac{|D\psi|}{|D\theta|} \right) (x) \geq \left(\frac{|D\theta|}{|D\psi|} + \frac{|D\psi|}{|D\theta|} \right) (x)$$

with equality only when $D\psi \cdot D\theta = 0$.

²A function $\theta : \overline{B_1} \setminus B_\epsilon \rightarrow \mathbb{R}/2\pi\mathbb{Z}$ is an element of $C^1(\overline{B_1} \setminus B_\epsilon; \mathbb{R}/2\pi\mathbb{Z})$ iff given any point $x \in \overline{B_1} \setminus B_\epsilon$ there exists an open neighborhood ω_x of x , relative to $\overline{B_1} \setminus B_\epsilon$, and a representative of $\theta \pmod{2\pi}$ that lies in $C^1(\omega_x; \mathbb{R})$. Notice that the globally defined derivative of $\theta \in C^1(\overline{B_1} \setminus B_\epsilon; \mathbb{R}/2\pi\mathbb{Z})$, $D\theta$, lies in $C^0(\overline{B_1} \setminus B_\epsilon; \mathbb{R}^2)$.

Proof. We calculate

$$D\phi = (J\phi) D\theta^T,$$

which immediately leads to the statements about $\text{Range}(D\phi)$ and $\text{Ker}(D\phi)$, and which also gives

$$D\theta = (D\phi)^T (J\phi) .$$

As a consequence

$$\begin{aligned} \det D\Phi &= \det (\phi D\psi^T + (J\phi) D\theta^T) \exp(2\psi) \\ &= \det (D\psi, D\theta) |\Phi|^2 \\ &= |D\theta| |D\psi| \sin \left(\widehat{D\psi, D\theta} \right) |\Phi|^2 . \end{aligned}$$

Here we have used that $\det D\Phi \neq 0$, since Φ is a bijective diffeomorphism of $\overline{B_1} \setminus B_\epsilon$ onto $\overline{B_1} \setminus B_{\frac{1}{2}}$; consequently $\det (D\psi, D\theta) \neq 0$ and $|D\psi| |D\theta| > 0$ and $\sin \left(\widehat{D\psi, D\theta} \right)$ (and $\widehat{D\psi, D\theta}$) is well-defined. It now follows that

$$\begin{aligned} \text{trace } \Phi_* [I] (\Phi(x)) &= \frac{\left(|D\theta|^2 + |D\psi|^2 \right)}{|D\theta| |D\psi| \left| \sin \left(\widehat{D\psi, D\theta} \right) \right|} (x) \\ &= \frac{1}{\left| \sin \left(\widehat{D\psi, D\theta} \right) \right|} \left(\frac{|D\theta|}{|D\psi|} + \frac{|D\psi|}{|D\theta|} \right) (x) \geq \frac{|D\theta|}{|D\psi|} (x) + \frac{|D\psi|}{|D\theta|} (x) , \end{aligned}$$

with equality if and only if $D\psi$ is normal to $D\theta$, and therefore in the kernel of $D\phi$. \square

3. THE RADIAL TRANSFORMATION CASE

For the general case of a radial transformation $\phi = \frac{x}{|x|}$, and $\psi = f(|x|)$. Then $D\theta = \frac{1}{|x|} J \frac{x}{|x|}$ and $D\psi = f'(|x|) \frac{x}{|x|}$. The transformation

$$\Phi = \exp(\psi)\phi$$

is a bijective C^1 diffeomorphism of $\overline{B_1} \setminus B_\epsilon$ onto $\overline{B_1} \setminus B_{\frac{1}{2}}$ with

$$\Phi|_{C_1} = Id , \text{ and } \Phi(C_\epsilon) = C_{\frac{1}{2}} ,$$

if and only if

$$f(\epsilon) = -\log 2 , f(1) = 0 , \text{ and } f \in C^1([\epsilon, 1]) \text{ with } f'(r) > 0 \text{ for all } r \in [\epsilon, 1].$$

In this case, $\sin \left(\widehat{D\psi, D\theta} \right) = 1$, and

$$\text{trace } \Phi_* [I] (\Phi(x)) = \frac{1}{|x| f'(|x|)} + |x| f'(|x|) .$$

Proposition 3. *Suppose $1 \leq p < \infty$, and let I_p denote the energy*

$$I_p(f) := \int_{B_1 \setminus B_\epsilon} (\text{trace } \Phi_* [I])^p (\Phi(x)) dx = 2\pi \int_\epsilon^1 \left(\frac{1}{r f'(r)} + r f'(r) \right)^p r dr ,$$

with values in $(0, \infty)$, defined on the convex set

$$\mathcal{C} = \left\{ f \in C^1([\epsilon, 1]) : f' > 0 , f(\epsilon) = -\log 2 , f(1) = 0 \right\} .$$

Then

- I_p has a unique minimizer, f_p , in \mathcal{C} .
- f_p lies in $C^\infty([\epsilon, 1])$, and is the unique solution in \mathcal{C} to the Euler–Lagrange equation

$$\left(\left(\frac{1}{rf_p'(r)} + rf_p'(r) \right)^{p-1} \left(-\frac{1}{(f_p')^2} + r^2 \right) \right)' = 0 \text{ in } [\epsilon, 1] . \quad (\text{E-L})$$

Proof. We start by establishing (part of) the last statement concerning the existence of a unique solution to the Euler–Lagrange equation (E-L). By integration, any C^1 solution to (E-L) must satisfy

$$G(rf_p'(r)) = \frac{C}{r^2}$$

for some constant C , with the function $G : \mathbb{R}_+ \rightarrow \mathbb{R}$ given by

$$G(t) = \left(\frac{1}{t} + t \right)^{p-1} \left(-\frac{1}{t^2} + 1 \right) .$$

Now suppose $1 < p < \infty$. A simple calculation shows that G is monotonically increasing, with $G(1) = 0$, $\lim_{t \rightarrow 0^+} G(t) = -\infty$ and $\lim_{t \rightarrow \infty} G(t) = \infty$. $G^{-1} : \mathbb{R} \rightarrow \mathbb{R}_+$ is thus well defined, and f_p has the form

$$f_p(r) = \int_\epsilon^r f_p'(t) dt - \log 2 = \int_\epsilon^r t^{-1} G^{-1} \left(\frac{C}{t^2} \right) dt - \log 2 ,$$

for some constant C . The constant C must be chosen so that f_p satisfies the boundary condition $f_p(1) = 0$. As $C \rightarrow \int_\epsilon^1 t^{-1} G^{-1} \left(\frac{C}{t^2} \right) dt - \log 2$ is continuous and monotonically increasing, with

$$\int_\epsilon^1 t^{-1} G^{-1} \left(\frac{C}{t^2} \right) dt - \log 2 \rightarrow \begin{cases} |\log \epsilon| - \log 2 > 0 & \text{when } C \rightarrow 0 \\ -\log 2 < 0 & \text{when } C \rightarrow -\infty \end{cases} ,$$

it follows immediately that there exists a unique value $C_0 < 0$ for which the boundary condition $f_p(1) = 0$ is satisfied. This shows the uniqueness of the solution to the Euler–Lagrange equation in \mathcal{C} . Furthermore, the formula

$$f_p(r) = \int_\epsilon^r t^{-1} G^{-1} \left(\frac{C_0}{t^2} \right) dt - \log 2$$

clearly gives rise to a C^∞ function in \mathcal{C} which solves the equation (E-L), thus establishing the existence. A slightly modified argument works for $p = 1$, and in that case we find the (even more) explicit formula

$$f_1 : r \rightarrow \log \left(\frac{3r + \sqrt{9r^2 + 16(2-\epsilon)\left(\frac{1}{2} - \epsilon\right)}}{4(2-\epsilon)} \right) .$$

We now proceed to show that f_p is the unique minimizer of I_p in \mathcal{C} . Since the function $(0, \infty) \ni x \rightarrow \left(\frac{1}{x} + x\right)^p \in (0, \infty)$ is strictly convex, it follows immediately that I_p is strictly convex on \mathcal{C} . Now suppose there existed a function $g \in \mathcal{C}$ with $I_p(g) < I_p(f_p)$. The convexity of the functional I_p implies that

$$\frac{d}{d\tau} \Big|_{\tau=0} I_p(f_p + \tau(g - f_p)) \leq I_p(g) - I_p(f_p) < 0 ,$$

or

$$\int_{\epsilon}^1 \left(\frac{1}{r f'(r)} + r f'(r) \right)^{p-1} \left(-\frac{1}{(f'_p)^2} + r^2 \right) (g - f_p)' dr < 0 ,$$

in contradiction with the fact that f_p is a solution to the Euler-Lagrange equation (E-L). This verifies that f_p is a minimizer of I_p in \mathcal{C} . The fact that the minimizer is unique follows immediately from the strict convexity of I_p . \square

Remark. The logarithmic amplitude f_1 gives rise to the transformation

$$\Phi_1 = \left(\frac{3|x| + \sqrt{9|x|^2 + 16(2-\epsilon)\left(\frac{1}{2}-\epsilon\right)}}{4(2-\epsilon)} \right) \frac{x}{|x|} .$$

We compute

$$I_1(f_1) = 2\pi \int_{\epsilon}^1 \left(\frac{1}{f'_1(r)} + r^2 f'_1(r) \right) dr = 2\pi \left(1 - \epsilon^2 + \frac{2}{3}(2\epsilon - 1)^2 \right) .$$

By comparison, the radial affine transformation

$$\Phi_{ra} = \left(\frac{|x| - 1}{2(1-\epsilon)} + 1 \right) \frac{x}{|x|} ,$$

with logarithmic amplitude

$$f_{ra}(r) = \log \left(\frac{r-1}{2(1-\epsilon)} + 1 \right) .$$

has

$$I_1(f_{ra}) = 2\pi \int_{\epsilon}^1 \left(\frac{1}{f'_{ra}(r)} + r^2 f'_{ra}(r) \right) dr = 2\pi \left(1 - \epsilon^2 + \ln 2 (2\epsilon - 1)^2 \right) \geq I_1(f_1) .$$

Equality occurs only when $\epsilon = \frac{1}{2}$ (when the associated transformations are both the identity). \square

Turning to maximum norm, we consider the minimization

$$\mathcal{I}_{\infty} = \inf_{f \in \mathcal{C}_{[\epsilon,1]}} \sup \left(\frac{1}{r f'(r)} + r f'(r) \right) .$$

We note that

$$\begin{aligned} \mathcal{I}_{\infty} &= \inf_{K>1} \left\{ \frac{1}{K} + K : \exists f \in \mathcal{C} \text{ with } \sup_{r \in [\epsilon,1]} \left\{ \frac{1}{r f'(r)} + r f'(r) \right\} \leq \frac{1}{K} + K \right\} \\ &\geq \inf_{K>1} \left\{ \frac{1}{K} + K : \exists f \in \mathcal{C} \text{ with } \frac{1}{K} |\log r| \leq |f(r)| \right\} \\ &\geq \inf \left\{ \frac{1}{K} + K : \frac{|\log \epsilon|}{\log 2} \leq K \right\} = \frac{\log 2}{|\log \epsilon|} + \frac{|\log \epsilon|}{\log 2} . \end{aligned}$$


Here we have used that, if $f \in \mathcal{C}$ and if $K > 1$, then

$$\begin{aligned} \frac{1}{r f'(r)} + r f'(r) \leq \frac{1}{K} + K \text{ in } (\epsilon, 1) &\implies \frac{1}{Kr} \leq f'(r) \leq \frac{K}{r} \text{ in } (\epsilon, 1) \\ &\implies \frac{1}{K} |\log r| \leq |f(r)| \leq K |\log r| \text{ in } (\epsilon, 1) . \end{aligned}$$

On the other hand, the function

$$(3.1) \quad f_\infty(r) = \frac{\log 2}{|\log \epsilon|} \log r$$

lies in \mathcal{C} , and has $I_\infty(f_\infty) = \frac{\log 2}{|\log \epsilon|} + \frac{|\log \epsilon|}{\log 2}$. It now follows immediately that f_∞ is a minimizer of I_∞ in \mathcal{C} . The following graph shows the logarithmic amplitudes f_{ra} (dashed orange line), $f_1, f_2, f_3, f_5, f_8, f_{13}$ and f_∞ (solid lines from red to blue), for $\epsilon = 1/100$.



4. OPTIMALITY OF RADIAL TRANSFORMS

We now return to the general, two dimensional case. By introducing $u = \psi$ and $V = -JD\theta$ in the formula

$$\text{trace } \Phi_* [I] (\Phi(x)) = \frac{|D\psi|^2 + |D\theta|^2}{\det(D\psi, D\theta)}(x),$$

we obtain

$$\text{trace } \Phi_* [I] (\Phi(x)) = \frac{|Du|^2 + |V|^2}{Du \cdot V}(x).$$

Similarly, by introducing $u = \theta$ and $V = JD\psi$, we obtain

$$\text{trace } \Phi_* [I] (\Phi(x)) = \frac{|Du|^2 + |V|^2}{Du \cdot V}(x).$$

We thus notice that the problem of minimizing

$$I_p(\Phi) = \int_{B_1 \setminus B_\epsilon} (\text{trace } \Phi_* [I])^p (\Phi(x)) dx$$

with respect to ψ given θ , and with respect to θ , given ψ merely differs by a change of the convex test set for u (essentially relating to boundary conditions). Let $\arg \in C^\infty(\overline{B_1} \setminus B_\epsilon; \mathbb{R}/2\pi\mathbb{Z})^3$ denote the standard argument function. We introduce the convex sets

$$\begin{aligned} \mathcal{C}_\theta &= C^{2,\alpha}(\overline{B_1} \setminus B_\epsilon; \mathbb{R}/2\pi\mathbb{Z}) \cap \{ u|_{C_1} = \arg \} \text{ and} \\ \mathcal{C}_\psi &= C^{2,\alpha}(\overline{B_1} \setminus B_\epsilon; \mathbb{R}) \cap \{ u|_{C_\epsilon} = -\log 2, u|_{C_1} = 0 \}, \end{aligned}$$

³The space $C^\infty(\overline{B_1} \setminus B_\epsilon; \mathbb{R}/2\pi\mathbb{Z})$ is defined as $\{ u \in C^1(\overline{B_1} \setminus B_\epsilon; \mathbb{R}/2\pi\mathbb{Z}) : Du \in C^\infty(\overline{B_1} \setminus B_\epsilon; \mathbb{R}^2) \}$. Similarly $C^{2,\alpha}(\overline{B_1} \setminus B_\epsilon; \mathbb{R}/2\pi\mathbb{Z}) = \{ u \in C^1(\overline{B_1} \setminus B_\epsilon; \mathbb{R}/2\pi\mathbb{Z}) : Du \in C^{1,\alpha}(\overline{B_1} \setminus B_\epsilon; \mathbb{R}^2) \}$

for some fixed $\alpha > 0$.

Proposition 4. *Given $\mathcal{C} = \mathcal{C}_\psi$ and a fixed $V \in C^0(\overline{B_1} \setminus B_\epsilon; \mathbb{R}^2)$, or $\mathcal{C} = \mathcal{C}_\theta$ and a fixed $V \in C^0(\overline{B_1} \setminus B_\epsilon; \mathbb{R}^2)$, and given $n \geq 1$, we introduce*

$$\mathcal{C}_n = \left\{ u \in \mathcal{C} : Du \cdot V \geq \frac{1}{n} \text{ and } \|u\|_{C^{2,\alpha}(\overline{B_1} \setminus B_\epsilon)} \leq n \right\}.$$

Suppose $\mathcal{C}_{N_0} \neq \emptyset$, for some $N_0 \geq 1$. Given any $1 \leq p < \infty$, the functional $F_p : \mathcal{C}_n \rightarrow \mathbb{R}$, $n \geq N_0$, defined by

$$u \rightarrow F_p(u) = \int_{B_1 \setminus B_\epsilon} \left(\frac{|Du|^2 + |V|^2}{Du \cdot V} \right)^p dx$$

is strictly convex, continuous, and attains its infimum on \mathcal{C}_n at a unique minimizer. If the unique minimizer, u , lies in $\text{int}(\mathcal{C}_n)$ ⁴, then it satisfies the associated Euler-Lagrange equation

$$(4.1) \quad \text{div} \left(\left(\frac{|Du|^2 + |V|^2}{Du \cdot V} \right)^{p-1} \left(\frac{2Du}{Du \cdot V} - \frac{|Du|^2 + |V|^2}{(Du \cdot V)^2} V \right) \right) = 0 \text{ in } B_1 \setminus B_\epsilon,$$

and in the case $\mathcal{C} = \mathcal{C}_\theta$, the additional boundary condition

$$(4.2) \quad \left(\frac{2Du}{Du \cdot V} - \frac{|Du|^2 + |V|^2}{(Du \cdot V)^2} V \right) \cdot \frac{x}{|x|} = 0 \text{ on } C_\epsilon.$$

Conversely, if there exists a solution to equation (4.1) (and equation (4.2) in case $\mathcal{C} = \mathcal{C}_\theta$) which lies in $\mathcal{C} \cap \{Du \cdot V > 0 \text{ on } \overline{B_1} \setminus B_\epsilon\}$, then, for some $N \geq 1$, this is the unique minimizer of F_p in \mathcal{C}_n , for any $n \geq N$. Consequently this u is also the unique minimizer of F_p in $\mathcal{C} \cap \{Du \cdot V > 0 \text{ on } \overline{B_1} \setminus B_\epsilon\}$.

For the proof of Proposition 4 we shall need the following lemma.

Lemma 5. *For any $1 \leq p < \infty$, and any $A > 0$, the function $G_p[A] : (0, \infty) \times \mathbb{R} \rightarrow \mathbb{R}_+$, given by*

$$(x, y) \rightarrow \left(\frac{A}{x} + \frac{x}{A} + \frac{x}{A} \left(\frac{y}{x} \right)^2 \right)^p$$

is convex. Furthermore,

$$G_p[A](x, y) - \frac{2A^4}{(A^2 + M^2)^3} (x^2 + y^2)$$

is convex on $B_M = \{(x, y) : x^2 + y^2 < M^2\}$.

Proof. The function $x \rightarrow \frac{A}{x} + \frac{x}{A}$ is strictly convex and positive valued on $(0, \infty) \times \mathbb{R}$. The map $(x, y) \rightarrow \frac{1}{A} \frac{y^2}{x}$ is convex and positive on $(0, \infty) \times \mathbb{R}$. Indeed, its Hessian has eigenvalues 0 and $\frac{2}{A} \frac{x^2 + y^2}{x^3}$. The sum of two convex (and positive valued) functions is convex (and positive valued), and the composition of it with $z \rightarrow z^p$, a monotonically increasing and convex function on $(0, \infty)$, results in a convex (positive valued) function.

To establish the second assertion, we compute lower bounds for $D^2 G_p[A]$. It is a fact that the lowest eigenvalue of a symmetric positive definite matrix is bounded

⁴The interior is formed relative to \mathcal{C}_ψ or \mathcal{C}_θ with the $C^{2,\alpha}$ topology, respectively.

below by the quotient of the determinant over the trace. We compute that for $p \geq 1$,

$$\frac{\det(D^2 G_p[A])}{\text{tr}(D^2 C_p[A])} > \frac{4p}{p+1} G_p[A] \frac{A^4}{(A^2 + x^2 + y^2)^3} \geq 4 \frac{A^4}{(A^2 + x^2 + y^2)^3}.$$

In particular, on the ball $B_M = \{(x, y) : x^2 + y^2 < M^2\}$ we have

$$D^2 G_p[A](x, y) > \frac{4A^4}{(A^2 + M^2)^3} I.$$

This immediately leads to the second assertion of the lemma. \square

We are now ready for the proof of Proposition 4.

Proof. Given $u \in \mathcal{C}_n$, we define

$$P_V(Du) = Du \cdot \frac{V}{|V|}, \text{ and } P_{V^\perp}(Du) = Du \cdot \frac{JV}{|V|}.$$

Then

$$\begin{aligned} \left(\frac{|Du|^2 + |V|^2}{Du \cdot V} \right)^p &= \left(\frac{|V|}{P_V(Du)} + \frac{P_V(Du)}{|V|} + \frac{P_V(Du)}{|V|} \left(\frac{P_{V^\perp}(Du)}{P_V(Du)} \right)^2 \right)^p \\ &= G_p[|V|](P_V(Du), P_{V^\perp}(Du)). \end{aligned}$$

Note that $\mathcal{C}_{N_0} \neq \emptyset$ implies $\inf |V| > 0$. On \mathcal{C}_n , $|P_V(Du)|^2 + |P_{V^\perp}(Du)|^2 \leq n^2$, and therefore for any $u, v \in \mathcal{C}_n$, $n \geq N_0$, and any $\tau \in [0, 1]$

$$\begin{aligned} &G_p[|V|](P_V(D(\tau u + (1-\tau)v)), P_{V^\perp}(D(\tau u + (1-\tau)v))) \\ &\leq \tau G_p[|V|](P_V(Du), P_{V^\perp}(Du)) + (1-\tau) G_p[|V|](P_V(Dv), P_{V^\perp}(Dv)) \\ &\quad - \tau(1-\tau) K |D(u-v)|^2, \end{aligned}$$

with

$$K = \frac{2 \inf |V|^4}{(n^2 + \sup |V|^2)^3} > 0.$$

For $u, v \in \mathcal{C}_n$, and $\tau \in [0, 1]$, we thus get

$$F_p(\tau u + (1-\tau)v) \leq \tau F_p(u) + (1-\tau) F_p(v) - \tau(1-\tau) K \int_{B_1 \setminus B_\epsilon} |D(u-v)|^2 dx,$$

and so F_p is strictly convex on \mathcal{C}_n . In regards to continuity, let u_m be a sequence in \mathcal{C}_n with $u_m \rightarrow u$ in the C^1 topology. Then the functions

$$x \rightarrow G_p[|V|](P_V(Du_m), P_{V^\perp}(Du_m))(x)$$

are measurable, non negative, uniformly bounded, and converge pointwise to the function

$$x \rightarrow G_p[|V|](P_V(Du), P_{V^\perp}(Du))(x).$$

Thanks to the Lebesgue Dominated Convergence Theorem, this implies

$$\lim F_p(u_m) = F_p(u).$$

Since \mathcal{C}_n is compact with respect to the C^1 topology, the C^1 continuity of F_p implies the existence of a minimizer. The convexity of \mathcal{C}_n and the strict convexity of F_p

yields the uniqueness of the minimizer. A computation shows that for any $u \in \mathcal{C}_n$, F_p is Gâteaux-differentiable at u , and its differential is given by

$$\begin{aligned} & \langle DF_p(u), h \rangle \\ &= \int_{B_1 \setminus B_\epsilon} p \left(\frac{|Du|^2 + |V|^2}{Du \cdot V} \right)^{p-1} \left(\frac{2Du}{Du \cdot V} - \frac{|Du|^2 + |V|^2}{(Du \cdot V)^2} V \right) \cdot Dh \, dx, \end{aligned}$$

for $h \in C^1$. Note that $u \in \mathcal{C}_n$ is the unique minimizer if and only if for all $v \in \mathcal{C}_n$ there holds

$$(4.3) \quad \langle DF_p(u), v - u \rangle \geq 0.$$

If the minimizer lies in the interior of \mathcal{C}_n , equation (4.3) implies

$$\langle DF_p(u), h \rangle = 0$$

for all $h \in C^{2,\alpha} \cap \{h = 0 \text{ on } C_\epsilon \text{ and } C_1\}$, if $\mathcal{C} = \mathcal{C}_\psi$, and for all $h \in C^{2,\alpha} \cap \{h = 0 \text{ on } C_1\}$, if $\mathcal{C} = \mathcal{C}_\theta$; in other words, u satisfies the Euler-Lagrange equation (4.1) (or equation (4.1) and equation (4.2) when $\mathcal{C} = \mathcal{C}_\theta$). Conversely, if $w \in \mathcal{C} \cap \{Du \cdot V > 0 \text{ on } \overline{B_1} \setminus B_\epsilon\}$ satisfies equation (4.1) (and equation (4.2) if $\mathcal{C} = \mathcal{C}_\theta$), then, for some N , it lies in \mathcal{C}_n for all $n \geq N$, and it satisfies $\langle DF_p(w), v - w \rangle = 0$ (in particular ≥ 0) for all $v \in \mathcal{C}_n$; w is thus the unique minimizer of F_p in \mathcal{C}_n for any $n \geq N$. It follows immediately that w is a minimizer of F_p in $\mathcal{C} \cap \{Du \cdot V > 0 \text{ on } \overline{B_1} \setminus B_\epsilon\}$. The uniqueness of this minimizer follows from the strict convexity of F_p on \mathcal{C}_n for any n . \square

Corollary 6. *A global $C^{2,\alpha}$ minimizer (ψ, θ) of I_p , subject to $\psi = -\log 2$ at $|x| = \epsilon$, $\psi = 0$ and $\theta = \arg$ at $|x| = 1$, and $\det(D\psi, D\theta) > 0$ on $\overline{B_1} \setminus B_\epsilon$, satisfies*

$$\operatorname{div} \left(\left(\frac{|D\psi|^2 + |D\theta|^2}{\det(D\psi, D\theta)} \right)^p \left(\frac{2D\psi}{|D\psi|^2 + |D\theta|^2} + \frac{JD\theta}{\det(D\psi, D\theta)} \right) \right) = 0,$$

and

$$\operatorname{div} \left(\left(\frac{|D\psi|^2 + |D\theta|^2}{\det(D\psi, D\theta)} \right)^p \left(\frac{2D\theta}{|D\psi|^2 + |D\theta|^2} - \frac{JD\psi}{\det(D\psi, D\theta)} \right) \right) = 0.$$

Furthermore,

$$\left(\frac{|D\psi|^2 + |D\theta|^2}{\det(D\psi, D\theta)} JD\psi - 2D\theta \right) \cdot \frac{x}{|x|} = 0 \text{ on } \{|x| = \epsilon\}.$$

Proof. The ψ component of this global minimizer automatically lies in $\operatorname{int}(\mathcal{C}_n)$ with $\mathcal{C} = \mathcal{C}_\psi$ and $V = -JD\theta$ for some n , and it is a minimizer of F_p in \mathcal{C}_n . The first equation of this corollary is now simply the Euler-Lagrange equation (4.1) for such a minimizer. Similarly, the θ component of this global minimizer lies in $\operatorname{int}(\mathcal{C}_n)$ with $\mathcal{C} = \mathcal{C}_\theta$ and $V = JD\psi$ for some n , and is a minimizer of F_p in \mathcal{C}_n . The two last equations of this corollary are simply the Euler-Lagrange equation (4.1) and the boundary condition (4.2) satisfied by such a minimizer. \square

Corollary 7. *Let f_p be the function introduced in Proposition 3. The transformation $x \rightarrow f_p(|x|) \frac{x}{|x|}$, or rather the function pair $(f_p(|x|), \arg(x))$ satisfies the three Euler-Lagrange equations from Corollary 6. As a consequence*

$$(4.4) \quad I_p(f_p(|x|) \frac{x}{|x|}) \leq I_p(\psi(x) \frac{x}{|x|}),$$




FIGURE 4.1. Illustration of the conclusions of Corollary 7.

for any $\psi \in \mathcal{C}_\psi \cap \{D\psi(x) \cdot \frac{x}{|x|} > 0 \text{ on } \overline{B_1} \setminus B_\epsilon\}$. The last two Euler-Lagrange equations from Corollary 6 are actually satisfied by any pair $(f(|x|), \arg(x))$, with $f \in \{f \in C^{2,\alpha}([\epsilon, 1]) : f' > 0, f(\epsilon) = -\log 2, f(1) = 0\}$. As a consequence

$$(4.5) \quad I_p(f_p(|x|)\frac{x}{|x|}) \leq I_p(f(|x|)\frac{x}{|x|}) \leq I_p(f(|x|)\phi(x)),$$


for any $\phi(x) = (\cos(\theta(x)), \sin(\theta(x))^t)$, with $\theta \in \mathcal{C}_\theta \cap \{D\theta \cdot J\frac{x}{|x|} > 0 \text{ on } \overline{B_1} \setminus B_\epsilon\}$ and any $f \in \{f \in C^{2,\alpha}([\epsilon, 1]) : f' > 0, f(\epsilon) = -\log 2, f(1) = 0\}$.

Proof. Direct calculations verify that the first Euler-Lagrange equation from Corollary 6 is satisfied by $(f_p(|x|), \arg(x))$, and that the last two Euler-Lagrange equations from Corollary 6 are satisfied by any pair $(f(|x|), \arg(x))$, with $f \in \{f \in C^{2,\alpha}([\epsilon, 1]) : f' > 0, f(\epsilon) = -\log 2, f(1) = 0\}$. The inequality (4.4) now follows immediately from the last statement in Proposition 4 in the case $\mathcal{C} = \mathcal{C}_\psi$ and $V = -JD \arg(x) = \frac{1}{|x|}\frac{x}{|x|}$. The first inequality in (4.5) is a direct consequence of (4.4). The second inequality follows from the last statement in Proposition 4 in the case $\mathcal{C} = \mathcal{C}_\theta$ and $V = JDf(|x|) = f'(|x|)J\frac{x}{|x|}$. \square

5. OPTIMAL CLOAKS FOR SIMPLY CONNECTED DOMAINS

So far our study has focused on the situation where the cloaks are constructed from diffeomorphisms of the classical annulus $\overline{B_1} \setminus B_\epsilon$ to the classical annulus $\overline{B_1} \setminus B_{\frac{1}{2}}$, and the corresponding push-forwards of the identity matrix. In a more general setting, one could consider instead three simply connected domains, $\omega_\epsilon \subset \omega_{\frac{1}{2}} \subset \Omega$ containing the origin (where ω_ϵ is comparable to B_ϵ) and a bijective diffeomorphism $\Psi_\epsilon : \overline{\Omega} \setminus \omega_\epsilon \rightarrow \overline{\Omega} \setminus \omega_{\frac{1}{2}}$, such that $\Psi_\epsilon = Id$ on $\partial\Omega$ and $\Psi_\epsilon(\partial\omega_\epsilon) = \partial\omega_{\frac{1}{2}}$. As before, the material parameters of the cloak would be the push-forward of I by Ψ_ϵ . Any smooth globally minimizing transformation would still satisfy the Euler-Lagrange equations of Corollary 6, if we continue to use the energy I_p .

The goal of this section is to show that for general geometries one should (naturally) not expect the optimal transformations to be radial. As we demonstrate this, we also derive a process for the construction of optimal transformations (based on a slightly revised energy). Suppose Ω is a bounded, smooth, simply connected domain containing the origin. Due to the Riemann Mapping Theorem, there exists

FIGURE 5.1. Cloaking by mapping where $\Omega = \sinh(B_1)$, with $\epsilon = 1/10$.

a unique (complex) analytic map Ψ such that $\Psi(0) = 0$, $D\Psi(0) = aI$ for some $a > 0$ and Ψ is a one-to-one mapping from $\overline{\Omega}$ onto $\overline{B_1}$. By the maximum modulus principle $\min\{|x| : x \in \overline{\Omega}\} \leq 1/a \leq \max\{|x| : x \in \overline{\Omega}\}$. Set $\omega_\epsilon = \Psi^{-1}(B_\epsilon)$, and $\omega_{\frac{1}{2}} = \Psi^{-1}(B_{\frac{1}{2}})$. By construction, $0 \in \omega_\epsilon \subset \omega_{\frac{1}{2}} \subset \Omega$. Provided ϵ is small enough, ω_ϵ is approximately $B_{\frac{\epsilon}{a}}$, in the sense that

$$\forall x \in C_\epsilon \quad \left| \Psi^{-1}(x) - \frac{x}{a} \right| \leq \frac{1}{2} \frac{\max_{\overline{B_{1/2}}} |D^2\Psi^{-1}|}{a} \epsilon^2 .$$

Given $\Phi_\epsilon \in C^1(\overline{B_1} \setminus B_\epsilon; \overline{B_1} \setminus B_{\frac{1}{2}})$ a (possibly optimal) bijective diffeomorphism with $\Phi_\epsilon|_{C_1} = Id$, and $\Phi_\epsilon(C_\epsilon) = C_{\frac{1}{2}}$, we define

$$(5.1) \quad \Psi_\epsilon := \Psi^{-1} \circ \Phi_\epsilon \circ \Psi .$$

Figure 5.1 shows some of the ‘‘rays’’ of the map Ψ_ϵ (Φ_ϵ being radial) in the case $\Psi^{-1} = \sinh$, $\Omega = \sinh(B_1)$, $\omega_{\frac{1}{2}} = \sinh(B_{\frac{1}{2}})$ and $\omega_\epsilon = \sinh(B_\epsilon)$. The green curves on the left are mapped to proper subsets of themselves, shown as red curves on the right. Clearly the transformation Ψ_ϵ is no longer radial.

For any $x \in \partial\Omega$, $\Psi(x)$ lies on C_1 , and thus $\Phi_\epsilon \circ \Psi(x) = \Psi(x)$. It follows that $\Psi_\epsilon(x) = x$, in other words: $\Psi_\epsilon = Id$ on $\partial\Omega$. Similarly, we obtain that $\Psi_\epsilon(\partial\omega_\epsilon) = \partial\omega_{\frac{1}{2}}$. $(\Psi_\epsilon)_*[I]$ therefore produces an approximate cloak (with same approximate invisibility as that of $(\Phi_\epsilon)_*[I]$). From composition of transformations we obtain

$$(\Psi_\epsilon)_*[I] = (\Psi^{-1})_*[(\Phi_\epsilon)_*[\Psi_*[I]]] .$$

Lemma 8. *There holds*

$$\text{trace}(\Psi_\epsilon)_*[I] = (\text{trace}(\Phi_\epsilon)_*[I]) \circ \Psi .$$

Proof. Since Ψ is conformal, $D\Psi = \gamma Q$ with γ a positive scalar and Q an orthogonal matrix. We are in 2d, and so this implies

$$\Psi_* [I] (y) = \frac{(D\Psi)(D\Psi)^T}{|\det D\Psi|} \circ \Psi^{-1} (y) = I .$$

Similarly,

$$\begin{aligned} (\Psi^{-1})_* [A] (x) &= \frac{(D\Psi^{-1}) A (D\Psi^{-1})^T}{|\det D\Psi^{-1}|} \circ \Psi (x) \\ &= Q^T (x) A (\Psi (x)) Q (x) , \end{aligned}$$

where we have used that $(D\Psi^{-1})(\Psi(x)) = (D\Psi)^{-1}(x) = \frac{1}{\gamma}Q^T(x)$. In summary, we conclude that $(\Psi_\epsilon)_* [I]$ is given by the formula

$$(\Psi_\epsilon)_* [I] (x) = Q^T (x) (\Phi_\epsilon)_* [I] (\Psi (x)) Q (x) ,$$

and the statement about the traces follows. \square

If Φ_ϵ is a transformation which minimizes the anisotropy of $(\Phi_\epsilon)_* [I]$, using the measure I_p for some $1 \leq p < \infty$, then it follows immediately from Lemma 8 above that Ψ_ϵ minimizes anisotropy of $(\Psi_\epsilon)_* [I]$, using the slightly modified measure

$$\tilde{I}_p (\Psi_\epsilon) = \int_{\Omega \setminus \omega_\epsilon} (\text{trace}(\Psi_\epsilon)_* [I])^p (\Psi_\epsilon(x)) |\det \Psi(x)| dx .$$

A similar statement holds for $p = \infty$. In that case there is no change in the measure of anisotropy.

Acknowledgements The research of MSV was partially supported by NSF Grant DMS-12-11330. Part of this work was carried out while MSV was visiting the University of Copenhagen and the Danish Technical University. This visit was made possible through support from the Nordea Foundation and the Otto Mønsted Foundation. This study also contributes to the IdEx Université de Paris ANR-18-IDEX-0001.

REFERENCES

1. H. Ammari, H. Kang, H. Lee, and M. Lim, *Enhancement of near cloaking using generalized polarization tensors vanishing structures. I: The conductivity problem*, Commun. Math. Phys. **317** (2013), no. 1, 253–266.
2. A. Greenleaf, Y. Kurylev, M. Lassas, and G. Uhlmann, *Invisibility and inverse problems*, Bull. Am. Math. Soc., New Ser. **46** (2009), no. 1, 55–97.
3. A. Greenleaf, M. Lassas, and G. Uhlmann, *On nonuniqueness for Calderón’s inverse problem*, Math. Res. Lett. **10** (2003), no. 5-6, 685–693.
4. R. Griesmaier and M. S. Vogelius, *Enhanced approximate cloaking by optimal change of variables*, Inverse Probl. **30** (2014), no. 3, 17, Id/No 035014.
5. H. Heumann and M. S. Vogelius, *Analysis of an enhanced approximate cloaking scheme for the conductivity problem*, Asymp. Anal. **87** (2014), 223–246.
6. R. V. Kohn, H. Shen, M. S. Vogelius, and M. I. Weinstein, *Cloaking via change of variables in electric impedance tomography*, Inverse Probl. **24** (2008), no. 1, 21, Id/No 015016.
7. G. W. Milton and N-A. P. Nicorovici, *On the cloaking effects associated with anomalous localized resonance*, Proceedings: Mathematical, Physical and Engineering Sciences **462** (2006), no. 2074, 3027–3059.
8. J. B. Pendry, D. Schurig, and D. R. Smith, *Controlling electromagnetic fields*, Science **312** (2006), no. 5781, 1780–1782.

UNIVERSITÉ DE PARIS AND SORBONNE UNIVERSITÉ, CNRS, LABORATOIRE JACQUES-LOUIS
LIONS (LJLL), F-75006 PARIS, FRANCE

Email address: `yves.capdeboscq@u-paris.fr`

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, NEW BRUNSWICK, NEW JERSEY
08901, USA

Email address: `vogelius@math.rutgers.edu`