



**HAL**  
open science

## Pyridoxal kinase and poly(ADP-ribose) affect the immune microenvironment of locally advanced cancers

Adrien Joseph, Juncheng Pan, Judith Michels, Guido Kroemer, Maria Castedo

► **To cite this version:**

Adrien Joseph, Juncheng Pan, Judith Michels, Guido Kroemer, Maria Castedo. Pyridoxal kinase and poly(ADP-ribose) affect the immune microenvironment of locally advanced cancers. *OncoImmunology*, 2021, 10 (1), pp.1950954. 10.1080/2162402x.2021.1950954 . hal-03342400


**HAL Id: hal-03342400**

**<https://hal.sorbonne-universite.fr/hal-03342400v1>**

Submitted on 13 Sep 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## Pyridoxal kinase and poly(ADP-ribose) affect the immune microenvironment of locally advanced cancers

Adrien Joseph, Juncheng Pan, Judith Michels, Guido Kroemer & Maria Castedo

To cite this article: Adrien Joseph, Juncheng Pan, Judith Michels, Guido Kroemer & Maria Castedo (2021) Pyridoxal kinase and poly(ADP-ribose) affect the immune microenvironment of locally advanced cancers, *Oncolmmunology*, 10:11, 1950954, DOI: [10.1080/2162402X.2021.1950954](https://doi.org/10.1080/2162402X.2021.1950954)

To link to this article: <https://doi.org/10.1080/2162402X.2021.1950954>



© 2021 The Author(s). Published with license by Taylor & Francis Group, LLC.



Published online: 08 Jul 2021.



Submit your article to this journal [↗](#)



Article views: 308





View related articles [↗](#)



View Crossmark data [↗](#)

## Pyridoxal kinase and poly(ADP-ribose) affect the immune microenvironment of locally advanced cancers

Adrien Joseph <sup>a,b,c</sup>, Juncheng Pan<sup>a,b,c</sup>, Judith Michels<sup>d</sup>, Guido Kroemer <sup>a,b,e,f,g,h</sup>, and Maria Castedo<sup>a,b</sup>

<sup>a</sup>Equipe 11 labellisée par la Ligue contre le Cancer, Université de Paris, Sorbonne Université, INSERM U1138, Centre de Recherche des Cordeliers, Paris, France; <sup>b</sup>Metabolomics and Cell Biology Platforms, Gustave Roussy Cancer Campus, Villejuif, France; <sup>c</sup>Faculté de médecine, Université de Paris Saclay, Kremlin Bicêtre, France; <sup>d</sup>Department of Medical Oncology, Gustave Roussy Cancer Campus, Villejuif, France; <sup>e</sup>Paris Descartes University, Sorbonne Paris Cité, Paris, France; <sup>f</sup>Pôle de Biologie, Hôpital Européen Georges Pompidou, Assistance Publique-Hôpitaux de Paris, Paris, France; <sup>g</sup>Suzhou Institute for Systems Medicine, Chinese Academy of Medical Sciences, Suzhou, China; <sup>h</sup>Karolinska Institute, Department of Women's and Children's Health, Karolinska University Hospital, Stockholm, Sweden

### ABSTRACT

Malignant cells adapt to the hostile tumor microenvironment by escaping from, or actively suppressing, anticancer immune responses. In the past, we reported that reduced synthesis of active vitamin B6 (due to downregulation of pyridoxal kinase) or overactivation of poly(ADP-ribose) polymerase confers resistance to chemotherapy with cisplatin. Recently, we found that these prognostically adverse alterations in oncometabolism also correlate with the rarefaction of immune effectors in the tumor bed.

### ARTICLE HISTORY

Received 28 June 2021  
Revised 29 June 2021  
Accepted 29 June 2021

### KEYWORDS

Tumor microenvironment; immune escape; metabolism; vitamin B; immunotherapy; CD8+ T cells; dendritic cells; survival; cervical cancer; non-small cell lung cancer

The immune system is known to play a major role in cancer immunosurveillance and response to chemotherapy.<sup>1,2</sup> Recently, immune checkpoint blockade therapy has created a paradigm shift in antineoplastic treatments, highlighting the importance of focusing on the tumor microenvironment. Cancer cell-intrinsic properties, and in particular metabolic features, are known to impact the tumor immune environment.

In the past, we reported that chemotherapy with cisplatin leads to the down-regulation of vitamin B6 activating enzyme pyridoxal kinase (PDXK)<sup>3</sup> and the up-regulation of the enzymatic activity of poly(ADP-ribose) (PAR) polymerase 1 (PARP1)<sup>4</sup> in tumor cells, and that these metabolic alterations affect the prognosis of non-small cell lung cancer (NSCLC) patients.<sup>5,6</sup> The activity of PARP1 requires a supply of nicotinamide (vitamin B3) adenine dinucleotide. Thus, both of these enzymes participate in vitamin B metabolism.

Recently, we dove further into the mechanisms underlying the interplay between cancer metabolism, immunosurveillance and prognosis,<sup>7</sup> and investigated the role of these two vitamin B metabolism-related markers on the immune microenvironment.


In locally advanced cervical carcinoma (LACC) patients, for whom cisplatin is still part of the standard of care, we observed that tumor infiltration by CD8<sup>+</sup> T lymphocytes and DC-LAMP<sup>+</sup> cells were both associated with an improved prognosis. Tumors scarcely infiltrated by both CD8<sup>+</sup> and DC-LAMP<sup>+</sup> cells had a particularly dismal prognosis. We also described

a negative correlation between CD8 density and PAR expression and a positive correlation between DC-LAMP density and PDXK expression in LACC.

These findings were reproduced in a cohort of NSCLC patients, in which the detection of high levels of CD8<sup>+</sup> or DC-LAMP<sup>+</sup> cells, were independent prognostic factors, and in which we observed similar associations between PAR and CD8 infiltration (negative correlation) and between PDXK and DC-LAMP (positive correlation). In accordance with these findings, a PAR<sup>high</sup>/PDXK<sup>low</sup> status was associated with dismal NSCLC prognosis.

In a final step, we used a heterotopic mouse model of NSCLC, in which we injected Lewis lung cancer (LLC) cells subcutaneously into the flank of C57BL/6 mice. These cells had been previously exposed to increasing concentrations of cisplatin for several weeks and exhibited cisplatin resistance, together with enhanced PARP1 activity, as demonstrated by upregulation of PAR in immunoblot analyses. Immunofluorescence cytometry revealed that tumors formed by PAR<sup>high</sup> LLC cells exhibited a reduced infiltration by CD8<sup>+</sup> T lymphocytes as compared to tumors formed by parental LLC cells. In addition, PAR<sup>high</sup> tumors were less infiltrated by subpopulations of myeloid cells (CD45<sup>+</sup>CD11b<sup>+</sup>Ly6G<sup>-</sup>Ly6C<sup>low/intermediate</sup>) and activated dendritic cells (CD45<sup>+</sup>CD11c<sup>+</sup>MHCII<sup>+</sup>, non-significant trend).

In essence, in our recent article, we show that two enzymes, PDXK and PARP1, both implicated in the vitamin B metabolism, affect the immune infiltrate in two different malignant diseases, LACC and NSCLC. Our NSCLC mouse



**Figure 1.** The interplay between oncometabolism, immune infiltrate and prognosis. In non-small cell lung cancer (NSCLC) and locally advanced cervical carcinoma (LACC), PAR level correlates negatively with CD8 T cells infiltration, whereas PDXK level correlates positively with DC-LAMP cells infiltration. Infiltration by each of these cell types is associated with a good prognosis in NSCLC and LACC, whereas PAR<sup>high</sup>/PDXK<sup>low</sup> status indicates dismal prognosis. Cisplatin-resistant LLC cells, which exhibit high level of poly (ADP- ribose), form tumors scarcely infiltrated by CD8<sup>+</sup> T lymphocytes after injection in C57BL/6 mice. Items from this figure are provided for illustrative purposes only.

model supports a causal relationship between these metabolic features and immune tonus, demonstrating that the upregulation of PARP1 activity required for cisplatin resistance leads to a rarefaction of CD8<sup>+</sup> T cell infiltration, exactly as this is observed in cancer patients. Based on these results, we hypothesize that PARP1 and PDXK activity within tumor cells can modulate the tumor immune environment. Thus, metabolic alterations of cancer cells associated with cisplatin resistance may have an impact on anticancer immunosurveillance (Figure 1).

These findings potentially pave the way for therapeutic interventions. Thus, we previously showed in an orthotopic mouse model of NSCLC that the addition of pyridoxine (phosphorylated by PDXK) to standard cisplatin treatment exhibited hyperadditive therapeutic effects only in the context of an intact immune system.<sup>8</sup> Additionally, exogenous supply of nicotinamide adenine dinucleotide (NAD), which is consumed by overactivated PARP1, potently stimulates immunosurveillance in the context of hormone-induced breast cancer.<sup>9</sup> PARP inhibitors are currently being tested in association with immune checkpoint inhibitors (e.g. NCT02571725 and NCT03598270). Our findings, along with others showing an increased infiltration of CD8<sup>+</sup> T cells after PARP inhibition,<sup>10</sup> support the hypothesis that PARP1 activity and immune infiltration are mechanistically linked, and provide a rationale for such combination therapies.

To conclude, metabolic features of cancer cells, such as PDXK and PAR levels, can impact the immune infiltrate, thereby affecting the clinical course of LACC and NSCLC. Although the interplay between tumor metabolism, immune infiltrate and prognosis remains to be explored and the detailed mechanisms underlying these findings remain to be

discovered, we believe that this area of research has the potential to elucidate new patient-relevant facets of the cancer-immune dialogue.

### Disclosure statement

This study contributes to the IdEx Université de Paris ANR-18-IDEX-0001.

### Funding

GK is supported by the Ligue contre le Cancer (équipe labellisée); Agence Nationale de la Recherche (ANR) – Projets blancs; ANR under the frame of E-Rare-2, the ERA-Net for Research on Rare Diseases; AMMICa US23/CNRS UMS3655; Association pour la recherche sur le cancer (ARC); Association “Ruban Rose”; Cancéropôle Ile-de-France; Chancellerie des universités de Paris (Legs Poix), Fondation pour la Recherche Médicale (FRM); a donation by Elior; European Research Area Network on Cardiovascular Diseases (ERA-CVD, MINOTAUR); Gustave Roussy Odyssey, the European Union Horizon 2020 Project Oncobiome; Fondation Carrefour; High-end Foreign Expert Program in China [GDW20171100085], Institut National du Cancer (INCa); Inserm (HTE); Institut Universitaire de France; LeDucq Foundation; the LabEx Immuno-Oncology (ANR-18-IDEX-0001); the RHU Torino Lumière; the Seerave Foundation; the SIRIC Stratified Oncology Cell DNA Repair and Tumor Immune Elimination (SOCRATE); and the SIRIC Cancer Research and Personalized Medicine (CARPEM). AJ is supported by a grant from Fondation ARC pour la Recherche sur le Cancer. This study contributes to the IdEx Université de Paris ANR-18-IDEX-0001.

### ORCID

Adrien Joseph  <http://orcid.org/0000-0002-5278-8966>  
Guido Kroemer  <http://orcid.org/0000-0002-9334-4405>

## References

1. Galluzzi L, Humeau J, Buqué A, Zitvogel L, Kroemer G. Immunostimulation with chemotherapy in the era of immune checkpoint inhibitors. *Nat Rev Clin Oncol*. 2020 Dec;17(12):725–741.
2. Zitvogel L, Perreault C, Finn OJ, Kroemer G. Beneficial autoimmunity improves cancer prognosis. *Nat Rev Clin Oncol*. 2021 May 11. doi:10.1038/s41571-021-00508-x.
3. Galluzzi L, Vitale I, Senovilla L, Olaussen KA, Pinna G, Eisenberg T, Goubar A, Martins I, Michels J, Kratassiouk G, et al. Prognostic impact of vitamin B6 metabolism in lung cancer. *Cell Rep*. 2012 Aug 30;2(2):257–269. doi:10.1016/j.celrep.2012.06.017.
4. Michels J, Vitale I, Galluzzi L, Adam J, Olaussen KA, Kepp O, Senovilla L, Talhaoui I, Guegan J, Enot DP, et al. Cisplatin resistance associated with PARP hyperactivation. *Cancer Res*. 2013 Apr 1;73(7):2271–2280. doi:10.1158/0008-5472.CAN-12-3000.
5. Michels J, Adam J, Goubar A, Obrist F, Damotte D, Robin A, Alifano M, Vitale I, Olaussen KA, Girard P, et al. Negative prognostic value of high levels of intracellular poly(ADP-ribose) in non-small cell lung cancer. *Ann Oncol Off J Eur Soc Med Oncol*. 2015 Dec;26(12):2470–2477. doi:10.1093/annonc/mdv393.
6. Galluzzi L, Vacchelli E, Michels J, Garcia P, Kepp O, Senovilla L, Vitale I, Kroemer G. Effects of vitamin B6 metabolism on oncogenesis, tumor progression and therapeutic responses. *Oncogene*. 2013 Oct 17;32(42):4995–5004. doi:10.1038/onc.2012.623.
7. Joseph A, Juncheng P, Mondini M, Labaied N, Loi M, Adam J, Lafarge A, Astesana V, Obrist F, Klein C, et al. Metabolic features of cancer cells impact immunosurveillance. *J ImmunoTher Cancer*. 2021;9:e002362. doi:10.1136/jitc-2021-002362.
8. Aranda F, Bloy N, Pesquet J, Petit B, Chaba K, Sauvat A, Kepp O, Khadra N, Enot D, Pfirschke C, et al. Immune-dependent antineoplastic effects of cisplatin plus pyridoxine in non-small-cell lung cancer. *Oncogene*. 2015 Jun 4;34(23):3053–3062. doi:10.1038/onc.2014.234.
9. Buqué A, Bloy N, Perez-Lanzón M, Iribarren K, Humeau J, Pol JG, Levesque S, Mondragon L, Yamazaki T, Sato A, et al. Immunoprophylactic and immunotherapeutic control of hormone receptor-positive breast cancer. *Nat Commun*. 2020 Jul 30;11(1):3819. doi:10.1038/s41467-020-17644-0.
10. Sen T, Rodriguez BL, Chen L, Corte CMD, Morikawa N, Fujimoto J, Cristea S, Nguyen T, Diao L, Li L, et al. Targeting DNA damage response promotes antitumor immunity through STING-mediated T-cell activation in small cell lung cancer. *Cancer Discov*. 2019;9(5):646–661. doi:10.1158/2159-8290.CD-18-1020.