



HAL
open science

Toward Pediatric T Lymphoblastic Lymphoma Stratification Based on Minimal Disseminated Disease and NOTCH1/FBXW7 Status

Amélie Trinquand, Adriana Plesa, Chrystelle Abdo, Fabien Subtil, Nathalie Aladjidi, Charlotte Rigaud, Aurore Touzart, Ludovic Lhermitte, Arnaud Petit, Katell Michaux, et al.

► **To cite this version:**

Amélie Trinquand, Adriana Plesa, Chrystelle Abdo, Fabien Subtil, Nathalie Aladjidi, et al.. Toward Pediatric T Lymphoblastic Lymphoma Stratification Based on Minimal Disseminated Disease and NOTCH1/FBXW7 Status. *HemaSphere*, 2021, 5 (10), pp.e641. 10.1097/hs9.0000000000000641 . hal-03350319

HAL Id: hal-03350319

<https://hal.sorbonne-universite.fr/hal-03350319v1>

Submitted on 21 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Toward Pediatric T Lymphoblastic Lymphoma Stratification Based on Minimal Disseminated Disease and *NOTCH1*/*FBXW7* Status

Amélie Trinquand^{1,2}, Adriana Plesa³, Chrystelle Abdo¹, Fabien Subtil⁴, Nathalie Aladjidi⁵, Charlotte Rigaud⁶, Aurore Touzart¹, Ludovic Lhermitte¹, Arnaud Petit⁷, Katell Michaux⁸, Charlotte Jung⁸, Catherine Chassagne-Clement⁹, Vahid Asnafi¹, Yves Bertrand⁸, Nathalie Garnier⁸, Elizabeth Macintyre¹

Correspondence: Elizabeth Macintyre (elizabeth.macintyre@aphp.fr).

Abstract

While outcome for pediatric T lymphoblastic lymphoma (T-LL) has improved with acute leukemia-type therapy, survival after relapse remains rare. Few prognostic markers have been identified: *NOTCH1* and/or *FBXW7* (*N/F*) mutations identify good prognosis T-LL and high-level minimal disseminated disease (MDD) is reported to be of poor prognosis. We evaluated MDD and/or MRD status by 8-color flow cytometry and/or digital droplet PCR in 82 pediatric T-LL treated according to the EURO-LB02 prednisone reference arm. Both techniques gave identical results for values $\geq 0.1\%$, allowing compilation. Unlike historical studies, an MDD threshold of 1% had no prognostic significance. The 54% (42/78) of patients with MDD $\geq 0.1\%$ had a relatively favorable outcome (5-y overall survival [OS] 97.6% versus 80.6%, $P = 0.015$, 5-y event-free-survival [EFS] 95.2% versus 80.6%, $P = 0.049$). MDD lower than 0.1% had no impact in *N/F* mutated T-LL, but identified the *N/F* germline patient with a high risk of relapse. Combining oncogenetic and MDD status identified 86% of patients ($n = 49$) with an excellent outcome and 14% of *N/F* germline/MDD $< 0.1\%$ patients ($n = 8$) with poor prognosis (5y-OS 95.9% versus 37.5%, $P < 0.001$; 5y-EFS 93.9% versus 37.5%, $P < 0.001$). If confirmed by prospective studies, MDD and *N/F* mutational status would allow identification of a subset of patients who merit consideration for alternative front-line treatment.

Introduction

T-cell lymphoblastic lymphoma (T-LL) and T-cell acute lymphoblastic leukemias (T-ALL) are both characterized by the proliferation of malignant immature T-cell precursors but differ by the extent of bone marrow (BM) involvement, which is (arbitrarily) $< 25\%$, assessed by morphology, in T-LL. Despite the improvement of current therapy for children, achieving an event-free survival (EFS) at 5 years of 75%–85%,^{1–3} the survival rate of refractory or relapsed LL remains very poor, at

10%–30%.^{4,5} Relatively favorable results are obtained in the rare T-LL cases who achieve second remission and can undergo hematopoietic stem cell transplantation.⁶ The early identification of poor risk T-LL is thus mandatory.

Contrary to T-ALL,^{7,8} prognostic factors in T-LL are rare, partly due to its low incidence and difficulty in obtaining diagnostic material. Retrospective analyses identified the absence of *NOTCH1* and/or *FBXW7* (*N/F*) mutations and biallelic T-cell receptor-gamma (TRG) deletions to be associated to unfavorable outcome^{9–13} and loss of heterozygosity at chromosome 6q (LOH6q) to an increased relapse risk.^{11,14}

In T-ALL, one of the strongest prognostic factors is minimal residual disease (MRD).⁸ MRD analysis in peripheral blood (PB) and BM give comparable results.¹⁵ In T-LL, high-level minimal disseminated disease (MDD) at diagnosis and MRD positivity by multiparameter flow cytometry (MFC) and/or quantitative polymerase chain reaction (qPCR) for TR rearrangements identified high-risk patients,^{16–18} justifying intensification of MDD $\geq 1\%$ patients in the Children's Oncology Group (COG) AALL0434 trial.¹⁹ This trial showed no difference in outcome when comparing high-risk (MDD $> 1\%$) to standard-risk subjects who were assigned the same COG ABFM C-MTX regimen. The authors suggested that the prognostic impact of MDD $> 1\%$ may have been abrogated by the COG ABFM C-MTX regimen, with higher dose, continuous asparaginase exposure.

qPCR is a well-established tool for MRD detection, notably in leukemia, but has limitations in lymphoma because of the need for a reference standard curve, based on tumor-specific target serial dilutions from diagnostic DNA with known infiltration, usually assessed by MFC.²⁰ Evaluating tumour infiltration is more difficult in diagnostic tissue samples. Droplet digital PCR (ddPCR) is based on sample compartmentalization in single-ol

¹Université de Paris and Institut Necker-Enfants Malades, Laboratoire d'Onco-Hématologie, AP-HP Hôpital Necker-Enfants Malades, Paris, France

²National Children's Research Centre at Children's Health Ireland, Dublin, Ireland

³Laboratoire d'Hématologie, Hospices Civiles de Lyon, France

⁴Service de Biostatistiques des Hospices Civiles de Lyon, France

⁵Unité d'Hématologie et Cancérologie pédiatrique/CEREVANCE, CHU Bordeaux, France

⁶Oncologie Pédiatrique, Institut Gustave Roussy, Villejuif, France

⁷Service d'Hématologie et d'Oncologie pédiatrique, AP-HP, Hôpital Armand Trousseau, Sorbonne Université, Paris, France

⁸Institut d'Hématologie et d'Oncologie Pédiatrique, Hospices Civiles de Lyon, France

⁹Département de Biopathologie, Centre Leon Berard, Lyon, France

Supplemental digital content is available for this article.

Copyright © 2021 the Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the European Hematology Association. This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

HemaSphere (2021) 5:10(e641).

<http://dx.doi.org/10.1097/HS9.0000000000000641>.

Received: 16 June 2021 / Accepted: 12 August 2021

droplets. An independent end-point PCR reaction allows absolute quantification, using Poisson statistics. Studies indicate that ASO-specific ddPCR is particularly adapted for lymphomas^{21–23} but ddPCR has not, to date, been evaluated in T-LL.

In France, pediatric T-LL was treated on the EURO-LB02 trial until its premature arrest due to excessive toxicity on January 1, 2008²⁴ and from then onward, according to the reference prednisone arm, with 24 months treatment. The present study compared T cell Receptor (TR) ddPCR and MFC to assess the clinical impact of MDD at diagnosis in patients on this reference arm. Since the recently opened European multicenter prospective pediatric LBL2018 protocol (NCT04043494) stratifies patients on their *N/F* mutation status, we also evaluated the correlation between these oncogenetic markers and MDD on outcome. Unexpectedly, patients with MDD positivity above 0.1% in PB/BM have a relatively favorable prognosis, essentially within the poor prognosis *N/F* germline (*N/F*^{GL}) subgroup.

Patients and methods

Patients and samples

Between December 2004 and December 2007, pediatric T-LL patients were treated in France on the EURO-LB02 European trial, derived from the Berlin-Frankfurt-Münster BFM90 protocol, with CNS prophylaxis by high-dose systemic Methotrexate (HD-MTX) in the interim phase instead of prophylactic central nervous system irradiation (NCT00275106).²⁴ The protocol included 2 randomizations (R) for T-LL: prednisone versus dexamethasone during induction (R1) and the total treatment duration of 18 versus 24 months (R2). Since January 1, 2008, pediatric T-LLs were treated according to the reference prednisone arm, with 24 months treatment according to EICNHL group recommendations. Asparaginase treatment comprised E Coli native asparaginase (10,000 UI/m² per dose, 8 doses in induction, 4 doses during reinduction). All clinical data for French patients treated in Société Française des Cancers de l'Enfant (SFCE) centers were registered centrally (Lyon). As per the declaration of Helsinki, the Lyon ethics committee approved the study and the signed informed consent was obtained for all patients.

Centralized prospective evaluation of oncogenetics on infiltrated diagnostic material in Paris-Necker and PB and BM MDD and MRD by 8-color multiparameter FC (MFC) in Lyon was set up in 2008. No qPCR MDD/MRD was planned initially, but leftover cells from FC were stored as pellets in Lyon, whenever possible, and later transferred to the Paris molecular platform.

MDD/MRD assessment by MFC

From 2008 to 2017, fresh PB and BM samples were collected at diagnosis and PB at day (d) 33 (end of induction). MFC analysis was performed with an 8-color panel on a FACS CANTOII with DIVA software (BD Bioscience), as described.^{25,26} Antibodies used are described in Supplemental Digital Material and Methods, <http://links.lww.com/HS/A197>. To exclude residual erythrocytes, dead cells, debris, platelet aggregates and doublets, a live gate was adjusted on forward scatter (FSC)/SSC and CD45/SSC. Positive MDD/MRD was defined as a cluster of >10 cells expressing at least 2 leukemia aberrant immunophenotype (LAIP) and SSC characteristics identified at diagnosis, amongst at least 500,000 viable cells (Supplemental Digital Figure S1B, <http://links.lww.com/HS/A197>). MFC sensitivity of 0.01% normal cells was possible in virtually all samples.

MDD/MRD assessment by ddPCR

To identify patient-specific clonotypic markers, genomic DNA was extracted from diagnostic tissue or effusions from

62 patients using a QIAampDNA mini kit (Qiagen Co., Hilden, Germany), TR clonality (TRD, TRB, and TRG) was assessed by one-step Next-Generation Sequencing using 100 ng DNA, EuroClonality-NGS amplicon primers²⁷ and Vidjil software analysis.²⁸ At least one allele-specific (ASO) CDR3 primer, selected on the basis of maximal clonotype amplification and minimal nonspecific positivity in normal PBL DNA, was designed for each patient. Thirty-eight patients with fresh or cryopreserved MDD samples (34 BM and 18 PB) and 18 patients with d33 MRD frozen samples (6 BM and 17 PB) were analyzed retrospectively (Paris-Necker). Quantification method is described in Supplemental Digital Material and Methods, <http://links.lww.com/HS/A197>.

NOTCH1/FBXW7 mutations

The mutational screening of 61 tumour samples (effusions, mediastinal mass, and lymph node) with at least 20% infiltration was performed by capture NGS, as described.²⁹ The functionally silent SNVs of *NOTCH1* and *FBXW7* were not reported as mutations. Mutational status was not used to modify therapy.

Statistical analysis

Comparison of the 2 MDD/MRD quantification methods was performed by performing Spearman's rank correlation coefficient. Group comparison for categorical and continuous variables was performed with Fisher's exact and Mann-Whitney U tests, respectively. Overall survival (OS) was calculated from the date of prephase initiation to the last follow-up date by censoring alive patients. Events accounting for EFS were a nonresponse on day 33 (defined as >5% blasts in the bone marrow and/or blasts in the cerebrospinal fluid) and/or <35% tumor regression), relapse, secondary malignancy, or death from any cause. Survival analysis was performed using the Kaplan-Meier method, and the curves were compared using the log-rank test. Statistical analysis was performed with the R software, version 4.0.2. All *P* values were two-sided, with *P* <0.05 denoting statistical significance.

Results

From January 2008 to December 2017, 205 pediatric T-LL were diagnosed in 30 SFCE centers (Flow Chart, Figure 1). Eighty-two T-LL patients, who did not differ from the overall group, had MDD and/or MRD analysis. MDD status was assessed for 78 patients, MRD status for 59, both for 55. Clinical and biological characteristics are detailed in Table 1 and Supplemental Digital Table S1, <http://links.lww.com/HS/A195>. As per the EURO-LB02 protocol, patients who received systemic corticosteroids for more than 8 days within 2 months before or at the time of MDD evaluation were excluded, although lower dose oral corticotherapy cannot be excluded for all patients. Two patients died in remission of toxic complications. Twelve patients diagnosed between 2008 and 2010 were previously published for *N/F* status.¹²

MDD assessment by MFC

Sixty-two pediatric T-LL patients had MDD quantification by 8-color MFC from 98 samples (23 PB only, 3 BM only, 36 both) (flow chart, Figure 1). As described for MRD,^{25,26} optimal combinations were those containing TdT, CD99, CD1a, and CD34, in addition to a CD3/cCD3/CD5 backbone. It was possible to compare pathological MDD phenotypes with those of the tumor in only 8 patients for whom a pleural or pericardiac effusion was

Flow chart




Figure 1. Flow chart of MDD/MRD study in French pediatric T-LL from 2008 to 2017. MDD = minimal disseminated disease; MRD = minimal residual disease; T-LL = T lymphoblastic lymphoma.

transmitted to Lyon for central MFC. Despite this, a pathological population of at least 0.01% was identified in 81/102 PB/BM samples (51/62 patients), with a median infiltration of 0.2% (range 0.01%–15%), when analyzing at least 500,000 viable cells (Figure 2A). Infiltration was similar in PB and BM MDD/MRD samples in the 36 patients with dual analysis (Supplemental Digital Figure S1A, <http://links.lww.com/HS/A198>).

MDD quantitation by ddPCR

To evaluate ddPCR MDD/MRD, TR immunogenetic status was evaluated in diagnostic DNA from 62 infiltrated tumor samples by genescan multiplex clonality analysis³⁰ and/or

EuroClonality amplicon NGS for TRD, TRG, and TRB (VDJ and DJ) rearrangements.²⁷ Clonal rearrangement of at least one TR was identified in all but one case, for which MDD analysis was performed by MFC, with TRG clonality identified in 92% (57/62), TRB VDJ ± DJ in 84% (52/62), TRB DJ only in 8% (5/62), and TRD in 50% (30/60). At least one ASO CDR3 primer was developed for each of the 41 T-LL with MDD/MRD samples and first tested by ddPCR on diagnostic tissue samples. ddPCR quantified infiltration in the 22 pleural/pericardiac effusions transmitted to the Necker molecular platform varied from 8% to 100%, median 44% and from 14% to 80%, median 56%, in tissue samples. As such, diagnostic liquid and tissue T-LL samples cannot be presumed to be 100% infiltrated at the DNA level, even if histological interpretation suggests massive

Table 1.

Clinical and Biological Characteristics of the MDD/MRD Cohort and the MDD Cohort According to MDD at 0.1% Threshold

	MDD/MRD cohort (n = 82)	MDD cutoff 0.1% (n = 78)		P
		< 0.1 % (n = 36)	≥0.1 % (n = 42)	
Clinical features				
Median age (y, range)	10.1 (0.5–17.7)	11.15 (4.3–16.9)	9.55 (0.5–17.7)	0.0798
Gender (M, %)	62 (76%)	26 (72%)	33 (79%)	0.515
Stage (n)	(n = 80)			
I/II	8 (10%)	4 (11%)	4 (10%)	0.818
III	55 (69%)	30 (86%)	22 (54%)	0.0027
IV	17 (21%)	1 (3%)	15 (36%)	0.0003
CNS involvement (yes, %)	6/79 (8%)	0/35 (3%)	4/40 (10%)	0.0545
BM involvement (yes, %)	14/79 (18%)	0	13/41 (32%)	0.0002
Relapse (yes, %)	7 (8.5%)	5 (14%)	1 (2%)	0.0512
5y-OS (% , 95% CI)	89.0% [0.825, 0.961]	80.6% [0.686, 0.946]	97.6% [0.931, 1.000]	0.015
5y-EFS (% , 95% CI)	87.8% [0.810, 0.952]	80.6% [0.686, 0.946]	95.2% [0.890, 1.000]	0.049
Biological features				
LDH, xN (average, range)	2.5 (0.6–15.2)	3.1 (0.6–15.2)	1.9 (0.6–6.2)	0.0145
NOTCH1/FBXW7 mutation (yes, %)	42/61 (69%)	19/27 (70%)	20/30 (67%)	0.764
MDD ≥ 0.1 %	42/78 (54%)	/	/	
MRD pos/BQL	9/59 (15%)	3/25 (12%)	5/30 (17%)	0.625

Significant P values, <0.05, are indicated in bold.

/ = not applicable; 5y-OS = 5-y overall survival; 5y-EFS = 5-y event-free survival; BM = bone marrow; BQL = below quantitative level by ddPCR; CNS = central nervous system; MDD = minimal disseminated disease; MRD = minimal residual disease at day 33.




Figure 2. MDD quantification assays. (A) Flow cytometry gating strategy for MDD. Cells were first gated on viable cells, then CyCD3+/SSC, surf CD3+/SSC, Cy CD3+/surf CD3-, then gated by the expression of CD5, TdT, and CD99. Back-gated blasts/MDD are stained black. MRD level = events MDD/events of total viable nucleated cells MDD LAIP (CyCD3+/surfCD3-CD99+TdT+). Top: MDD positive at 0.4%, bottom: MDD undetectable, sensitivity 0.1%. (B) MFC and ddPCR gave comparable quantitative results above 0.1% (Spearman's rank correlation coefficient, kappa = 0.93 [IC 95%: 0.81–1.00]). Sixty-one samples from 27 patients were evaluated (PB, BM, or effusions at diagnosis and/or follow-up). The single BQL positive result by ddPCR was arbitrarily shown at 0.001%. (C) Representation of MDD quantification according to each assay (MFC for 62 patients, ddPCR for 38 patients; higher value of MDD retained when PB and BM performed) and for all patients (highest MDD value retained regardless of the assay or sample). BM = bone marrow; BQL = below quantitative level; ddPCR = droplet digital PCR; MDD = minimal disseminated disease; MFC = multiparameter flow cytometry; MRD = minimal residual disease; PB = peripheral blood; T-LL = T lymphoblastic lymphoma.

involvement by tumor. This is relevant if tumor DNA is used to design qPCR calibration curves for ASO IG/TR MDD/MRD strategies.

ddPCR comparison with MFC

For MDD ddPCR quantification, we first evaluated cell pellets frozen after prospective MFC analysis. A total of 61 samples (18 PB MDD, 19 BM MDD, 5 effusions at diagnosis, 16 PB MRD and 3 BM MRD at d33) from 27 patients were evaluated by both methods (Figure 2B). MFC and ddPCR showed good correlation above the threshold of 0.1% (Spearman's rank correlation coefficient, kappa = 0.93 [IC 95%: 0.81–1.00]). ddPCR gave quantifiable positivity in 10/61 samples (plus one BQL result) with no detectable infiltration by MFC, in keeping with reported sensitivity limits of 0.01% for MFC, compared to 0.001% by ddPCR, at least in MCL.²² For 7 samples (4 patients with 4 MRD and 3 MDD samples) with ddPCR quantification between 0.1% and 0.01%, MFC raw data review confirmed the absence of detectable T-lymphoblasts, albeit in patients whose tumour immunophenotype had not been assessed. In 5 samples (3 patients with 4 MDD and 1 MRD samples), ddPCR

was positive/BQL below the sensitivity of MFC (0.01%). Taken together, these results demonstrate that ddPCR is an appropriate method for MDD/MRD assessment in T-LL, that MFC and ddPCR can be compiled for values above 0.1% and that ddPCR is more sensitive for low level positivity than MFC, as practiced here (0.5–1 million cells analyzed).

MDD infiltration at diagnosis

We extended our ddPCR study to all frozen PB/BM MDD/MRD samples for which a tumour sample with a clonal TR was also available (Figure 2C). In total, 41 patients were evaluated by ddPCR, including 38 with MDD quantification. ASO targets were TRB for 28 patients, TRD for 7 and TRG for 6. Only 2 patients had enough DNA to be quantified on 2 targets, when the most positive result was retained. All samples met the defined ddPCR criteria.²² The vast majority of MDD (34/38, 89%) were quantifiably positive, with 7 samples in the 0.001%–0.01% range and only 1 positive BQL (Figure 2C). Infiltration was similar in PB and BM in the 14 patients with dual analysis (Supplemental Digital Figure S1B, <http://links.lww.com/HS/A198>).

When combined with MFC results, 78 patients had MDD evaluation by MFC, ddPCR or both (Figure 2C). For each patient, the highest MDD value was retained, regardless of the sample type or assay used. We observed a continuum of infiltration in PB/BM, with 10% (8/78; 5 by MFC only, 3 by ddPCR \pm MFC) of patients having undetectable disease, 36% (28/78) positivity $<0.1\%$, 26% (20/78) positive $\geq 0.1\%$, and $<1\%$ and 28% (22/78) $\geq 1\%$. Overall, 54% of patients were MDD positive above 0.1%, at which level MFC and ddPCR were totally concordant.

MDD $\geq 0.1\%$ identifies patients with a good response to EURO-LB02 prednisone/24-month therapy

To evaluate the prognostic relevance of MDD status, we first showed that the patients with MDD analysis were representative of the overall cohort treated with the prednisone/24-month treatment regimen (Supplemental Digital Table S1, <http://links.lww.com/HS/A195>). Applying the historically proposed 1% MDD threshold^{16,17} to the present series demonstrated no prognostic value (Supplemental Digital Figure S2A and B, <http://links.lww.com/HS/A195>). In contrast, when using a MDD cut-off of 0.1%, above which there was an absolute correlation between MFC and ddPCR, patients with MDD $\geq 0.1\%$ had a favorable 5-year OS of 97.6% compared to 80.6% for patients with MDD $<0.1\%$ (HR 0.11, 95% confidence interval [CI]: 0.01–0.93, $P = 0.015$ and a favorable 5-year EFS (95% versus 84%, HR 0.24, 95% confidence interval [CI]: 0.05–1.13, $P = 0.049$) (Figure 3A and B).

To investigate whether there were clinical differences in MDD (cutoff 0.1%) high and low/negative patients, we compared clinical and biological characteristics of the 2 groups (Table 1). Sex ratios were comparable. As expected, Stage IV and BM involvement were more common in patients with MDD $>0.1\%$ (Figure 3C). Patients with MDD $>0.1\%$ had significantly lower LDH values and tended to be younger and have more CNS involvement at diagnosis (Table 1).

The prognostic impact of MDD status is restricted to *NOTCH1/FBXW7* germline T-LL

Since pediatric T-LL patients with *NOTCH1/FBXW7* germline status (N/F^{GL}) are randomized for intensification in the

LBL2018 protocol, we determined MDD impact as a function of N/F status. 21/82 (26%) patients were not evaluated due to absence of sufficiently infiltrated diagnostic material. 19/61 (31%) were N/F^{GL} and 42/61 N/F^{mut} (69%). The level of MDD was comparable in the 3 groups (Supplemental Digital Figure S3A, <http://links.lww.com/HS/A195>). As expected, N/F^{GL} patients had worse outcomes (Supplemental Digital Figure S3B and C, <http://links.lww.com/HS/A195>).

Using the MDD 0.1% threshold, MDD status had no prognostic impact in N/F^{mut} cases (5-y OS: 95% for MDD $\geq 0.1\%$ versus 94.7% for MDD $<0.1\%$; 5-y EFS: 90% versus 94.7%, respectively; Figure 4A and B). In contrast, among N/F^{GL} cases, 8/18 (44%) were MDD $<0.1\%$ and had inferior prognosis (5-y OS of 37.5% versus 100% for MDD $\geq 0.1\%$, $P < 0.001$; a 5-y EFS of 37.5% versus 100%, respectively, $P < 0.001$; Figure 4A and B). Details of the patients who progressed or relapsed are shown in Supplemental Digital Table S2, <http://links.lww.com/HS/A196>.

Overall, a molecular classifier combining MDD assessment and N/F genotype identified striking differences in outcome between the 8/57 (14%) high-risk patients (N/F^{GL} and MDD $<0.1\%$) and low-risk patients (all others) with 5-year OS at 37.5% versus 95.9%, HR 20.70, 95% CI: 3.98–107.50, $P < 0.001$; 5-year EFS of 37.5% versus 93.9% HR 14.27, 95% CI: 3.36–60.71, $P < 0.001$, respectively (Figure 4C and D). Taken together, these results demonstrate that combined MDD and oncogenetic evaluation at diagnosis is likely to allow the identification of patients who should be considered for alternative treatment front line, given the very poor survival of relapsed pediatric T-LL.

Prognostic impact of MRD

MRD at the end of induction is a fundamentally important prognostic factor in T-ALL. We therefore evaluated MRD at d33 in 59 patients with available samples (41 by MFC only, 2 by ddPCR only, 16 by both; flow chart, Figure 1). One N/F^{mut} case (N° 6134, Supplemental Digital Table S2, <http://links.lww.com/HS/A196>) remained persistently MRD positive by ddPCR 4 months after diagnosis, so was switched to a high-risk T-ALL regimen with Nelarabine, and is in CR 3 years later. Samples from 9 patients (15%) were positive or BQL (2 were MFC+/ddPCR+, 4 ddPCR+/MFC- and 3 MFC+/ddPCRnd), and 50 (85%) were undetectable, with a sensitivity of 0.01%. Details of the patients with pos/BQL MRD are shown in Supplemental Digital Table S2,




Figure 3. Outcomes of patients according to MDD at 0.1% threshold and MDD quantification according to clinical stage. (A) 5-year overall survival, Kaplan–Meier curve according to MDD at 0.1% threshold, (B) 5-year event-free survival, Kaplan–Meier curve according to MDD at 0.1% threshold, (C) MDD quantification according to clinical stage. Percentage of T-cell lymphoblastic lymphoma cells in BM or PB according to disease stage in the 78 patients of the study. Horizontal bars indicate the median value in each group. BM = bone marrow; MDD = minimal disseminated disease; PB = peripheral blood.




Figure 4. Outcomes of patients according to *NOTCH1*/*FBXW7* in combination with MDD at 0.1% threshold in pediatric T-LL. (A) 5-y OS, Kaplan-Meier curve according to *N/F* and MDD 0.01% status, (B) 5-y EFS, Kaplan-Meier curve according to *N/F* and MDD 0.01% status, (C) 5-y OS comparing the 14% of *N/F^{GL}* and MDD $< 0.1\%$ patients with all others, (D) 5-y EFS comparing the 14% of *N/F^{GL}* and MDD $< 0.1\%$ patients with all others. 5y-OS = 5-y overall survival; 5y-EFS = 5-y event-free survival; ddPCR = droplet digital PCR; MDD = minimal disseminated disease; OS = overall survival; T-LL = T lymphoblastic lymphoma.

<http://links.lww.com/HS/A196>. Prognostic evaluation showed an inferior outcome in MRD positive patients, with 4-year OS of 77.8% for MRD positive/BQL versus 92.0% for MRD undetectable, HR 0.31, 95% CI: 0.06–1.71, $P = 0.157$; a 4-year EFS of 66.7% versus 92.0%, respectively, HR 0.31, 95% CI: 0.06–1.71, $P = 0.015$, respectively (Figure 5A and B). MRD alone did not predict the majority of the 6 relapses/progression, whose clinico-biological characteristics are detailed in Supplemental Digital Table S2, <http://links.lww.com/HS/A196>. The small sample size and the mixture of techniques used preclude reliable analysis of MRD significance within MDD or *N/F* defined subsets, but only 2/7 relapsing patients tested had MDD $\ge 0.1\%$ and only 2/7 tested at diagnosis had *N/F^{mut}* T-LL, suggest that this should be addressed specifically and prospectively in a collaborative study.

Discussion

T-LL prognosis has greatly improved with ALL treatment strategies, achieving more than 85% OS. In contrast, refractory patients and those who relapse, predominantly within the first 2 years, have a dismal outcome, so their early identification is essential for timely therapeutic adaptation. We show that poor prognosis pediatric T-LL is virtually restricted to patients with

MDD levels in PB/BM below 0.1% and absence of *NOTCH1*/*FBXW7* mutations at diagnosis. These results contradict prior data suggesting that MDD positivity is associated with poor prognosis.

We performed MDD and MRD quantification by MFC and ddPCR, following ddPCR approaches developed for Mantle Cell Lymphoma, which also frequently disseminates to PB and BM.²² Results were perfectly concordant for positivity levels above 0.1%, allowing compilation. ddPCR was, however, more likely to detect lower level positivity, since 10/32 samples were positive/BQL by ddPCR, but negative by MFC, compared to only 3 which were MFC positive (all at 0.01%) but ddPCR negative. In keeping with this, a disproportionate number of positive MRD samples were detected by ddPCR. As such, either technique is acceptable for detection of high level positivity, but evaluation of levels below 0.1% should either be performed by molecular clonotype quantification or by MFC of higher cell numbers than the 0.5–1 million events analyzed here. This is increasingly practiced in ALL MRD evaluation, but it is noteworthy that MFC is not recommended in isolation to determine MRD negativity in the European ALL-Together trial (EUDRACT 2018-001795-38). MDD evaluation by MFC may also be challenging without characterization of the tumour phenotype and should only be assessed by experienced reference platforms with sufficient T-LL recruitment.




Figure 5. Outcomes of patients according to MRD at day 33 in 59 pediatric T-LL patients. (A) 5y-OS according to MRD analysis in blood and/or bone marrow at end of induction. (B) 5y-EFS according to MRD analysis in blood and/or bone marrow at end of induction. 5y-OS = 5-y overall survival; 5y-EFS = 5-y event-free survival; MRD = minimal residual disease; OS = overall survival; T-LL = T lymphoblastic lymphoma.

Despite a comparable incidence of patients with high level MDD values, we did not confirm the inferior outcome of high level MDD of the COG-A5971 trial.¹⁶ On the contrary, when using a lower MDD cutoff at 0.1% (10^{-3}), patients with circulating T lymphoblasts responded relatively well to ALL-type therapy. This is in keeping with the, at least, comparable outcome of Stage IV (5-y EFS 88%) compared to stage III (EFS 78%) patients in the EURO-LB02 trial.²⁴ Reasons for this discrepancy are not evident, but may result from differences in treatment compared to A5971.¹⁶ This is in keeping with the loss of prognostic significance of MDD in the COG AALL0434 trial,¹⁹ interpreted to result from a modified interim phase with escalating intravenous methotrexate (Capizzi MTX)/pegasparaginase regimens. However, the EURO-LB02 used a HD-MTX regime similar to A5971, while achieving similar results to AALL0434.²⁴ In light of our data, it would be interesting to compare MDD outcome in AALL0434 using a 0.1% cutoff in patients with known *N/F* status, although comparisons are hampered by use of different use of MFC and ddPCR for MDD assessment. MDD low/negative values here could be due to prior steroids, since only systemic corticotherapy was an exclusion criterion, but this seems unlikely to explain the poor prognosis, since corticosenitivity is usually associated with a relatively good outcome.

Given the relatively favorable outcome of *N/F^{mut}* in pediatric T-LL,^{11,12} this parameter is used to randomize patients for treatment intensification in the LBL2018 trial. The incidence of *N/F^{mut}* is slightly higher in the present cohort of patients compared to our initial assessment (66% versus 55%)¹² at least partly due to the replacement of Sanger screening by next-generation-sequencing capture screening for *N/F* status.²⁹

The impact of MDD status was strikingly different in *N/F^{mut}* and *N/F^{GL}* patients. Whereas there was no prognostic impact in *N/F^{mut}* patients, MDD below 0.1% clearly identified *N/F^{GL}* patients at risk of relapse. These observations suggest that significant dissemination to PB/BM may correlate with sensitivity to ALL-type therapy. Conversely, purely tissue-based disease with low/undetectable MDD negativity may be less sensitive. One therapeutic possibility for the latter group might be proteasome inhibitors, given the demonstration in the COG AALL1231 trial that Bortezomib was of benefit for de novo pediatric T-LL but not T-ALL.³¹ Anti-CD38 immunotherapy could also be considered.

Our data encourage addition of MDD assessment for patients in the LBL2018 trial, preferably by central ddPCR or qPCR, given the superior sensitivity compared to flow cytometry, as practiced here, and since it is better adapted to retrospective analysis.

MRD evaluation in a limited number of patients, predominantly by MFC, confirmed previous reports for poor outcome. It failed, however, to detect the majority of relapses and will be difficult to undertake in purely nodal, MDD low/negative patients. Since a disproportionate number of positive results were from the minority of samples assessed by ddPCR, this technique would seem preferable to MFC for MRD assessment. Although it will be possible to perform MRD evaluation in the majority of patients, its value relative to combined oncogenetic/MDD stratification should be evaluated within *N/F* defined subgroups, as should its prognostic value compared to imaging evaluation. Whether patients with MDD low/negative status should be screened for MRD emergence (presuming that tumor immunogenotype/phenotype can be determined from diagnostic tissue) is unclear, but our data illustrate that this is unlikely to be as useful as in T-ALL, somewhat complicating inclusion of T-LL in MRD-driven T-ALL protocols.

In practical terms, pediatric T-LLs with *N/F^{mut}* respond very well to current standard protocols and MDD status adds little/nothing to their risk assessment. Most would be easily MRD accessible, since 82% demonstrated at least 0.01% PB/BM clonal dissemination at diagnosis, but their excellent outcome will make it difficult to demonstrate added value of MRD stratification, and MRD kinetics may differ in T-LL and T-ALL. Although the small number of relapsing cases precludes definitive assessment, of the 6 *N/F* evaluated relapses with MRD evaluation, only two were MRD positive (Supplemental Digital Table S2, <http://links.lww.com/HS/A196>). Among *N/F^{GL}* patients, MDD <0.1% status represents a promising means of rapidly identifying patients with a very high relapse risk, if confirmed in other T-LL trials with larger patient numbers. Approximately 80% would be MRD accessible, and the impact of MRD in this small *N/F^{GL}*, MDD low/neg. subgroup (8/57, 14%) should also be evaluated prospectively, although this will require large patient numbers and universal availability of diagnostic tissue for baseline molecular and immunophenotypic assessment. It will also be important to evaluate other oncogenetic markers in these patients,¹³ who merit consideration for alternative front line therapy.

Acknowledgements

The authors thank the SFCE and the investigators, the patients and their families from the 24 SFCE-centers involved for providing patient material and data: Anne Lutin, Isabelle Pelier, Nathalie Cheik, Marianna Dapris, Justyna Kanold, H el ene Pacquement, Florent

Neumann, Dominique Plantaz, Brigitte Nelken, Charlotte Oudot, Catherine Curtillet, Claudine Schmitt, Marie Laure Couec, Maryline Poiree, Thierry Leblanc, Arnaud Petit, Frederic Millot, Gregory Guimard, Virginie Gandemer, Nimrod Buchbinder, Audrey David, Veronique Minard.

Disclosures

The authors have no conflicts of interest to disclose.

Sources of funding

This study was supported by La Ligue Nationale de Recherche sur le Cancer (Enfants, Adolescents et Cancer), by Enfants et Santé and by the SFCE. The Trousseau team was supported by the Recherche en Hémopathies Malignes de l'Enfant (RHME). The Necker LL RELYE (Réseau des lymphomes de l'Enfant) tissue bank is supported by IMAGINE for Margo via the SFCE, the Institut National du Cancer and the CARPEM SIRIC.

References

- Hofmans M, Suci S, Ferster A, et al. Results of successive EORTC-CLG 58 881 and 58 951 trials in paediatric T-cell acute lymphoblastic leukaemia (ALL). *Br J Haematol*. 2019;186:741–753.
- Bergeron C, Coze C, Segura C, et al; Société Française d'Oncologie Pédiatrique (SFOP), France. Treatment of childhood T-cell lymphoblastic lymphoma-long-term results of the SFOP LMT96 trial. *Pediatr Blood Cancer*. 2015;62:2150–2156.
- Uyttebroeck A, Suci S, Laureys G, et al. Treatment of childhood T-cell lymphoblastic lymphoma according to the strategy for acute lymphoblastic leukaemia, without radiotherapy: long term results of the EORTC CLG 58881 trial. *Eur J Cancer*. 2008;44:840–846.
- Schmidt E, Burkhardt B. Lymphoblastic lymphoma in childhood and adolescence. *Pediatr Hematol Oncol*. 2013;30:484–508.
- Michaux K, Bergeron C, Gandemer V, et al. Relapsed or refractory lymphoblastic lymphoma in children: results and analysis of 23 patients in the EORTC 58951 and the LMT96 protocols. *Pediatr Blood Cancer*. 2016;63:1214–1221.
- Gross TG, Hale GA, He W, et al. Hematopoietic stem cell transplantation for refractory or recurrent non-Hodgkin lymphoma in children and adolescents. *Biol Blood Marrow Transplant*. 2010;16:223–230.
- Trinquand A, Tanguy-Schmidt A, Ben Abdelali R, et al. Toward a NOTCH1/FBXW7/RAS/PTEN-based oncogenetic risk classification of adult T-cell acute lymphoblastic leukemia: a Group for Research in Adult Acute Lymphoblastic Leukemia study. *J Clin Oncol*. 2013;31:4333–4342.
- Petit A, Trinquand A, Chevret S, et al. Oncogenetic mutations combined with MRD improve outcome prediction in pediatric T-cell acute lymphoblastic leukemia. *Blood*. 2018;131:289–300.
- Balbach ST, Makarova O, Bonn BR, et al. Proposal of a genetic classifier for risk group stratification in pediatric T-cell lymphoblastic lymphoma reveals differences from adult T-cell lymphoblastic leukemia. *Leukemia*. 2016;30:970–973.
- Basso K, Mussolin L, Lettieri A, et al. T-cell lymphoblastic lymphoma shows differences and similarities with T-cell acute lymphoblastic leukemia by genomic and gene expression analyses. *Genes Chromosomes Cancer*. 2011;50:1063–1075.
- Bonn BR, Rohde M, Zimmermann M, et al. Incidence and prognostic relevance of genetic variations in T-cell lymphoblastic lymphoma in childhood and adolescence. *Blood*. 2013;121:3153–3160.
- Callens C, Baleyrier F, Lengline E, et al. Clinical impact of NOTCH1 and/or FBXW7 mutations, FLASH deletion, and TCR status in pediatric T-cell lymphoblastic lymphoma. *J Clin Oncol*. 2012;30:1966–1973.
- Khanam T, Sandmann S, Seggewiss J, et al. Integrative genomic analysis of pediatric T-cell lymphoblastic lymphoma reveals candidates of clinical significance. *Blood*. 2021;137:2347–2359.
- Burkhardt B, Bruch J, Zimmermann M, et al. Loss of heterozygosity on chromosome 6q14-q24 is associated with poor outcome in children and adolescents with T-cell lymphoblastic lymphoma. *Leukemia*. 2006;20:1422–1429.
- van der Velden VH, Jacobs DC, Wijkhuijs AJ, et al. Minimal residual disease levels in bone marrow and peripheral blood are comparable in children with T cell acute lymphoblastic leukemia (ALL), but not in precursor-B-ALL. *Leukemia*. 2002;16:1432–1436.
- Coustan-Smith E, Sandlund JT, Perkins SL, et al. Minimal disseminated disease in childhood T-cell lymphoblastic lymphoma: a report from the children's oncology group. *J Clin Oncol*. 2009;27:3533–3539.
- Mussolin L, Buldini B, Lovisa F, et al. Detection and role of minimal disseminated disease in children with lymphoblastic lymphoma: The AIEOP experience. *Pediatr Blood Cancer*. 2015;62:1906–1913.
- Stark B, Avigad S, Luria D, et al. Bone marrow minimal disseminated disease (MDD) and minimal residual disease (MRD) in childhood T-cell lymphoblastic lymphoma stage III, detected by flow cytometry (FC) and real-time quantitative polymerase chain reaction (RQ-PCR). *Pediatr Blood Cancer*. 2009;52:20–25.
- Hayashi RJ, Winter SS, Dunsmore KP, et al. Successful outcomes of newly diagnosed T lymphoblastic lymphoma: results from Children's Oncology Group AALL0434. *J Clin Oncol*. 2020;38:3062–3070.
- van der Velden VH, Cazzaniga G, Schrauder A, et al. Analysis of minimal residual disease by Ig/TCR gene rearrangements: guidelines for interpretation of real-time quantitative PCR data. *Leukemia*. 2007;21:604–611.
- Cavalli M, De Novi LA, Della Starza I, et al. Comparative analysis between RQ-PCR and digital droplet PCR of BCL2/IGH gene rearrangement in the peripheral blood and bone marrow of early stage follicular lymphoma. *Br J Haematol*. 2017;177:588–596.
- Drandi D, Alcantara M, Benmaad I, et al. Droplet digital PCR quantification of mantle cell lymphoma follow-up samples from four prospective trials of the European MCL Network. *HemaSphere*. 2020;4:e347.
- Drandi D, Ferrero S, Ladetto M. Droplet digital PCR for minimal residual disease detection in mature lymphoproliferative disorders. *Methods Mol Biol*. 2018;1768:229–256.
- Landmann E, Burkhardt B, Zimmermann M, et al. Results and conclusions of the European Intergrup EURO-LB02 trial in children and adolescents with lymphoblastic lymphoma. *Haematologica*. 2017;102:2086–2096.
- Fossat C, Roussel M, Arnoux I, et al. Methodological aspects of minimal residual disease assessment by flow cytometry in acute lymphoblastic leukemia: a French multicenter study. *Cytometry B Clin Cytom*. 2015;88:21–29.
- Garand R, Beldjord K, Cavé H, et al. Flow cytometry and IG/TCR quantitative PCR for minimal residual disease quantitation in acute lymphoblastic leukemia: a French multicenter prospective study on behalf of the FRALLE, EORTC and GRAALL. *Leukemia*. 2013;27:370–376.
- Brüggemann M, Kotrová M, Knecht H, et al. Standardized next-generation sequencing of immunoglobulin and T-cell receptor gene recombinations for MRD marker identification in acute lymphoblastic leukaemia; a EuroClonality-NGS validation study. *Leukemia*. 2019;33:2241–2253.
- Duez M, Giraud M, Herbert R, et al. Vidjil: a web platform for analysis of high-throughput repertoire sequencing. *PLoS One*. 2016;11:e0166126.
- Bond J, Graux C, Lhermitte L, et al. Early response-based therapy stratification improves survival in adult early thymic precursor acute lymphoblastic leukemia: a group for research on adult acute lymphoblastic leukemia study. *J Clin Oncol*. 2017;35:2683–2691.
- van Dongen JJ, Langerak AW, Brüggemann M, et al. Design and standardization of PCR primers and protocols for detection of clonal immunoglobulin and T-cell receptor gene recombinations in suspect lymphoproliferations: report of the BIOMED-2 concerted action BMH4-CT98-3936. *Leukemia*. 2003;17:2257–2317.
- Teachey DT, Devidas M, Wood BL, et al. Cranial radiation can be eliminated in most children with T-cell acute lymphoblastic leukemia (T-ALL) and bortezomib potentially improves survival in children with T-cell lymphoblastic lymphoma (T-L): results of Children's Oncology Group (COG) trial AALL. *Blood*. 2020;136(Suppl 1):11–12.