



HAL
open science

Effect of 2020 containment strategies on trauma workflow in Ile-de-France region: another benefit of lockdown?

Jean-Denis Moyer, Paer-Selim Abback, Sophie Hamada, Thibault Martinez, Marie Werner, Arthur James

► To cite this version:

Jean-Denis Moyer, Paer-Selim Abback, Sophie Hamada, Thibault Martinez, Marie Werner, et al.. Effect of 2020 containment strategies on trauma workflow in Ile-de-France region: another benefit of lockdown?. *Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine*, 2021, 29 (1), 10.1186/s13049-021-00918-x . hal-03353752

HAL Id: hal-03353752

<https://hal.sorbonne-universite.fr/hal-03353752>

Submitted on 24 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

COMMENTARY

Open Access



Effect of 2020 containment strategies on trauma workflow in *Ile-de-France* region: another benefit of lockdown?

Jean-Denis Moyer^{1*} , Paer-Selim Abback¹, Sophie Hamada², Thibault Martinez³, Marie Werner⁴, Arthur James⁵ and the Traumabase® Group

Abstract

Background: During the SARS-CoV-2 pandemic, the French Government imposed various containment strategies, such as severe lockdown (SL) or moderate lockdown (ML). The aim of this study was to evaluate the effect of both strategies on severe trauma admissions and ICU capacity in *Ile-de-France* region (Paris Area).

Main text: We conducted a multicenter cohort-based observational study from 1st January 2017 to 31st December 2020, including all consecutive trauma patients admitted to the trauma centers of *Ile-de-France* region participating in the national registry (Traumabase®). Two periods were defined, the “non-pandemic period” (NPP) from 2017 to 2019, and the “pandemic period” (PP) concerning those admitted in 2020. The number of ICU beds released during 2020 pandemic period (overall period, SL and ML) was estimated by multiplying difference in trauma admissions by the median length of stay during the same week of pandemic period (ICU day-beds in 2020).

A 15% yearly reduction of trauma patients was observed during the PP, associated with the release of 6422 ICU day-beds in 2020. During SL and ML, the observed decrease in trauma admission was respectively 49 and 39% compared with similar dates of the NPP. The number of beds released was 1531 days-beds in SL and 679 day-beds in ML. Those reductions respectively accounted for 4.5 and 6.0% of the overall ICU admission for COVID-19 in *Ile-de-France*.

Conclusion: The lockdown strategies during pandemic resulted in a reduction of severe trauma admissions. In addition to the social distancing effect, lockdown strategies freed up an important number of ICU beds in trauma centers, available for severe COVID-19 patients.

Keywords: COVID-19, Trauma, France, Traumacenter, ICU capacity

Background

Because of the first and the second waves of SARS-CoV-2 pandemic in 2020, a large number of patients were admitted to French intensive care units (ICU) [1, 2]. As a consequence of this massive influx of patients, ICU capacity had to be increased considerably. In order to

absorb the SARS-CoV-2 burden and to limit the ICU overflow, the French government, as well as many others in the world, imposed strict containment strategies during the spring and fall waves. The Covitrauma study recently published in this journal as confirmed that the first lockdown reduced severe trauma admissions by 50% [3], indicating an indirect benefit of the lockdown by reducing non SARS-CoV-2 ICU emergency admissions. The aim of this commentary was to complete our first study through the evaluation of the effect of

* Correspondence: Jean-denis.moyer@aphp.fr

¹Department of Anesthesiology and Critical Care, Beaujon Hospital, DMU Parabol, AP-HP.Nord, 100 boulevard du General Leclerc, F92110 Paris, Cligny, France

Full list of author information is available at the end of the article



© The Author(s). 2021 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

containment strategies on severe trauma admissions and ICU capacity in *Ile-de-France* region (Paris Area).

Main text

We conducted a multicenter cohort-based observational study from 1st January 2017 to 31th December 2020. During each of these 4 years, all consecutive trauma admitted to the trauma centers of *Ile-de-France* region (12.2 million of inhabitants) participating in the national registry (Traumabase[®]) were included (5 teaching hospitals and one military teaching hospital). Patients were categorized in two groups according to their year of admission, allocating those admitted within years 2017, 2018 and 2019 to the “non-pandemic period” group and those admitted in 2020 to the “pandemic period” group. Each variable result over the non-pandemic period were summarized using its **median** (and interquartile) or number (and relative percentage).

During the pandemic period, two types of lockdown strategies were applied in *Ile de France* area:

- Severe lockdown (SL) from week 12 to week 19 **in 2020**: leaving home was only allowed for essential purchases, medical care, exercise (1-h session per

day), and work journeys if work from home was impossible.

- Moderate lockdown (ML) from week 45 to week 51 **in 2020**: leaving home was only allowed for essential purchases, medical care, exercise (1-h session per day), and work journeys (work in factories and public works were allowed). Kindergarten, schools and high schools were opened.

The number of ICU beds released during 2020 pandemic period (overall period, SL and ML) was estimated by multiplying difference in trauma admissions (difference between median number of trauma admissions each week during the pandemic and the non-pandemic period) by the median length of stay during the same week of pandemic period (ICU day-beds in 2020).

Data for SARS-CoV-2 patients admitted in ICU in *Ile-de-France* region are available on open access data [4].

Over the 4 year-period, 11,230 trauma patients were admitted in the participating trauma centers, 8749 during the non-pandemic period (average 2916 per year) and 2481 during the pandemic period, representing a 15% yearly reduction during the pandemic period (Table 1 and Fig. 1). This observed 15% reduction

Table 1 Epidemiology of patients admitted to Trauma centers during pandemic period (2020) and non-pandemic period (2017–2019)

Variable	Period					
	Overall year		Severe Lockdown		Moderate lockdown	
	Non-pandemic	Pandemic	Non-pandemic	Pandemic (SL)	Non-pandemic	Pandemic (ML)
Number of admitted patients	2916 ^b	2481	466 ^b	237	312 ^b	189
Age (Years)	36 [25, 52]	34 [23, 50] ^a	37 [25, 53]	35 [26, 49]	36 [25, 54]	33 [23, 48] ^a
Gender (Men)	2279 ^b (78.5)	2002 (82.2) ^a	364 ^b (78.2)	196 (83.1)	235 ^b (76.1)	147 (78.2)
Mechanism of injury distribution						
Penetrating trauma	364 ^b (12.5)	387 (15.6) ^a	55 ^b (11.8)	36 (15.2) ^a	39 ^b (12.4)	34 (18.0) ^a
Road traffic accident	1566 ^b (53.7)	1452 (50.5) ^a	249 ^b (53.3)	93 (39.9) ^a	170 ^b (54.7)	91 (48.1) ^a
Fall from height	606 ^b (20.8)	546 (22.0) ^a	97 ^b (20.9)	86 (36.6) ^a	55 ^b (17.8)	52 (27.5) ^a
Other	381 ^b (13.1)	296 (11.9) ^a	64 ^b (13.7)	12 (9.3) ^a	47 ^b (15.2)	12 (6.3) ^a
ISS	11 [5, 21]	11 [5, 20]	11 [5, 22]	12 [5, 20]	12 [5, 22]	13 [8, 22]
SAPS 2	20 [13, 34]	20 [13, 31]	21 [13, 34]	21 [13, 30]	21 [13, 36]	20 [13, 32]
Hemorrhagic shock	176 ^b (6.1)	141 (5.8)	33 ^b (7.1)	11 (4.8)	19 ^b (6.2)	11 (5.8)
Traumatic brain injury	707 ^b (24.4)	516 (21.7) ^a	108 ^b (23.4)	44 (19.4)	80 ^b (26.2)	30 (16.5)
Surgery in the first 24 h	1452 ^b (49.8)	1226 (49.4)	240 ^b (51.6)	127 (53.6)	140 ^b (45.5)	96 (50.8)
In-hospital mortality	832 ^b (9.6)	171 (7.5) ^a	133 ^b (9.6)	10 (4.6)	101 ^b (10.9)	14 (8.1)
ICU Length of stay (days)	2 [2, 6]	2 [2, 5]	2 [2, 6]	2 [2, 5]	2 [2, 7]	2 [2, 6]
Predicted mortality (TRISS, mean, SD) (%)	8.5 (19.1)	7.4 (17.9)	8.4 (19.3)	7.7 (19.2)	9.2 (19.8)	9.1 (20.2)

Non-pandemic period: 2017–2018–2019. Pandemic period: 2020

^a highlight differences with p -value ≤ 0.01 between the pandemic and the non-pandemic period

^b For the non-pandemic period, we divided number of categorical variables by 3 in order make a comparison possible

Data are n (%) or median [Q1, Q3]

ICU Intensive Care Unit, ISS Injury Severity Score, SAPS 2 Simplified Acute Physiology Score, TRISS Trauma Related Injury Severity Score

SL severe lockdown; ML moderate lockdown

between both periods was associated with the release of 6422 ICU day-beds in 2020. During SL and ML, the observed decrease in trauma admission was respectively 49 and 39% compared with similar dates of the non-pandemic period (Fig. 1 and Table 1). During the 8 weeks of SL and the 6 weeks of ML, the number of beds released by the reduction of the trauma activity was 1531 days-beds and 679 day-beds respectively. Those reductions of trauma admission respectively accounted for 4.5 and 6.0% of the overall ICU admission required for COVID-19 over the same period of time in *Ile-de-France*.

The characteristics of the population during pandemic and non-pandemic period are described Table 1. Trauma severity assessed by Injury Severity Score did not differ during both periods (11 [5, 21] vs 10 [5, 21] $p = 0.68$). Mechanisms of injury differed markedly between both period: road traffic accidents decreased during both lockdown whereas proportion of penetrating trauma and fall from height significantly increased (but not the absolute numbers of event) (Table 1).

These results highlight how trauma epidemiology in Paris Area has been impacted during the pandemic period and especially during the SL and the ML. The national lockdowns imposed by authorities to control the pandemic reduced severe trauma incidence. This study suggests that lockdown benefice on SARS-CoV-2

pandemic management is not only related to the prospect of controlling virus circulation. The reduction of trauma center workflow during lockdown freed up an important number of ICU beds for critical care patients and this effect seems to be more pronounced with a severe lockdown rather than a moderate one. This “collateral effect” of lockdown partly protected our health care system when facing the overwhelming and sudden surge in the number of patients in need of urgent treatment, it allowed to reallocate these free ICU beds to SARS-CoV-2 patients and probably contributed to maintain a part of scheduled major surgical activity such as cancer, cardio-vascular surgery or transplantation. This important observation might help public health stakeholders when preparing for potential future pandemic waves.

Furthermore, during this period, trauma centers did not restrict their admission policy, as indicated by the similar median ISS across the study periods. The proportion of patients operated within the first 24 h remained identical suggesting timely access to complex trauma surgery was not impeded. The trauma system analyzed in this study provides adequate and appropriate care equivalent to non-pandemic period, while participating significantly to the 2020 COVID-19 surge.

Study limitations first include a lack of specific data on SARS-CoV-2 admission in Trauma centers. However, the official occupation rate of ICU in Paris area during




Fig. 1 Weekly number of patients admitted to Trauma center and to *Ile-de-France* Intensive Care Unit for SARS-CoV2 infection in pandemic period (2020), compared with patients admitted to Trauma center during non-pandemic period (average of 2017–2019). SL: severe lockdown, ML: moderate lockdown

pandemic waves was over 100% during SL and ML compared with ICU standard capacity. Secondly, this study only represents urban territory and should not be transposed in rural territories. Thirdly, data for non-trauma centers are not available; then the effect of lockdown on ICU beds release in non-trauma center could not be analyzed.

Conclusion

The lockdown strategies during pandemic resulted in a reduction of severe trauma admissions. This reduction did not appear to be driven by modification of severe trauma admission policy or quality of care, although surge capacity may have impacted both. In addition to the social distancing effect, lockdown strategies freed up an important number of ICU beds in trauma centers, available for severe COVID-19 patients.

Abbreviations

ICU: Intensive care unit; SL: Severe lockdown; ML: Moderate lockdown

Financial support

None.

Authors' contributions

All authors have made substantial contributions to this work and have approved the final version of the manuscript. Data collection and quality control: All Authors. Concept and design: JDM, AJ. Statistical analysis: AJ. Interpretation of data: JDM, AJ, PSA, SH. Writing original draft: JDM, AJ, SH, PSA, TM, MW. Writing review and editing: all authors.

Availability of data and materials

The data that support the findings of this study are available from the Traumabase® group but restrictions apply to the availability of these data, which were used under license for the current study, and so are not publicly available. All data are however available from the authors upon reasonable request and with permission of the Traumabase® group. COVID-19 ICU related data are freely available (<https://www.data.gouv.fr/fr/datasets/r/41b9bd2a-b5b6-4271-8878-e45a8902ef00>).

Declarations

Competing of interests

No conflict of interest.

Ethics approval and consent to participate

Given the retrospective nature of the study and according to French law, the TraumaBase group obtained approval for this study, including use of data retrospectively, from the Institutional Review Board (Comité de Protection des Personnes, Paris VI, Pitié, president Pr Laurent Lacapelle, Bâtiment de la Force, 47 Boulevard de l'Hôpital, 75651 Paris Cedex 13, 28 November 2012) which is equivalent to an Ethical Committee and from the Advisory Committee for Information Processing in Health Research (Comité consultatif sur le traitement de l'information en matière de recherche dans le domaine de la santé, autorisation 11.305bis) and from the National Commission for Data Protection (Commission Nationale de l'Informatique et des Libertés, autorisation 911461).

Consent for publication

Not applicable.

Author details

¹Department of Anesthesiology and Critical Care, Beaujon Hospital, DMU Parabol, AP-HP.Nord, 100 boulevard du General Leclerc, F92110 Paris, Clichy, France. ²Department of Anesthesiology and Critical Care, Hôpital Européen Georges Pompidou, AP-HP, Paris, France. ³Intensive Care Unit, Percy Military

Teaching Hospital, 101 Avenue Henri Barbusse 92140, Clamart, Val de Grace Academy, Place Alphonse Laveran, 75005 Paris, France. ⁴Department of Anesthesiology and Critical Care, AP-HP, Bicêtre Hôpitaux Universitaires Paris-Sud, Université Paris Saclay, Le Kremlin Bicêtre, France. ⁵GRC 29, AP-HP, DMU DREAM, Department of Anaesthesiology and critical care, Pitié-Salpêtrière Hospital, Sorbonne University, Paris, France.

Received: 24 June 2021 Accepted: 10 July 2021

Published online: 14 September 2021

References

1. Lefrant J-Y, Fischer M-O, Potier H, Degryse C, Jaber S, Muller L, et al. A national healthcare response to intensive care bed requirements during the COVID-19 outbreak in France. *Anaesth Crit Care Pain Med* Published online October 5, 2020;(6):709–15. <https://doi.org/10.1016/j.accpm.2020.09.007>.
2. COVID-ICU Group on behalf of the REVA Network and the COVID-ICU Investigators. Clinical characteristics and day-90 outcomes of 4244 critically ill adults with COVID-19: a prospective cohort study. *Intensive Care Med*. 2021;47(1):60–73. <https://doi.org/10.1007/s00134-020-06294-x>.
3. Moyer J-D, James A, Gakuba C, et al. Impact of the SARS-COV-2 outbreak on epidemiology and management of major trauma in France: a registry-based study (the COVITRAUMA study). *Scand J Trauma Resusc Emerg Med*. 2021; 29(1):51. <https://doi.org/10.1186/s13049-021-00864-8>.
4. Données hospitalières relatives à l'épidémie de COVID-19. <https://www.data.gouv.fr/fr/datasets/donnees-hospitalieres-relatives-a-lepidemie-de-covid-19/>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

