



HAL
open science

A new strange Archaeorthoptera from the Moscovian of Avion (France) (Insecta, Polyneoptera)

André Nel, Patrick Roques

► To cite this version:

André Nel, Patrick Roques. A new strange Archaeorthoptera from the Moscovian of Avion (France) (Insecta, Polyneoptera). *Historical Biology*, 2021, 34 (9), pp.1769-1771. 10.1080/08912963.2021.1978082 . hal-03356138

HAL Id: hal-03356138

<https://hal.sorbonne-universite.fr/hal-03356138>

Submitted on 27 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Historical Biology

2 Manuscript DOI: 10.1080/08912963.2021.1978082

3

4 A new strange Archaeorthoptera from the Moscovian of Avion (France) (Insecta, Polyneoptera)

5

6 André Nel^a and Patrick Roques^b

7

8 ^aInstitut de Systématique, Évolution, Biodiversité (ISYEB) Muséum national d'Histoire
9 naturelle, CNRS, Sorbonne Université, EPHE, Université des Antilles, CP50, 57 rue Cuvier
10 75005 Paris, France

11 ^b2 Chemin des Processions, 93360 Neuilly Plaisance, France.

12

13 anel@mnhn.fr; <https://orcid.org/0000-0002-4241-7651>

14 patrick.roques93@wanadoo.fr; <https://orcid.org/0000-0002-8721-9763>

15

16 ABSTRACT

17 A new archaeorthopteran genus and species, *Avionugonioneura jouaulti*, is described from the
18 Moscovian of the Avion in Northern France. It shows several strong similarities, with the two
19 genera *Contracladus* (Pennsylvanian) and *Nugonioneura* (early Permian), **viz. a simple vein M**
20 **emerging anteriorly from M+CuA+CuPa and an anteriorly branched CuA+CuPa**, which are
21 putative synapomorphies, suggesting a possible attribution to the family Nugonioneuridae. This
22 discovery confirms the very important diversity of the Archaeorthoptera in the outcrop of
23 Avion.

24

25 KEYWORDS

26 Insecta; Polyneoptera; ?Nugonioneuridae; early late Carboniferous; diversity

27

28 **Introduction**

29 For a Moscovian outcrop, The Konservat-Lagerstätte of Avion (Northern France) is especially
30 rich in polyneopteran insects of the superorder Archaeorthoptera (so-called ‘orthopteroid’
31 insects), with no less than eight genera and nine species: *Aviogramma gracilis* Prokop et al.,
32 2014 (plus another Caloneurodea), *Aviocladus pectinatus* Prokop et al., 2014, *Aviohapaloptera*
33 *bethouxi* Prokop et al., 2014, *Aviologus duquesnei* Coty et al., 2014, *Cacurgus avionensis*
34 Schubnel et al., 2020, *Beloatta duquesnei* Nel et al., 2021, cf. *Tococladus* sp. (Coty et al. 2014;
35 Prokop et al. 2014; Schubnel et al. 2020; Nel et al. 2020).

36 Here we describe a further new genus and species belonging to a family different of
37 those previously recorded, confirming the impressive diversity of the clade in this outcrop.

38

39 **Material and methods**

40 The fossil was found by one of us (P.R.) in the slag heap of Avion. It was examined under a
41 Nikon SMZ1500 and photographed with an AmScope camera MU900.

42 We follow the wing venation terminology of Béthoux and Nel (2002).

43 Wing vein terminology: CuA, cubitus anterior; CuP, cubitus posterior; CuPa, anterior branch
44 of CuP; CuPb, posterior branch of CuP; M, median vein; PCu, postcubital vein; RA, radius
45 anterior; RP, radius posterior; ScP, subcosta posterior.

46

47 *Systematic palaeontology*

48 Superorder Archaeorthoptera Béthoux and Nel, 2002

49 Family ?Nugonioneuridae Carpenter, 1976

50 *Avionugonioneura* gen. nov.

51 Zoobank xxxx

52

53 *Type species*

54 *Avionugonioneura jouaulti* sp. nov.

55

56 *Diagnosis*

57 Forewing characters only. CuPa ending into M+CuA and not in a free CuA; no CuPa β , vein
58 M+CuA+CuPa anteriorly pectinate; ScP ending into costa; short posterior branches of
59 CuA+CuPa; five simple branches of RP; M simple and strictly parallel to branches of RP and
60 to first anterior branch of CuA+CuPa; area between RA and RP broad with two rows of cells.

61

62 *Etymology*

63 Named after the outcrop of Avion and the genus name *Avionugonioneura*. Gender feminine.

64

65 *Avionugonioneura jouaulti* sp. nov.

66 (Fig. 1)

67 Zoobank xxxx

68

69 *Etymology*

70 Named after our friend and colleague Corentin Jouault, for his contribution to
71 Palaeoentomology.

72

73 *Type material*

74 MNHN.F.A71353 (two nearly complete forewings overlapped at rest), stored at the Muséum
75 National d'Histoire Naturelle, Paris, France.

76

77 *Type locality*

78 ‘Terril N°7’, Avion, Pas-de-Calais, France.

79

80 *Stratigraphic occurrence*

81 Moscovian (Westphalian C/D equivalent to Bolsovian/Asturian).

82

83 *Diagnosis*

84 As for the genus by monotypy.

85

86 *Description*

87 Forewing elongate, 16.0 mm long, 5.0 mm wide; ScP reaching anterior wing margin at 8.7 mm
88 from wing base; costal area rather narrow, 0.6 mm wide with no visible veinlet; a series of
89 simple short crossveins in area between ScP and R; RP separating from RA 4.4 mm from wing
90 base; area between RA and RP broad, 0.7 mm wide, with two rows of cells in its distal part,
91 apically narrower; RA with a series of short, more or less curved anterior veinlets between it
92 and anterior wing margin; apex of RA 2.4 mm of wing apex; RP with five posterior branches,
93 three of them being forked again distally, parallel to M; RA and RP approximate in apical parts;
94 a common stem R+M+CuA; M+CuA separating from R 2.5 mm from wing base, straight; M
95 distally simple and straight, separating from CuA+CuPa 0.4 mm distal of point of fusion of
96 CuPa with M+CuA; CuA+CuPa with at least an anterior simple branch parallel to M, and two
97 weak posterior subapical branches; concave CuPa curved, elongate, 2.0 mm long, ending into
98 M+CuA; concave CuPb straight and simple in its preserved part; convex PCu basally curved
99 and simple in its preserved part; anal veins not preserved.

100

101 **Discussion**

102 This fossil is a pair of forewings in life position. It belongs to the Archaeorthoptera because of
103 the following characters (Béthoux and Nel 2002): CuP differentiated into concave CuPa and
104 CuPb; a common stem R+M+CuA, from which M+CuA separates distally; CuPa ending into
105 M+CuA. *Avionugonioneura* gen. nov. does not fall into the Panorthoptera because CuPa is not
106 differentiated into two branches CuPa α and CuPa β ; MA1 and MA2 cannot be differentiated. It
107 has a very peculiar vein M+CuA+CuPa, with two strong and simple anterior branches, which
108 is rarely found among the Archaeorthoptera but also present in the Paoliida for CuA only
109 (Prokop et al. 2014a). Similar pectinations are found in the Cnemidolestodea sensu Béthoux
110 (2005), but with the important difference in the first branch being itself forked several times.
111 Another difference to the members of Cnemidolestodea is the vein CuPa ending into M+CuA
112 in *Avionugonioneura* gen. nov., while it ends into CuA in the Cnemidolestodea. Two
113 Cnemidolestodea are already known from Avion, viz. *Aviocladus* Prokop et al., 2014 that
114 differs from *Avionugonioneura* gen. nov. in ScP ending into R, and base of RP in a much distal
115 position (Prokop et al. 2014b); and *Piesbergopterus avionensis* Nel and Roques, 2021 that
116 differs from *Avionugonioneura* gen. nov. in the presence of a specialized area between an
117 anterior branch of M and a posterior branch of the same vein (Nel and Roques, 2021). Few
118 cases of Archaeorthoptera with CuPa ending into M+CuA and not in a free CuA are known. It
119 is the case for *Bruaylogus magnificus* Coty et al., 2014, but this taxon has a CuPa β as
120 panorthopteran synapomorphy, and no anterior pectination of CuA+CuPa (Coty et al. 2014; Gu
121 et al. 2017). *Nugonioneura* Tillyard, 1937 (*N. problematica* Tillyard, 1937 known from the
122 Lower Permian of Elmo in Kansas, U.S.A.) (Tillyard 1937; Aristov 2020, p. 6, Figs 3, 4) and
123 *Contracladus impar* Dvořák et al., 2021 (Pennsylvanian of Piesberg, Germany) also have no
124 CuPa β , a simple M anteriorly emerging from vein M+CuA+CuPa, and an anterior branch of
125 CuA+CuPa parallel to M, as in *Avionugonioneura* gen. nov. Both differ from

126 *Avionugonioneura* gen. nov. in ScP ending into RA, the presence of an angle in M below base
127 of RP so that these two veins approximate, no posterior branch of CuA+CuPa, shorter anterior
128 branches of CuA+CuPa. *Nugonioneura* has only three simple branches of RP instead of five
129 with some being forked again more distally in *Avionugonioneura* gen. nov. and *Contracladus*.
130 M is distally forked in *Contracladus*, instead of being simple as in *Avionugonioneura* gen. nov.
131 and *Nugonioneura*.

132 As a conclusion, *Contracladus*, *Nugonioneura*, and *Avionugonioneura* gen. nov. can
133 only be considered as Archaeorthoptera of uncertain affinities. A new phylogenetic analysis
134 will be necessary to clarify their positions, even if they possibly all belong to the same
135 archaeorthopteran family Nugonioneuridae that would be characterized by the putative
136 apomorphies: ‘CuPa ending into M+CuA’, a ‘simple M anteriorly emerging from
137 M+CuA+CuPa’, and ‘presence of a simple anterior branch of CuA+CuPa’.

138

139 Remarks. Coty et al. (2014: 464, fig. 1) described and figured a wing fragment they attributed
140 to a ‘cf. *Tococladus* sp.’ (Cnemidolestodea Tococladidae). It strongly resemble the wing apex
141 of *Avionugonioneura* gen. nov. in the number of branches of RP, simple M and anterior
142 branches of CuA+CuPa, and RA and RP strongly approximating distally. The only difference
143 is the area between RA and RP narrower in the putative ‘cf. *Tococladus* sp.’ than in
144 *Avionugonioneura* gen. nov. Numerous Palaeozoic wings of Polyneoptera have very similar
145 patterns of venations in their distal parts. They can be discriminated and attributed to precise
146 clades only on the basis of the basal halves of the wings.

147 Rasnitsyn and Aristov (2021: 151) indicated that in *Tococladus garrici* Béthoux et al., 2003,
148 the vein ‘CuPa’ is convex, after the original photograph in Béthoux et al. (2003: fig. 4),
149 suggesting that the attribution of the Tococladidae to the Archaeorthoptera could be not well-
150 grounded. But the original photograph shows a CuPa of ‘intermediate’ convexity compared to

151 those of the convex M+CuA and the concave stem of CuP. Also Béthoux et al. (2003) did not
152 indicate anything about the convexity of this vein. This uncertainty in the determination of
153 the convexity of the vein CuPa is due to fossilization, but it is clear that this vein has a diameter
154 and shape different of the more distal vein CuA+CuPa, supporting that it is a branch of CuP
155 and not CuA. Thus the argument of Rasnitsyn and Aristov (2021) does not stand.

156

157 **Acknowledgements**

158 We sincerely thank Mr. Stéphane Carlier and Mr. Patrick Wypych, Eiffage Route Nord Est, for
159 the kind authorization to collect fossil insects in the terril of Avion.

160

161 **References**

162 Aristov DS. 2020. Revision of the genera *Heteroptilon*, *Nugonioneura* and *Opistocladus* from
163 the Lower Permian of USA (Insecta: Cnemidolestida: Tillyardembiidae, Nugonioneuridae
164 and Parmapteridae). *Far Eastern Entomologist* 401:1–9.

165 Béthoux, O. 2005. Cnemidolestodea (Insecta): an ancient order reinstated. *Journal of*
166 *Systematic Palaeontology* 3:403–408. <https://doi.org/10.1017/S147720190500163X>

167 Béthoux O, Nel A. 2002. Venation pattern and revision of Orthoptera sensu nov. and sister
168 groups. *Phylogeny of Palaeozoic and Mesozoic Orthoptera sensu nov.* *Zootaxa* 96:1–88.
169 <https://doi.org/10.11646/zootaxa.96.1.1>

170 Béthoux O, Nel A, Galtier J, Lapeyrie J, Gand G. 2003. A new species of Tococladidae
171 Carpenter, 1966 from the Permian of France (Insecta: Archaeorthoptera). *Geobios* 36:275–
172 283. [https://doi.org/10.1016/S0016-6995\(03\)00030-5](https://doi.org/10.1016/S0016-6995(03)00030-5)

173 Carpenter FM. 1976. The Lower Permian insects of Kansas. Part 12. Protorthoptera
174 (continued), Neuroptera, additional Palaeodictyoptera, and families of uncertain position.
175 *Psyche* 83:336–376. <https://doi.org/10.1155/1976/932123>

176 Coty D, Háva J, Prokop J, Roques P, Nel A. 2014. New archaeorthopteran insects from the Late
177 Carboniferous of the Nord and Pas-de-Calais basins in northern France (Insecta:
178 Cnemidolestodea, Panorthoptera). *Zootaxa* 3878:462–470.
179 <https://doi.org/10.11646/zootaxa.3878.5.4>

180 Dvořák T, Pecharová M, Leipner A, Nel A, Prokop J. 2021. New archaeorthopteran insects
181 from the Pennsylvanian of Piesberg reveal unexpected mosaic of morphological traits and
182 colouration pattern of the tegmina. *Historical Biology*
183 <https://doi.org/10.1080/08912963.2020.1867127>

184 Gu J-J, Béthoux O, Ren D. 2017. A new, rare and distinctive species of Panorthoptera (Insecta,
185 Archaeorthoptera) from the Upper Carboniferous of Xiaheyuan (Ningxia, China). *Fossil*
186 *Record* 20:253–257. <https://doi.org/10.5194/fr-20-253-2017>

187 Nel A, Garrouste R, Roques P. 2020. The first representative of the archaeorthopteran family
188 Eoblattidae in the Konservat-Lagerstätte of Avion (France) (Insecta: Polyneoptera).
189 *Palaeoentomology* 3:552–555. <https://doi.org/10.11646/palaeoentomology.3.6.3>

190 Nel A, Roques P. 2021. The second species of the cnemidolestid genus *Piesbergopterum* from
191 the Upper Carboniferous of Avion, Northern France (Archaeorthoptera: Cnemidolestidae).
192 *Palaeoentomology* 4:323–325. <https://doi.org/10.11646/palaeoentomology.4.4.6>

193 Prokop J, Krzeminski W, Krzeminska E, Hörnschemeyer T, Ilger J-M, Brauckmann C,
194 Grandcolas P, Nel A. 2014a. Late Palaeozoic Paoliida is the sister group of Dictyoptera
195 (Insecta: Neoptera). *Journal of Systematic Palaeontology* 12:601–622.
196 <https://doi.org/10.1080/14772019.2013.823468>

197 Prokop J, Roques P, Nel A. 2014b. New non-holometabolous insects from Pennsylvanian of
198 Avion locality in Pas-de-Calais, France (Insecta: ‘Exopterygota’). *Alcheringa* 38:155–169.
199 <https://doi.org/10.1080/03115518.2014.848620>

200 Rasnitsyn AP, Aristov DS. 2021. Review of the Family Tococladidae Carpenter (Insecta:
201 Eoblattida) from the Permian of North America and Europe. Paleontological Journal
202 55:151–156. <https://doi.org/10.1134/S003103012102012X>

203 Schubnel T, Roberts D, Roques P, Garrouste R, Desutter-Grandcolas L, Nel A. 2020.
204 Moscovian fossils shed light on the enigmatic polyneopteran families Cacurgidae and
205 Eoblattidae (Insecta: ‘Eoblattida’, Archaeorthoptera). Journal of Systematic Palaeontology
206 18:499–511. <https://doi.org/10.1080/14772019.2019.1627595>

207 Tillyard RJ. 1937. Kansas Permian insects. 17. The order Megasecoptera and additions to the
208 Palaeodictyoptera, Odonata, Protoperlaria, Copeognatha and Neuroptera. American
209 Journal of Science (5) 33:81–110. <https://doi.org/10.2475/ajs.s5-33.194.81>

210

211 Figure 1. *Avionugonioneura jouaulti* gen. et sp. nov., holotype MNHN.F.A71353. (A)
212 photograph of habitus; (B) forewing reconstruction, arrowhead: M+CuA, arrow: CuPa. Scale
213 bars = 1 mm.

214

