



HAL
open science

Denseness results in the theory of algebraic fields

Sylvy Anscombe, Philip Dittmann, Arno Fehm

► **To cite this version:**

Sylvy Anscombe, Philip Dittmann, Arno Fehm. Denseness results in the theory of algebraic fields. *Annals of Pure and Applied Logic*, 2019, 172 (8), pp.102973. 10.1016/j.apal.2021.102973 . hal-03373671

HAL Id: hal-03373671

<https://hal.sorbonne-universite.fr/hal-03373671>

Submitted on 11 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

DENSENESS RESULTS IN THE THEORY OF ALGEBRAIC FIELDS

SYLVY ANSCOMBE, PHILIP DITTMANN AND ARNO FEHM

ABSTRACT. We study when the property that a field is dense in its real and p -adic closures is elementary in the language of rings and deduce that all models of the theory of algebraic fields have this property.

1. INTRODUCTION

The completions of the field of rational numbers are \mathbb{R} and the \mathbb{Q}_p for prime numbers p , and completions of number fields are accordingly finite extensions of \mathbb{R} and \mathbb{Q}_p . The importance of these completions in number theory (for example in the form of local-global principles) and the success of local algebra has motivated the attempt to apply these concepts also to other fields. Here the problem arises that the completion of an ordered field or a (suitably) valued field need in general not share any of the crucial properties of \mathbb{R} or the finite extensions of the \mathbb{Q}_p . This has led to the definitions of real closures and p -adic closures (of p -rank 1, corresponding to \mathbb{Q}_p , and of higher p -rank corresponding to finite extensions of \mathbb{Q}_p) of arbitrary fields with orderings respectively so-called p -valuations. But here the converse problem arises: These closures are not complete, but do they at least satisfy other crucial properties of the completions - in particular, is the field dense in its closures?

This question has been answered positively for several classes of fields, in particular for fields that satisfy a local-global principle for rational points on varieties: Prestel [Pre81] shows that pseudo-real closed (PRC) fields are dense in all their real closures, and the analogue for pseudo- p -adically closed (PpC) fields was proven in [Gro87]. The most general result in this direction is by Pop [Pop03], who proves that arbitrary so-called pseudo classically closed (PCC) fields are dense in all their real closures and all their p -adic closures (of arbitrary p -rank). Ershov [Ers92] has a similar result for so-called RC_π -fields.

In this note we study when denseness in real and p -adic closures is an elementary property. We show that this is not always the case (Section 7), but we also prove some general principles that allow us to exhibit a large class of fields dense in all their real and p -adic closures, and which likely includes all the classes listed in the previous paragraph:

Theorem 1.1. *Every model of the theory of algebraic fields of characteristic zero is dense in all its real closures and all its p -adic closures (of arbitrary p -rank). Equivalently (see Proposition 3.4), the completion of any such model with respect to any ordering is real closed, and the completion of any such model with respect to any p -valuation is p -adically closed.*

Here, by an algebraic field of characteristic zero we mean an algebraic extension of \mathbb{Q} . Note that this result applies in particular to elementary extensions of number fields and even non-standard number fields in the most general sense (see [Che75]). By results of Jarden [Jar88, Jar91] every PRC field and every PpC field is a model of the theory of

algebraic fields and this suggests that also arbitrary PCC fields might be of that kind. The only other result about models of the theory of algebraic fields we are aware of concerns their cohomological dimension [Cha90]. To appreciate the non-triviality of Theorem 1.1 note that algebraic fields are dense in their henselization with respect to *any* valuation (not just with respect to p -valuations), which however no longer holds true for arbitrary models of the theory of algebraic fields, see Remark 6.12.

While Theorem 1.1 is interesting for its own sake, our motivation for it came from a concrete application in the area of definable valuation rings based on [AF17], as we explain in Section 6. As one consequence we obtain:

Corollary 1.2. *Let F be an algebraic extension of \mathbb{Q} properly contained in the algebraic part of \mathbb{R} or of some \mathbb{Q}_p . Then there exists an existential formula $\phi(x)$ in the language of rings which defines in $F((t))$ the valuation ring $F[[t]]$. Equivalently, there exists a polynomial $g \in \mathbb{Z}[X_1, \dots, X_n]$ such that the projection of the zero set of g in $F((t))$ onto the first coordinate is precisely $F[[t]]$.*

Although Theorem 1.1 is a result in the model theory of fields, its proof relies heavily on recent number theoretic results: While in the case of orderings it can build on Euler's well-known Four Squares Theorem and Siegel's generalization to number fields, which implies the finiteness of the Pythagoras number for algebraic fields, in the p -adic case we make use of the recent p -adic analogue of Siegel's theorem [ADF19b] for the so-called p -Pythagoras number, where the squares are replaced by a certain rational function, the Kochen operator. The other crucial observation involved is that for algebraic fields, denseness in real or p -adic closures implies a certain uniform denseness statement, which then allows an axiomatization. In fact we show more generally that denseness in real or p -adic closures transfers to elementarily equivalent fields as soon as Pythagoras number respectively p -Pythagoras number are finite. In this generality the equivalence of denseness and uniform denseness is less obvious and builds on our approximation theorems from [ADF19a].

2. STATEMENT OF GENERAL RESULT

We recall definitions from [Feh13] that we will use, see also [PR84] for basic notions regarding p -valuations. Let F be a field of characteristic 0. A **prime** \mathfrak{P} of F is either an ordering on F , in which case we also denote it by $\leq_{\mathfrak{P}}$, or an equivalence class of p -valuations on F (of arbitrary p -rank), in which case we write $v_{\mathfrak{P}}$ for a representative of \mathfrak{P} which has \mathbb{Z} as smallest non-trivial convex subgroup of the value group. However, for brevity, we will freely say ‘ \mathfrak{P} is a p -valuation’ to mean that \mathfrak{P} is an equivalence class of p -valuations. We denote by $\mathcal{O}_{\mathfrak{P}}$ the valuation ring of $v_{\mathfrak{P}}$, if \mathfrak{P} is a p -valuation, or the positive cone of $\leq_{\mathfrak{P}}$, if \mathfrak{P} is an ordering.¹ The set of all primes of F is denoted $\mathcal{S}(F)$. For example, we write $\mathcal{S}(\mathbb{Q}) = \{2, 3, \dots, \infty\}$. A **closure** of (F, \mathfrak{P}) is a real closure if \mathfrak{P} is an ordering, or is a p -adic closure if \mathfrak{P} is a p -valuation. Thus the closure of (F, \mathfrak{P}) is unique up to isomorphism over F if and only if \mathfrak{P} is an ordering or \mathfrak{P} is a p -valuation such that $v_{\mathfrak{P}}(F^\times)$ is a \mathbb{Z} -group (see Section 5 for the definition of a \mathbb{Z} -group). Note also that every real closed field and every p -adically closed field carries a unique prime, which induces a canonical topology.

We consider two conditions on a set $S \subseteq \mathcal{S}(F)$:

Denseness is the condition

¹Note that in the case of orderings this definition agrees with [Feh13] but disagrees with a definition we made in [ADF19a], where for an ordering \mathfrak{P} we put $\mathcal{O}_{\mathfrak{P}} = \{x \in F : -1 \leq_{\mathfrak{P}} x \leq_{\mathfrak{P}} 1\}$.

D_S For every monic $g \in F[X]$ and all $\mathfrak{P} \in S$ for which g has a zero in some closure of (F, \mathfrak{P}) , for every $a \in F^\times$ there exists $x \in F$ with $1 - g(x)^2 a^{-2} \in \mathcal{O}_{\mathfrak{P}}$.

Uniform denseness is the condition

UD_S For every monic $g \in F[X]$ and every $a \in F^\times$ there exists $x \in F$ such that for all $\mathfrak{P} \in S$ for which g has a zero in some closure of (F, \mathfrak{P}) , we have $1 - g(x)^2 a^{-2} \in \mathcal{O}_{\mathfrak{P}}$.

Lemma 2.1. UD_S implies D_S .

Proof. This is trivial. \square

The following lemma is a variant of [Ers01, Proposition 1.6.1]. For the proof we use the *completion* of a field with respect to a valuation or ordering. This can be defined in terms of transfinite Cauchy sequences, see [EP05, Theorem 2.4.3] respectively [Hau67] (written $\overline{F}_{X(F)}$ in the latter). Many alternative definitions are also available in the literature, see for instance [Bou07, III §6, No 8] and [Bae70].

Lemma 2.2. F satisfies D_S if and only if F is dense in every closure at elements of S .

Proof. Let $\mathfrak{P} \in S$. First note that if \mathfrak{P} is a p -valuation, then $1 - g(x)^2 a^{-2} \in \mathcal{O}_{\mathfrak{P}}$ if and only if $v_{\mathfrak{P}}(g(x)) \geq v_{\mathfrak{P}}(a)$, and if \mathfrak{P} is an ordering, then $1 - g(x)^2 a^{-2} \in \mathcal{O}_{\mathfrak{P}}$ if and only if $|g(x)| \leq_{\mathfrak{P}} |a|$. Now let (F', \mathfrak{P}') be a closure of (F, \mathfrak{P}) . Of course, if F is dense in F' , then any zero $y \in F'$ of $g \in F[X]$ can be approximated arbitrarily well by elements $x \in F$.

Conversely, we suppose that F satisfies D_S . If an irreducible polynomial $g \in F[X]$ has a zero y in F' then we may find $x \in F$ with $v_{\mathfrak{P}}(g(x))$ arbitrarily large if \mathfrak{P} is a p -valuation, or with $|g(x)|$ arbitrarily small if \mathfrak{P} is an ordering. Then g has a zero in the completion \hat{F} of F with respect to \mathfrak{P} : this follows from [EP05, Theorem 2.4.5] in the case of valuations, and is proved analogously in the case of orderings, by factoring g into linear and quadratic factors over a real closure of \hat{F} . Since this holds for all such g , and F'/F is algebraic, there is an F -embedding of F' into \hat{F} , see [FJ08, Lemma 20.6.3(a)]. As F' carries a unique prime, this embedding is in fact an isometry, and so F is dense also in F' . \square

The main idea of the proof of the generalization of Theorem 1.1 is that for sufficiently nice sets of primes S , D_S in fact implies UD_S , which in turn is an elementary statement given a predicate for the holomorphy domain:

Fix a number field K and a pair $\tau = (e, f) \in \mathbb{N}^2$, where \mathbb{N} is the set of positive integers. For a prime \mathfrak{p} of K and an extension F/K , we denote by $\mathcal{S}_{\mathfrak{p}}^{\tau}(F)$ the set of primes of F lying above \mathfrak{p} , where in the case of p -valuations we restrict to relative initial ramification e' at most e and relative residue degree f' dividing f . The pair $\tau' = (e', f')$ is called the **relative type** of \mathfrak{P} , and we write $\tau' \leq \tau$ if and only if $e' \leq e$ and $f' | f$. The **holomorphy domain** corresponding to $\mathcal{S}_{\mathfrak{p}}^{\tau}(F)$ is

$$R_{\mathfrak{p}}^{\tau}(F) = \bigcap_{\mathfrak{P} \in \mathcal{S}_{\mathfrak{p}}^{\tau}(F)} \mathcal{O}_{\mathfrak{P}}.$$

We adopt the convention that the relative type $\tau = (1, 1)$ can always be omitted from any notation. We write $\mathcal{S}_{\mathfrak{p}}^{\tau, \mathbb{Z}}(F)$ for the set of those $\mathfrak{P} \in \mathcal{S}_{\mathfrak{p}}^{\tau}(F)$ that are either orderings, or are p -valuations for which $v_{\mathfrak{P}}(F^\times)$ is a \mathbb{Z} -group. We write $\mathcal{S}_{\mathfrak{p}}^{=\tau}(F)$ for the set of those $\mathfrak{P} \in \mathcal{S}_{\mathfrak{p}}^{\tau}(F)$ that are either orderings, or are p -valuations of exact relative type τ . For $t, s \in F$ we denote by $\mathcal{S}_{\mathfrak{p}}^{=\tau}(F, t, s)$ the set of those $\mathfrak{P} \in \mathcal{S}_{\mathfrak{p}}^{=\tau}(F)$ that are either orderings, or are p -valuations for which t is a uniformizer and s is a unit whose residue generates the multiplicative group of the residue field. Finally, we let $\mathcal{S}_{\mathfrak{p}}^{=\tau, \mathbb{Z}}(F, t, s) = \mathcal{S}_{\mathfrak{p}}^{=\tau}(F, t, s) \cap \mathcal{S}_{\mathfrak{p}}^{\tau, \mathbb{Z}}(F)$.

To state the general results we denote by $\mathcal{L}_{\text{ring}} = \{+, \cdot, 0, 1\}$ the language of rings and we define $\mathcal{L}_R = \mathcal{L}_{\text{ring}} \cup \{R\}$, where R is a unary predicate, as well as expansions $\mathcal{L}_{\text{ring}}(K)$ and $\mathcal{L}_R(K)$ by constants for the elements of K . Then the following holds:

Proposition 2.3. $D_{\mathcal{S}_p^\tau(F)}$ holds if and only if $\text{UD}_{\mathcal{S}_p^{\tau'}(F,t,s)}$ holds for every $t, s \in F$ and every $\tau' \leq \tau$.

Proposition 2.4. There exists an $\mathcal{L}_R(K)$ -theory T_p^τ such that $(F, R_p^\tau(F)) \models T_p^\tau$ if and only if $\text{UD}_{\mathcal{S}_p^{\tau'}(F,t,s)}$ holds for every $t, s \in F$ and every $\tau' \leq \tau$.

Corollary 2.5. F is dense in every closure at elements of $\mathcal{S}_p^\tau(F)$ if and only if $(F, R_p^\tau(F)) \models T_p^\tau$.

We prove Proposition 2.3 in Section 4 and Proposition 2.4 in Section 5. In the case of algebraic fields (the setting of Theorem 1.1) we obtain an $\mathcal{L}_{\text{ring}}(K)$ -theory, since the holomorphy domain is uniformly definable. This part is carried out in Section 6, where we also draw some consequences of the results. In Section 7 we show that the \mathcal{L}_R -theory T_∞ cannot be replaced by an $\mathcal{L}_{\text{ring}}$ -theory with the same property. We start in Section 3 with some purely algebraic characterizations of denseness conditions.

3. DENSENESS IN HENSELIZATIONS, p -ADIC CLOSURES AND REAL CLOSURES

Let F be a field. For a valuation v on F , we write \mathcal{O}_v for its valuation ring, vF for its value group, and Fv for its residue field. A pair of valuations on F is **independent** if they induce distinct topologies on F . Likewise for a pair of orderings, or a pair consisting of an ordering and a valuation. In particular, this applies to primes of F . A valuation u on F is a **refinement** of v , and v is a **coarsening** of u , if $\mathcal{O}_u \subseteq \mathcal{O}_v$. If u is a coarsening or refinement of v , then u and v are **comparable**. Note that a p -valuation and a q -valuation, for primes p and q , are never comparable as soon as they are distinct.

Lemma 3.1. Let v be a valuation on F . The following are equivalent:

- (1) F is dense in a henselization of (F, v) (with respect to the valuation topology)
- (2) For every non-trivial coarsening w of v , the valuation \bar{v} induced by v on the residue field Fw is henselian.
- (3) For any separable monic polynomial $g \in \mathcal{O}_v[X]$ for which there exists $a_0 \in \mathcal{O}_v$ with $v(g(a_0)) > 0$ and $v(g'(a_0)) = 0$, the set $\{v(g(a)) : a \in F, g(a) \neq 0\}$ is unbounded in vF .

Proof. See [Ers01, Theorem 1.6.1] and [Ers01, Proposition 1.6.1]. □

Lemma 3.2. Let F' be a field extension of F with a valuation v' extending v . If F is dense in (F', v') , then $F'v' = Fv$ and $v'F' = vF$.

Proof. Given $x \in \mathcal{O}_{v'}$, any $y \in F$ with $v'(x - y) > 0$ has the same residue as x ; this proves the equality of residue fields. Given $x \in (F')^\times$, any $y \in F$ with $v'(x - y) > v'(x)$ has the same valuation as x ; this proves the equality of value groups. □

Lemma 3.3. Let v_1, v_2 be valuations on F . Suppose that F is dense in henselizations of both (F, v_1) and (F, v_2) . Then

- (1) v_1 and v_2 are comparable, or
- (2) v_1 and v_2 are independent, or
- (3) both Fv_1 and Fv_2 are algebraically closed.

Proof. Suppose that v_1 and v_2 are incomparable and dependent. Then both v_1 and v_2 are non-trivial. Let w be the finest common coarsening of v_1 and v_2 . Since v_1 and v_2 are dependent, w is also non-trivial, see [EP05, Theorem 2.3.4]. Since v_1 and v_2 are incomparable, w is a proper coarsening of both v_1 and v_2 . Thus v_1 and v_2 induce independent non-trivial valuations \bar{v}_1 and \bar{v}_2 on Fw . By (1) \Rightarrow (2) of Lemma 3.1, both \bar{v}_1 and \bar{v}_2 are henselian. Therefore, by F. K. Schmidt's theorem [EP05, Theorem 4.4.1], Fw is separably closed, and so both Fv_1 and Fv_2 are algebraically closed, see [EP05, Theorem 3.2.11]. \square

Let F now be a field of characteristic 0. Given a valuation u and an ordering \leq on F , we say that u is \leq -**convex** if its valuation ring \mathcal{O}_u is a convex subset of F with respect to \leq . We may now say that u is a **coarsening** of a prime $\mathfrak{P} \in \mathcal{S}(F)$ if u is a coarsening of $v_{\mathfrak{P}}$, in the usual sense for valuations, if \mathfrak{P} is a p -valuation, or if u is $\leq_{\mathfrak{P}}$ -convex, if \mathfrak{P} is an ordering.

Given an ordering \leq on F , we denote by v_{\leq} the valuation whose valuation ring is the convex hull of the prime field \mathbb{Q} with respect to \leq . Thus v_{\leq} is the finest valuation which is \leq -convex, and u is \leq -convex if and only if u is a coarsening of v_{\leq} . Note that \leq is archimedean if and only if v_{\leq} is the trivial valuation. Also note that if \leq is non-archimedean then \leq and v_{\leq} are dependent, whereas if \leq is archimedean then it is independent from all valuations and all other orderings.

If u is \leq -convex, then \leq induces an ordering on Fu , which we denote by \leq_u . In general, \leq_u does not uniquely determine \leq : by the Baer–Krull theorem [EP05, Theorem 2.2.5], there is a family of orderings on F each of which induces \leq_u on Fu .

We can now give an equivalent statement of denseness conditions in terms of completions.

Proposition 3.4. *Let F be a field.*

- (1) *If v is a valuation on F and (F', v') a henselization of (F, v) , then F is dense in (F', v') if and only if the completion (\hat{F}, \hat{v}) of (F, v) is henselian.*
- (2) *If v is a p -valuation on F and (F', v') a p -adic closure of (F, v) , then F is dense in (F', v') if and only if the completion (\hat{F}, \hat{v}) of (F, v) is p -adically closed.*
- (3) *If \leq is an ordering on F and (F', \leq') a real closure of (F, \leq) , then F is dense in (F', \leq') if and only if the completion $(\hat{F}, \hat{\leq})$ of (F, v) is real closed.*

Proof. For the first point, observe that if the completion (\hat{F}, \hat{v}) is henselian, then the henselization (F', v') embeds into \hat{F} , and therefore F is dense in it. For the converse direction we imitate an argument from the proof of [Ers01, Proposition 1.6.1]: if (F, v) is dense in its henselization, then (\hat{F}, \hat{v}) is in turn dense in its own henselization, as the condition from Lemma 3.1(3) transfers from (F, v) to (\hat{F}, \hat{v}) by approximating a separable polynomial in $\mathcal{O}_{\hat{v}}[X]$ by a separable polynomial in $\mathcal{O}_v[X]$. By completeness, this means that (\hat{F}, \hat{v}) must already be henselian.

The second point follows from the first by observing that by Lemma 3.2 F can only be dense in a p -adic closure if vF is a \mathbb{Z} -group, in which case the henselization of (F, v) is already a p -adic closure.

The third point is immediate from [Hau67, Satz 13, Satz 14]. \square

For lack of a convenient reference, we give a proof of the following simple lemma.

Lemma 3.5. *Let F'/F be any field extension.*

- (1) *If v' is a valuation on F' prolonging a valuation v on F , and if w is a coarsening of v , then there is a coarsening w' of v' which is a prolongation of w .*

- (2) If \leq' is an ordering on F' extending an ordering \leq on F , and if w is a coarsening of \leq , then there is a coarsening w' of \leq' which is a prolongation of w .

Proof. Recalling the correspondence between coarsenings of v and convex subgroups of vF , we denote by Δ the subgroup corresponding to w . In the analogous correspondence for coarsenings of v' , we choose w' to correspond to the convex hull of Δ in $v'F'$. Then w' is both a coarsening of v' and a prolongation of w . This proves (1). For (2), we apply (1) in the case $v = v_{\leq}$ and $v' = v_{\leq'}$. \square

Lemma 3.6. *Let \leq be an ordering on F and let v be a \leq -convex valuation. Then (F, \leq) is real closed if and only if*

- (1) vF is divisible,
- (2) (Fv, \leq_v) is real closed, and
- (3) (F, v) is henselian.

Proof. See [EP05, Theorem 4.3.7]. \square

Lemma 3.7. *Let \leq be an ordering on F and let v be a non-trivial \leq -convex valuation. Then F is dense in a real closure of (F, \leq) if and only if*

- (1) vF is divisible,
- (2) (Fv, \leq_v) is real closed, and
- (3) F is dense in a henselization of (F, v) .

Proof. (\implies) Let (F', \leq') be a real closure of (F, \leq) , and let (F^h, v^h) be a henselization of (F, v) . By Lemma 3.5(2), there is a valuation v' on F' which is a prolongation of v and which is a coarsening of $v_{\leq'}$, i.e. v' is \leq' -convex. As F is dense in (F', \leq') and v' induces the same topology as \leq' , F is also dense in (F', v') , in particular (by Lemmas 3.2 and 3.6) $Fv = F'v'$ is real closed and $vF = v'F'$ is divisible. Also by Lemma 3.6, (F', v') is henselian, hence (F^h, v^h) embeds into (F', v') over (F, v) , so F is dense also in (F^h, v^h) .

(\impliedby) By the Baer–Krull theorem, and since vF is divisible, \leq is the unique ordering on F which induces \leq_v on Fv . Any henselization (F^h, v^h) of (F, v) is an immediate extension, hence the value group $v^h F^h = vF$ is still divisible, and the residue field is still Fv . By the Baer–Krull theorem, there is a unique ordering \leq^h on F^h which induces \leq_v on Fv and obviously extends \leq . As \leq^h and v^h induce the same topology on F^h , F is dense in (F^h, \leq^h) . Applying Lemma 3.6 and assumptions (1) and (2), in fact (F^h, \leq^h) is real closed, hence is a real closure of (F, \leq) . \square

The next proposition is an analogue of Lemma 3.1(2) for non-archimedean orderings.

Proposition 3.8. *Let \leq be an ordering on F . Then F is dense in a real closure of (F, \leq) if and only if for all non-trivial \leq -convex valuations v on F we have*

- (1) vF is divisible and
- (2) (Fv, \leq_v) is real closed.

Proof. If F is dense in a real closure of (F, \leq) , then (1) and (2) follow from Lemma 3.7. For the converse, we suppose (1) and (2) hold. First observe that if \leq is archimedean then F is dense in every real closure of (F, \leq) . Thus we may suppose \leq to be non-archimedean, and so v_{\leq} is non-trivial. If w is any non-trivial \leq -convex valuation on F (equivalently, any non-trivial coarsening of v_{\leq}), then (Fw, \leq_w) is real closed, by (2). In particular $(Fw, \overline{v_{\leq}})$ is henselian, by Lemma 3.6. Since w was an arbitrary non-trivial coarsening of v_{\leq} , it follows from (2) \implies (1) of Lemma 3.1 that F is dense in a henselization of (F, v_{\leq}) . This verifies condition (3) of Lemma 3.7 for $v = v_{\leq}$. Therefore F is dense in a real closure of (F, \leq) , as required. \square

Proposition 3.9. *Let $S \subseteq \mathcal{S}(F)$. If D_S holds then the elements of S are pairwise independent.*

Proof. Let \mathfrak{P} and \mathfrak{Q} be distinct primes in S , and suppose that F is dense in every closure of (F, \mathfrak{P}) and (F, \mathfrak{Q}) . Note that if \mathfrak{P} is a p -valuation, this implies in particular that F is dense in a henselization of $(F, v_{\mathfrak{P}})$.

As a first case, suppose that \mathfrak{P} is a p -valuation and \mathfrak{Q} is a q -valuation, for prime numbers p and q . Since such valuations are incomparable, and the residue field of a p -valuation is finite, by Lemma 3.3 we have that \mathfrak{P} and \mathfrak{Q} are independent.

Next, suppose that \mathfrak{P} is an ordering and \mathfrak{Q} is a p -valuation. If $\leq_{\mathfrak{P}}$ is archimedean, then it is independent from all valuations and other orderings; in particular it is independent from $v_{\mathfrak{Q}}$. If $\leq_{\mathfrak{P}}$ is non-archimedean, then the finest $\leq_{\mathfrak{P}}$ -convex valuation $v_1 = v_{\leq_{\mathfrak{P}}}$ is non-trivial. By Lemma 3.7, F is dense in a henselization of (F, v_1) , $v_1 F$ is divisible, and (Fv_1, \leq_1) is real closed, where \leq_1 is the ordering induced on Fv_1 by $\leq_{\mathfrak{P}}$. Since a real closed field admits no p -valuations, v_1 and $v_{\mathfrak{Q}}$ are incomparable. Thus, by Lemma 3.3, v_1 and $v_{\mathfrak{Q}}$ are independent, and it follows that \mathfrak{P} and \mathfrak{Q} are independent.

Finally, suppose that \mathfrak{P} and \mathfrak{Q} are both orderings. If either is archimedean, then they are independent; thus we may assume that neither is archimedean, and so both $v_1 = v_{\leq_{\mathfrak{P}}}$ and $v_2 = v_{\leq_{\mathfrak{Q}}}$ are non-trivial. By Lemma 3.7, F is dense in henselizations of (F, v_1) and (F, v_2) , both $v_1 F$ and $v_2 F$ are divisible, and both (Fv_1, \leq_1) and (Fv_2, \leq_2) are real closed, where \leq_1 is the ordering induced on Fv_1 by $\leq_{\mathfrak{P}}$, and likewise \leq_2 is induced on Fv_2 by $\leq_{\mathfrak{Q}}$. We claim that v_1 and v_2 are incomparable. To see this: suppose without loss of generality that v_1 is a (not necessarily proper) refinement of v_2 . Then v_2 is $\leq_{\mathfrak{P}}$ -convex, and so $\leq_{\mathfrak{P}}$ induces an ordering \leq on Fv_2 . Since the real closed field Fv_2 admits a unique ordering, we must have $\leq = \leq_2$. Since $v_2 F$ is divisible, by the Baer–Krull theorem, we have $\leq_{\mathfrak{P}} = \leq_{\mathfrak{Q}}$, i.e. $\mathfrak{P} = \mathfrak{Q}$, which is a contradiction, proving the claim. Since neither Fv_1 nor Fv_2 is algebraically closed, and together with the claim, it follows from Lemma 3.3 that v_1 and v_2 are independent, and therefore \mathfrak{P} and \mathfrak{Q} are independent, as required. \square

4. UNIFORM DENSENESS

Let K be a number field, \mathfrak{p} a prime of K , $\tau = (e, f) \in \mathbb{N}^2$ and F an extension of K . If \mathfrak{p} is a p -valuation let $t_{\mathfrak{p}}$ be a uniformizer of $v_{\mathfrak{p}}$, and let $q_{\mathfrak{p}} = |Kv_{\mathfrak{p}}|$ be the size of the residue field. We equip the set $\mathcal{S}(F)$ of all primes of F with the topology with subsbasis of open-closed sets

$$\{\mathfrak{P} \in \mathcal{S}(F) : a \in \mathcal{O}_{\mathfrak{P}}\}, \quad a \in F.$$

This clearly makes $\mathcal{S}(F)$ a totally disconnected Hausdorff space.

Lemma 4.1. $\mathcal{S}_{\mathfrak{p}}^{\tau}(F)$ is compact.

Proof. [Feh13, Lemma 10.3] states that the subspace topology on $\mathcal{S}_{\mathfrak{p}}^{\tau}(F)$ coincides with the subspace topology obtained by declaring the $\{\mathfrak{P} \in \mathcal{S}(F) : a \in \mathcal{O}_{\mathfrak{P}}\}$ to be a subsbasis of open sets, and that this gives a compact space. \square

Lemma 4.2. Suppose that \mathfrak{p} is a p -valuation, let $\mathfrak{P} \in \mathcal{S}_{\mathfrak{p}}^{\tau}(F)$ and let $t, s \in F$. Then $\mathfrak{P} \in \mathcal{S}_{\mathfrak{p}}^{=\tau}(F, t, s)$ if and only if $t^e t_{\mathfrak{p}}^{-1} \in \mathcal{O}_{\mathfrak{P}}^{\times}$, $s \in \mathcal{O}_{\mathfrak{P}}^{\times}$ and $s^n - 1 \in \mathcal{O}_{\mathfrak{P}}^{\times}$ for all $n | q_{\mathfrak{p}}^f - 1$, $n \neq q_{\mathfrak{p}}^f - 1$.

Proof. This follows directly from the definitions. \square

Lemma 4.3. If \mathfrak{p} is a p -valuation, then for every $t, s \in F$, $\mathcal{S}_{\mathfrak{p}}^{=\tau}(F, t, s)$ is open-closed in $\mathcal{S}_{\mathfrak{p}}^{\tau}(F)$.

Proof. As each of the finitely many conditions in the previous lemma defines an open-closed subset of $\mathcal{S}_p^\tau(F)$, also $\mathcal{S}_p^{\tau, \mathbb{Z}}(F, t, s)$ is open-closed. \square

Next we quote two approximation results from [ADF19a]:

Proposition 4.4. *Suppose that \mathfrak{p} is a p -valuation. Let $S_1, \dots, S_n \subseteq \mathcal{S}_p^\tau(F)$ be disjoint compact sets, and $z_1, \dots, z_n \in F^\times$. If for every valuation v on F coarsening some prime in S_i and some prime in S_j , for some i, j , we have $v(z_i) = v(z_j)$, then there exists a $z \in F^\times$ with $v_{\mathfrak{p}}(z) = v_{\mathfrak{p}}(z_i)$ for all $\mathfrak{P} \in S_i$, for $i = 1, \dots, n$.*

Proof. This is an application of [ADF19a, Corollary 5.1], for which we verify the hypotheses. Condition (U) holds as shown in [ADF19a, Example 3.5(2,3)]. Condition (I) follows from the fact that distinct elements of $\mathcal{S}_p^\tau(F)$ are pairwise incomparable. \square

For the next proposition we will need the notion of a ball with respect to a prime \mathfrak{P} of F : if \mathfrak{P} is a p -valuation we write

$$B_{\mathfrak{P}}(y, z) := \{x \in F : v_{\mathfrak{P}}(x - y) > v_{\mathfrak{P}}(z)\},$$

whereas if \mathfrak{P} is an ordering we write

$$B_{\mathfrak{P}}(y, z) := \{x \in F : |x - y| <_{\mathfrak{P}} |z|\},$$

for $y \in F$ and $z \in F^\times$.

Proposition 4.5. *Let $S \subseteq \mathcal{S}_p^\tau(F)$ be compact, and suppose that the elements of S are pairwise independent. Let $\gamma_1, \dots, \gamma_m \in F(X_1, \dots, X_r)$ and write $\gamma_j = \frac{g_j}{h_j}$ with $g_j, h_j \in F[X_1, \dots, X_r]$. For each $j = 1, \dots, m$, let $y_j \in F$, $z_j \in F^\times$ be given. If for each $\mathfrak{P} \in S$ there exists $x_{\mathfrak{P}} \in F^r$ such that $h_j(x_{\mathfrak{P}}) \neq 0$ and $\gamma_j(x_{\mathfrak{P}}) \in B_{\mathfrak{P}}(y_j, z_j)$ for each j , then there exists $x \in F^r$ such that $h_j(x) \neq 0$ and $\gamma_j(x) \in B_{\mathfrak{P}}(y_j, z_j)$ for each j and $\mathfrak{P} \in S$.*

Proof. This is an application of [ADF19a, Corollary 7.8]. Note that $\mathcal{S}_p^\tau(F)$ satisfies condition (U) of that corollary by [ADF19a, Example 3.5(2,3)]. \square

For a set $S \subseteq \mathcal{S}(F)$ of primes of F and an $\mathcal{L}_R(F)$ -sentence φ we define

$$\varphi(S) := \{\mathfrak{P} \in S : (F, \mathcal{O}_{\mathfrak{P}}) \models \varphi\}$$

and

$$\varphi'(S) := \{\mathfrak{P} \in S : (F', \mathcal{O}_{\mathfrak{P}'}) \models \varphi \text{ for each closure } (F', \mathfrak{P}') \text{ of } (F, \mathfrak{P})\}.$$

Lemma 4.6. *If φ is a quantifier-free $\mathcal{L}_R(F)$ -sentence and $S \subseteq \mathcal{S}(F)$, then $\varphi(S) = \varphi'(S)$ is open-closed in S .*

Proof. Since the family of open-closed sets is closed under boolean operations we can reduce to atomic sentences $a = 0$ and $R(a)$, with $a \in F$. For $a = 0$ the statement is trivial, while for $R(a)$ it follows from the definition of the topology on $\mathcal{S}(F)$. \square

Lemma 4.7. *There exists a recursive map $\varphi \mapsto \varphi_p^\tau$ from $\mathcal{L}_R(K)$ -sentences to universal $\mathcal{L}_R(K)$ -sentences such that for every $\mathfrak{P} \in \mathcal{S}_p^{\tau, \mathbb{Z}}(F)$ and every closure (F', \mathfrak{P}') of (F, \mathfrak{P}) the following are equivalent:*

- (1) $(F', \mathcal{O}_{\mathfrak{P}'}) \models \varphi$
- (2) $(F', \mathcal{O}_{\mathfrak{P}'}) \models \varphi_p^\tau$
- (3) $(F, \mathcal{O}_{\mathfrak{P}}) \models \varphi_p^\tau$

Proof. This is a consequence of model completeness, see [Feh13, Prop. 8.2]. \square

Proposition 4.8. *Let $S \subseteq \mathcal{S}_p^{\tau, \mathbb{Z}}(F)$. Then $\varphi'(S)$ is open-closed in S for every $\mathcal{L}_R(F)$ -sentence φ .*

Proof. By Lemma 4.7 we can find a quantifier-free $\mathcal{L}_R(F)$ -formula $\psi_1(x)$, $x = (x_1, \dots, x_r)$, such that with $\varphi_1 \equiv \forall x \psi_1(x)$ we have $\varphi'(S) = \varphi_1'(S) = \varphi_1(S)$. Thus,

$$\varphi'(S) = \bigcap_{a \in F^r} \psi_1(a)(S).$$

Similarly, we find a quantifier-free $\mathcal{L}_R(F)$ -formula $\psi_2(x)$ with

$$S \setminus \varphi'(S) = (\neg\varphi)'(S) = \bigcap_{a \in F^r} \psi_2(a)(S).$$

By Lemma 4.6, each $\psi_i(a)(S)$ is in particular closed, so $\varphi'(S)$ is open-closed. \square

We are now able to conclude that denseness implies uniform denseness, for certain compact sets of primes.

Proposition 4.9. *For every compact subset $S \subseteq \mathcal{S}_p^{\tau}(F)$, D_S implies UD_S .*

Proof. Suppose that D_S holds. Then S is contained in $\mathcal{S}_p^{\tau, \mathbb{Z}}(F)$ by Lemma 3.2, and the elements of S are pairwise independent by Proposition 3.9. Let $g \in F[X]$ monic and $a \in F^\times$. By Proposition 4.8, the set S_g of $\mathfrak{P} \in S$ for which g has a zero in some (or equivalently, every) closure of (F, \mathfrak{P}) is open-closed in S , therefore in particular compact. For each $\mathfrak{P} \in S_g$, D_{S_g} implies that there exists $x_{\mathfrak{P}} \in F$ with $1 - g(x_{\mathfrak{P}})^2 a^{-2} \in \mathcal{O}_{\mathfrak{P}}$. Therefore, Proposition 4.5 (in the case $m = r = 1$) gives $x \in F$ with $1 - g(x)^2 a^{-2} \in \mathcal{O}_{\mathfrak{P}}$ for every $\mathfrak{P} \in S_g$. \square

Remark 4.10. Important special cases to which the previous proposition applies are:

- (1) $S = \mathcal{S}_p^{\tau}(F, t, s)$ for $t, s \in F$, which is compact by Lemma 4.3 and Lemma 4.1,
- (2) $S = \mathcal{S}_p(F)$, as $\mathcal{S}_p(F) = \mathcal{S}_p^{(1,1)}(F, t_p, 1)$.

Remark 4.11. We want to stress that we proved Proposition 4.9 only for compact sets S . This condition cannot be dropped. More precisely, if UD_S holds for some $S \subseteq \mathcal{S}_p^{\tau}(F)$, then S is relatively compact in $\mathcal{S}_p^{\tau}(F)$:

Indeed, by applying UD_S to the cyclotomic polynomial $g = \Phi_{q^f-1}$ and $a = t_p$ we find $s \in F$ such that the residue of s is a generator of the multiplicative group of the residue field of (F, \mathfrak{P}) , for every $\mathfrak{P} \in S$. Next, adapting a familiar argument (for example in [PR84, Lemma 3.5(ii)]), we may select finitely many polynomials $g_1, \dots, g_n \in K(s)[X]$ in order that for each $\mathfrak{P} \in S$ there exists $i \in \{1, \dots, n\}$ such that there is a uniformizer of (F', \mathfrak{P}') with minimal polynomial g_i over $K'(s)$, where (F', \mathfrak{P}') and (K', \mathfrak{p}') are closures of (F, \mathfrak{P}) and (K, \mathfrak{p}) , respectively. Moreover, any element of F' sufficiently close to a root of g_i is a uniformizer of \mathfrak{P}' . Applying UD_S to each g_i , there are elements $t_i \in F$ such that S is contained in the finite union $\bigcup_{i=1}^n \mathcal{S}_p^{\tau}(F, t_i, s)$, which is compact by Remark 4.10.

The following example shows that $\mathcal{S}_p^{\tau}(F)$ itself is in general not compact.

Example 4.12. Fix $K = \mathbb{Q}$ and a prime number p . We start with a quadratic number field K_0 in which p is split. Now in each step construct K_{i+1} as a quadratic extension of K_i which satisfies the following three conditions:

- (1) one of the two primes above p of type $(1, 1)$ is inert,
- (2) the other prime of type $(1, 1)$ is split, and
- (3) all primes above p of type $(1, 2)$ are split.

The existence of such K_{i+1} is a special case of the Grunwald–Wang theorem. If F is the union of the K_i , then the p -valuations on F correspond to branches in the tree of primes constructed and are all in $S_F := \mathcal{S}_p^{(1,2)}(F)$. Out of these, exactly one will have type $(1, 1)$,

and all the others have type $\tau := (1, 2)$. Now $S'_F := \mathcal{S}_p^{\tau}(F)$ is not compact, as the open-closed cover $S'_F = \bigcup_{s \in F} \mathcal{S}_p^{\tau}(F, p, s)$ has no finite subcover: If $S'_F = \bigcup_{j=1}^n \mathcal{S}_p^{\tau}(F, p, s_j)$ then there exists i with $s_j \in K_i$ for all j , and there are $\mathfrak{P} \in S_F$ of type τ whose restriction to K_i is of type $(1, 1)$, so $\mathfrak{P} \notin \mathcal{S}_p^{\tau}(F, p, s_j)$ for all j . Assuming that p is odd for simplicity, one can in fact check that F satisfies D_{S_F} but not UD_{S_F} for the polynomial $g = X^2 - c$, where c is an integer which is not a square modulo p .

In this situation we have $\mathcal{S}_p^{\tau, \mathbb{Z}}(F) = \mathcal{S}_p^{\tau}(F)$, which additionally shows that Proposition 4.8 does not hold for $\mathcal{S}_p^{\tau, \mathbb{Z}}(F)$ instead of $\mathcal{S}_p^{\tau, \mathbb{Z}}(F)$ (take φ to be an \mathcal{L}_R -sentence asserting that the residue field has p^2 elements).

Proof of Proposition 2.3. For every $t, s \in F$ and $\tau' \leq \tau$, the set $\mathcal{S}_p^{\tau'}(F, t, s)$ is compact by Remark 4.10, so $D_{\mathcal{S}_p^{\tau'}(F)}$ implies $UD_{\mathcal{S}_p^{\tau'}(F, t, s)}$ by Proposition 4.9. Conversely,

$$\mathcal{S}_p^{\tau}(F) = \bigcup_{\tau' \leq \tau} \bigcup_{t, s \in F} \mathcal{S}_p^{\tau'}(F, t, s),$$

and $UD_{\mathcal{S}_p^{\tau'}(F, t, s)}$ implies $D_{\mathcal{S}_p^{\tau'}(F, t, s)}$ (Lemma 2.1). \square

5. AXIOMATIZING UNIFORM DENSENESS

Let again K be a number field, \mathfrak{p} a prime of K and $\tau = (e, f) \in \mathbb{N}^2$. As above, if \mathfrak{p} is a p -valuation let $t_{\mathfrak{p}}$ be a uniformizer of $v_{\mathfrak{p}}$, and let $q_{\mathfrak{p}} = |Kv_{\mathfrak{p}}|$ be the size of the residue field. When working in a first-order language expanding $\mathcal{L}_{\text{ring}}$ in the following, we freely use multiplicative inversion as though it were represented by a function symbol, as it can be replaced by either a universal or existential definition for multiplicative inversion. This replacement thus does not affect the quantifier complexity of any formula that is not itself quantifier free.

We will use the following result from [Feh13] to axiomatize quantifiers over real and p -adic closures.

Proposition 5.1. *There exists a recursive map $\varphi(\mathbf{x}) \mapsto \hat{\varphi}_{\mathfrak{p}}^{\tau}(\mathbf{x})$ from \mathcal{L}_R -formulas to $\mathcal{L}_R(K)$ -formulas such that for every extension F/K and tuple $\mathbf{a} \in F^m$ the following holds:*

- (1) *If $(F', \mathcal{O}_{\mathfrak{p}'}) \models \varphi(\mathbf{a})$ holds for all closures (F', \mathfrak{P}') of (F, \mathfrak{P}) with $\mathfrak{P} \in \mathcal{S}_p^{\tau}(F)$, then $(F, R_{\mathfrak{p}}^{\tau}(F)) \models \hat{\varphi}_{\mathfrak{p}}^{\tau}(\mathbf{a})$.*
- (2) *If $(F, R_{\mathfrak{p}}^{\tau}(F)) \models \hat{\varphi}_{\mathfrak{p}}^{\tau}(\mathbf{a})$, then $(F', \mathcal{O}_{\mathfrak{p}'}) \models \varphi(\mathbf{a})$ for all closures (F', \mathfrak{P}') of (F, \mathfrak{P}) with $\mathfrak{P} \in \mathcal{S}_p^{\tau, \mathbb{Z}}(F)$.*

Proof. We can copy the proof of [Feh13, Proposition 8.4], except that we note that [Feh13, Lemma 8.3] actually only applies [Feh13, Proposition 8.2] and can therefore take an \mathcal{L}_R -formula instead of an $\mathcal{L}_{\text{ring}}$ -formula, and that we omit the step of replacing the occurrences of $x \in R$ by the formula $\theta_{R, \mathfrak{p}}^{\tau'}(x)$, which makes the application of [Feh13, Proposition 6.6] and therefore the assumption that F satisfies $T_{R, \mathfrak{p}}^{\tau'}$ unnecessary. \square

In order to use this result, we first deal with the value groups being \mathbb{Z} -groups and then use Proposition 5.1 to axiomatize uniform denseness.

Recall that an ordered abelian group G is a \mathbb{Z} -group if it is elementarily equivalent to \mathbb{Z} ; equivalently, if it is discrete (i.e. it has a convex subgroup isomorphic to \mathbb{Z} , which we will also denote by \mathbb{Z}) and satisfies $G = nG + \{0, \dots, n-1\}$ for all $n \in \mathbb{N}$. So, given a valuation v on a field F which has a uniformizer, the value group vF is a \mathbb{Z} -group if and only if for each $n \in \mathbb{N}$ and for each $y \in F^{\times}$ there exists $x \in F^{\times}$ such that $v(yx^n) \in \{0, \dots, n-1\} \subseteq \mathbb{Z}$, or, equivalently, there exists $x \in F$ such that $v(yx^n) \in \mathbb{Z}$.

We want to axiomatize the class of fields F/K with $\mathcal{S}_p^\tau(F) = \mathcal{S}_p^{\tau, \mathbb{Z}}(F)$. In both Lemma 5.2 and Proposition 5.3 we assume that \mathfrak{p} is a p -valuation.

Lemma 5.2. *For any $n \in \mathbb{N}$, there exists a polynomial $\phi_n \in \mathbb{Z}[X_1, \dots, X_n]$ such that for any field F/K , elements $x_1, \dots, x_n \in F$, and $\mathfrak{P} \in \mathcal{S}_p^\tau(F)$, we have $v_{\mathfrak{P}}(\phi_n(x_1, \dots, x_n)) = 0$ if and only if $\min(v_{\mathfrak{P}}(x_1), \dots, v_{\mathfrak{P}}(x_n)) = 0$.*

Proof. Let $g \in \mathbb{Z}[X]$ be a monic polynomial whose reduction has no zero in the extension of Kv_p of degree f ; for instance we may take g to be a lift of an irreducible polynomial over \mathbb{F}_p of sufficiently large degree. Let $\phi_2(X, Y) = Y^{\deg g} g(X/Y) \in \mathbb{Z}[X, Y]$ be the homogenization. For any F/K , $x, y \in F$ and $\mathfrak{P} \in \mathcal{S}_p^\tau(F)$, we have $v_{\mathfrak{P}}(g(x)) = \deg(g) \min(v_{\mathfrak{P}}(x), 0)$, and therefore $v_{\mathfrak{P}}(\phi_2(x, y)) = \deg(g) \min(v_{\mathfrak{P}}(x), v_{\mathfrak{P}}(y))$; in particular ϕ_2 is as desired for $n = 2$. For higher n , one uses the polynomial $\phi_2(X_1, \phi_2(X_2, \phi_2(X_3, \dots) \cdot \dots))$. For $n = 1$ we may take $\phi_1 = X_1$. \square

Proposition 5.3. *There exists an $\mathcal{L}_R(K)$ -theory $T_{\mathbb{Z}, p}^\tau$ such that for every extension F/K , we have $(F, R_p^\tau(F)) \models T_{\mathbb{Z}, p}^\tau$ if and only if $\mathcal{S}_p^\tau(F) = \mathcal{S}_p^{\tau, \mathbb{Z}}(F)$.*

Proof. Let $\nu_{p, n}^\tau$ be the $\mathcal{L}_R(K)$ -sentence

$$\forall y \neq 0 \exists x_0, \dots, x_{n-1} R^\times(\phi_n(y^{e!} t_p^0 x_0^n, \dots, y^{e!} t_p^{n-1} x_{n-1}^n)),$$

where $R^\times(x)$ is short for $R(x) \wedge R(x^{-1})$, and let $T_{\mathbb{Z}, p}^\tau = \{\nu_{p, n}^\tau : n > 0\}$.

Consider a field F/K with $(F, R_p^\tau(F)) \models T_{\mathbb{Z}, p}^\tau$ and let $\mathfrak{P} \in \mathcal{S}_p^\tau(F)$. Given $y \in F^\times$ and $n > 0$, and letting $n' = e!n$, there exist $i \in \{0, \dots, n' - 1\}$ and $x \in F$ with $v_{\mathfrak{P}}(y^{e!} t_p^i x^{n'}) = 0$ by using $(F, R_p^\tau(F)) \models \nu_{p, n'}$ and Lemma 5.2. Therefore $v_{\mathfrak{P}}(y^{e!} x^{n'}) = e!v_{\mathfrak{P}}(yx^n) \in \mathbb{Z}$ and thus $v_{\mathfrak{P}}(yx^n) \in \mathbb{Z}$. Hence $\mathfrak{P} \in \mathcal{S}_p^{\tau, \mathbb{Z}}(F)$. This proves $\mathcal{S}_p^\tau(F) = \mathcal{S}_p^{\tau, \mathbb{Z}}(F)$.

For the converse direction, let F/K be a field with $\mathcal{S}_p^\tau(F) = \mathcal{S}_p^{\tau, \mathbb{Z}}(F)$, and fix $n > 1$ and $y \in F^\times$. We shall find $x_0, \dots, x_{n-1} \in F$ such that for any $\mathfrak{P} \in \mathcal{S}_p^\tau(F)$ we have $v_{\mathfrak{P}}(x_i^n) \geq -e!v_{\mathfrak{P}}(y) - i v_{\mathfrak{P}}(t_p)$ for any $i \in \{0, \dots, n - 1\}$, with equality for some i . We see that then $v_{\mathfrak{P}}(y^{e!} t_p^i x_i^n) \geq 0$ for all i , with equality for some i , hence $\nu_{p, n}^\tau$ is satisfied by Lemma 5.2.

It remains to prove that such x_0, \dots, x_{n-1} can always be found, for which we use Proposition 4.4. For every \mathfrak{P} , write $e_{\mathfrak{P}} = v_{\mathfrak{P}}(t_p)$ for the relative ramification index. Let $i_{\mathfrak{P}} \in \{0, \dots, n - 1\}$ such that $i_{\mathfrak{P}} \equiv -\frac{e!}{e_{\mathfrak{P}}} v_{\mathfrak{P}}(y) \pmod{n}$, choose $x_{\mathfrak{P}}^{(i_{\mathfrak{P}})}$ with $n v_{\mathfrak{P}}(x_{\mathfrak{P}}^{(i_{\mathfrak{P}})}) = -e!v_{\mathfrak{P}}(y) - i_{\mathfrak{P}} e_{\mathfrak{P}}$. For $i \in \{0, \dots, n - 1\} \setminus \{i_{\mathfrak{P}}\}$, choose $x_{\mathfrak{P}}^{(i)}$ with

$$-e!v_{\mathfrak{P}}(y) - i e_{\mathfrak{P}} \leq n v_{\mathfrak{P}}(x_{\mathfrak{P}}^{(i)}) \leq -e!v_{\mathfrak{P}}(y) - (i - n) e_{\mathfrak{P}}.$$

For every \mathfrak{P} , the set

$$V_{\mathfrak{P}} = \{\mathfrak{Q} : n v_{\mathfrak{Q}}(x_{\mathfrak{P}}^{(i_{\mathfrak{P}})}) = -v_{\mathfrak{Q}}(y^{e!} t_p^{i_{\mathfrak{P}}}) \text{ and } -v_{\mathfrak{Q}}(y^{e!} t_p^i) \leq n v_{\mathfrak{Q}}(x_{\mathfrak{P}}^{(i)}) \leq -v_{\mathfrak{Q}}(y^{e!} t_p^{i-n}) \text{ for } i \neq i_{\mathfrak{P}}\}$$

is an open-closed neighbourhood of \mathfrak{P} in $\mathcal{S}_p^\tau(F)$. Since $\mathcal{S}_p^\tau(F)$ is compact by Lemma 4.1, the $V_{\mathfrak{P}}$ are compact and there are finitely many $\mathfrak{P}_1, \dots, \mathfrak{P}_m$ such that the corresponding $V_{\mathfrak{P}_j}$ cover $\mathcal{S}_p^\tau(F)$, and we may choose open-closed and hence compact subsets $S_j \subseteq V_{\mathfrak{P}_j}$ that partition $\mathcal{S}_p^\tau(F)$.

By Proposition 4.4, we can find x_0, \dots, x_{n-1} such that $v_{\mathfrak{Q}}(x_i) = v_{\mathfrak{Q}}(x_{\mathfrak{P}_j}^{(i)})$ for all $\mathfrak{Q} \in S_j$, all $i \in \{0, \dots, n - 1\}$, and all $j \in \{1, \dots, m\}$; the hypothesis of the proposition is satisfied since for any valuation v which is a proper coarsening of $v_{\mathfrak{Q}}$ for some prime $\mathfrak{Q} \in S_j$ we have $v(x_{\mathfrak{P}_j}^{(i)}) = -\frac{e!}{n} v(y)$, and the right-hand side is independent of j . These x_i are as desired by construction. \square

We return to the setting of an arbitrary prime \mathfrak{p} of K .

Lemma 5.4. *There exists an $\mathcal{L}_R(K)$ -formula $\chi_{\mathfrak{p}}^{\tau}(t, s)$ such that for every extension F/K , every $\mathfrak{P} \in \mathcal{S}_{\mathfrak{p}}^{\tau}(F)$, and every $t, s \in F$, the following are equivalent:*

- (1) $\mathfrak{P} \in \mathcal{S}_{\mathfrak{p}}^{\tau}(F, t, s)$
- (2) $(F, \mathcal{O}_{\mathfrak{P}}) \models \chi_{\mathfrak{p}}^{\tau}(t, s)$
- (3) $(F', \mathcal{O}_{\mathfrak{P}'}) \models \chi_{\mathfrak{p}}^{\tau}(t, s)$ for some closure (F', \mathfrak{P}') of (F, \mathfrak{P})
- (4) $(F', \mathcal{O}_{\mathfrak{P}'}) \models \chi_{\mathfrak{p}}^{\tau}(t, s)$ for every closure (F', \mathfrak{P}') of (F, \mathfrak{P})

Proof. If \mathfrak{p} is an ordering, then $\mathcal{S}_{\mathfrak{p}}^{\tau}(F, t, s) = \mathcal{S}_{\mathfrak{p}}^{\tau}(F)$, for all t, s . Thus we may take $\chi_{\mathfrak{p}}^{\tau}(t, s)$ to be $t = t$. Otherwise, \mathfrak{p} is a p -valuation, and we let $\chi_{\mathfrak{p}}^{\tau}(t, s)$ be the $\mathcal{L}_R(K)$ -formula

$$R^{\times}(t^e t_{\mathfrak{p}}^{-1}) \wedge R^{\times}(s) \wedge \bigwedge_{n|q_{\mathfrak{p}}^f - 1, n \neq q_{\mathfrak{p}}^f - 1} R^{\times}(s^n - 1).$$

The equivalence of (1) and (2) follows from Lemma 4.2. For the equivalence of (2), (3) and (4) note that $\chi_{\mathfrak{p}}^{\tau}$ is equivalent both to an existential and a universal $\mathcal{L}_R(K)$ -formula, and if (F', \mathfrak{P}') is a closure of (F, \mathfrak{P}) , then $\mathcal{O}_{\mathfrak{P}'}^{\times} \cap F = \mathcal{O}_{\mathfrak{P}}^{\times}$. \square

Proposition 5.5. *There is a recursive theory $T_{\text{UD}, \mathfrak{p}}^{\tau}$ in the language $\mathcal{L}_R(K)$ such that for every extension F/K the following holds:*

- (1) *If F satisfies $\text{UD}_{\mathcal{S}_{\mathfrak{p}}^{\tau}(F, t, s)}$ for every $t, s \in F$, then $(F, R_{\mathfrak{p}}^{\tau}(F)) \models T_{\text{UD}, \mathfrak{p}}^{\tau}$.*
- (2) *If $(F, R_{\mathfrak{p}}^{\tau}(F)) \models T_{\text{UD}, \mathfrak{p}}^{\tau}$, then F satisfies $\text{UD}_{\mathcal{S}_{\mathfrak{p}}^{\tau, \mathbb{Z}}(F, t, s)}$ for all $t, s \in F$.*

Proof. For a n -tuple \mathbf{c} denote by $g_{\mathbf{c}}$ the monic polynomial of degree n with coefficients \mathbf{c} . Let $\varphi(\mathbf{c}, x, a, t, s)$ be the $\mathcal{L}_R(K)$ -formula

$$[\chi_{\mathfrak{p}}^{\tau}(t, s) \wedge (\exists y)(g_{\mathbf{c}}(y) = 0)] \rightarrow R(1 - g_{\mathbf{c}}(x)^2 a^{-2}).$$

Proposition 5.1 gives us an $\mathcal{L}_R(K)$ -formula $\hat{\varphi}_{\mathfrak{p}}^{\tau}(\mathbf{c}, x, a, t, s)$. Let ψ_n be the sentence

$$\forall \mathbf{c} \forall a \neq 0 \forall t \forall s \exists x \hat{\varphi}_{\mathfrak{p}}^{\tau}(\mathbf{c}, x, a, t, s)$$

with \mathbf{c} of length n . Then $T_{\text{UD}, \mathfrak{p}}^{\tau} = \{\psi_n : n \in \mathbb{N}\}$ is the required $\mathcal{L}_R(K)$ -theory, by Lemma 5.4. \square

Proof of Proposition 2.4. Let $T_{\mathfrak{p}}^{\tau} = T_{\mathbb{Z}, \mathfrak{p}}^{\tau} \cup \bigcup_{\tau' \leq \tau} T_{\text{UD}, \mathfrak{p}}^{\tau'}$.

If $(F, R_{\mathfrak{p}}^{\tau}(F)) \models T_{\mathfrak{p}}^{\tau}$, then in particular $(F, R_{\mathfrak{p}}^{\tau'}(F)) \models T_{\text{UD}, \mathfrak{p}}^{\tau'}$ for every $\tau' \leq \tau$, so F satisfies $\text{UD}_{\mathcal{S}_{\mathfrak{p}}^{\tau', \mathbb{Z}}(F, t, s)}$ for every $t, s \in F$ by Proposition 5.5. Since also $(F, R_{\mathfrak{p}}^{\tau}(F)) \models T_{\mathbb{Z}, \mathfrak{p}}^{\tau}$, we have that $\mathcal{S}_{\mathfrak{p}}^{\tau, \mathbb{Z}}(F, t, s) = \mathcal{S}_{\mathfrak{p}}^{\tau}(F, t, s)$ by Proposition 5.3.

Conversely, if $\text{UD}_{\mathcal{S}_{\mathfrak{p}}^{\tau'}(F, t, s)}$ holds for every $t, s \in F$ and $\tau' \leq \tau$, then in particular $\text{D}_{\mathcal{S}_{\mathfrak{p}}^{\tau}(F)}$ holds, hence $\mathcal{S}_{\mathfrak{p}}^{\tau}(F) = \mathcal{S}_{\mathfrak{p}}^{\tau, \mathbb{Z}}(F)$ since (in the case of p -valuations) the value group of a prime in $\mathcal{S}_{\mathfrak{p}}^{\tau}(F)$ is the same as the value group of a corresponding closure by Lemma 3.2, and the latter is a \mathbb{Z} -group. Thus by Proposition 5.3 and Proposition 5.5, $(F, R_{\mathfrak{p}}^{\tau}(F)) \models T_{\mathfrak{p}}^{\tau}$. \square

Remark 5.6. A closer inspection shows that the theory $T_{\mathfrak{p}}^{\tau}$ thus obtained has quantifier complexity $\forall \exists \forall$. However, it is not difficult to see that if $(F_n, R_{\mathfrak{p}}^{\tau}(F_n))_{n \in \mathbb{N}}$ is a chain of models of $T_{\mathfrak{p}}^{\tau}$, then also $F = \bigcup_{n \in \mathbb{N}} F_n$ satisfies $\text{D}_{\mathcal{S}_{\mathfrak{p}}^{\tau}(F)}$ and so $(F, R_{\mathfrak{p}}^{\tau}(F))$ is a model of $T_{\mathfrak{p}}^{\tau}$. Therefore, by general principles, $T_{\mathfrak{p}}^{\tau}$ is equivalent to an $\forall \exists$ -theory.

Remark 5.7. Corollary 2.5 shows that if $\text{D}_{\mathcal{S}_{\mathfrak{p}}^{\tau}(F)}$ holds and $(E, R_{\mathfrak{p}}^{\tau}(E)) \equiv (F, R_{\mathfrak{p}}^{\tau}(F))$, as $\mathcal{L}_R(K)$ -structures, then also $\text{D}_{\mathcal{S}_{\mathfrak{p}}^{\tau}(E)}$ holds. In Section 7 we show—at least in the case $K = \mathbb{Q}$ and $\mathfrak{p} = \infty$ —that this statement no longer holds true in $\mathcal{L}_{\text{ring}}$ instead of \mathcal{L}_R , i.e. if F is dense in all its real closures and $E \equiv F$, as $\mathcal{L}_{\text{ring}}$ -structures, it does *not* follow that

also E is dense in all its real closures. In other words, the property $D_{\mathcal{S}_p^\tau}$ is in general not preserved under elementary equivalence in the language of rings.

Remark 5.8. However, we observe that the property $D_{\mathcal{S}_p^\tau}$ does pass down elementary extensions in the language of rings. Let $F \preceq F^*$ be an $\mathcal{L}_{\text{ring}}$ -elementary extension and suppose that $D_{\mathcal{S}_p^\tau(F^*)}$ holds. Let $\mathfrak{P} \in \mathcal{S}_p^\tau(F)$, and let w be a non-trivial valuation on F which is a coarsening of \mathfrak{P} . There exists an ultrapower F^{**} of F into which we may $\mathcal{L}_{\text{ring}}(F)$ -elementarily embed F^* . Identifying embeddings with inclusions, we have $F \preceq F^* \preceq F^{**}$. Since F^{**} is an ultrapower of F , we may let w^{**} and \mathfrak{P}^{**} be the natural extensions of w and \mathfrak{P} to F^{**} , respectively. By Łos's theorem, we have

$$(F, \mathcal{O}_w, \mathcal{O}_{\mathfrak{P}}) \preceq (F^{**}, \mathcal{O}_{w^{**}}, \mathcal{O}_{\mathfrak{P}^{**}}),$$

where these are viewed as structures in a language $\mathcal{L}_{R,S}(K)$ which is the expansion of $\mathcal{L}_{\text{ring}}(K)$ by two unary predicates R and S . In particular: \mathfrak{P}^{**} is also a prime of F^{**} , lying above \mathfrak{p} , and of the same exact type $\tau' \leq \tau$ as \mathfrak{P} ; and w^{**} is a non-trivial valuation on F^{**} coarsening \mathfrak{P}^{**} . Next, we let w^* and \mathfrak{P}^* denote the restrictions of w^{**} and \mathfrak{P}^{**} to F^* , respectively. Then

$$(\dagger) \quad (F, \mathcal{O}_w, \mathcal{O}_{\mathfrak{P}}) \text{ is existentially closed in } (F^*, \mathcal{O}_{w^*}, \mathcal{O}_{\mathfrak{P}^*})$$

as $\mathcal{L}_{R,S}(K)$ -structures. Again \mathfrak{P}^* is of the same exact type as \mathfrak{P} , and w^* is a coarsening of \mathfrak{P}^* . In particular $\mathfrak{P}^* \in \mathcal{S}_p^\tau(F^*)$.

We break into two cases. First, suppose that \mathfrak{p} is a p -valuation. Then $v_{\mathfrak{P}}$ induces a p -valuation $\bar{v}_{\mathfrak{P}}$ on Fw , and $v_{\mathfrak{P}^*}$ induces a p -valuation $\bar{v}_{\mathfrak{P}^*}$ on F^*w^* . In fact $\bar{v}_{\mathfrak{P}}$ is the restriction of $\bar{v}_{\mathfrak{P}^*}$ to Fw . By assumption, F^* is dense in a closure of (F^*, \mathfrak{P}^*) ; and thus $v_{\mathfrak{P}^*}F^*$ is a \mathbb{Z} -group and $(F^*w^*, \bar{v}_{\mathfrak{P}^*})$ is henselian, the latter by (1) \Rightarrow (2) of Lemma 3.1. From (\dagger) , it follows that $v_{\mathfrak{P}}F$ is existentially closed in $v_{\mathfrak{P}^*}F^*$, thus $v_{\mathfrak{P}}F$ is also a \mathbb{Z} -group. Also from (\dagger) , it follows that Fw is existentially closed in F^*w^* . In particular, Fw is relatively algebraically closed in F^*w^* , and so $\bar{v}_{\mathfrak{P}}$ is henselian. By (2) \Rightarrow (1) of Lemma 3.1, and since w was an arbitrary non-trivial valuation coarsening $v_{\mathfrak{P}}$, F is dense in a henselization of (F, \mathfrak{P}) , which is a p -adic closure. Note that, by ignoring w altogether, this argument shows that the property $\mathcal{S}_p^\tau = \mathcal{S}_p^{\tau, \mathbb{Z}}$ passes down elementary extensions in the language of rings.

If \mathfrak{p} is an ordering, on the other hand, then \mathfrak{P} induces an ordering \leq_w on Fw , and \mathfrak{P}^* induces an ordering \leq_{w^*} on F^*w^* . Again, \leq_w is the restriction of \leq_{w^*} to Fw . Again, by assumption, F^* is dense in a closure of (F^*, \mathfrak{P}^*) ; and thus w^*F^* is divisible and (F^*w^*, \leq_{w^*}) is real closed, by Proposition 3.8. From (\dagger) , it follows that wF is existentially closed in w^*F^* , and that (Fw, \leq_w) is existentially closed in (F^*w^*, \leq_{w^*}) . Thus wF is divisible and (Fw, \leq_w) is real closed. By Proposition 3.8, and since w was an arbitrary non-trivial $\leq_{\mathfrak{P}}$ -convex valuation, F is dense in a real closure of (F, \mathfrak{P}) . Thus, in either case, $D_{\mathcal{S}_p^\tau(F)}$ holds.

Remark 5.9. It is not hard to see that denseness of F in the closure of one specific prime \mathfrak{P} of F is an elementary property in \mathcal{L}_R where R interprets $\mathcal{O}_{\mathfrak{P}}$: If \mathfrak{P} is a valuation then (3) of Lemma 3.1 gives an elementary characterization of denseness of (K, v) in its henselization, and one axiomatizes that a henselization is a closure by demanding that the value group be a \mathbb{Z} -group. If \mathfrak{P} is an ordering then a similar characterization is given in [Hau67, Satz 13].

6. ALGEBRAIC FIELDS AND APPLICATIONS

Recall that the Pythagoras number of a field F is the smallest n such that every sum of squares in F is a sum of at most n squares. We denote the Pythagoras number of F

by $\pi_\infty(F)$. Let $\Phi_{\infty,n}$ be the $\mathcal{L}_{\text{ring}}$ -theory stating that every sum of $n+1$ squares is a sum of n squares, and let $\varphi_{\infty,n}(x)$ be an $\mathcal{L}_{\text{ring}}$ -formula defining the sums of n squares. As the sums of squares of F are precisely the elements of $R_\infty(F)$, the following is obvious:

Lemma 6.1. *For every field F , and every n , we have $\varphi_{\infty,n}(F) \subseteq R_\infty(F)$. Moreover, the following are equivalent:*

- (1) $\varphi_{\infty,n}(F) = R_\infty(F)$
- (2) $\pi_\infty(F) \leq n$.
- (3) $F \models \Phi_{\infty,n}$

For a p -valuation \mathfrak{p} on a number field K , and $\tau \in \mathbb{N}^2$, [ADF19b] defines a p -adic analogue $\pi_{\mathfrak{p}}^\tau(F)$ of the Pythagoras number, as follows. For each F/K , we first define an increasing chain $(R_{\mathfrak{p},n}^\tau(F))_{n \in \mathbb{N}}$ of subsets of F , as in [ADF19b, Section 2.2], such that

$$R_{\mathfrak{p}}^\tau(F) = \bigcup_{n \in \mathbb{N}} R_{\mathfrak{p},n}^\tau(F).$$

Then the (\mathfrak{p}, τ) -Pythagoras number of F is

$$\pi_{\mathfrak{p}}^\tau(F) := \inf \{n \in \mathbb{N} \mid R_{\mathfrak{p}}^\tau(F) = R_{\mathfrak{p},n}^\tau(F)\} \in \mathbb{N} \cup \{\infty\}.$$

In the language of [ADF19b, Section 3], for each n , the map $F \mapsto R_{\mathfrak{p},n}^\tau(F)$ is a 1-dimensional diophantine family, by [ADF19b, Example 3.8]. That is, for each n , there is an existential $\mathcal{L}_{\text{ring}}(K)$ -formula $\varphi_{\mathfrak{p},n}^\tau(x)$ such that $\varphi_{\mathfrak{p},n}^\tau(F) = R_{\mathfrak{p},n}^\tau(F)$, for all F/K . Let $\Phi_{\mathfrak{p},n}^\tau$ be the universal-existential $\mathcal{L}_{\text{ring}}(K)$ -theory consisting of the sentences

$$\forall x [\phi_{\mathfrak{p},m}^\tau(x) \rightarrow \phi_{\mathfrak{p},n}^\tau(x)],$$

for all $m \geq n$. Just as for orderings, the following is now obvious.

Lemma 6.2. *If F/K is an extension, then $\varphi_{\mathfrak{p},n}^\tau(F) \subseteq R_{\mathfrak{p}}^\tau(F)$, and the following are equivalent:*

- (1) $\varphi_{\mathfrak{p},n}^\tau(F) = R_{\mathfrak{p}}^\tau(F)$
- (2) $\pi_{\mathfrak{p}}^\tau(F) \leq n$.
- (3) $F \models \Phi_{\mathfrak{p},n}^\tau$

Let \mathfrak{p} now be an arbitrary prime of K . For a class of fields \mathcal{F} we let $\pi_{\mathfrak{p}}^\tau(\mathcal{F}) = \sup_{F \in \mathcal{F}} \pi_{\mathfrak{p}}^\tau(F)$.

Lemma 6.3. *For a class of fields \mathcal{F} let \mathcal{F}^* denote its closure under elementary equivalence, direct limits and ultraproducts. Then $\pi_{\mathfrak{p}}^\tau(\mathcal{F}^*) = \pi_{\mathfrak{p}}^\tau(\mathcal{F})$.*

Proof. This is clear, since $\Phi_{\mathfrak{p},n}^\tau$ is a $\forall\exists$ -theory. □

Let $T_{\mathfrak{p},n}^\tau$ denote the union of $\Phi_{\mathfrak{p},n}^\tau$ and $T_{\mathfrak{p}}^\tau$, with all occurrences of R replaced with $\varphi_{\mathfrak{p},n}^\tau(x)$.

Proposition 6.4. *$F \models T_{\mathfrak{p},n}^\tau$ if and only if $\pi_{\mathfrak{p}}^\tau(F) \leq n$ and $D_{\mathcal{S}_{\mathfrak{p}}^\tau(F)}$ holds.*

Proof. This follows from Corollary 2.5 together with Lemma 6.2. □

Corollary 6.5. *If $\pi_{\mathfrak{p}}^\tau(F) = n < \infty$ and F is dense in all closures at elements of $\mathcal{S}_{\mathfrak{p}}^\tau(F)$, then every F' which is $\mathcal{L}_{\text{ring}}(K)$ -elementarily equivalent to F is dense in all closures at elements of $\mathcal{S}_{\mathfrak{p}}^\tau(F')$.*

We denote by \mathcal{A} the class of algebraic fields of characteristic zero, i.e. fields $F \subseteq \mathbb{Q}^{\text{alg}}$. We denote by $\text{Th}(\mathcal{A}) = \bigcap_{F \in \mathcal{A}} \text{Th}_{\mathcal{L}_{\text{ring}}}(F)$ the $\mathcal{L}_{\text{ring}}$ -theory of \mathcal{A} .

Proposition 6.6. *For each prime p of \mathbb{Q} and pair τ of positive integers, there exists $n_p^\tau \in \mathbb{N}$ such that $\pi_p^\tau(F) \leq n_p^\tau$ for each field $F \in \mathcal{A}$.*

Proof. For F in the class \mathcal{K} of number fields, the statement is [ADF19b, Thm 1.2]. By Lemma 6.3 it then holds for the class \mathcal{K}^* , which contains \mathcal{A} . \square

Corollary 6.7. *Every $F' \models \text{Th}(\mathcal{A})$ is dense in all its real closures and all its p -adic closures (of arbitrary p -rank).*

Proof. Let $K = \mathbb{Q}$ and let T be the union of $T_{p,n_p^\tau}^\tau$ for all primes p of \mathbb{Q} and all $\tau \in \mathbb{N}^2$. Then every $F \in \mathcal{A}$ is a model of T : For every p and τ , every $\mathfrak{P} \in \mathcal{S}_p^\tau(F)$ is either an archimedean ordering or a valuation with value group \mathbb{Z} , so F is dense in a closure of (F, \mathfrak{P}) . By Proposition 6.4, $F \models T_{p,n_p^\tau}^\tau$.

Therefore $T \subseteq \text{Th}(\mathcal{A})$ and so also every $F' \models \text{Th}(\mathcal{A})$ is a model of $T_{p,n_p^\tau}^\tau$ for every $p \in \mathcal{S}(\mathbb{Q})$ and $\tau \in \mathbb{N}^2$. Let \mathfrak{P} be a prime of F' . If p is the restriction of \mathfrak{P} to \mathbb{Q} and τ denotes the relative type of \mathfrak{P} over p , then $\mathfrak{P} \in \mathcal{S}_p^\tau(F')$. As $F' \models T_{p,n_p^\tau}^\tau$, Proposition 6.4 gives that $\text{D}_{\mathcal{S}_p^\tau(F')}$ holds. In particular, F' is dense in every closure at \mathfrak{P} . \square

Remark 6.8. After this work was finished we realized that for p -adic closures this corollary can in fact also be deduced from the literature in a different way: For each $F \in \mathcal{A}$, the ring $R_p^\tau(F)$ satisfies Darnière’s *Block Approximation Property (BAP)* [Dar98, II.2 p. 29] and is dense in all of its henselizations, i.e. F is dense in the quotient fields of henselizations of the localizations of $R_p^\tau(F)$ at maximal ideals, which in this case are precisely the p -adic closures of F . As this class of rings is elementary by [Dar98, II.5 Prop. 5], and $\varphi_{p,n_p^\tau}^\tau$ defines $R_p^\tau(F')$ in every model F' of $\text{Th}(\mathcal{A})$, $R_p^\tau(F')$ will be dense in all its henselizations. However, also in this case the quotient fields of these henselizations are precisely the p -adic closures of F' : Indeed, for a henselian p -valued field E with value group Γ , the ℓ -cohomological dimension is $1 + \dim_{\mathbb{F}_\ell} \Gamma/\ell\Gamma$, in particular this is bigger than 2 for some prime number ℓ if Γ is not a \mathbb{Z} -group, i.e. if E is not p -adically closed, and by [Cha90, p. 129 Theorem], every model F' of $\text{Th}(\mathcal{A})$ has (virtual) cohomological dimension at most 2, and therefore so has every algebraic extension E of F' .

Remark 6.9. By Proposition 4.9 and Remark 4.10, $\text{D}_{\mathcal{S}_p^\tau(F)}$ implies the uniform version $\text{UD}_{\mathcal{S}_p^\tau(F,t,s)}$, for each t, s . We note that the proof of this fact generalizes easily to allow the prime p to vary within a given finite set S of primes of K . More precisely, if we denote $\mathcal{S}_S^\tau(F) = \bigcup_{p \in S} \mathcal{S}_p^\tau(F)$ and $\mathcal{S}_S^{\tau}(F, t, s) = \bigcup_{p \in S} \mathcal{S}_p^{\tau}(F, t, s)$; then $\text{D}_{\mathcal{S}_S^\tau(F)}$ implies $\text{UD}_{\mathcal{S}_S^{\tau}(F,t,s)}$, for each t, s .

In particular, every model F of the theory of algebraic fields satisfies $\text{UD}_{\mathcal{S}_S^{\tau}(F,t,s)}$, for every t, s , and every finite set S . However, for models of the common theory of number fields one can say more: if F^* is a model of the common $\mathcal{L}_{\text{ring}}(K)$ -theory of number fields extending K , and S is a finite set of primes of K , then F^* satisfies $\text{UD}_{S'}$ where $S' = \bigcup_{\tau \in \mathbb{N}^2} \mathcal{S}_S^\tau(F^*)$.

Remark 6.10. As we already observed in the introduction, our denseness result has long been known for PRC fields and PpC fields, which are models of the theory of algebraic fields. In a sense the archetypal result of this kind is the Frey–Prestel theorem [FJ08, Proposition 11.5.3] – every PAC field is dense with respect to any non-trivial valuation in its algebraic closure –, from which one in particular deduces that henselizations of PAC fields are separably closed.

There are also other fields in the literature which are dense in all their real and p -adic closures in a non-trivial way, for instance coming from other local–global principles which are not subsumed by Pop’s notion of PCC fields: for example, the regularly closed fields

of [HP84] are dense in their real and p -adic closures, but they are not always models of the theory of algebraic fields.

We now draw a few simple consequences from Corollary 6.7.

Corollary 6.11. *Let F be a model of the theory of algebraic fields, and let v be a non-trivial valuation on F . If Fv is formally real, respectively formally p -adic, then Fv is real closed, resp. p -adically closed.*

Proof. If Fv has a prime \mathfrak{Q} , this is induced from a prime \mathfrak{P} of F of which v is a coarsening: In the case of p -valuations, this is simply composition of places, and in the case of orderings it follows from the Baer–Krull theorem. Now F is dense in every closure with respect to \mathfrak{P} by Corollary 6.7. If \mathfrak{Q} is an ordering, Proposition 3.8 implies that $(Fv, \leq_{\mathfrak{Q}})$ is real closed. If \mathfrak{Q} is a p -valuation, Lemma 3.1 implies that $(Fv, v_{\mathfrak{Q}})$ is henselian, and Lemma 3.2 implies that the value group $v_{\mathfrak{P}}F$, and therefore also its convex subgroup $v_{\mathfrak{Q}}(Fv)$, is a \mathbb{Z} -group. Hence $(Fv, v_{\mathfrak{Q}})$ is p -adically closed. \square

Remark 6.12. Algebraic fields are not just dense in all p -adic closures, which are the henselizations with respect to p -valuations, but in fact in *all* their henselizations. We note that this stronger property does *not* transfer to models of the theory of algebraic fields: In fact, already any proper elementary extension \mathbb{Q}^* of \mathbb{Q} has many valuations for which this fails, one example being the following: Like in any ordered field, as in Section 3, the convex closure \mathcal{O} of $\mathbb{Z} \subseteq \mathbb{Q}^*$ is the valuation ring of a valuation $v = v_{\leq}$ with residue field $F := \mathbb{Q}^*v \subseteq \mathbb{R}$. Since \mathbb{Q}^* is a proper elementary extension of \mathbb{Q} , the valuation v is non-trivial, and hence F is real closed by Corollary 6.11. Let \bar{w} be any non-trivial valuation on F and note that \bar{w} is necessarily non-henselian. Then \mathbb{Q}^* is not dense in a henselization of the composition $w = \bar{w} \circ v$, as w induces the non-henselian valuation \bar{w} on the residue field of its non-trivial coarsening v , see Lemma 3.1.

Corollary 6.13. *Let F be an algebraic field that admits a prime \mathfrak{P} such that (F, \mathfrak{P}) is neither real closed nor p -adically closed. If $F \prec F^*$ is an elementary extension and v a non-trivial valuation on F^* , then there is no embedding of F^*v into F^* .*

Proof. Suppose that $F^*v \subseteq F^*$. There is a prime \mathfrak{P}^* of F^* of the same type as \mathfrak{P} , i.e. either both $\mathfrak{P}, \mathfrak{P}^*$ are orderings, or both $\mathfrak{P}, \mathfrak{P}^*$ are p -valuations of the same exact type τ (see [PR84, Theorem 6.4] for the case of p -valuations). Restricting \mathfrak{P}^* to F^*v gives a prime \mathfrak{p} of F^*v , so F^*v is real or p -adically closed by Corollary 6.11. In particular, F^* contains a real or p -adically closed field, which implies that also $F = F^* \cap \mathbb{Q}^{\text{alg}}$ is real or p -adically closed, contradicting the assumption. \square

In the language of [AF17], this means that an algebraic field F as in the corollary does not have *embedded residue* when viewed as a \mathbb{Z} -field. By [AF17, Corollary 5.3], this immediately implies the following, where we write $\mathbb{Q}_{p,\text{alg}} = \mathbb{Q}_p \cap \mathbb{Q}^{\text{alg}}$, $\mathbb{R}_{\text{alg}} = \mathbb{R} \cap \mathbb{Q}^{\text{alg}}$:

Corollary 6.14. *Let F be a field with $F \not\subseteq \mathbb{R}_{\text{alg}}$ or $[F\mathbb{Q}_{p,\text{alg}} : \mathbb{Q}_{p,\text{alg}}] < \infty$ and $\mathbb{Q}_{p,\text{alg}} \not\subseteq F$ for some p . Then there exists an existential formula $\phi(x)$ in the language of rings which defines in $F((t))$ the valuation ring $F[[t]]$.*

We note that the assumption $F \not\subseteq \mathbb{R}_{\text{alg}}$ or $[F\mathbb{Q}_{p,\text{alg}} : \mathbb{Q}_{p,\text{alg}}] < \infty$ and $\mathbb{Q}_{p,\text{alg}} \not\subseteq F$ cannot be dropped: For example, the statement does not hold for $F = \mathbb{R}_{\text{alg}}$ or $F = \mathbb{Q}_{p,\text{alg}}$, see [AF17, Corollary 6.6].

7. DENSENESS IN REAL CLOSURES IN THE LANGUAGE OF RINGS

Let F be a field of characteristic zero. By Proposition 6.4, if $\pi_\infty(F) < \infty$, then denseness of F in its real closures transfers to every F' which is elementarily equivalent to F in the language of rings. In this section we adapt a construction of Prestel from [Pre78] to show that this can fail in the case $\pi_\infty(F) = \infty$. In fact we will see that denseness in real closures transfers in general not to all ultrapowers F^* of F .

Proposition 7.1. *There exists a field $K_\infty \subseteq \mathbb{R}$ which has a unique ordering, and for each $i \in \mathbb{N}$ has a valuation u_i with residue field $K_\infty u_i$ which is hilbertian of finite level² $s(K_\infty u_i) \geq 4^i$.*

Proof. We construct a chain of countable fields $K_0 \subseteq K_1 \subseteq K_2 \subseteq \dots$ and for each i extensions H_i/K_i and k_{i+1}/K_i and elements $d_i \in K_i$ such that

- (1) d_i is a sum of $2 \cdot 4^i + 1$ squares in K_i
- (2) K_i embeds into \mathbb{R} and has a unique ordering
- (3) H_i carries a henselian valuation v_i trivial on K_{i-1} with residue field $H_i v_i = K_i v_i = k_i$
- (4) k_i is hilbertian with $4^i \leq s(k_i) \leq 3 \cdot 4^i$
- (5) k_i carries a valuation w_i trivial on K_{i-1} with residue field contained in H_{i-1}

We begin with $K_{-1} = \mathbb{R}_{\text{alg}}$, $d_{-1} = 1$. Given K_{i-1} , H_{i-1} , k_{i-1} and d_{i-1} satisfying (1)-(5), let $F_i = K_{i-1}(x_{i,1}, \dots, x_{i,4^i})$ for some new indeterminates $x_{i,j}$, $L_i = F_i(y_i)$ for a new indeterminate y_i , $e_i = d_{i-1} + \sum_{j=1}^{4^i} x_{i,j}^2 \in F_i$ and $d_i = e_i + y_i^2 \in L_i$. Note that d_i is a sum of $2 \cdot 4^{i-1} + 1 + 4^i + 1 \leq 2 \cdot 4^i + 1$ squares. The field L_i carries a valuation v_i , coming from the prime ideal $(y_i^2 + e_i)$ of $F_i[y_i]$ (prime since e_i is totally positive), with residue field $k_i := F_i(\sqrt{-e_i})$; this k_i is hilbertian since it is a finitely generated transcendental extension of K_{i-1} , see [FJ08, Chapter 13]. Now e_i cannot be written as a sum of fewer than 4^i squares in F_i by a theorem of Cassels and Pfister [Lam05, Corollary IX.2.3], which implies that $4^i \leq s(k_i) < \infty$, cf [Lam05, p. 421 Exercise 5]. As e_i is a sum of fewer than $3 \cdot 4^i$ squares in F_i , $s(k_i) \leq 3 \cdot 4^i$.

We construct H_i as a henselization of L_i with respect to v_i , and K_i as the intersection of H_i with a real closure R_i of L_i with respect to an archimedean ordering. (Evidently L_i has an archimedean ordering since it can be embedded into \mathbb{R} , since K_{i-1} can be assumed.) By [Pre78, Lemma 1.3], K_i is then uniquely ordered. As $L_i \subseteq K_i \subseteq H_i$ we get that $H_i v_i = K_i v_i = L_i v_i = k_i$.

We now construct a valuation on k_i (for $i > 1$), trivial on K_{i-1} , with residue field contained in H_{i-1} . By Hensel's Lemma, $s(k_{i-1}) \leq 3 \cdot 4^{i-1}$ implies that also $s(H_{i-1}) \leq 3 \cdot 4^{i-1}$. Therefore we can find $z_{i,1}, \dots, z_{i,4^i} \in H_{i-1}$ such that $y_{i-1}^2 + z_{i,1}^2 + \dots + z_{i,4^i}^2 = 0$. On $F_i = K_{i-1}(x_{i,1}, \dots, x_{i,4^i})$, take a H_{i-1} -valued place³ w , trivial on K_{i-1} , under which each $x_{i,j}$ has residue $z_{i,j}$. Then the residue of $e_i = e_{i-1} + y_{i-1}^2 + x_{i,1}^2 + \dots + x_{i,4^i}^2$ is e_{i-1} , whose negative has a square root in H_{i-1} by Hensel's Lemma; therefore we can extend w to an H_{i-1} -valued place w_i on k_i .

Let $K_\infty = \bigcup_{i \in \mathbb{N}} K_i$. This field is naturally embedded into \mathbb{R} by using the embeddings of all the K_i , and it is uniquely ordered since all the K_i are, by (2). It remains to show that for each i_0 , there is a valuation on K_∞ with residue field k_{i_0} . For each $i > i_0$, we have a place $w_i \circ v_i : H_i \dashrightarrow k_i \dashrightarrow H_{i-1}$ trivial on K_{i-1} , by (3) and (5); by composition, we obtain a place $H_i \dashrightarrow k_{i_0}$ and therefore a place $K_i \dashrightarrow k_{i_0}$, with residue field exactly

²Recall that the *level* $s(F)$ of a field F is the least number of squares that sum to -1 .

³We use the same symbol for a valuation and the associated place.

k_{i_0} . The corresponding valuations on the K_i are all compatible, since $w_i \circ v_i$ is trivial on K_{i-1} ; therefore we can take an inductive limit of valuation rings to obtain a valuation u_{i_0} on K_∞ with residue field k_{i_0} . \square

Lemma 7.2. $D_{\mathcal{S}_\infty(F)}$ implies that for every $g \in F[X]$ of odd degree and every $\varepsilon \in F^\times$, there exists $x \in F$ with $\varepsilon^2 - g(x)^2$ a sum of squares in F .

Proof. By Proposition 2.3, $D_{\mathcal{S}_\infty(F)}$ implies $UD_{\mathcal{S}_\infty(F)}$. Since g has a zero in every real closure of F , $UD_{\mathcal{S}_\infty(F)}$ gives an $x \in F$ with $1 - g(x)^2\varepsilon^{-2} \in R_\infty(F)$. As $R_\infty(F)$ consists of the sums of squares of F , we get that $\varepsilon^2 - g(x)^2$ is a sum of squares. \square

Lemma 7.3. Let $s \geq 2$ and let v be a valuation on F whose residue field Fv has level $s(Fv) \geq s$. Let $g \in \mathcal{O}_v[X]$ for which the reduction $\bar{g} \in Fv[X]$ has no root in Fv . Then for no $x \in F$ and $\varepsilon \in F^\times$ with $v(\varepsilon) > 0$ is $\varepsilon^2 - g(x)^2$ a sum of $s - 1$ squares in F .

Proof. Suppose for contradiction that $\varepsilon^2 = g(x)^2 + y_1^2 + \cdots + y_{s-1}^2$, and look for terms of lowest valuation on the right-hand side. Since $v(\varepsilon^2) > 0 \geq v(g(x)^2)$ by the assumption on \bar{g} , and no non-trivial sum of s squares in Fv is zero, we obtain a contradiction. \square

Corollary 7.4. The field K_∞ is dense in all its real closures but has an elementary extension which is not.

Proof. As all orderings of K_∞ are archimedean, it is dense in its real closures.

Let $\varphi_n(a, \varepsilon)$ be a formula expressing that $\varepsilon \neq 0$ and with $g_a = X^3 - a$, there is no x such that $\varepsilon^2 - g_a(x)^2$ is a sum of $4^n - 1$ squares. Then $\{\varphi_1, \dots, \varphi_n\}$ is satisfiable in K_∞ : As $K_\infty u_n$ is hilbertian there exists $a \in \mathcal{O}_{u_n}$ such that $X^3 - \bar{a}$ has no zero in $K_\infty u_n$. By Lemma 7.3, for every $\varepsilon \in K_\infty^\times$ with $u_n(\varepsilon) > 0$, $K_\infty \models \varphi_n(a, \varepsilon)$.

Therefore, if $K_\infty \prec K^*$ is an \aleph_0 -saturated extension, $\{\varphi_1, \varphi_2, \dots\}$ is realized in K^* , i.e. there exists $0 \neq \varepsilon \in K^*$ and $a \in K^*$ such that $\varepsilon^2 - f_a(x)^2$ is not a sum of squares for any x . By Lemma 7.2 this shows that K^* is not dense in all its real closures. \square

ACKNOWLEDGEMENTS

Part of this work was done while all three authors were guests of the Institut Henri Poincaré. The authors would like to thank the IHP for funding and hospitality, and the organizers of the trimester ‘Model theory, combinatorics and valued fields’ for the invitation.

S. A. was supported by the Leverhulme Trust under grant RPG-2017-179. P. D. was supported by KU Leuven IF C14/17/083. A. F. was funded by the Deutsche Forschungsgemeinschaft (DFG) - 404427454.

REFERENCES

- [ADF19a] Sylvy Anscombe, Philip Dittmann and Arno Fehm. Approximation theorems for spaces of localities. [arXiv:1901.02632](https://arxiv.org/abs/1901.02632), 2019.
- [ADF19b] Sylvy Anscombe, Philip Dittmann and Arno Fehm. A p -adic analogue of Siegel’s theorem on sums of squares. To appear in *Mathematische Nachrichten*, 2019. [arXiv:1904.03466](https://arxiv.org/abs/1904.03466).
- [AF17] Sylvy Anscombe and Arno Fehm. Characterizing diophantine henselian valuation rings and valuation ideals. *Proc. London Math. Soc.* 115:293–322, 2017.
- [Bae70] Reinhold Baer. Dichte, Archimedizität und Starrheit geordneter Körper. *Math. Ann.*, 188:165–205, 1970.
- [Bou07] Nicolas Bourbaki. *Topologie générale: Chapitres 1 à 4*. Springer, 2007.
- [Cha90] Zoé Chatzidakis. On the cohomological dimension of non-standard number fields. *J. Pure Appl. Algebra* 69:121–133, 1990.

- [Che75] Gregory Cherlin. Ideals of integers in nonstandard number fields. In: D. H. Saracino and V. B. Weispfenning (eds.), *Model Theory and Algebra. A Memorial Tribute to Abraham Robinson.*, 1975.
- [Dar98] Luck Darnière. *Étude modèle-théorique d'anneaux satisfaisant un principe de Hasse non singulier*. Doctoral thesis, Université de Rennes 1, 1998.
- [EP05] Antonio J. Engler and Alexander Prestel. *Valued Fields*. Springer, 2005.
- [Ers92] Yuri Ershov. Relative regular closeness and π -valuations. *Algebra and Logic*, 31(6):342–360, 1992.
- [Ers01] Yuri L. Ershov. *Multi-Valued Fields*. Siberian School of Algebra and Logic. Springer, 2001.
- [Feh13] Arno Fehm. Elementary local-global principles for fields. *Annals Pure Appl. Logic* 164:989–1008, 2013.
- [FJ08] Michael D. Fried and Moshe Jarden. *Field Arithmetic*. Springer, Third Edition, 2008.
- [Gro87] Camilla Grob. *Die Entscheidbarkeit der Theorie der maximalen pseudo p -adisch abgeschlossenen Körper*. Dissertation, Konstanz, 1987.
- [Hau67] Kurt Hauschild. Cauchyfolgen höheren Typus in angeordneten Körpern. *Zeitschr. math. Logik und Grundlagen d. Math.*, 13:55–66, 1967.
- [HP84] Bernhard Heinemann and Alexander Prestel. Fields regularly closed with respect to finitely many valuations and orderings. In C. R. Riehm and I. Hambleton, editors, *Quadratic and Hermitian Forms. 1983 Conference on Quadratic Forms and Hermitian K -Theory held at McMaster University, Hamilton, Ontario*, volume 4 of *Canadian Mathematical Society Conference Proceedings*, pages 297–336. American Mathematical Society, 1984.
- [Jar88] Moshe Jarden. The algebraic nature of the elementary theory of PRC fields. *Manuscripta math.* 60(4):463–476, 1988.
- [Jar91] Moshe Jarden. Algebraic realization of p -adically projective groups. *Compositio Math.* 79(1):21–62, 1991.
- [Lam05] Tsit Yuen Lam. *Introduction to Quadratic Forms over Fields*. Amer. Math. Soc., 2005.
- [Pop03] Florian Pop. Classically projective groups and pseudo classically closed fields. In: F.-V. Kuhlmann, S. Kuhlmann, M. Marshall (Eds.), *Valuation Theory and Its Applications*, Fields Inst. Commun., vol.II, American Mathematical Society, 2003, pp. 251–283.
- [Pre78] Alexander Prestel. Remarks on the Pythagoras and Hasse number of real fields. *J. reine angew. Math.* 303/304:284–294, 1978.
- [Pre81] Alexander Prestel. Pseudo real closed fields. In R. B. Jensen and A. Prestel, editors, *Set Theory and Model Theory, Proceedings, Bonn 1979*, pages 127–156. Springer, 1981.
- [PR84] Alexander Prestel and Peter Roquette. *Formally p -adic fields*. Springer, 1984.

JEREMIAH HORROCKS INSTITUTE, UNIVERSITY OF CENTRAL LANCASHIRE, PRESTON PR1 2HE, UNITED KINGDOM

E-mail address: sanscombe@uclan.ac.uk

AFDELING ALGEBRA, KU LEUVEN, CELESTIJNENLAAN 200B, 3001 LEUVEN, BELGIUM

E-mail address: philip.dittmann@kuleuven.be

INSTITUT FÜR ALGEBRA, TECHNISCHE UNIVERSITÄT DRESDEN, 01062 DRESDEN, GERMANY

E-mail address: arno.fehm@tu-dresden.de