



**HAL**  
open science

## Earliest occurrence of Embiidae: A new genus from earliest Eocene Oise amber (Insecta: Embiodea)

Elsa Falières, Michael S Engel, André Nel

### ► To cite this version:

Elsa Falières, Michael S Engel, André Nel. Earliest occurrence of Embiidae: A new genus from earliest Eocene Oise amber (Insecta: Embiodea). *Comptes Rendus. Palevol*, 2021, 20 (38), pp.799-805. 10.5852/cr-palevol2021v20a38 . hal-03380464

**HAL Id: hal-03380464**

**<https://hal.sorbonne-universite.fr/hal-03380464>**

Submitted on 15 Oct 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

2020 0248

Earliest occurrence of Embiidae: A new genus from earliest Eocene Oise amber (Insecta: Embiodea)

Le plus ancien Embiidae : un nouveau genre de l'ambre éocène basal de l'Oise (Insecta: Embiodea)

Elsa Falières<sup>a</sup>, Michael S. Engel<sup>b</sup>, André Nel<sup>a,\*</sup>

<sup>a</sup> *Institut Systématique Evolution Biodiversité (ISYEB), Muséum national d'Histoire naturelle, CNRS, Sorbonne Université, Université des Antilles, EPHE, 57 rue Cuvier, CP 50, 75005 Paris, France*

<sup>b</sup> *Division of Entomology, Natural History Museum, and Department of Ecology & Evolutionary Biology, 1501 Crestline Drive – Suite 140, University of Kansas, Lawrence, Kansas 66045, USA*

#### ABSTRACT

A new genus and species of webspinner, *Galloembia raholai* gen. et sp. nov., is described from the earliest Eocene Oise amber of France. The species is the current oldest record of the family, a lineage previously known from only as far back as the middle Eocene Baltic amber and an exceptionally dubious Eocene-Oligocene impression fossil of the central United States. The genus is diagnosed and compared with other genera of Embiidae.

*Keywords:* Insecta; Embiodea; Embiidae; gen. et sp. nov.; oldest record

## RÉSUMÉ

Un nouveau genre et espèce d'embie, *Galloembia raholai* gen. et sp. nov., sont décrits de l'ambre éocène basal de l'Oise, France. Il s'agit du plus ancien représentant des Embiidae, une lignée connue de l'ambre de l'Eocène moyen de la Baltique et par un fossile en compression éocène-oligocène douteux du centre des États-Unis. Le genre est diagnostiqué et comparé aux autres genres d'Embiidae.

*Mots clés* : Insecta; Embiodea; Embiidae; gen. et sp. nov.; plus ancien fossile

\*Corresponding author. E-mail address: [anel@mnhn.fr](mailto:anel@mnhn.fr) (A. Nel).

## 1. Introduction

Webspinners (order Embiodea) are one of the smaller lineages of polyneopterous insects, with presently less than 500 described species. Species are gregarious and live in silken galleries which are spun from characteristic glands in the swollen probasitarsi. Females are apterous and spend much of their lives within these galleries tending to the developing brood, while males are usually fully winged. Although individuals are comparatively soft-bodied and therefore unlikely to preserve in many kinds of sediments, there are remains of Embiodea, albeit few in number, from as far back as the Jurassic. Aside from a single impression from the Eocene-Oligocene boundary of Florissant, Colorado, USA (Cockerell, 1908), the fossil record for the order is confined to inclusions in amber and its higher fidelity of preservation. There are well-documented Embiodea in mid-Cretaceous amber from Myanmar (Cockerell, 1919; Davis, 1939; Engel & Grimaldi, 2006; Engel et al., 2016), early Eocene amber of the Cambay Basin (Engel et al., 2011), middle Eocene amber of the Baltic region (Ross, 1956), and middle

Miocene amber of the Dominican Republic (Szumik, 1994, 1998; Ross, 2003). These fossil occurrences span a number of different families ranging from the putatively primitive Clothodidae to the more derived lineages of Scelembiidae and Anisembiidae. The family Embiidae is a diverse lineage of doubtful monophyly (e.g., Szumik, 1996; Miller et al., 2012; Szumik et al., 2019). Its fossil record is scarce, with a single genus and species in middle Eocene Baltic amber and the aforementioned Eocene-Oligocene fossil of dubious familial attribution. Herein we report the discovery of an early Eocene embiid preserved in Oise amber, which is also the currently oldest representative of the family and the first record of the order for this deposit.

## **2. Material and methods**

The fossil is embedded in a small, clear piece of amber. It was prepared using a diamond disk and examined using a Nikon binocular microscope SMZ 1500. Photographs were taken with an Amscope camera MU900, and the images processed using Adobe Photoshop CS6. The fossil is incomplete, head partial and fragmentary and lacking dorsal anterior part of the body lost at the amber surface. The ventral portion of the body is well visible. The wings are along the outer part of the fossil, with the venation clearly visible. The classification followed herein is that proposed by Miller et al. (2012), while the morphological terminology is adapted from that of Ross (2000). Abbreviations are as follows: HP hypandrium process; LCB left cercus basipodite; LC<sub>1</sub> basal left cercomere; LPPT left paraproct; LPPT-P hook of left paraproct; MA median vein; RC<sub>1</sub> right basal cercomere; 10LP process of left hemitergite; 10RP process of right hemitergite.

## **3. Systematic palaeontology**

Class Insecta Linnaeus, 1758

Order Embiodea Lameere, 1900

Family Embiidae Burmeister, 1839

Subfamily *Incertae sedis*

Genus ***Galloembia*** gen. nov.

*Type species. Galloembia raholai* sp. nov.

*Diagnosis.* Male: Moderate body size (as preserved), at least 7.6 mm. Metabasitarsus with two ventral bladders (or euplantulae) (medial and apical). Fully winged; MA forked well distad forewing midlength. Tenth tergite cleft to base (*i.e.*, hemitergites completely separated with basal membranous connection); right and left hemitergites with short processes; 10LP short, narrow, and arched with acutely rounded apex (10LP analogous to that of *Embia ramburi* Rimsky-Korsakov); 10RP exceptionally short, broad, and bluntly rounded; medial flap apparently absent (or scarcely developed); LPPT enlarged, with a short, subapical LPPT-P; HP poorly developed, apically blunt; LC<sub>1</sub> with large, prominent inner lobe subapical in position, lobe slightly echinulate on dorsal surface, otherwise remainder of LC<sub>1</sub> not echinulate; left and right cercomeres with abundant, elongate, erect, simple setae.

*Etymology.* The new generic name is a combination of the Greek words *Γαλλία* (*Gallia*, meaning, “France”, also of Latin derivation), and *ἐμβιός* (*embiós*, meaning, “lively” or “vivacious”; itself the root of the type genus of the family). The gender of the name is feminine.

***Galloembia raholai*** sp. nov.

(Figs 1-3)

*Type material.* Holotype, MNHN.F.A71313 (PA 17338), head only partly preserved, antennae and mouthparts missing, fore legs incomplete, thorax, abdomen, complete wings, mid and hind legs well preserved), deposited in the MNHN, Paris, France.

*Type locality and horizon.* Le Quesnoy, Chevrère, region of Creil, Oise department, France. earliest Eocene, - 53 Ma, ‘Sparnacian’, level MP7 of the mammal fauna of Dormaal.

*Etymology.* The specific epithet honors Pompeu Rahola, entomologist who first showed Embiodes to the senior author.

*Diagnosis.* As for the genus (*vide supra*).

*Description.* Male, fully winged; body slender, light brown, 7.6 mm long (as preserved); pronotum longer than wide, 1.18 mm long, 0.87 mm wide; mesonotum narrower and longer than pronotum, triangular; without visible longitudinal median sulcus; metanotum quadrate; fore legs only partly preserved; mesobasitarsus 0.16 mm long, mesotarsomere III 0.08 mm long; metabasitarsus 0.24 mm long, metatarsomere III 0.16 mm long; metabasitarsus with two ventral bladders; forewing 5.0 mm long; venation similar to that of *Embia* spp., in particular vein MA bifurcate; abdomen elongate cylindrical, 2.8 mm long, 0.84 wide; terminalia asymmetrical; tergite X completely divided medially, with inner base of left hemitergite well separated from right hemitergite by a membranous area and no median sclerite between; apical process of left hemitergite (10LP) short, narrow, arched, and acutely rounded apically; process of right hemitergite (10RP) broadly and bluntly rounded, short, and smooth, without apical hook; left paraproct (LPPT) enlarged, with short, subapical hook (LPPT-P); left cercomeres not fused; basal left cercomere (LC<sub>1</sub>) with a prominent, inner, subapical lobe, dorsal surface of lobe weakly echinulate, remainder of LC<sub>1</sub> not echinulate; right basal cercomere (RC<sub>1</sub>) elongate and slender.

#### **4. Discussion**

Although Embiidae are likely not a monophyletic group (e.g., Szumik, 1996; Szumik et al., 2008; Miller et al., 2012), the present fossil would generally fall among this assemblage of genera based on the characters outlined by Ross (2007) and Miller et al. (2012). Specifically,

*Galloembia* gen. nov. would be classified within Embiidae as currently circumscribed owing to the following attributes: male fully winged; metabasitarsus with two ventral bladders (medial and apical); forewing vein MA forked; terminalia strongly asymmetrical; tenth tergum completely divided medially, with distinct membranous area between hemitergites; inner base of left hemitergite well separated from right one by a membranous area, no median sclerite between the two halves; right basal cercomere elongate and slender; left cercomeres not fused; basal left cercomere with distinct inner, subapical lobe, with dorsal surface of lobe weakly echinulate.

Given its diversity and lack of homogeneity, there has been no modern attempt to provide a comprehensive revision of Embiidae. Accordingly, we provide comparisons between *Galloembia* and the various extant and fossil genera currently placed therein.

*Galloembia* differs from the embiid genera as follows:

- from *Acrosembia* Ross, 2006: simple, short, narrowly arched 10LP (elongate with inner talon in *Acrosembia*); short, bluntly rounded 10RP (projecting and hooked in *Acrosembia*) (Ross, 2006);
- *Apterembia* Ross, 1957: 10R shortened (10R with a sharp apex); LPPT as wide as long (LPPT wider than long) (Krauss, 1911: pl. 3, fig. 15; Esben-Petersen, 1920; Davis, 1939b: figs 1–10);
- *Arabembia* Ross, 1981; no basal lobe on LC<sub>1</sub> (presence of such a lobe); absence a small echinulations proximally on LC<sub>1</sub> (small echinulations on proximal part of LC<sub>1</sub>); a broader LPPT, as wide as long (LPPT longer than wide) (Ross, 1981: fig. 2);
- *Berlandembia* Davis, 1940: absence of small echinulations proximally on LC<sub>1</sub> (echinulations present in basal part of LC<sub>1</sub>); longer LPPT, as long as wide (LPPT wider than long) (Davis, 1939b: figs 3–5);

- *Chirembia* Davis, 1940: shorter 10LP (10LP with long processes); longer LPPT, as long as wide (LPPT wider than long) (Davis, 1940: figs 14, 17; Ross, 2006);
- *Cleomia* Stefani, 1953: only one median lobe of basal left cercomere (two such lobes); longer LPPT, as long as wide (LPPT wider than long) (Stefani, 1953: fig. 4e-f; Ross, 1966);
- *Dihibocercus* Enderlein, 1912: short apical process of left hemitergite (long apical process); only one median lobe of basal left cercomere (two such lobes); larger LPPT, as long as wide (wider than long) (Davis, 1939a: figs 6, 9);
- *Dinembia* Davis, 1939: absence of indentation on basal right cercomere (present in *Dinembia*); broad LPPT, as wide as long (longer than wide) (Davis, 1939b: figs 6, 7);
- *Donaconethis* Enderlein, 1909: absence of a basal lobe and small echinulations at base of left cercomere (structures present); left paraproct longer than wide (rounded); larger LPPT (LPPT small) (Ross, 1951: fig. 2);
- *Embia* Latreille, 1829: presence of two ventral bladders on metabasitarsus (medial bladder absent in *Embia*) (Ross, 1966, 2006);
- *Electroembia* Ross, 1956 (middle Eocene Baltic amber): more pronounced median lobe of basal left cercomere (basal inner margin straight); curved and shorter apical process of left hemitergite (sharp and straight process) (Ross, 1956: fig. 2; 1966: fig. 1);
- *Enveja* Navás, 1916: median lobe of basal left cercomere rounded (quadrangular); shape of left paraproct; large LPPT (LPPT very small) (Davis, 1939a: 491, figs 5, 6);
- *Leptembia* Krauss, 1911: crenulation only on median lobe of basal left cercomere (crenulation in basal inner part); median lobe of basal left cercomere larger than in *Leptembia* (Krauss, 1911: pl. 5, fig. 23; Ross, 2006);
- *Lithembia* Ross, 1984, *nomen nudum* (Eocene-Oligocene Nearctic, based on an impression fossil without any detail of body structures, terminalia, or most venational



elements (Ross, 1984). It is impossible to attribute this species, "*Embia*" *florissantensis* Cockerell, 1908, to the Embiidae, nor can it be meaningfully assigned to a genus. In fact, Ross (1984) was unable to give any characters to circumscribe the taxon, leaving the generic name as a *nomen nudum*;

- *Machadoembia* Ross, 1952: presence of two well developed ventral bladders on metabasitarsus (medial bladder absent in *Machadoembia*); vein MA forked well distal of middle of forewing (forked more basally); LPPT separated from LCB (fused) (Ross, 1952: 43–44);
- *Metembia* Davis, 1939: only one lobe one LC<sub>1</sub> (two lobes); large left paraproct (reduced); large LPPT (reduced) (Davis, 1939a: fig. 2–3; Ross, 1950: fig. 3);
- *Macrembia* Davis, 1940: LC<sub>1</sub> without a basal indentation (a basal indentation); large left paraproct (reduced); LPPT as long as wide (wide than long) (Davis, 1940: figs. 39–41);
- *Odontembia* Davis, 1939: only one median lobe one LC<sub>1</sub> (two lobes); large LPPT (reduced) (Davis, 1939a: figs 23, 25);
- *Oedembia* Ross, 2007: median lobe of LC<sub>1</sub> close to its apex (in a more basal position); right hemitergite not rounded (rounded) (Ross, 2007: fig. 1);
- *Parachirembia* Davis, 1940: exceptionally short apical process of right hemitergite (very long process); a larger left paraproct (narrow); large LPPT (very small) (Davis, 1940: figs 30, 34);
- *Parembia* Davis, 1939: smaller median lobe of LC<sub>1</sub>; apical process of left hemitergite shorter and less sharp; left paraproct with a more pronounced lobe; large LLPT (small) (Davis, 1939a: figs 23–27; Ross, 1981: fig. 1);

- *Parthenembia* Ross, 1960 (female material, male unknown): hardly comparable to our male fossil; nevertheless, in *Parthenembia*, the second bladder of metabasitarsus is medial, instead of being subapical as in *Galloembia* (Ross, 1960);
- *Pseudembia* Davis, 1939: absence of a basal lobe and presence of a larger median lobe on LC<sub>1</sub> (basal lobe present and median lobe reduced) (Davis, 1939a: figs. 9–19).

## **5. Concluding remark**

The Embioidea are living in the liter, generally under rather warm conditions, but some species are known under Mediterranean climates. Flying males can disperse. Our fossil was probably trapped in resin on the trunk of the producing tree. The order appears extremely rare in the Oise amber (one specimen for ca. 20000 insects), which is probably due to their cryptic way of life.

## **Acknowledgments**

We are sincerely thankful to Thomas Schubnel for photographing the fossil, and to two anonymous reviewers for their comments on the manuscript.

## **References**

- Davis, C., 1939a. Taxonomic notes of the order Embioptera. 6–10. Proceedings of the Linnean Society of New South Wales 64, 474–495.
- Davis, C., 1939b. Taxonomic notes of the order Embioptera. 11–14. Proceedings of the Linnean Society of New South Wales 64, 559–575.
- Davis, C., 1940. Taxonomic notes on the order Embioptera. 15. The genus *Rhagadochir* Enderlein and genera convergent to it. Proceedings of the Linnean Society of New South Wales 65, 171–191.

- Esben-Petersen, T., 1920. New species of neuropterous insects from South Africa (Ephemera, Megaloptera, Embiida). *Annals of the South African Museum* 17, 499–506.
- Krauss, H.A., 1911. Monographie der Embien. *Zoologica* 60, 1–78.
- Miller, K.B., Hayashi, C., Whiting, M.F., Svenson, G.J., Edgerly, J.S., 2012. The phylogeny and classification of Embioptera (Insecta). *Systematic Entomology* 37, 550–570.
- Navás, L., 1916. Neurópteros nuevos o poco conocidos (Séptima serie). *Memorias de la Real Academia de Ciencias y Artes de Barcelona* (3) 12, 219–243.
- Ross, E.S., 1950. The Embiidae of India (Embioptera). *The Wasmann Journal of Biology* 8, 133–153.
- Ross, E.S., 1951. Three new African Embiidae. *Annals and Magazine of Natural History* (12) 4, 381–389.
- Ross, E.S., 1952. The Embioptera of Angola. *Publicacoes Culturais Companhia de Diamantes de Angola* 14, 41–54.
- Ross, E.S., 1956. A new genus of Embioptera from Baltic amber. *Mitteilungen aus dem Geologischen Staatsinstitut in Hamburg* 25, 76–81.
- Ross, E.S., 1960. Parthenogenetic African Embioptera. *The Wasmann Journal of Biology* 18, 297–304.
- Ross, E.S., 1966. The Embioptera of Europe and the Mediterranean region. *Bulletin of the British Museum, (Natural History), Entomology* 17, 273–326.
- Ross, E.S., 1981. Insects of Saudi Arabia. Embiidina. *Fauna of Saudi Arabia* 3, 201–212.
- Ross, E.S., 1984. A synopsis of the Embiidina of the United States. *Proceedings of the Entomological Society of Washington* 86, 82–93.
- Ross, E.S., 2000. EMBIA. Contributions to the biosystematics of the insect order Embiidina. Part 1. Origin, relationships and integumental anatomy of the insect order Embiidina. *Occasional Papers of the California Academy of Sciences* 149, 1–53.

Ross, E.S., 2006. The insect order Embiidina of northeastern Africa and the Red Sea region. *Fauna of Arabia* 22, 287–343.

Ross, E.S., 2007. The Embiidina of eastern Asia, part I. *Proceedings of the California Academy of Sciences* 58, 23–30.

Stefani, R., 1953. Nuovi Embiotteri della Sardegna. *Bollettino della Societa Entomologica Italiana* 83, 84–98.

Szumik, C.A., 1996. The higher classification of the order Embioptera: a cladistic analysis. *Cladistics* 12, 41–64.

Szumik, C., Juárez, M.L., Ramirez, M.J., Goloboff, P., Pereyra, V.V., 2019. Implications of the tympanal hearing organ and ultrastructure of chaetotaxy for the higher classification of Embioptera. *American Museum Novitates* 3933, 1–31.

**Fig. 1.** *Galloembia raholai* gen. et sp. nov., holotype MNHN.F.A71313, habitus. A. Dorsal view. B. Ventral view. Scale bars: 1 mm.

**Fig. 1.** *Galloembia raholai* gen. et sp. nov., holotype MNHN.F.A71313, habitus. A. Vue dorsale. B. Vue ventrale. Barres d'échelles : 1 mm.

**Fig. 2.** *Galloembia raholai* gen. et sp. nov., holotype MNHN.F.A71313, male terminalia. A. Ventral view. B. Dorsal view. Scale bars: 1 mm.

**Fig. 2.** *Galloembia raholai* gen. et sp. nov., holotype MNHN.F.A71313, terminalia mâle. A. Vue ventrale. B. Vue dorsale. Barres d'échelles : 1 mm.

**Fig. 3.** *Galloembia raholai* gen. et sp. nov., holotype MNHN.F.A71313. A. Metabasitarsus in ventral view, arrows indicate position of bladders. B. Forewing. Scale bars: 0.2 mm (A), 2 mm (B).

**Fig. 3.** *Galloembia raholai* gen. et sp. nov., holotype MNHN.F.A71313. A. Métabasitarse, vue ventrale, la fleche indique la position des coussinets. B. Aile antérieure. Barres d'échelles : 0,2 mm (A), 2 mm (B).

**A**



**B**







