



HAL
open science

First Detection in West Africa of a Mutation That May Contribute to Artemisinin Resistance *Plasmodium* *falciparum*

Hui Zhao, Liang Pi, Luyi Zhao, Yucheng Qin, Weilin Zeng, Zheng Xiang, Qi
Yang, Maohua Pan, Xinxin Li, Chunyan Zou, et al.

► **To cite this version:**

Hui Zhao, Liang Pi, Luyi Zhao, Yucheng Qin, Weilin Zeng, et al.. First Detection in West Africa of a Mutation That May Contribute to Artemisinin Resistance *Plasmodium falciparum*. *Frontiers in Genetics*, 2021, 12, pp.701750. 10.3389/fgene.2021.701750 . hal-03409089

HAL Id: hal-03409089

<https://hal.sorbonne-universite.fr/hal-03409089>

Submitted on 29 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



First Detection in West Africa of a Mutation That May Contribute to Artemisinin Resistance *Plasmodium falciparum*

Hui Zhao^{1†}, Liang Pi^{1†}, Luyi Zhao^{1†}, Yucheng Qin², Weilin Zeng¹, Zheng Xiang¹, Qi Yang¹, Maohua Pan², Xinxin Li¹, Chunyan Zou³, Xi Chen¹, Wei Zhao¹, Yuxin Lu², Yanrui Wu⁴, Mengxi Duan¹, Xun Wang¹, Xiaosong Li¹, Dominique Mazier⁵, Yaming Huang^{1,6} and Zhaoqing Yang^{1*}

OPEN ACCESS

Edited by:

Toshihiro Mita,
Juntendo University, Japan

Reviewed by:

Billy Ngasala,
Muhimbili University of Health and
Allied Sciences, Tanzania
Emanuele Micaglio,
IRCCS Policlinico San Donato, Italy
Moritoshi Iwagami,
National Center for Global Health and
Medicine, Japan

*Correspondence:

Zhaoqing Yang
zhaoqingy92@hotmail.com

[†]These authors have contributed
equally to this work

Specialty section:

This article was submitted to
Genetics of Common and Rare
Diseases,
a section of the journal
Frontiers in Genetics

Received: 01 May 2021

Accepted: 31 August 2021

Published: 08 October 2021

Citation:

Zhao H, Pi L, Zhao L, Qin Y, Zeng W,
Xiang Z, Yang Q, Pan M, Li X, Zou C,
Chen X, Zhao W, Lu Y, Wu Y, Duan M,
Wang X, Li X, Mazier D, Huang Y and
Yang Z (2021) First Detection in West
Africa of a Mutation That May
Contribute to Artemisinin Resistance
Plasmodium falciparum.
Front. Genet. 12:701750.
doi: 10.3389/fgene.2021.701750

¹ Department of Pathogen Biology and Immunology, Kunming Medical University, Kunming, China, ² Shanglin County People's Hospital, Guangxi, China, ³ Guangxi Zhuang Autonomous Region People's Hospital, Nanning, China, ⁴ Department of Cell Biology & Genetics, Kunming Medical University, Kunming, China, ⁵ Sorbonne Université, INSERM, CNRS, Centre d'Immunologie et des Maladies Infectieuses, CIMI, Paris, France, ⁶ Guangxi Zhuang Autonomous Region Center for Disease Prevention and Control, Nanning, China

Background: The spread of drug resistance has seriously impacted the effective treatment of infection with the malaria parasite, *Plasmodium falciparum*. Continuous monitoring of molecular marker polymorphisms associated with drug resistance in parasites is essential for malaria control and elimination efforts. Our study describes mutations observed in the resistance genes *Pfkelch13*, *Pfcr1*, and *Pfmdr1* in imported malaria and identifies additional potential drug resistance-associated molecular markers.

Methods: Chinese patients infected in Africa with *P. falciparum* were treated with intravenous (IV) injections of artesunate 240–360 mg for 3–5 days while hospitalized and treated with oral dihydroartemisinin-piperaquine (DHP) for 3 days after hospital discharge. Blood samples were collected and PCR sequencing performed on genes *Pfkelch13*, *Pfcr1*, and *Pfmdr1* from all isolates.

Results: We analyzed a total of 225 patients from Guangxi, China with *P. falciparum* malaria acquired in Africa between 2016 and 2018. All patients were cured completely after treatment. The F446I mutation of the *Pfkelch13* gene was detected for the first time from samples of West African *P. falciparum*, with a frequency of 1.0%. Five haplotypes of *Pfcr1* that encode residues 72–76 were found, with the wild-type CVMNK sequence predominating (80.8% of samples), suggesting that the parasites might be chloroquine sensitive. For *Pfmdr1*, N86Y (13.1%) and Y184F (58.8%) were the most prevalent, suggesting that artemether-lumefantrine may not, in general, be a suitable treatment for the group.

Conclusions: For the first time, this study detected the F446I mutation of the *Pfkelch13* gene from Africa parasites that lacked clinical evidence of resistance. This study provides the latest data for molecular marker surveillance related to antimalarial drug resistance

genes *Pfkelch13*, *Pfprt*, and *Pfmdr1* imported from Africa, in Guangxi, China from Chinese migrate workers.

Clinical Trial Registration: ChiCTROPC17013106.

Keywords: *Plasmodium falciparum*, antimalarial drug, polymorphism, drug resistance genes *Pfkelch13*, *Pfprt*, *Pfmdr1*

INTRODUCTION

Malaria is a serious public health problem in tropical and subtropical areas of the world. There were ~229 million cases and 409,000 malaria-related deaths worldwide in 2019, with Africa shouldering over 94% of the global malaria burden (WHO, 2020). While five species of *Plasmodium* can cause human disease, *Plasmodium falciparum* is most notorious due to high levels of mortality and morbidity associated with infection. The recent decline in malaria prevalence is mainly due to the application of efficacious antimalarial drugs, early diagnosis by RDT, and distribution of long-lasting insecticide-treated nets (LLINs) in malaria-endemic areas. However, the emergence of drug resistance remains a persistent obstacle to control and eliminate malaria. Here we sought to better understand the geography of resistance alleles in hopes of guiding clinical decisions and support malaria surveillance and elimination programs.

Imported malaria is increasing in China owing to increasing global economic integration. Disease among migrant laborers attributable to *P. falciparum* has increased rapidly in recent years (Feng et al., 2014; Liu et al., 2014). From 2011 to 2016, 19,154 imported cases were reported in mainland China, most of which came from Africa (72%) (Lai et al., 2019). In the past, China suffered seriously from endemic malaria, but the occurrence of malaria has been reduced; 2017 marked the first time that no locally acquired cases were reported (Feng et al., 2018). The World Health Organization (WHO) deemed China to have achieved malaria elimination in 2021. Thus, imported disease has become the main challenge remaining to eliminate malaria from China. It is particularly important to continue to monitor malaria imported into China through molecular monitoring. Chinese migrant workers lack prior immunity; thus, most self-administer antimalarial drugs when working in endemic regions, often without medical guidance; such practices may induce drug resistance. Understanding the emergence of drug resistance therefore requires attention to the behaviors and consequences of such migrant workers (Yang et al., 2017; Wang et al., 2018, 2020; Yao et al., 2018; Li et al., 2019; She et al., 2020; Zhao et al., 2020; Shi et al., 2021).

Antimalarial drugs are principal tools for malaria control. In Africa in the 1990s, chloroquine (CQ) was used to treat malaria and saved tens of millions of lives (Trape, 2001). Chloroquine was once highly efficacious against uncomplicated *P. falciparum* infection. Widespread use engendered resistance (CQR) (Young et al., 1963; Fogh et al., 1979; Wellems and Plowe, 2001); the WHO then recommended artemisinin-based combination therapies (ACTs) as the first-line antimalarial treatment of choice. According to a WHO report, Côte d'Ivoire and Gabon used artesunate-amodiaquine as a first-line treatment for *P. falciparum*

in 2003 and artesunate-amodiaquine as a first-line treatment for *P. falciparum* in Cameroon and Liberia in 2004. Mozambique in 2004 used artemether-lumefantrine as a first-line treatment for *P. falciparum*. In Sierra Leone and Ghana, artemether-lumefantrine and artesunate-amodiaquine were used as first-line treatments for *P. falciparum* in 2004, and artemether-lumefantrine and artesunate-amodiaquine were used as first-line treatments for *P. falciparum* in Mali in 2007 (WHO, 2015). Artemether-lumefantrine is used widely in African countries.

Artemisinin resistance (ART-R) has emerged in Africa (Lu et al., 2017a), and its appearance is a major threat to malaria control efforts including those efforts aimed at treating imported malaria. Mutations in the Kelch propeller gene (*Pfkelch13*) have, elsewhere, been linked to reduced efficacy of artemisinin. Particular mutations in the *Pfkelch13* gene discovered to play an important role in ART-R include Y439H, F446I, M476I, R539T, I543T, and C580Y increasing ring-stage survival (RSA) (Straimer et al., 2015; Mita et al., 2016; Zhang et al., 2017; Wang et al., 2018). Mutations that have been detected in Africa samples include T149S, K189T/N, P441S, S459T, D464E, C469F, T474I, A481T, K503E, R539T, R561H, P574L, A578S, C580Y, V589I, V603E, E612K, Q613E, R622I, V650E, G665S, A675V, V692L, and N694K, but not F446I (Conrad et al., 2014a; Torrentino-Madamet et al., 2014; Tacoli et al., 2016; Ikeda et al., 2018; Wang et al., 2018, 2020; Yao et al., 2018; Li et al., 2019; Ocan et al., 2019; She et al., 2020).

Alleles of the *P. falciparum* chloroquine resistance transporter gene (*Pfprt*) have been identified as molecular markers of CQR (Warhurst, 2001; Arieu et al., 2014). These include mutations in amino acids 72–76 (Awasthi and Das, 2013) with K76T being the most common change found in CQR parasites. The *P. falciparum* multidrug resistance protein-1 (*Pfmdr1*) has also been associated with resistance to multiple antimalarial drugs. Mutations at residues 86, 184, 1034, 1042, and 1246 have been shown to be associated with resistance to CQ, amodiaquine (AQ), quinine (QN), mefloquine (MQ), halofantrine (HF), and artemisinin (ART) (Duraisingh et al., 1997; Sidhu et al., 2005; Humphreys et al., 2007; Lekostaj et al., 2008; Tinto et al., 2008; Folarin et al., 2011).

In Africa, artemisinin resistance has emerged, and its appearance threatens current efforts to control malaria (Lu et al., 2017b). Antimalarial drug use among Chinese laborers may contribute to drug pressures. Therefore, monitoring the molecular markers of drug resistance in imported malaria cases complements studies in endemic African populations, enriching the picture of polymorphisms that may contribute to drug resistance. Here, we describe the distribution of *Pfkelch13*, *Pfprt*, and *Pfmdr1* from *P. falciparum* isolates imported from Africa to Guangxi, South China from 2016 to 2018.

MATERIALS AND METHODS

Sample Collection and Treatment

All samples were collected between 2016 and 2018 from Chinese migrant workers that had returned from Africa and who were diagnosed at the Guangxi Shanglin Hospital with uncomplicated *falciparum* malaria. The diagnosis was based on the microscopy of Giemsa-stained blood smears. Venous blood samples (2–5 ml) were collected before treatment. Prospective cases with complex travel histories were excluded from the study (Zhao et al., 2020). All uncomplicated patients were treated with intravenous (IV) injections of artesunate, on day 0 with 120 mg, and days 1, 2, 3, and 4 with 60 mg on each subsequent day during hospitalization. Upon resolution of symptoms, the patients were discharged, and 3 days of oral treatment with dihydroartemisinin and piperazine phosphate tablets (dihydroartemisinin-piperazine, DHP, dihydroartemisinin 40 mg, and piperazine phosphate 320 mg for each tablet) was prescribed, on day 1, day 2, and day 3 with 4 pills, 2 pills, and 2 pills, respectively.

This study was approved by the Ethics Review Committees of the Institutional Review Board (IRB) of Shanglin Hospital. All participants supplied written informed consent. This study was registered on the Chinese Clinical Trial Registry with Registration Number ChiCTROPC17013106.

DNA Extraction and PCR

Parasite genomic DNA was isolated from 2 to 5 ml of venous blood using the High Pure PCR Template Preparation Kit (Roche Biotech Co., Ltd., Germany). The template DNA was amplified by Nest-PCR-sequence of the *Pfkelch13* gene (Gene ID: PF3D7_1343700; Wang et al., 2015), a 145-bp fragment of the *Pfcrtr* gene (Gene ID: PF3D7_0709000; Zhou et al., 2016), and two fragments (N1: 526 bp and N2: 799 bp) of the *Pfmdr1* gene (Gene ID: PF3D7_0523000; Li et al., 2015), including the entire sequence of the *Pfkelch13* gene, *Pfcrtr* mutations at codons 72–76, and *Pfmdr1* mutations at codons 86, 130, 184, 1034, 1042, 1109, and 1246. PCR products were purified, sequenced by the Sangon Biotech Co Ltd (Kunming, China), and analyzed using DNASTAR (DNASTAR Inc., Madison, WI, USA).

Statistical Analyses

Experimental data were analyzed by SPSS 19.0 (SPSS Inc. Chicago, IL, USA). The variation tendency for haplotypes and the frequency mutations of *Pfkelch13*, *Pfcrtr*, and *Pfmdr1* over the study period was evaluated by chi-square test; the trend among the groups from patients from different regions was assessed by chi-square test with $p < 0.05$ being considered significant.

RESULTS

Epidemiologic Date of Cases and Clinical Outcome

A total of 225 uncomplicated *P. falciparum* cases that returned from Africa to Guangxi Province between 2016 and 2018 were analyzed in this study. None of the patients had traveled to any other areas prior to, or after, spending time in Africa (Zhao et al., 2020). Among the infected individuals, most (77.8%, 175/225)

had spent time in West Africa, with 20.4% (46/225) and 1.8% (4/225) having been in Central and South Africa, respectively (Supplementary Table 1). The majority of the patients who returned from West Africa had been in either Kumasi or Asankragua in Ghana. The patients' clinical information is included in Supplementary Table 2. Fever was resolved in 80% of the patients within 2 days of treatment. Patients were discharged on days 3, 4, and 5 (15, 70, and 15%, respectively). Six months after discharge, the patients reported no additional fever.

Prevalence of Mutations in *Pfkelch13*, *Pfcrtr*, and *Pfmdr1* Genes

The complete *Pfkelch13* gene was successfully amplified and sequenced from 92.4% (208/225) of the samples. The sequences were deposited in GenBank (MK877250–MK877457). The distribution of *Pfkelch13* mutations is shown in Figure 1 and Tables 1, 2.

The wild-type allele accounted for 25.5% of the samples and was found in higher frequency in samples from patients who had been in areas other than West Africa ($p < 0.01$). A total of 19 single nucleotide polymorphisms (SNPs), including 9 synonymous and 10 non-synonymous changes, were observed in the gene, of which 6 non-synonymous (T149S, K189T/N, R255K, F446I, G533S, and V589I) had been reported previously (Wang et al., 2015; Boussaroque et al., 2016; Yao et al., 2018) while 4 (G112E, L258M, A569V, and G665S) were new (Table 1). One, two, or three asparagine insertions at codon 142 were found in multiple samples (13.00%, 27/208). The non-synonymous mutations at K189T/N or T149S were most prevalent, being found in 50.9, 5.3, and 1.9% of the samples, respectively (Figure 1). In this study, the mutation F446I was observed in two of the isolates from Ghana (Figure 1 and Table 2, GenBank accession numbers MK877371 and MK877381).

From the 225 samples, the *Pfcrtr* and *Pfmdr1* genes were successfully sequenced from 193 (85.8%) to 204 (90.7%), respectively. The sequences were deposited in GenBank (MZ572562–MZ572754, MZ572358–MZ572561, MZ577609–MZ577812). The mutations found in *Pfcrtr* and *Pfmdr1* are indicated in Tables 3–5. For *Pfcrtr*, there were no polymorphisms at position 72 or 73. Mutations leading to changes at positions M74I and N75E were found in the same abundance, making up 15.5% (30/193) of the total samples analyzed, while mutations impacting residue 76 were most frequent (17.6%; Table 3). Five haplotypes within *Pfcrtr* residues 72–76 were identified, including CVMNK (wild type), CVIET, CVIEK, CVMNT (mutant types), and CVM/I N/E K/T (mixed types) with prevalence of 80.8% (156/193), 13.5% (26/193), 1.6% (3/193), 3.6% (7/193), and 0.5% (1/193), respectively (Table 4). For *Pfmdr1*, the prevalence of polymorphisms impacting positions 184 and 86 totaled 58.8 and 13.1%, respectively. Other, less frequent mutations included D89G (0.5%), F1226Y (3.9%), and D1246Y (2.4%). No mutations at position 1034 or 1042 were detected (Table 3). Eleven haplotypes with differences at residues 86, 184, and 1246 were found, including wild type (NYD), mutation haplotypes (YYD, NFD, YFD, and YYY), and mixed haplotypes (N/Y YD, N/Y FD, N Y/F D, Y Y/F D, N/Y Y/F D, and Y Y/F D/Y), accounting




FIGURE 1 | Prevalence of Pfk13 mutations from studied isolates. *The difference is statistically significant between 2016 and 2018.

TABLE 1 | Prevalence rate of wild and mutant *Pfkelch13* gene from studied isolates.

Mutants	No. (%) of samples from Western Africa	No. (%) of samples from Non-Western Africa	No. (%) total	p-value
Wild strain	32 (20)	21 (43.7)	53 (25.5)	<0.001*
N insertion	3 (1.9)	1 (2.1)	4 (1.9)	0.93
NN insertions	21 (13.1)	1 (2.1)	22 (10.6)	0.03*
NNN insertions	1 (2.1)	0 (0.0)	1 (0.5)	0.58
G112 E	2 (1.2)	0 (0.0)	2 (1.0)	0.44
T149 S	2 (1.2)	2 (4.2)	4 (1.9)	0.20
K189 T/N	93 (58.1)	24 (50)	117 (56.2)	0.32
R255 K	1 (0.6)	1 (2.1)	2 (1.0)	0.36
L258 M	0 (0.0)	1 (2.1)	1 (0.5)	0.07
F446 I	2 (1.2)	0 (0.0)	2 (1.0)	0.44
G533 S	1 (0.6)	0 (0.0)	1 (0.5)	0.58
A569 V	1 (0.6)	0 (0.0)	1 (0.5)	0.58
V589 I	1 (0.6)	0 (0.0)	1 (0.5)	0.58
G665 S	1 (0.6)	0 (0.0)	1 (0.5)	0.58

*The difference is statistically significant.

for 34.3, 4.4, 41.7, 2.9, 2.0, 0.5, 0.5, 10.7, 0.5, 2.0, and 0.5%, respectively (Table 4). Among the haplotypes from samples from migrants from West and Non-West Africa, the wild type (NYD) was found with similar frequency ($p > 0.05$). The NFD and YFD haplotypes show a significant difference among the groups ($p < 0.05$). The haplotype frequency of *Pfmdr1* YYD was markedly reduced during the time period of the study, accounting for 8.3% of the diversity in 2016 but undetected in 2018 ($p < 0.05$; Table 5). The distribution of all *Pfmdr1* haplotypes is shown in Table 5.

DISCUSSION

In our study, several antimalarial molecular markers (*Pfkelch13*, *Pfprt*, *Pfmdr1*) were used to monitor drug resistance in imported

P. falciparum cases returning from Africa to Shanglin County of Guangxi Province during 2016–2018. Resistance to artemisinin, a widely used front-line antimalarial drug, is increasing. Mutations in the *Pfkelch13* gene have served as useful molecular markers for ART-R (Ariey et al., 2014). To date, more than 200 non-synonymous mutations in the *Pfkelch13* gene have been reported (MalariaGEN Plasmodium falciparum Community Project, 2016; WHO, 2018), and several validated resistance mutations in the *Pfkelch13* gene have been described that are associated with delayed parasite clearance *in vitro* or *in vivo*, or both. These include F446I, N458Y, M476I, Y493H, R539T, I543T, P553L, R561H, and C580Y (Amaratunga et al., 2014a,b; Ariey et al., 2014; Ashley et al., 2014; Huang et al., 2015; Straimer et al., 2015; Takala-Harrison et al., 2015; WHO, 2018). These changes have been found in distinct geographic regions. For example,

TABLE 2 | *Pfkelch13* mutations from studied isolates.

Mutation	Type	Source countries
N insertion	NS	Ghana (<i>N</i> = 3) Cameroon (<i>N</i> = 1)
NN insertions	NS	Ghana (<i>N</i> = 20) Cameroon (<i>N</i> = 1) Liberia (<i>N</i> = 1)
NNN insertions	NS	Ghana (<i>N</i> = 1)
L119L*	S	Ghana (<i>N</i> = 3) Liberia (<i>N</i> = 1)
T149S	NS	Ghana (<i>N</i> = 2) Cameroon (<i>N</i> = 1) Congo (<i>N</i> = 1)
K189T	NS	Ghana (<i>N</i> = 81) Cameroon (<i>N</i> = 16) Congo (<i>N</i> = 3) Central African Republic (<i>N</i> = 2) Mozambique (<i>N</i> = 1) Sierra Leone (<i>N</i> = 1) Mali (<i>N</i> = 1) Gabon (<i>N</i> = 1)
K189N	NS	Ghana (<i>N</i> = 8) Mozambique (<i>N</i> = 1) Liberia (<i>N</i> = 2)
R255K	NS	Liberia (<i>N</i> = 1) Cameroon (<i>N</i> = 1)
L258M*	NS	Central African Republic (<i>N</i> = 1)
Y288Y*	S	Ghana (<i>N</i> = 1)
F446I	NS	Ghana (<i>N</i> = 2)
T456C*	S	Ghana (<i>N</i> = 1)
C469C	S	Ghana (<i>N</i> = 2)
Y493Y*	S	Ghana (<i>N</i> = 1)
G496G*	S	Ghana (<i>N</i> = 2)
G533S	NS	Ghana (<i>N</i> = 1)
A569V*	NS	Ghana (<i>N</i> = 1)
V589I	NS	Ghana (<i>N</i> = 1)
A627A	S	Cote d'Ivoire (<i>N</i> = 1)
G665S*	NS	Ghana (<i>N</i> = 1)
A676A*	S	Sierra Leone (<i>N</i> = 1) Cameroon (<i>N</i> = 1)
G690G*	S	Ghana (<i>N</i> = 1) Mali (<i>N</i> = 1)

S, synonymous mutations; NS, non-synonymous mutations.

*Mutation was unreported before.

TABLE 3 | Polymorphisms of *Pfprt* and *Pfmdr1* in studied isolates.

Gene	Codons position	No. of isolates	Prevalence of mutation (%)
<i>Pfprt</i> (<i>N</i> = 193)	M74I	30	15.5
	N75E	30	15.5
	K76T	34	17.6
<i>Pfmdr1</i> (<i>N</i> = 204)	N86Y	27	13.1
	D89G*	1	0.5
	Y184F	120	58.8
	F1226Y	8	3.9
	D1246Y	5	2.4

*Mutation was unreported before.

F446I, R539T, I543T, P553L, and C580Y have been found in high prevalence in the Greater Mekong Subregion (GMS) (Wang et al., 2015), while the most frequent mutation found along the Thailand–Myanmar border and in Cambodia, Thailand, and Laos is the C580Y variant (Ariey et al., 2014; Menard et al., 2016; Phyo et al., 2016; Imwong et al., 2017). Previous research has shown that the mutations at residues 539 and 580 were also observed in migrant workers returning from Ghana (Feng et al.,

2015a), and other mutations were detected from African samples, including T149S, K189T/N, P574L, Q613E, and A675V (Conrad et al., 2014a; Torrentino-Madamet et al., 2014; Tacoli et al., 2016; Ikeda et al., 2018). In this study, two Ghanaian samples with mutations known to convey ART-R phenotypes were found at residue 446 of the *Pfkelch13* gene, marking the first time this genotype has been identified in samples from West Africa. As a validated resistance mutation in the *Pfkelch13*, the mutant F446I is mainly found proximal to the China–Myanmar border and has been associated with high RSA (Zhang et al., 2017; Wang et al., 2018) and delayed parasite clearance (Feng et al., 2015b; Huang et al., 2015; Wang et al., 2018). It has not been detected before in Africa.

The infections of patients treated in this study with IV artesunate followed by oral DHP resolved 3–5 days thereafter, suggesting that ART-R succeeded even against parasites bearing the F446I mutation. The clinical success may relate to the use of IV treatment with artesunate, which is typically reserved for severe cases but which was used for all patients in this study. Treatment modality deserves further consideration as an explanation for clinical success in these instances (Wang et al., 2019). It may also be true that F446I-bearing parasites remained sensitive to artemisinin owing to the presence (or absence) of additional mutations relevant to drug sensitivity. Among all samples, the highest frequency of mutations was K189T/N, T149S, or an insertion of N, NN, and NNN, which is consistent with a previous report from Dakar (Torrentino-Madamet et al., 2014). However, there is no definitive evidence that the mutations at 189 and 149 are related to ART-R (Miotto et al., 2015).

Only three residues (amino acids 74, 75, and 76) were found to be mutated in the *Pfprt* gene, of which 76 had the highest frequency of change with 17.6% prevalence. No mutations were found at positions 72 and 73, which was consistent with previous studies of *P. falciparum* imported to Shandong Province, China (Xu et al., 2018). Previous work has shown that the *Pfprt* 76T mutation was the main marker for CQR (Fidock et al., 2000), while research conducted in Nigeria demonstrated an association between the 76T mutation in the *Pfprt* gene and reduced susceptibility to artemether *in vitro* (Bustamante et al., 2012). In addition, increased frequency of the 76T allele raises concerns about the use of ACTs because it has been found after using artemether-lumefantrine (AL) and dihydroartemisinin-piperaquine (DHAP) in Burkina Faso (Baraka et al., 2015). We also found five haplotypes of the *Pfprt* gene at positions 72–76, including CVMNK (wild type), CVIET, CVIEK, CVMNT (mutation type), and CVM/I N/E K/T (multiple clonal infection). The prevalence of the *Pfprt* wild-type haplotype was high (80.8%), which is consistent with a previous report of the prevalence of this allele in Ghana (90%; Osarfo et al., 2018). This result was also similar to a recent report on imported malaria, in which the prevalence of wild-type *Pfprt* CQ sensitive parasites from different regions of Africa was dominant (Lu et al., 2017c). In this study, CVIET (13.5%) was more frequently observed than other mutant haplotypes, and this finding was similar in samples of malaria parasites acquired from either West Africa or Non-West African regions. The prevalence of molecular markers associated with CQ resistance has decreased since changes in treatment have

TABLE 4 | Distribution of *Pfcr*t and *Pfmdr*1 haplotypes from studied isolates.

Gene	Haplotype	No. (%) of samples from West Africa	No. (%) of samples from Non-West Africa	No. (%) total	p-value
<i>Pfcr</i> t Codon 72–76	CVMNK(wild type)	123 (81.5)	33 (78.6)	156 (80.8)	0.674
	CVIET	19 (12.6)	7 (16.7)	25 (13.5)	0.493
	CVIEK	2 (1.3)	1 (2.4)	3 (1.6)	0.624
	CVMNT	6 (3.9)	1 (2.4)	7 (3.6)	0.625
	CVM/I N/E K/T	1 (0.7)	0 (0.0)	1 (0.5)	0.597
	Total	151	42	193	
<i>Pfmdr</i> 1 Codon 86, 184, 1246	NYD(wild type)	50 (31.2)	20 (45.4)	70 (34.3)	0.079
	YYD	5 (3.1)	4 (9.1)	9 (4.4)	0.088
	NFD	78 (48.7)	7 (15.9)	85 (41.7)	<0.0001*
	YFD	2 (1.2)	4 (9.1)	6 (2.9)	0.006*
	YYY	3 (1.8)	1 (2.3)	4 (2.0)	0.866
	N/Y YD	1 (0.6)	0 (0.0)	1 (0.5)	0.326
	N/Y FD	1 (0.6)	0 (0.0)	1 (0.5)	0.599
	N Y/F D	15 (9.4)	7 (15.9)	22 (10.7)	0.216
	Y Y/F D	1 (0.6)	0 (0.0)	1 (0.5)	0.599
	N/Y Y/F D	3 (1.9)	1 (2.3)	4 (2.0)	0.866
	Y Y/F D/Y	1 (0.6)	0 (0.0)	1 (0.5)	0.599
	Total	160	44	204	

*The difference is statistically significant.

TABLE 5 | Haplotypes of *Pfcr*t and *Pfmdr*1 from studied isolates^a.

Gene	Haplotype ^b	Year No. (%)			p-value
		2016	2017	2018	
<i>Pfcr</i> t	CVMNK	61 (84.7)	60 (75.0)	35 (85.4)	0.818
	CVIET	11 (15.3)	11 (13.8)	4 (9.8)	0.426
	CVIEK	0 (0.0)	3 (3.7)	0 (0.0)	0.708
	CVMNT	0 (0.0)	6 (7.5)	1 (2.4)	0.274
	CVM/I N/E K/T	0 (0.0)	0 (0.0)	1 (2.4)	0.120
	Total	72	80	41	
<i>Pfmdr</i> 1	NYD	23 (27.4)	33 (44.6)	14 (30.4)	0.209
	YYD	7 (8.3)	2 (2.7)	0 (0.0)	0.019 ^c
	NFD	38 (45.2)	26 (35.1)	21 (45.6)	0.831
	YFD	4 (4.8)	1 (1.4)	1 (2.2)	0.315
	YYY	2 (2.4)	2 (2.7)	0 (0.0)	0.414
	N/Y YD	0 (0.0)	1 (1.4)	0 (0.0)	0.810
	N/Y FD	0 (0.0)	0 (0.0)	1 (2.2)	0.126
	N Y/F D	10 (11.9)	5 (6.7)	7 (15.2)	0.749
	Y Y/F D	0 (0.0)	1 (1.4)	0 (0.0)	0.810
	N/Y Y/F D	0 (0.0)	3 (4.0)	1 (2.2)	0.209
	Y Y/F D/Y	0 (0.0)	0 (0.0)	1 (2.2)	0.125
	Total	84	74	46	

^aThe haplotypes were constructed considering codon positions 72–76 of *Pfcr*t and codon positions 86, 184, and 1246 of *Pfmdr*1.

^bAmino acid mutation is in bold type.

^cThe difference is statistically significant.

been enacted (Conrad et al., 2014b). For example, in Malawi and Tanzania, *Pfcr*t alleles associated with CQ sensitivity have reemerged after CQ withdrawal (Kublin et al., 2003; Laufer et al., 2006; Mohammed et al., 2013). This suggests that CQ could

possibly be reintroduced as an effective antimalarial drug in the future (Duah et al., 2013; Lu et al., 2017c).

Mutations in the *Pfmdr*1 gene at residues 86, 184, 1034, 1042, and 1246 have been reported to be associated with resistance to

multiple antimalarial drugs. In our study, we found five non-synonymous mutations at positions 86, 89, 184, 1226, and 1246. The main mutation sites were 86 and 184, being found in 13.1 and 58.8% of the samples, respectively. As far as we know, D89G, which we detected at a frequency of 0.5%, has not been described before as a resistance allele. Its role in drug resistance remains to be confirmed. The F1226Y mutation has previously been detected in imported isolates from Africa (Yao et al., 2018). Together, our data and the literature have revealed that at least 11 haplotypes are possible at positions 86, 184, and 1246 in the *Pfmdr1* gene. The high-frequency N86, 184F, and D1246 alleles have been previously reported to be correlated with selection *in vivo* with AL (Vinayak et al., 2010; Baliraine and Rosenthal, 2011; Thomsen et al., 2011, 2013). A mutation N86Y in *Pfmdr1* increases sensitivity to lumefantrine and mefloquine (Menard and Dondorp, 2017). In Africa, the treatment of uncomplicated malaria has changed from a CQ-based approach to one based on AL and DHAP (Duah et al., 2013; Gadalla et al., 2013). The difference of NFD and YFD that we have observed among imported samples from West Africa and Non-West Africa may be related to subtle differences in treatment schemes (and thereby selective pressure) among the countries in Africa.

In conclusion, this study detected the prevalence of polymorphisms in *Pfkelch13*, *Pfcr1*, and *Pfmdr1* genes in Chinese migrant workers returning to Guangxi Province. Mutations at residue 446 in the *Pfkelch13* gene, which is associated with ART-R (Wang et al., 2018), was detected for the first time in authentic malaria samples from West Africa in our study, but it did not coincide with clinical evidence of ART-R. *Pfcr1* gene polymorphisms were rare, and the wild-type haplotype was more frequently identified, suggesting that CQ could be used as an effective antimalarial drug in the future. The frequencies of *Pfmdr1* N86Y and Y184F alleles were most abundant, while the D1246Y allele was low. These observations suggest that AL may not be an effective choice for treatment.

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

AUTHOR CONTRIBUTIONS

HZ, LP, and LZ: conceptualization, investigation, writing—original draft, and visualization. YQ: validation

REFERENCES

- Amaratunga, C., Witkowski, B., Dek, D., Try, V., Khim, N., Miotto, O., et al. (2014a). *Plasmodium falciparum* founder populations in western Cambodia have reduced artemisinin sensitivity *in vitro*. *Antimicrob. Agents Chemother.* 58, 4935–4937. doi: 10.1128/AAC.03055-14
- Amaratunga, C., Witkowski, B., Khim, N., Menard, D., and Fairhurst, R. M. (2014b). Artemisinin resistance in *Plasmodium falciparum*. *Lancet Infect. Dis.* 14, 449–450. doi: 10.1016/S1473-3099(14)70777-7
- Ariey, F., Witkowski, B., Amaratunga, C., Beghain, J., Langlois, A. C., Khim, N., et al. (2014). A molecular marker of artemisinin-resistant *Plasmodium falciparum* malaria. *Nature* 505, 50–55. doi: 10.1038/nature12876
- Ashley, E. A., Dhorda, M., Fairhurst, R. M., Amaratunga, C., Lim, P., Suon, S., et al. (2014). Spread of artemisinin resistance in *Plasmodium falciparum* malaria. *N. Engl. J. Med.* 371, 411–423. doi: 10.1056/NEJMoa1314981
- Awasthi, G., and Das, A. (2013). Genetics of chloroquine-resistant malaria: a haplotypic view. *Mem. Inst. Oswaldo Cruz* 108, 947–961. doi: 10.1590/0074-0276130274
- and investigation. WZe: investigation, methodology, supervision. WZe and MP: validation and data curation. ZX: data curation. YW: visualization and supervision. DM: writing—review and editing. YH: resources: project administration and conceptualization. ZY: conceptualization, methodology, and funding acquisition. QY, XinL, CZ, YL, WZh, MD, XiaL, and XW: resources and validation. All authors contributed to the article and approved the submitted version.

FUNDING

This study was supported by the National Science Foundation of China (31860604 and U1802286), major science and technology projects of Yunnan Province (2018ZF0081), and the International Science and Technology Cooperation–Yunnan International Science and Technology Cooperation Base (202003AE140004). YL was funded by grants of the Science and Technology Bureau Programs for Science and Technology Development of Nanning (ZC20153012). YQ was funded by grants of Guangxi Zhuang Autonomous Region Health Commission of Scientific Research Project (Z20190892). CZ was funded by grants of the Youth Fund Project of People's Hospital of Guangxi Zhuang Autonomous Region, China (No. QN2017-10). WZe was under the support of the Education Department Fund of Yunnan Province (2019J1184). XC and ZX were under sponsoring from the Yunnan Applied Basic Research Projects–Union Foundation [2018FE001-190 and 2019FE001 (-015)]. YW was supported by the Hundred-Talent Program of Kunming Medical University (60117190439), Foundation of the Education Department of Yunnan Province (2018JS151), and the Innovation Experiment Project of Yunnan Province (202010678064) and Kunming Medical University (2020JXD014).

ACKNOWLEDGMENTS

The authors thank all the patients for volunteering to participate in the study.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fgene.2021.701750/full#supplementary-material>

- Baliraine, F. N., and Rosenthal, P. J. (2011). Prolonged selection of *pfmdr1* polymorphisms after treatment of *falciparum* malaria with artemether-lumefantrine in Uganda. *J. Infect. Dis.* 204, 1120–1124. doi: 10.1093/infdis/jir486
- Baraka, V., Tinto, H., Valea, I., Fitzhenry, R., Delgado-Ratto, C., Mbonye, M. K., et al. (2015). In vivo selection of *Plasmodium falciparum* Pfcrt and Pfdm1 variants by artemether-lumefantrine and dihydroartemisinin-piperaquine in Burkina Faso. *Antimicrob. Agents Chemother.* 59, 734–737. doi: 10.1128/AAC.03647-14
- Boussaroque, A., Fall, B., Madamet, M., Camara, C., Benoit, N., Fall, M., et al. (2016). Emergence of mutations in the K13 propeller gene of *Plasmodium falciparum* isolates from Dakar, Senegal, in 2013–2014. *Antimicrob. Agents Chemother.* 60, 624–627. doi: 10.1128/AAC.01346-15
- Bustamante, C., Folarin, O. A., Gbotosho, G. O., Batista, C. N., Mesquita, E. A., Brindeiro, R. M., et al. (2012). *In vitro*-reduced susceptibility to artemether in *P. falciparum* and its association with polymorphisms on transporter genes. *J. Infect. Dis.* 206, 324–332. doi: 10.1093/infdis/jis359
- Conrad, M. D., Bigira, V., Kapisi, J., Muhindo, M., Kanya, M. R., Havlir, D. V., et al. (2014a). Polymorphisms in K13 and falcipain-2 associated with artemisinin resistance are not prevalent in *Plasmodium falciparum* isolated from Ugandan children. *PLoS ONE* 9:e105690. doi: 10.1371/journal.pone.0105690
- Conrad, M. D., Leclair, N., Arinaitwe, E., Wanzira, H., Kakuru, A., Bigira, V., et al. (2014b). Comparative impacts over 5 years of artemisinin-based combination therapies on *Plasmodium falciparum* polymorphisms that modulate drug sensitivity in Ugandan children. *J. Infect. Dis.* 210, 344–353. doi: 10.1093/infdis/jiu141
- Duah, N. O., Matrevi, S. A., De Souza, D. K., Binnah, D. D., Tamakloe, M. M., Opoku, V. S., et al. (2013). Increased *pfmdr1* gene copy number and the decline in *pfcr* and *pfmdr1* resistance alleles in Ghanaian *Plasmodium falciparum* isolates after the change of anti-malarial drug treatment policy. *Malar. J.* 12:377. doi: 10.1186/1475-2875-12-377
- Duraisingh, M. T., Drakeley, C. J., Muller, O., Bailey, R., Snounou, G., Targett, G. A., et al. (1997). Evidence for selection for the tyrosine-86 allele of the *pfmdr1* gene of *Plasmodium falciparum* by chloroquine and amodiaquine. *Parasitology* 114 (Pt 3), 205–211. doi: 10.1017/S0031182096008487
- Feng, J., Li, J., Yan, H., Feng, X., and Xia, Z. (2015a). Evaluation of antimalarial resistance marker polymorphism in returned migrant workers in China. *Antimicrob. Agents Chemother.* 59, 326–330. doi: 10.1128/AAC.04144-14
- Feng, J., Yan, H., Feng, X. Y., Zhang, L., Li, M., Xia, Z. G., et al. (2014). Imported malaria in China, 2012. *Emerging Infect. Dis.* 20, 1778–1780. doi: 10.3201/eid2010.140595
- Feng, J., Zhang, L., Huang, F., Yin, J. H., Tu, H., Xia, Z. G., et al. (2018). Ready for malaria elimination: zero indigenous case reported in the People's Republic of China. *Malar. J.* 17:315. doi: 10.1186/s12936-018-2444-9
- Feng, J., Zhou, D., Lin, Y., Xiao, H., Yan, H., and Xia, Z. (2015b). Amplification of *pfmdr1*, *pfcr*, *pvmdr1*, and K13 propeller polymorphisms associated with *Plasmodium falciparum* and *Plasmodium vivax* isolates from the China-Myanmar border. *Antimicrob. Agents Chemother.* 59, 2554–2559. doi: 10.1128/AAC.04843-14
- Fidock, D. A., Nomura, T., Talley, A. K., Cooper, R. A., Dzekunov, S. M., Ferdig, M. T., et al. (2000). Mutations in the *P. falciparum* digestive vacuole transmembrane protein Pfcrt and evidence for their role in chloroquine resistance. *Mol. Cell* 6, 861–871. doi: 10.1016/S1097-2765(05)00077-8
- Fogh, S., Jepsen, S., and Effersoe, P. (1979). Chloroquine-resistant *Plasmodium falciparum* malaria in Kenya. *Trans. R. Soc. Trop. Med. Hyg.* 73, 228–229. doi: 10.1016/0035-9203(79)90220-7
- Folarin, O. A., Bustamante, C., Gbotosho, G. O., Sowunmi, A., Zalis, M. G., Oduola, A. M., et al. (2011). *In vitro* amodiaquine resistance and its association with mutations in *pfcr* and *pfmdr1* genes of *Plasmodium falciparum* isolates from Nigeria. *Acta Trop.* 120, 224–230. doi: 10.1016/j.actatropica.2011.08.013
- Gadalla, N. B., Abdallah, T. M., Atwal, S., Sutherland, C. J., and Adam, I. (2013). Selection of *pfdhfr/pfdhps* alleles and declining artesunate/sulphadoxine-pyrimethamine efficacy against *Plasmodium falciparum* eight years after deployment in eastern Sudan. *Malar. J.* 12:255. doi: 10.1186/1475-2875-12-255
- Huang, F., Takala-Harrison, S., Jacob, C. G., Liu, H., Sun, X., Yang, H., et al. (2015). A Single mutation in K13 predominates in Southern China and is associated with delayed clearance of *Plasmodium falciparum* following artemisinin treatment. *J. Infect. Dis.* 212, 1629–1635. doi: 10.1093/infdis/jiv249
- Humphreys, G. S., Merinopoulos, I., Ahmed, J., Whitty, C. J., Mutabingwa, T. K., Sutherland, C. J., et al. (2007). Amodiaquine and artemether-lumefantrine select distinct alleles of the *Plasmodium falciparum* *mdr1* gene in Tanzanian children treated for uncomplicated malaria. *Antimicrob. Agents Chemother.* 51, 991–997. doi: 10.1128/AAC.00875-06
- Ikeda, M., Kaneko, M., Tachibana, S.-I., Balikagala, B., Sakurai-Yatsushiro, M., Yatsushiro, S., et al. (2018). Artemisinin-resistant *Plasmodium falciparum* with high survival rates, Uganda, 2014–2016. *Emerging Infect. Dis.* 24, 718–726. doi: 10.3201/eid2404.170141
- Imwong, M., Suwannasin, K., Kunasol, C., Sutawong, K., Mayxay, M., Rekol, H., et al. (2017). The spread of artemisinin-resistant *Plasmodium falciparum* in the Greater Mekong subregion: a molecular epidemiology observational study. *Lancet Infect. Dis.* 17, 491–497. doi: 10.1016/S1473-3099(17)30048-8
- Kublin, J. G., Cortese, J. F., Njunju, E. M., Mukadam, R. A., Wirima, J. J., Kazembe, P. N., et al. (2003). Reemergence of chloroquine-sensitive *Plasmodium falciparum* malaria after cessation of chloroquine use in Malawi. *J. Infect. Dis.* 187, 1870–1875. doi: 10.1086/375419
- Lai, S., Sun, J., Ruktanonchai, N. W., Zhou, S., Yu, J., Routledge, I., et al. (2019). Changing epidemiology and challenges of malaria in China towards elimination. *Malar. J.* 18:107. doi: 10.1186/s12936-019-2736-8
- Laufer, M. K., Thesing, P. C., Eddington, N. D., Masonga, R., Dzinjalumala, F. K., Takala, S. L., et al. (2006). Return of chloroquine antimalarial efficacy in Malawi. *N. Engl. J. Med.* 355, 1959–1966. doi: 10.1056/NEJMoa062032
- Lekostaj, J. K., Amoah, L. E., and Roepe, P. D. (2008). A single S1034C mutation confers altered drug sensitivity to PfMDR1 ATPase activity that is characteristic of the 7G8 isoform. *Mol. Biochem. Parasitol.* 157, 107–111. doi: 10.1016/j.molbiopara.2007.09.008
- Li, J., Chen, J., Xie, D., Eyi, U. M., Matesa, R. A., Obono, M. M. O., et al. (2015). Molecular mutation profile of Pfcrt and Pfdm1 in *Plasmodium falciparum* isolates from Bioko Island, Equatorial Guinea. *Infect. Genet. Evol.* 36, 552–556. doi: 10.1016/j.meegid.2015.08.039
- Li, J., Shi, Y., Zhang, W., Yan, H., Lin, K., Wei, S., et al. (2019). K13-propeller gene polymorphisms of *Plasmodium falciparum* and the therapeutic effect of artesunate among migrant workers returning to Guangxi, China (2014–2017). *Malar. J.* 18:349. doi: 10.1186/s12936-019-2984-7
- Liu, Y., Hsiang, M. S., Zhou, H., Wang, W., Cao, Y., Gosling, R. D., et al. (2014). Malaria in overseas labourers returning to China: an analysis of imported malaria in Jiangsu Province, 2001–2011. *Malar. J.* 13:29. doi: 10.1186/1475-2875-13-29
- Lu, F., Culleton, R., and Cao, J. (2017a). Artemisinin-Resistant *Plasmodium falciparum* in Africa. *N. Engl. J. Med.* 377:306. doi: 10.1056/NEJMc1705789
- Lu, F., Culleton, R., Zhang, M., Ramaprasad, A., Von Seidlein, L., Zhou, H., et al. (2017b). Emergence of indigenous artemisinin-resistant *Plasmodium falciparum* in Africa. *N. Engl. J. Med.* 376, 991–993. doi: 10.1056/NEJMc1612765
- Lu, F., Zhang, M., Culleton, R. L., Xu, S., Tang, J., Zhou, H., et al. (2017c). Return of chloroquine sensitivity to Africa? Surveillance of African *Plasmodium falciparum* chloroquine resistance through malaria imported to China. *Parasit. Vectors* 10:355. doi: 10.1186/s13071-017-2298-y
- MalariaGEN *Plasmodium falciparum* Community Project (2016). Genomic epidemiology of artemisinin resistant malaria. *Elife* 5:e08714. doi: 10.7554/eLife.08714
- Menard, D., and Dondorp, A. (2017). Antimalarial drug resistance: a threat to malaria elimination. *Cold Spring Harb. Perspect. Med.* 7:a025619. doi: 10.1101/cshperspect.a025619
- Menard, D., Khim, N., Beghain, J., Adegnik, A. A., Shafiu-Alam, M., Amodu, O., et al. (2016). A worldwide map of *Plasmodium falciparum* K13-propeller polymorphisms. *N. Engl. J. Med.* 374, 2453–2464. doi: 10.1056/NEJMoa1513137
- Miotto, O., Amato, R., Ashley, E. A., Macinnis, B., Almagro-Garcia, J., Amaratunga, C., et al. (2015). Genetic architecture of artemisinin-resistant *Plasmodium falciparum*. *Nat. Genet.* 47, 226–234. doi: 10.1038/ng.3189
- Mita, T., Tachibana, S., Hashimoto, M., and Hirai, M. (2016). *Plasmodium falciparum* kelch 13: a potential molecular marker for tackling artemisinin-resistant malaria parasites. *Expert Rev. Anti Infect. Ther.* 14, 125–135. doi: 10.1586/14787210.2016.1106938
- Mohammed, A., Ndaru, A., Kalinga, A., Manjurano, A., Mosha, J. F., Mosha, D. F., et al. (2013). Trends in chloroquine resistance marker, Pfcrt-K76T

- mutation ten years after chloroquine withdrawal in Tanzania. *Malar. J.* 12:415. doi: 10.1186/1475-2875-12-415
- Ocan, M., Akena, D., Nsoby, S., Kanya, M. R., Senono, R., Kinengyere, A. A., et al. (2019). K13-propeller gene polymorphisms in *Plasmodium falciparum* parasite population in malaria affected countries: a systematic review of prevalence and risk factors. *Malar. J.* 18:60. doi: 10.1186/s12936-019-2701-6
- Osarfo, J., Tagbor, H., Magnussen, P., and Alifrangis, M. (2018). Molecular markers of *Plasmodium falciparum* drug resistance in parasitemic pregnant women in the middle forest Belt of Ghana. *Am. J. Trop. Med. Hyg.* 98, 1714–1717. doi: 10.4269/ajtmh.18-0009
- Phyo, A. P., Ashley, E. A., Anderson, T. J. C., Bozdech, Z., Carrara, V. I., Sriprawat, K., et al. (2016). Declining efficacy of artemisinin combination therapy against *P. falciparum* malaria on the thai-myanmar border (2003-2013): the role of parasite genetic factors. *Clin. Infect. Dis.* 63, 784–791. doi: 10.1093/cid/ciw388
- She, D., Wang, Z., Liang, Q., Lu, L., Huang, Y., Zhang, K., et al. (2020). Polymorphisms of *pfprt*, *pfmdr1*, and K13-propeller genes in imported *falciparum* malaria isolates from Africa in Guizhou province, China. *BMC Infect. Dis.* 20:513. doi: 10.1186/s12879-020-05228-8
- Shi, S. M., Shi, T. Q., Chen, S. B., Cui, Y. B., Kassegne, K., Okpeku, M., et al. (2021). Genome-wide scans for ghanaiian *Plasmodium falciparum* Genes under selection from local and chinese host populations. *Front. Cell. Infect. Microbiol.* 11:630797. doi: 10.3389/fcimb.2021.630797
- Sidhu, A. B., Valderramos, S. G., and Fidock, D. A. (2005). *pfmdr1* mutations contribute to quinine resistance and enhance mefloquine and artemisinin sensitivity in *Plasmodium falciparum*. *Mol. Microbiol.* 57, 913–926. doi: 10.1111/j.1365-2958.2005.04729.x
- Straimer, J., Gnädig, N. F., Witkowski, B., Amaratunga, C., Duru, V., Ramadani, A. P., et al. (2015). Drug resistance. K13-propeller mutations confer artemisinin resistance in *Plasmodium falciparum* clinical isolates. *Science* 347, 428–431. doi: 10.1126/science.1260867
- Tacoli, C., Gai, P. P., Bayingana, C., Sift, K., Geus, D., Ndoli, J., et al. (2016). Artemisinin resistance-associated K13 polymorphisms of *Plasmodium falciparum* in Southern Rwanda, 2010-2015. *Am. J. Trop. Med. Hyg.* 95, 1090–1093. doi: 10.4269/ajtmh.16-0483
- Takala-Harrison, S., Jacob, C. G., Arze, C., Cummings, M. P., Silva, J. C., Dondorp, A. M., et al. (2015). Independent emergence of artemisinin resistance mutations among *Plasmodium falciparum* in Southeast Asia. *J. Infect. Dis.* 211, 670–679. doi: 10.1093/infdis/jiu491
- Thomsen, T. T., Ishengoma, D. S., Mmbando, B. P., Lusungu, J. P., Vestergaard, L. S., Theander, T. G., et al. (2011). Prevalence of single nucleotide polymorphisms in the *Plasmodium falciparum* multidrug resistance gene (*Pfmdr-1*) in Korogwe District in Tanzania before and after introduction of artemisinin-based combination therapy. *Am. J. Trop. Med. Hyg.* 85, 979–983. doi: 10.4269/ajtmh.2011.11-0071
- Thomsen, T. T., Madsen, L. B., Hansson, H. H., Tomas, E. V., Charlwood, D., Bygbjerg, I. C., et al. (2013). Rapid selection of *Plasmodium falciparum* chloroquine resistance transporter gene and multidrug resistance gene-1 haplotypes associated with past chloroquine and present artemether-lumefantrine use in Inhambane District, southern Mozambique. *Am. J. Trop. Med. Hyg.* 88, 536–541. doi: 10.4269/ajtmh.12-0525
- Tinto, H., Guekoun, L., Zongo, I., Guiguemde, R. T., D'alessandro, U., and Ouedraogo, J. B. (2008). Chloroquine-resistance molecular markers (*Pfprt* T76 and *Pfmdr-1* Y86) and amodiaquine resistance in Burkina Faso. *Trop. Med. Int. Health* 13, 238–240. doi: 10.1111/j.1365-3156.2007.01995.x
- Torrentino-Madamet, M., Fall, B., Benoit, N., Camara, C., Amalvict, R., Fall, M., et al. (2014). Limited polymorphisms in k13 gene in *Plasmodium falciparum* isolates from Dakar, Senegal in 2012-2013. *Malar. J.* 13:472. doi: 10.1186/1475-2875-13-472
- Trape, J. F. (2001). The public health impact of chloroquine resistance in Africa. *Am. J. Trop. Med. Hyg.* 64, 12–17. doi: 10.4269/ajtmh.2001.64.12
- Vinayak, S., Alam, M. T., Sem, R., Shah, N. K., Susanti, A. I., Lim, P., et al. (2010). Multiple genetic backgrounds of the amplified *Plasmodium falciparum* multidrug resistance (*pfmdr1*) gene and selective sweep of 184F mutation in Cambodia. *J. Infect. Dis.* 201, 1551–1560. doi: 10.1086/651949
- Wang, J., Huang, Y., Zhao, Y., Ye, R., Zhang, D., and Pan, W. (2018). Introduction of F446I mutation in the K13 propeller gene leads to increased ring survival rates in *Plasmodium falciparum* isolates. *Malar. J.* 17:248. doi: 10.1186/s12936-018-2396-0
- Wang, J., Xu, C., Liao, F. L., Jiang, T., Krishna, S., and Tu, Y. (2019). A temporizing solution to “artemisinin resistance”. *N. Engl. J. Med.* 380, 2087–2089. doi: 10.1056/NEJMp1901233
- Wang, X., Ruan, W., Zhou, S., Huang, F., Lu, Q., Feng, X., et al. (2020). Molecular surveillance of *Pfprt* and k13 propeller polymorphisms of imported *Plasmodium falciparum* cases to Zhejiang Province, China between 2016 and 2018. *Malar. J.* 19:59. doi: 10.1186/s12936-020-3140-0
- Wang, Z., Shrestha, S., Li, X., Miao, J., Yuan, L., Cabrera, M., et al. (2015). Prevalence of K13-propeller polymorphisms in *Plasmodium falciparum* from China-Myanmar border in 2007-2012. *Malar. J.* 14:168. doi: 10.1186/s12936-015-0672-9
- Warhurst, D. C. (2001). A molecular marker for chloroquine-resistant *falciparum* malaria. *N. Engl. J. Med.* 344, 299–302. doi: 10.1056/NEJM200101253440411
- Wellems, T. E., and Plowe, C. V. (2001). Chloroquine-Resistant Malaria. *J. Infect. Dis.* 184, 770–776. doi: 10.1086/322858
- WHO (2015). *World Malaria Report*. WHO.
- WHO (2018). *World Malaria Report 2018*. Geneva: World Health Organization.
- WHO (2020). *World Malaria Report 2020*. Geneva: World Health Organization.
- Xu, C., Wei, Q., Yin, K., Sun, H., Li, J., Xiao, T., et al. (2018). Surveillance of antimalarial resistance *Pfprt*, *Pfmdr1*, and *Pfkelch13* polymorphisms in African *Plasmodium falciparum* imported to Shandong Province, China. *Sci. Rep.* 8:12951. doi: 10.1038/s41598-018-31207-w
- Yang, C., Zhang, H., Zhou, R., Qian, D., Liu, Y., Zhao, Y., et al. (2017). Polymorphisms of *Plasmodium falciparum* k13-propeller gene among migrant workers returning to Henan Province, China from Africa. *BMC Infect. Dis.* 17:560. doi: 10.1186/s12879-017-2634-z
- Yao, Y., Wu, K., Xu, M., Yang, Y., Zhang, Y., Yang, W., et al. (2018). Surveillance of genetic variations associated with antimalarial resistance of *Plasmodium falciparum* isolates from returned migrant workers in Wuhan, Central China. *Antimicrob. Agents Chemother.* 62:e02387-17. doi: 10.1128/AAC.02387-17
- Young, M. D., Contacos, P. G., Stitche, J. E., and Willar, J. W. (1963). Drug resistance in *Plasmodium Falciparum* from Thailand. *Am. J. Trop. Med. Hyg.* 12, 305–314. doi: 10.4269/ajtmh.1963.12.305
- Zhang, J., Feng, G. H., Zou, C. Y., Su, P. C., Liu, H. E., and Yang, Z. Q. (2017). Overview of the improvement of the ring-stage survival assay: a novel phenotypic assay for the detection of artemisinin-resistant *Plasmodium falciparum*. *Zool. Res.* 38, 317–320. doi: 10.24272/j.issn.2095-8137.2017.075
- Zhao, L., Pi, L., Qin, Y., Lu, Y., Zeng, W., Xiang, Z., et al. (2020). Widespread resistance mutations to sulfadoxine-pyrimethamine in malaria parasites imported to China from Central and Western Africa. *Int. J. Parasitol. Drugs Drug Resist.* 12, 1–6. doi: 10.1016/j.ijpddr.2019.11.002
- Zhou, R. M., Zhang, H. W., Yang, C. Y., Liu, Y., Zhao, Y. L., Li, S. H., et al. (2016). Molecular mutation profile of *pfprt* in *Plasmodium falciparum* isolates imported from Africa in Henan province. *Malar. J.* 15:265. doi: 10.1186/s12936-016-1306-6

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Zhao, Pi, Zhao, Qin, Zeng, Xiang, Yang, Pan, Li, Zou, Chen, Zhao, Lu, Wu, Duan, Wang, Li, Mazier, Huang and Yang. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.