



HAL
open science

Mapping the molecular and cellular complexity of cortical malformations

Esther Klingler, Fiona Francis, Denis Jabaudon, Silvia Cappello

► **To cite this version:**

Esther Klingler, Fiona Francis, Denis Jabaudon, Silvia Cappello. Mapping the molecular and cellular complexity of cortical malformations. *Science*, 2021, 371 (6527), pp.eaba4517. 10.1126/science.aba4517 . hal-03425482

HAL Id: hal-03425482

<https://hal.sorbonne-universite.fr/hal-03425482>

Submitted on 10 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **TITLE**

2

3 Mapping the molecular and cellular complexity of cortical malformations

4

5

6 **AUTHORS AND AFFILIATIONS**

7 Esther Klingler¹, Fiona Francis^{2,3,4}, Denis Jabaudon^{1,5*} & Silvia Cappello^{6*}

8

9 ¹Department of Basic Neurosciences, University of Geneva, Geneva, Switzerland ; ²INSERM
10 U 1270, Paris, France ; ³Sorbonne University, UMR-S 1270, F-75005 Paris, France ; ⁴Institut
11 du Fer à Moulin, Paris, France ; ⁵Clinic of Neurology, Geneva University Hospital, Geneva,
12 Switzerland ; ⁶Max Planck Institute of Psychiatry, 80804 Munich, Germany.

13 * co-corresponding authors

14

15

16

1 **ABSTRACT**

2 The cerebral cortex is an intricate structure underlying human features such as language and
3 cognition. Cortical functions rely on specialized neurons, which emerge during development
4 from complex molecular and cellular interactions. Neurodevelopmental disorders occur when
5 one or several of these steps are incorrectly executed. While a number of causal genes and
6 disease phenotypes have been identified, the sequence of events linking molecular disruption
7 to clinical expression mostly remains obscure. Here, focusing on human malformations of
8 cortical development, we illustrate how complex interactions at genetic, cellular and circuit
9 levels combinatorially contribute to diversity and variability in disease phenotypes. Through
10 select examples and an online resource, we propose that a multi-level assessment of disease
11 processes is key to identify points of vulnerability and develop novel therapeutic strategies.

12

1 MAIN TEXT

2 The cerebral cortex, or neocortex, is critical to key behavioral functions in mammals, including
3 language, sociability, and fine motor skills. This brain structure consists of several dozens of
4 specialized types of neurons organized across layers and areas, which are generated during
5 development by the divisions of diverse progenitor cells. Newborn neurons undergo sequential
6 molecular programs that drive their specific local and long-range circuit connectivity and adult
7 function (1). The complexity of corticogenesis is staggering and, while necessary for proper
8 cortical function to emerge, the myriads of molecular and cellular developmental processes
9 involved also provide multiple points of vulnerability for “malformations of cortical
10 development” (2), here termed “cortical malformations”. These are usually first detected
11 through their clinical expression, including developmental delay with various combinations of
12 intellectual and motor disabilities, often associated with seizures. Despite the toll on patients
13 and their caregivers, only limited treatments exist and while a number of causal genes have
14 been identified, the sequence of events linking molecular disruption with clinical expression
15 mostly remains obscure.

16 Here, to interrogate cortical malformations and highlight potential points of intervention,
17 we first present basic principles of neocortical development and highlight vulnerable cellular
18 compartments and processes, with particular focus on neurogenesis and neuron migration.
19 Second, we present different “levels” of developmental processes, from genes and gene
20 products to cells, circuits, and clinical expression. Focusing on cortical malformations, we
21 illustrate how complex interactions within and across these levels may account for variable
22 disease patterns. Finally, we propose a framework integrating these different levels of
23 organization, towards better understanding and treatment of the disease, as well as an online
24 resource (<http://genebrowser.unige.ch/humous/>) to identify and compare genetic and cellular
25 developmental processes in mice, human embryos, and human organoids.

26 Neocortical development

27 During embryonic neocortical development, neurons are not born at the place where they will
28 reside in adulthood, but instead migrate relatively long distances to reach their destination. Not
29 only are cortical neurons born at different places, but distinct types of neurons are born at
30 different times, over several days in mice (from embryonic day (E) 11 to 17 (3)) and several
31 weeks in humans (from post-conception week (pcw) 8 to 24 (4)) (Fig. 1A). Generation of the
32 cortex results from billions of cells undergoing four key sequential and partially overlapping

1 processes: (1) progenitor division and neurogenesis, (2) migration, (3) neuritogenesis (*i.e.*
2 extension of axon and dendrites) and (4) synaptogenesis (Fig. 1A-C). Cortical malformations
3 occur when one or several of these developmental events are improperly executed (Fig. 1B);
4 they typically manifest as macroscopic anatomical defects visible with brain imaging
5 techniques such as magnetic resonance imaging (MRI) (Fig. 2). Morphological manifestations
6 include microcephaly (decreased brain size), lissencephaly (disrupted cortical folding and
7 lamination), polymicrogyria (numerous small cortical folds), and heterotopias (abnormally
8 positioned cells in periventricular or subcortical regions) (5). In other cases, more subtle defects
9 are visible only in pathology specimens, *e.g.* in focal cortical dysplasias (abnormal cortical
10 lamination) (2).

11 From E8.5-E9.5 in mice and pcw4 in human (4), progenitors of the dorsal pallium, at the
12 rostral end of the neural tube, which initially form a single-layered neuroepithelium, undergo
13 self-replicating divisions to increase in numbers before generating apical radial glia (aRG, also
14 called apical progenitors) and forming an expanded germinal zone called the ventricular zone
15 (3). Around E11.5 in mice and pcw8 in human, aRGs start dividing asymmetrically to give rise
16 to their first neuronal progeny (Fig.1A) (4). Different types of excitatory cortical neurons are
17 then sequentially born between E11.5 and E17.5 in mice (pcw8-24 in human) and distribute to
18 form six radially-organized layers in an inside-out manner: deep layer neurons, which project
19 subcortically, are born first while superficial layer neurons, which project intracortically, are
20 born last (Fig. 1A) (1). Of note, inhibitory GABAergic neurons, which are not discussed in
21 detail in this review, are born in a distinct germinal zone, the ventral pallium, and migrate at
22 later timepoints into the cortex to form local circuits with glutamatergic neurons (6).

23 *Radial cortical organization* – The laminar organization of the adult neocortex tightly relies
24 on the radial polarity of progenitors and their daughter cells during development. Accordingly,
25 aRGs are highly polarized cells with a radial process extending to the pial surface, necessary
26 to guide the daughter neuron migration (Fig. 1A). Adherens junctions at the apical (*i.e.* towards
27 the ventricle) pole of aRGs are critical to this polarity, as they maintain a cohesive ventricular
28 zone and regulate the balance between proliferation and differentiation through cell-cell
29 interactions (3). Mutations of genes coding for adherens junction proteins, such as
30 α/β -catenins and N-cadherin, indirectly disrupt cell divisions, leading to changes in cortical
31 size and folding, as well as neuronal heterotopias in mice and human (7-10). Similarly,
32 mutations in the genes coding for cell-cell contact proteins such as DCHS1 and FAT4 disrupt
33 aRG polarity and neuronal migration, resulting in human periventricular heterotopias (11).

1 Another notable determinant of aRG polarity is the primary cilium, a specialized organelle
2 which transduces signals from ventricular cerebrospinal fluid (Fig. 1A) (3). Mutations in genes
3 coding for cilium proteins (“ciliopathy” genes, *e.g.* *BBS1*, *BBS7*, *BBS10* and *TMEM216*) affect
4 progenitor proliferation and newborn neuron migration in mice (12) and mutations in the
5 microtubule-associated protein encoding gene *EML1*, which causes neuronal heterotopia in
6 mice and humans, are likewise thought to act by destabilization of the primary cilium (13, 14).

7 *Cycling of progenitor cells* – The ability to undergo cell cycle is a central feature of progenitors.
8 A number of mutations can perturb this process, either by affecting cell divisions, or through
9 premature differentiation into neurons. For example, mutations of genes coding for proteins
10 controlling mitotic spindle organization (Fig. 1A) typically lead to premature differentiation
11 and / or aRG apoptosis, resulting in reduced neurogenesis, and manifesting in humans as
12 microcephaly with or without lissencephaly (*e.g.* *ASPM*, *CENPJ*, *WDR62* (15)). Of note, both
13 premature differentiation into neurons as well as excessive self-replication of progenitors may
14 in principle both result in decreased neuronal output: in one case the progenitor pool is
15 prematurely depleted while in the other, progenitors fail to give rise to neurons. Divisions of
16 aRGs are initially rapid and then slow down as corticogenesis unfolds (from 8 hours at early
17 stages to 18 hours at the end of corticogenesis in mice, and from 23 to 54 hours in primates
18 (16, 17)). The slowing down of cell cycle as corticogenesis proceeds may *per se* confer new
19 properties to daughter cells as low-affinity DNA ligands such as transcription factors are more
20 likely to bind when target sites are available for longer times. Hence, disruption of cell cycle
21 not only affects the number of neurons being generated (typically decreasing neuronal output)
22 but also the relative proportions of daughter neuron types, since distinct types are sequentially
23 produced at different stages of corticogenesis (18).

24 *Diversity of progenitor cells* – aRGs not only generate neurons, but also a distinct type of
25 progenitor, termed intermediate progenitors (also called basal progenitors), which delaminate
26 from the ventricular zone and form a new germinal layer, the subventricular zone (Fig. 1A).
27 Intermediate progenitors are more neurogenic than aRGs, and act to boost neuronal production
28 (19). Consistent with such a function, human mutations affecting intermediate progenitors,
29 such as mutations in *EOMES* (also known as *TBR2*), perturb gyri formation and result in
30 microcephaly, lissencephaly or polymicrogyria (20). In most gyrencephalic species (*i.e.*
31 species with a folded neocortex), including primates, additional progenitor types exist,
32 including basal radial glia (also called outer radial glia), which form a distinct proliferative
33 sub-compartment, the outer subventricular zone (Fig. 1A). Basal radial glia express distinctive

1 genes (*e.g. TMEM14B (21)*) and morphologies (*e.g. many have a basal process but no apical*
2 *process*), and are thought to underlie the disproportionate expansion in superficial layers
3 occurring during gyrification (22).

4 *Neuronal migration* – Following cell division, while undergoing a complex series of
5 morphological changes, postmitotic daughter neurons migrate radially along aRG basal
6 processes away from the ventricular zone to the cortical plate, where they settle in their
7 appropriate final positions. The radial migration of neurons relies on the extension of a leading
8 process, with the leading edge responding to environmental cues. The neuron’s centrosome
9 moves within the leading process allowing somal translocation and locomotion (Fig. 1A) (23).
10 The proper execution of this migratory sequence is key to building the correct cortical
11 architecture and relies tightly on microtubule and actin cytoskeleton-associated proteins.
12 Accordingly, mutations in the genes coding for these proteins have dramatic effects on cortical
13 structure (5, 15), and account for a disproportionate fraction of cortical malformations, as will
14 be detailed in a later section.

15 Preplate splitting, which occurs at E11 in mouse (human pcw9-10) (24), is a critical event
16 during corticogenesis and evolution (Fig. 1A). The glycoprotein Reelin secreted by Cajal-
17 Retzius cells present in the marginal zone plays an important role in these processes. Preplate
18 splitting defects consistently give rise to abnormal cortex development due to aberrant cortical
19 neuron position. Indeed, lamination defects in *Reeler* mice, mutant for the *Reln* gene, not only
20 reflect abnormal neuronal migration, but also abnormal splitting of the preplate into a
21 superficial marginal zone (which becomes layer 1) and a transient deep layer subplate
22 (sometimes called layer 6b) (Fig. 1A). In humans, mutations in *RELN* cause autosomal
23 recessive lissencephaly (25). As discussed in a later section, despite the severe cortical
24 lamination defects in *Reeler* mouse mutants, long-range cortical input/output connectivity
25 appears largely preserved (26, 27). *Pomgnt2* mouse mutants, a model of “cobblestone”
26 lissencephaly caused by mutation in the *POMGNT2* gene in human, also show preplate splitting
27 defects, since abnormal clusters of subplate neurons are observed in superficial layers (28).
28 The subplate itself plays an important role in the radial migration of the neurons which traverse
29 it and, later, in circuit assembly: subplate neurons make transient synaptic connections with
30 migrating neurons, which primes their radial migration (29), and also act as a scaffold for
31 thalamocortical and inhibitory circuit maturation (30, 31). Although mutations specifically
32 affecting this process have not been reported in humans, the subplate is usually visible in

1 human fetal MRIs; in the future, it may be possible to reveal subtle subplate defects underlying
2 more significant circuit dysfunction.

3 *Building circuits* – Neuritogenesis and synaptogenesis are critical for circuit assembly and a
4 bottleneck for cortical function. They have been discussed in detail in recent reviews (32) and
5 will only be briefly covered here. Axon-dendrite polarization starts soon after neurons are born,
6 when they still are migrating (33). Axon and dendrite growth relies on their growth cones which
7 are highly dynamic actin-rich protrusions sensitive to environmental guidance cues (Fig. 1A).
8 As is the case for migration, axonal guidance and extension thus heavily rely on cytoskeletal
9 function. Cortical malformations can be accompanied by anomalies in axonal tracts, most
10 notably in the form of atrophy or agenesis the corpus callosum, which connects both
11 hemispheres in human (34). Other anatomically less prominent axonal pathway defects are
12 likely to occur, and the use of MRI techniques such as diffusion tensor imaging (35) may in
13 the future further address this possibility. Synaptic dysfunction, including inhibitory synaptic
14 dysfunction, is thought to underlie many if not most neuropsychiatric disorders in humans (36),
15 and like migration and neuritogenesis, synaptogenesis involves a highly organized
16 cytoskeleton (Fig. 1A). Neuropsychiatric disorders are not generally associated with salient
17 MRI defects, yet some level of anatomical dysfunction likely exists. For example, patches of
18 cortical disorganization and white matter defects (particularly in the corpus callosum) have
19 been reported in patients with autism spectrum disorder (37, 38). Similarly, decreased neuronal
20 density have been reported in neuropathological specimens of patients with schizophrenia (39).
21 The extent to which these anatomical findings drive the clinical phenotype or are secondary to
22 another underlying process remains unclear.

23 **Sources of complexity in cortical malformations**

24 Cortical malformations have a broad range of clinical expression, including seizures,
25 intellectual disabilities, autistic features, failure to reach developmental milestones, and non-
26 neurological co-morbidities (e.g. dysmorphic facial and musculoskeletal features, skin
27 abnormalities, cardiovascular defects) (5). These features alone often do not allow
28 identification of the cause of the disease, because genetically diverse defects lead to largely
29 overlapping clinical pictures, and impairment of single genes can lead to multiple clinical
30 defects. Thus, while an increasing number of causal genes is being identified and a diversity of
31 anatomical and clinical phenotypes have been recognized (15), the sequence of events linking
32 molecular disruption to final outcome remains mostly out of reach. In the following section,
33 we will highlight potential sources of variability which have precluded establishment of clear

1 causal relationships across genes and molecules, cell types and circuits, and clinical expression
2 (Fig. 3A).

3 *Genetic and molecular dysfunction* – Neurodevelopmental disorders can result from single
4 point mutations in critical genes (*i.e.* monogenic mutations; Fig. 3B), or from DNA sequence
5 variations at multiple loci that together increase disease probability. Convergent, divergent, or
6 mixed relationships between genetic defect(s) and disease phenotype(s) may occur (Fig. 3C-
7 F), and penetrance of the genetic mutation may be incomplete and variable (*i.e.* the mutation
8 is not expressed at the clinical level (Fig. 3G)), together complexifying causal analyses.
9 Understanding this cross-level organization is critical in therapeutic terms: for example,
10 “hidden hubs” represent convergent points of pharmacological intervention across disease
11 phenotypes, while disorders with “hidden complexity” would require distinct treatments
12 despite similar clinical presentations (Fig. 3E, F). The contribution of monogenic causes varies
13 across conditions; in the case of lissencephaly for example, 20% of cases could not be
14 associated with a single genetic cause (40), which may reflect yet un-identified gene mutations,
15 mutations in non-coding regions (41), polygenic or multifactorial inheritance, modulated by
16 the influence of non-genetic factors (*e.g.* infectious or toxic), or somatic mosaicism (42). While
17 monogenic causes appear to predominate in cortical malformations, neuropsychiatric disorders
18 such as autism spectrum disorders and schizophrenia instead appear to mostly involve
19 polygenic processes including through chromosomal deletions (39). Of note, in chromosomal
20 deletions, disease-causing genes need not to be in the affected chromosomal region as local
21 DNA conformation changes may in principle epigenetically alter gene accessibility in other
22 parts of the genome. Although not discussed in detail here, some genetic mutations occur in a
23 clonal manner, *i.e.* do not affect the whole brain but only the progeny of some cells with
24 somatic mutations. Somatic mutations have been clearly established as a cause of focal cortical
25 dysplasia and hemi-megalencephaly (hypertrophy of a single hemisphere) (42), but whether
26 clonal processes are involved in other disorders, and in particular whether at least some
27 psychiatric disorders may result from mosaic synaptic dysfunction, is unknown (43).
28 Overlapping clinical pictures can emerge from mutations in genes which are not apparently
29 functionally correlated. For example, over 200 *de novo* variants have been identified in a cohort
30 of patients with periventricular heterotopia, suggesting that many genes will be associated with
31 this condition, with a high level of convergence in the final outcome (*e.g.* mismigration in the
32 case of heterotopias) (44).

1 Complexifying the situation, the aforementioned developmental processes (*i.e.*
2 neurogenesis, migration, axo- and synaptogenesis) often involve redundant molecular
3 mechanisms. For example, the cyclin-dependent kinase CDK5 (whose gene mutation is
4 associated with lissencephaly in human (45)) is critical for mitosis in mouse progenitors, but
5 also in their daughter neurons for migration and gyrification, synapse formation and even long-
6 term plasticity in adults (Fig. 1C) (46, 47). Similarly, many, if not most, axon guidance
7 molecules also have earlier roles in neuron migration and even neuron generation, and/or in
8 synapse and spine formation and maturation (48-50). For example, the Ephrin receptor EphB2
9 both guides corpus callosum axons and stabilizes synaptic contacts (51, 52). Gene mutations
10 may therefore affect sequential cellular processes in parallel. Such versatility in protein
11 function likely contributes to the complexity and variability of disease processes.

12 *Cytoskeletal perturbations* – Given their involvement in multiple cellular processes, from
13 cytokinesis to migration and neuritogenesis, mutations in genes coding for cytoskeletal proteins
14 are particularly prevalent causes of cortical malformations. Strikingly however, quite different
15 anatomical defects can result from mutations in genes coding for apparently related proteins.
16 For example, in “tubulinopathies”, mutations in a tubulin-encoding gene, *TUBA1A*, generally
17 decreases gyrification in humans, while disruption of another tubulin, *TUBB3* often gives
18 polymicrogyria (53). Microtubule-related proteins can behave similarly, as mutations in the
19 gene coding for doublecortin (*DCX*, which stabilizes microtubules and regulates certain
20 molecular motors during neuronal migration (54)), or *LIS1* (which regulates dynein and is
21 required for interkinetic migration of aRGs as well as neuronal migration (55)), variably result
22 in either subcortical heterotopias or lissencephaly in humans (5). In the case of *DCX*, which is
23 located on the X chromosome, mutations in women typically lead to heterotopia while in men
24 it is associated with lissencephaly, suggesting dose-dependent effects (56).

25 Post-transcriptional and post-translational dysregulations are also a cause of cortical
26 malformations. For example, abnormal alternative splicing can occur in the Filamin A actin-
27 binding transcript (*Flna*), causing periventricular heterotopias in humans (57). Inhibition of a
28 neuronal “poison exon” by the Polypyrimidine Tract Binding Protein PTBP1 is necessary to
29 maintain neural progenitor identity. In the absence of PTBP1 function, de-repression of this
30 poison exon leads to decreased expression of *FLNA* and precocious neuronal differentiation
31 (Fig. 4A). Mutations of proteins modifying the conformation and activity of other proteins,
32 like kinases (*e.g.* PI₃K/AKT pathway (58)), glycosylases (59) and tubulin chaperones /

1 modifiers (*e.g.* Tubulin Folding Cofactor D (*TBCD*) (60)) are further post-translational causes
2 of cortical malformations.

3 Obviously, not all cortical malformations are the result of disrupted cytoskeletal function.
4 Other classes of proteins, such as transcription factors, are also involved (*e.g.* ARX, PAX6,
5 EOMES and FOXC1 whose gene mutations lead to severe cortical malformations in humans
6 (Fig. 2) and mice (20, 61-63)). Such examples are, however, comparatively rare despite their
7 critical function in orchestrating developmental genetic programs. One explanation for this
8 apparent paradox is that many transcription factors may be “too” critical for corticogenesis,
9 such that loss of their function is embryonically lethal.

10 *Disrupted spatio-temporal choreographies* – Normal gene expression is tightly regulated in
11 space and time during development. Hence, abnormal dynamic regulation of gene expression
12 is likely a critical factor in the pathogenesis of cortical malformations. Supporting this
13 possibility, *Flna* mutant mice do not display neuronal migration defects, while acute RNAi-
14 mediated knockdown of *Flna* in rats reproduces the periventricular heterotopias similar to those
15 observed in human patients (64). Also, constitutive *Dcx* mutant mice have a fairly normal
16 neocortex and behavior (65), while acute RNAi-mediated knockdown of *Dcx* causes
17 subcortical band heterotopia (66) (Fig. 4B). In principle, while dysfunction of early-onset genes
18 cascade onto a broad array of subsequent cellular processes and may lead to more severe
19 phenotypes, early-onset defects may also be more easily compensated for than defects affecting
20 late differentiation stages.

21 Spatial regulation of gene expression is also important, particularly in the cortex where
22 functionally specialized areas exist. In many cases, malformations are focal or regionalized
23 rather than diffuse, although what determines their location is usually unknown. Cortical
24 malformations often show antero-posterior gradients (40). For example, *FLNA* mutations
25 typically cause frontal-predominant heterotopias (67), and *GPR56* mutations frontal-
26 predominant polymicrogyria (68). In contrast, *CEP85L* shows higher expression in posterior
27 cortical regions and mutations in this gene lead to posterior-predominant polymicrogyria (69).
28 Some affected genes, however, are ubiquitously expressed (*e.g.* *GPR56* (70)); interactions with
29 locally expressed binding partners or specific developmental dynamics may here account for
30 the stereotypic location of the cortical defect. In some cases, such as *FLNA*, *GPR56* or *EOMES*,
31 spatial restriction of gene expression in regions underlying gyri or sulci (71) could further
32 contribute to impairments in cortical folding. Finally, somatic mutations may also account for

1 the focal nature of a defect (*e.g.* PI₃K pathway (72)), although what determines area location
2 is often unknown (42).

3 *Abnormal cell types* – Cell types are the end-result of the dynamic spatio-temporal expression
4 of genes during development. Cortical malformations typically affect multiple types of neurons
5 and/or progenitors, which is in contrast with neurodegenerative diseases, where single cell-
6 types (*e.g.* dopaminergic neurons in Parkinson disease) are predominantly affected. The
7 apparent lack of cell type specific impairments may reflect that differentiation processes are
8 mostly generic across neuron types, and that cell type identity mostly emerges from the
9 combinatorial interactions of multiple genes rather than single master regulators (73). In some
10 cases (mainly focal cortical dysplasia, autism spectrum disorder, schizophrenia and epilepsy),
11 specific defects have also been reported in inhibitory neurons, astrocytes and oligodendrocytes
12 (43, 74) (Fig. 4C). Failure in astrocyte-dependent removal of extracellular debris may lead to
13 a chronically pro-inflammatory environment, potentially increasing susceptibility to disease,
14 but the extent to which reported changes in non-neuronal cells are reactive to neuronal loss is,
15 however, often unclear (75). Of note, since astrocytes are born towards the end of neurogenesis
16 (likely from the progeny of aRGs), mutations that affect aRG divisions are probably associated
17 with a decreased number of astrocytes, although this has not been systematically assessed.

18 Depending on progenitor types and stage of corticogenesis, daughter neurons differ and the
19 aberrant generation of one of these progenitor types could in principle affect specific progenies.
20 For example, mutations affecting intermediate progenitor function could preferentially affect
21 superficial layers, since such progenitors become more and more prevalent as corticogenesis
22 unfolds and superficial layer neurons are born last. Recently, electrophysiological properties
23 of progenitors have been shown to contribute to the sequential generation of neuronal types
24 (76) and may thus contribute to cortical malformations. Accordingly, mutation in the sodium
25 channel Nav1.3 (SCN3A) gene affects outer subventricular zone progenitors and causes focal
26 polymicrogyria in humans, potentially reflecting excess or altered superficial neuron
27 production (77).

28 *Abnormal circuits* – Although neuronal heterotopia likely affects circuit assembly, neuronal
29 identity and basic circuit properties are often conserved despite abnormal neuronal position. In
30 mice, ectopic neurons usually still express their proper laminar markers and display conserved
31 long-range projections (27, 78, 79) (Fig. 4D). Similarly, L5-type neurons in L4 retain their
32 connectivity, as do L4-type neurons in L2/3 (80, 81) and ectopic neurons still attract the proper
33 subtypes of GABAergic interneurons with whom they make synaptic connections (as if they

1 were normally positioned in the cortex), suggesting that local connectivity might likewise be
2 preserved (82). How GABAergic interneuron migration is affected in microcephaly or other
3 cortical malformations has not been systematically examined (but see (83)), but disruption of
4 this process could provide a cellular/circuit explanation for the seizures seen in some patients.
5 In the *Reeler* mouse, although distinct layers are no longer visible, L4-type neurons still receive
6 proper innervation from the thalamus and somatotopic innervation of the cortex is relatively
7 preserved (26, 27) (Fig. 4D). Supporting spared circuit function, perception, learning and
8 memory in *Reeler* mutant mice are little affected by disrupted cortical lamination (27), although
9 more cortex-targeted behavioral tests would likely reveal some level of dysfunction. Together,
10 these results suggest that neuron identity emerges largely cell-autonomously, such that long-
11 distance projections (and possibly microcircuits) may be relatively spared in neuronal
12 heterotopias.

13 *Behavior and clinical features* – As a consequence of the number and complexity of the
14 molecular and cellular processes involved in cortical disorders, the clinical manifestations are
15 heterogeneous, and include intellectual disabilities, epilepsy, and a broad spectrum of
16 neuropsychiatric manifestations. What determines this spectrum is unknown, but postnatal,
17 environmental factors and overall genetic background may contribute (84). Epilepsy often
18 occurs in the context of cortical malformations, but which factors determine seizure
19 susceptibility is unclear. In mouse models of cortical heterotopia, the ectopic and isotopic
20 cortices remain interconnected, which may contribute to abnormal neuronal activation and
21 seizures seen in human patients (85, 86). In some cases, abnormal electrical activity was
22 initiated within or adjacent to the heterotopia while in others, seizures were initiated from the
23 overlying cortex. This suggests either that aberrantly located neurons are capable of generating
24 dysfunctional circuits at a distance, or that the overlying homotopic cortex is abnormal (87).
25 Like other clinical features, intellectual functions are also variably affected, spanning from
26 essentially normal functions to severe disability, in which case seizures are often associated
27 (88). In the absence of additional biomarkers, this relatively generic array of symptoms and
28 signs, which have historically been the main criterion for disease classification, thus offer
29 limited opportunities for a mechanistic understanding of the disease process.

30 **Towards a multi-level assessment of neurodevelopmental disorders**

31 The examples above highlight several levels (*i.e.* genetic, molecular, cellular, circuit and
32 behavioral) within and across which combinatorial interactions may occur during cortical
33 malformations and preclude a causal understanding of the disease process (Figs. 3-4).

1 Understanding the processes at play at each of these levels for individuals is key to provide
2 patients and their families with prognostic indicators (including biomarkers and, more broadly,
3 recognized “endophenotypes”) and, in the longer term, therapeutic perspectives.

4 Towards these two aims, the use of animal models, including mouse, ferret and non-human
5 primates (the latter being gyrencephalic and thus closer to humans) is important, in particular,
6 at some point, to assess the therapeutic effect of approaches on behavior, including through
7 insertion of human mutations via gene editing technologies (47). *In vitro* human models are
8 also emerging as a promising and feasible avenue: somatic cells can now be reprogrammed
9 into induced pluripotent stem cells (iPSCs) and serve as a basis for the generation of 3D
10 organoids and assembloids of the brain (89, 90). Organoids have been used to study a variety
11 of neurodevelopmental disorders including cortical malformations (*e.g.* *CDK5RAP2* mutation
12 leading to cobblestone lissencephaly in humans) (91). Despite their limitations in terms of
13 reproducibility and restriction to early stages of corticogenesis (gyrification is for example
14 challenging to study in these models) (92), organoids are poised to become broadly used in
15 personalized medicine by giving access to the neurons of the patient under study.

16 Recent advances in single-cell transcriptomics now allow the dynamic developmental
17 expression of genes in emerging cell types to be assessed (93, 94) (Fig. 5A), potentially linking
18 cellular diversity and characteristics with circuits and behavioral repertoires. Comparison of
19 gene expression across brain development in mouse, monkey, human and primate- / human-
20 derived brain organoids is particularly important to identify perturbed processes and select the
21 appropriate study model. As a first step in this direction, here we provide an online resource
22 (<http://genebrowser.unige.ch/humous/>) compiling transcriptional maps across development
23 and neuron differentiation for mouse embryos, human embryos and human-derived organoids
24 (94-96). As an example of an application of such a database, combinatorial analysis of arrays
25 of genes implicated in microcephaly, lissencephaly or polymicrogyria using this resource
26 confirms that microcephaly predominantly affects progenitor function, while lissencephaly and
27 polymicrogyria affect postmitotic neurons (Fig. 5B). In the latter two cases, human data
28 emphasizes that progenitors may also be affected, highlighting the value of *trans*-species
29 comparisons (see arrowheads in Fig. 5B).

30 Multi-omics approaches, in which transcriptomics are combined with genomics,
31 epigenomics and proteomics will provide an opportunity to further unravel the cell type
32 specific processes at play and distinguish core molecular / cellular disease processes from
33 idiosyncratic or stochastic variability. Comparing the distribution of affected processes (at any

1 level of analysis) across patients will help to build a spectrum of abnormal states and predict
2 vulnerable processes (Fig. 5C). Integrative approaches including electrophysiological,
3 imaging, and clinical and biological data from patients, for example using state-of-the art
4 artificial intelligence algorithms (97, 98), may allow the bridging of DNA mutation(s) to gene
5 expression, cellular, anatomy and circuit consequences (Fig. 5D). This will be instrumental for
6 a pathogenic classification of diseases, an essential step for a more precise patient
7 stratification and for the design of personalized diagnostic and therapeutic tools. Although this
8 level of integration may seem futuristic, high-throughput techniques and analytical tools are
9 increasingly available, paving the road for such strategies in a realistically close future.

10
11
12

1 REFERENCES

- 2
- 3 1. D. Jabaudon, Fate and freedom in developing neocortical circuits. *Nature*
4 *Communications*. **8**, 16042–9 (2017).
 - 5 2. M. Severino *et al.*, Definitions and classification of malformations of cortical
6 development: practical guidelines. *Brain*. **32**, 1123–21 (2020).
 - 7 3. E. Taverna, M. Götz, W. B. Huttner, The cell biology of neurogenesis: toward an
8 understanding of the development and evolution of the neocortex. *Annu. Rev. Cell Dev. Biol.*
9 **30**, 465–502 (2014).
 - 10 4. J. C. Silbereis, S. Pochareddy, Y. Zhu, M. Li, N. Šestan, The cellular and molecular
11 landscapes of the developing human central nervous system. *Neuron*. **89**, 248–268 (2016).
 - 12 5. R. Guerrini, W. B. Dobyns, Malformations of cortical development: clinical features and
13 genetic causes. *The Lancet Neurology*. **13**, 710–726 (2014).
 - 14 6. L. Lim, D. Mi, A. Llorca, O. Marín, Development and functional diversification of cortical
15 interneurons. *Neuron*. **100**, 294–313 (2018).
 - 16 7. W.-H. Lien, O. Klezovitch, T. E. Fernandez, J. Delrow, V. Vasioukhin, AlphaE-catenin
17 controls cerebral cortical size by regulating the hedgehog signaling pathway. *Science*. **311**,
18 1609–1612 (2006).
 - 19 8. M. Kadowaki *et al.*, N-cadherin mediates cortical organization in the mouse brain.
20 *Developmental Biology*. **304**, 22–33 (2007).
 - 21 9. A. E. Schaffer *et al.*, Biallelic loss of human CTNNA2, encoding α N-catenin, leads to
22 ARP2/3 complex overactivity and disordered cortical neuronal migration. *Nat Genet*. **50**,
23 1093–1101 (2018).
 - 24 10. A. Winczewska-Wiktor *et al.*, A de novo CTNNB1 nonsense mutation associated with
25 syndromic atypical hyperekplexia, microcephaly and intellectual disability: a case report. *BMC*
26 *Neurology*, 1–6 (2016).
 - 27 11. J. Klaus *et al.*, Altered neuronal migratory trajectories in human cerebral organoids derived
28 from individuals with neuronal heterotopia. *Nat. Med*. **25**, 561–568 (2019).

- 1 12. J. Guo *et al.*, Developmental disruptions underlying brain abnormalities in ciliopathies.
2 *Nature Communications*. **6**, 7857–13 (2015).
- 3 13. M. Kielar *et al.*, Mutations in *Eml1* lead to ectopic progenitors and neuronal heterotopia
4 in mouse and human. *Nat Neurosci*. **17**, 923–933 (2014).
- 5 14. A. Uzquiano *et al.*, Mutations in the heterotopia gene *Eml1/EML1* severely disrupt the
6 formation of primary cilia. *CellReports*. **28**, 1596–1611.e10 (2019).
- 7 15. D. M. Romero, N. Bahi-Buisson, F. Francis, Genetics and mechanisms leading to human
8 cortical malformations. *Seminars in Cell and Developmental Biology*. **76**, 33–75 (2018).
- 9 16. V. S. Caviness, T. Takahashi, Proliferative events in the cerebral ventricular zone. *Brain*
10 *Dev*. **17**, 159–163 (1995).
- 11 17. C. Dehay, H. Kennedy, Cell-cycle control and cortical development. *Nat Rev Neurosci*. **8**,
12 438–450 (2007).
- 13 18. L.-J. Pilaz *et al.*, Prolonged mitosis of neural progenitors alters cell fate in the developing
14 brain. **89**, 83–99 (2016).
- 15 19. A. Pontious, T. Kowalczyk, C. Englund, R. F. Hevner, Role of intermediate progenitor
16 cells in cerebral cortex development. *Dev Neurosci*. **30**, 24–32 (2008).
- 17 20. L. Baala *et al.*, Homozygous silencing of T-box transcription factor *EOMES* leads to
18 microcephaly with polymicrogyria and corpus callosum agenesis. *Nat Genet*. **39**, 454–456
19 (2007).
- 20 21. J. Liu *et al.*, The primate-specific gene *TMEM14B* marks outer radial glia cells and
21 promotes cortical expansion and folding. *Stem Cell*. **21**, 635–649.e8 (2017).
- 22 22. J. H. Lui, D. V. Hansen, A. R. Kriegstein, Development and evolution of the human
23 neocortex. *Cell*. **146**, 18–36 (2011).
- 24 23. C. G. Silva, E. Peyre, L. Nguyen, Cell migration promotes dynamic cellular interactions
25 to control cerebral cortex morphogenesis. *Nat Rev Neurosci*. **20**, 318–329 (2019).
- 26 24. Z. Molnár *et al.*, Comparative aspects of cerebral cortical development. *European Journal*
27 *of Neuroscience*. **23**, 921–934 (2006).

- 1 25. S. E. Hong *et al.*, Autosomal recessive lissencephaly with cerebellar hypoplasia is
2 associated with human RELN mutations. *Nat Genet.* **26**, 93–96 (2000).
- 3 26. V. S. Caviness, Patterns of cell and fiber distribution in the neocortex of the reeler mutant
4 mouse. *J. Comp. Neurol.* **170**, 435–447 (1976).
- 5 27. R. J. Wagener, C. David, S. Zhao, C. A. Haas, J. F. Staiger, The somatosensory cortex of
6 reeler mutant mice shows absent layering but intact formation and behavioral activation of
7 columnar somatotopic maps. *Journal of Neuroscience.* **30**, 15700–15709 (2010).
- 8 28. N. Nakagawa, H. Yagi, K. Kato, H. Takematsu, S. Oka, Ectopic clustering of Cajal-
9 Retzius and subplate cells is an initial pathological feature in Pomgnt2-knockout mice, a model
10 of dystroglycanopathy. *Sci Rep.* **5**, 11163 (2015).
- 11 29. C. Ohtaka-Maruyama *et al.*, Synaptic transmission from subplate neurons controls radial
12 migration of neocortical neurons. *Science.* **360**, 313–317 (2018).
- 13 30. A. Hoerder Suabedissen, Z. Molnár, Development, evolution and pathology of neocortical
14 subplate neurons. *Nat Rev Neurosci.* **16**, 133–146 (2015).
- 15 31. P. O. Kanold, R. Deng, X. Meng, The integrative function of silent synapses on subplate
16 neurons in cortical development and dysfunction. *Front. Neuroanat.* **13**, 41 (2019).
- 17 32. C. R. Cadwell, A. Bhaduri, M. A. Mostajo-Radji, M. G. Keefe, T. J. Nowakowski,
18 Development and arealization of the cerebral cortex. *Neuron.* **103**, 980–1004 (2019).
- 19 33. C. Xu *et al.*, Radial glial cell-neuron interaction directs axon formation at the opposite side
20 of the neuron from the contact site. *J. Neurosci.* **35**, 14517–14532 (2015).
- 21 34. T. J. Edwards, E. H. Sherr, A. J. Barkovich, L. J. Richards, Clinical, genetic and imaging
22 findings identify new causes for corpus callosum development syndromes. *Brain.* **137**, 1579–
23 1613 (2014).
- 24 35. M. Wahl, A. J. Barkovich, P. Mukherjee, Diffusion imaging and tractography of
25 congenital brain malformations. *Pediatr Radiol.* **40**, 59–67 (2010).
- 26 36. P. Monteiro, G. Feng, SHANK proteins: roles at the synapse and in autism spectrum
27 disorder. *Nat Rev Neurosci.* 1–11 (2017).

- 1 37. R. Stoner *et al.*, Patches of disorganization in the neocortex of children with autism. *N*
2 *Engl J Med.* **370**, 1209–1219 (2014).
- 3 38. D. Dimond *et al.*, Reduced white matter fiber density in autism spectrum disorder. *Cereb.*
4 *Cortex.* **29**, 1778–1788 (2019).
- 5 39. M. Li *et al.*, Integrative functional genomic analysis of human brain development and
6 neuropsychiatric risks. *Science.* **362**, eaat7615–15 (2018).
- 7 40. N. Di Donato *et al.*, Analysis of 17 genes detects mutations in 81% of 811 patients with
8 lissencephaly. *Genet. Med.* **20**, 1354–1364 (2018).
- 9 41. E. Perenthaler, S. Yousefi, E. Niggli, T. S. Barakat, Beyond the Exome: The non-coding
10 genome and enhancers in neurodevelopmental disorders and malformations of cortical
11 development. *Front. Cell. Neurosci.* **13**, 352 (2019).
- 12 42. A. M. D’Gama, C. A. Walsh, Somatic mosaicism and neurodevelopmental disease. *Nat*
13 *Neurosci.* **21**, 1504–1514 (2018).
- 14 43. D. Velmeshev *et al.*, Single-cell genomics identifies cell type-specific molecular changes
15 in autism. *Science.* **364**, 685–689 (2019).
- 16 44. E. L. Heinzen *et al.*, De novo and inherited private variants in MAP1B in periventricular
17 nodular heterotopia. *PLoS Genet.* **14**, e1007281 (2018).
- 18 45. D. Magen *et al.*, Autosomal recessive lissencephaly with cerebellar hypoplasia is
19 associated with a loss-of-function mutation in CDK5. *Hum. Genet.* **134**, 305–314 (2015).
- 20 46. A. H. Hawasli *et al.*, Cyclin-dependent kinase 5 governs learning and synaptic plasticity
21 via control of NMDAR degradation. *Nat Neurosci.* **10**, 880–886 (2007).
- 22 47. Y. Shinmyo *et al.*, Folding of the cerebral cortex requires CDK5 in upper-layer neurons in
23 gyrencephalic mammals. *Cell Reports.* **20**, 2131–2143 (2017).
- 24 48. H. Blockus, A. Chédotal, The multifaceted roles of Slits and Robos in cortical circuits:
25 from proliferation to axon guidance and neurological diseases. *Current Opinion in*
26 *Neurobiology.* **27**, 82–88 (2014).

- 1 49. W. D. Andrews, M. Barber, J. G. Parnavelas, Slit-Robo interactions during cortical
2 development. *Journal of Anatomy*. **211**, 188–198 (2007).
- 3 50. E. Arbeille *et al.*, Cerebrospinal fluid-derived Semaphorin3B orients neuroepithelial cell
4 divisions in the apicobasal axis. *Nature Communications*, 1–14 (2019).
- 5 51. M. B. Dalva *et al.*, EphB receptors interact with NMDA receptors and regulate excitatory
6 synapse formation. *Cell*. **103**, 945–956 (2000).
- 7 52. M. S. Kayser, M. J. Nolt, M. B. Dalva, EphB receptors couple dendritic filopodia motility
8 to synapse formation. *Neuron*. **59**, 56–69 (2008).
- 9 53. N. Bahi-Buisson, M. Cavallin, Tubulinopathies overview (available at
10 <https://www.ncbi.nlm.nih.gov/books/>).
- 11 54. F. Francis *et al.*, Doublecortin is a developmentally regulated, microtubule-associated
12 protein expressed in migrating and differentiating neurons. *Neuron*. **23**, 247–256 (1999).
- 13 55. J.-W. Tsai, K. H. Bremner, R. B. Vallee, Dual subcellular roles for LIS1 and dynein in
14 radial neuronal migration in live brain tissue. *Nat Neurosci*. **10**, 970–979 (2007).
- 15 56. R. Guerrini, T. Filippi, Neuronal migration disorders, genetics, and epileptogenesis. *J*
16 *Child Neurol*. **20**, 287–299 (2005).
- 17 57. X. Zhang *et al.*, Cell-type-specific alternative splicing governs cell fate in the developing
18 cerebral cortex. *Cell*. **166**, 1147–1162.e15 (2016).
- 19 58. L. A. Jansen *et al.*, PI3K/AKT pathway mutations cause a spectrum of brain malformations
20 from megalencephaly to focal cortical dysplasia. *Brain*. **138**, 1613–1628 (2015).
- 21 59. A. R. Nickolls, C. G. Bönnemann, The roles of dystroglycan in the nervous system:
22 insights from animal models of muscular dystrophy. *Dis Model Mech*. **11** (2018),
23 doi:10.1242/dmm.035931.
- 24 60. E. Flex *et al.*, Biallelic mutations in TBCD, encoding the tubulin folding cofactor D,
25 perturb microtubule dynamics and cause early-onset encephalopathy. *Am. J. Hum. Genet*. **99**,
26 962–973 (2016).

- 1 61. M. Kato *et al.*, Mutations of ARX are associated with striking pleiotropy and consistent
2 genotype-phenotype correlation. *Hum. Mutat.* **23**, 147–159 (2004).
- 3 62. L. K. Davis *et al.*, Pax6 3' deletion results in aniridia, autism and mental retardation. *Hum.*
4 *Genet.* **123**, 371–378 (2008).
- 5 63. M. Pringsheim *et al.*, Structural brain anomalies in patients with FOXP1 syndrome and in
6 Foxg1^{+/-} mice. *Ann Clin Transl Neurol.* **6**, 655–668 (2019).
- 7 64. A. Carabalona *et al.*, A glial origin for periventricular nodular heterotopia caused by
8 impaired expression of Filamin-A. *Human Molecular Genetics.* **21**, 1004–1017 (2012).
- 9 65. J. C. Corbo *et al.*, Doublecortin is required in mice for lamination of the hippocampus but
10 not the neocortex. *Journal of Neuroscience.* **22**, 7548–7557 (2002).
- 11 66. S. Sahu *et al.*, Spontaneous epileptiform activity in a rat model of bilateral subcortical band
12 heterotopia. *Epilepsia.* **60**, 337–348 (2019).
- 13 67. E. Parrini *et al.*, Periventricular heterotopia: phenotypic heterogeneity and correlation with
14 Filamin A mutations. *Brain.* **129**, 1892–1906 (2006).
- 15 68. X. Piao *et al.*, G protein-coupled receptor-dependent development of human frontal cortex.
16 *Science.* **303**, 2033–2036 (2004).
- 17 69. A. Kodani *et al.*, Posterior neocortex-specific regulation of neuronal migration by CEP85L
18 identifies maternal centriole-dependent activation of CDK5. *Neuron.* **106**, 246–255 (2020).
- 19 70. B.-I. Bae *et al.*, Evolutionarily dynamic alternative splicing of GPR56 regulates regional
20 cerebral cortical patterning. *Science.* **343**, 764–768 (2014).
- 21 71. C. De Juan Romero, C. Bruder, U. Tomasello, J. M. Sanz Anquela, V. Borrell, Discrete
22 domains of gene expression in germinal layers distinguish the development of gyrencephaly.
23 *The EMBO Journal.* **34**, 1859–1874 (2015).
- 24 72. G. M. Mirzaa *et al.*, Characterisation of mutations of the phosphoinositide-3-kinase
25 regulatory subunit, PIK3R2, in perisylvian polymicrogyria: a next-generation sequencing
26 study. *The Lancet Neurology.* **14**, 1182–1195 (2015).

- 1 73. L. Telley, D. Jabaudon, A mixed model of neuronal diversity. *Nature*. **555**, 452–454
2 (2018).
- 3 74. X. Jin *et al.*, In vivo Perturb-Seq reveals neuronal and glial abnormalities associated with
4 Autism risk genes. *BioRxiv*. **10**, 2233–37 (2019).
- 5 75. M. Kielbinski, K. Gzielo, Z. Soltys, Review: Roles for astrocytes in epilepsy: insights from
6 malformations of cortical development. *Neuropathol Appl Neurobiol*. **42**, 593–606 (2016).
- 7 76. I. Vitali *et al.*, Progenitor hyperpolarization regulates the sequential generation of neuronal
8 subtypes in the developing neocortex. *Cell*. **174**, 1264–1276.e15 (2018).
- 9 77. R. S. Smith *et al.*, Sodium channel SCN3A (NaV1.3) regulation of human cerebral cortical
10 folding and oral motor development. *Neuron*. **99**, 905–913.e7 (2018).
- 11 78. F. Schottler, D. Couture, A. Rao, H. Kahn, K. S. Lee, Subcortical connections of
12 normotopic and heterotopic neurons in sensory and motor cortices of the tish mutant rat. *J*
13 *Comp. Neurol*. **395**, 29–42 (1998).
- 14 79. A. Croquelois *et al.*, Characterization of the HeCo mutant mouse: a new model of
15 subcortical band heterotopia associated with seizures and behavioral deficits. *Cereb. Cortex*.
16 **19**, 563–575 (2009).
- 17 80. A. De la Rossa *et al.*, In vivo reprogramming of circuit connectivity in postmitotic
18 neocortical neurons. *Nat Neurosci*. **16**, 193–200 (2013).
- 19 81. E. Klingler, *et al.*, A translaminal genetic logic for the circuit identity of intracortically-
20 projecting neurons. *Curr Biol.*, **29**, 332–339 (2019).
- 21 82. S. Lodato *et al.*, Excitatory projection neuron subtypes control the distribution of local
22 inhibitory interneurons in the cerebral cortex. *Neuron*. **69**, 763–779 (2011).
- 23 83. M. Thom, L. Martinian, J. G. Parnavelas, S. M. Sisodiya, Distribution of cortical
24 interneurons in grey matter heterotopia in patients with epilepsy. *Epilepsia*. **45**, 916–923
25 (2004).
- 26 84. S. S. Jeste, D. H. Geschwind, Disentangling the heterogeneity of autism spectrum disorder
27 through genetic findings. *Nat Rev Neurol*. **10**, 74–81 (2014).

- 1 85. R. Mai *et al.*, A neuropathological, stereo-EEG, and MRI study of subcortical band
2 heterotopia. *Neurology*. **60**, 1834–1838 (2003).
- 3 86. J. A. Christodoulou *et al.*, Integration of gray matter nodules into functional cortical
4 circuits in periventricular heterotopia. *Epilepsy Behav.* **29**, 400–406 (2013).
- 5 87. S. V. Kothare *et al.*, Seizure onset from periventricular nodular heterotopias: depth-
6 electrode study. *Neurology*. **51**, 1723–1727 (1998).
- 7 88. P. Chiurazzi, F. Pirozzi, Advances in understanding - genetic basis of intellectual
8 disability. *F1000Res.* **5** (2016), doi:10.12688/f1000research.7134.1.
- 9 89. E. Di Lullo, A. R. Kriegstein, The use of brain organoids to investigate neural development
10 and disease. *Nat Rev Neurosci.* **18**, 573–584 (2017).
- 11 90. S. P. Pasca, The rise of three-dimensional human brain cultures. *Nature.* **553**, 437–445
12 (2018).
- 13 91. M. A. Lancaster *et al.*, Cerebral organoids model human brain development and
14 microcephaly. *Nature.* **501**, 373–379 (2013).
- 15 92. S. Velasco *et al.*, Individual brain organoids reproducibly form cell diversity of the human
16 cerebral cortex. *Nature.* **570**, 523–527 (2019).
- 17 93. J. A. Farrell *et al.*, Single-cell reconstruction of developmental trajectories during
18 zebrafish embryogenesis. *Science.* **360**, eaar3131 (2018).
- 19 94. L. Telley *et al.*, Temporal patterning of apical progenitors and their daughter neurons in
20 the developing neocortex. *Science.* **364**, eaav2522–9 (2019).
- 21 95. T. J. Nowakowski *et al.*, Spatiotemporal gene expression trajectories reveal developmental
22 hierarchies of the human cortex. *Science.* **358**, 1318–1323 (2017).
- 23 96. A. A. Pollen *et al.*, Establishing cerebral organoids as models of human-specific brain
24 evolution. *Cell.* **176**, 743–756 (2019).
- 25 97. A. Esteva *et al.*, Dermatologist-level classification of skin cancer with deep neural
26 networks. *Nature*, 1–12 (2017).

1 98. N. Schwalbe, B. Wahl, Artificial intelligence and the future of global health. *The Lancet*.
2 **395**, 1579–1586 (2020).

3

4

1 **ACKNOWLEDGMENTS**

2 We thank members of the Jabaudon, Cappello and Francis laboratories for insightful
3 discussions and suggestions, Natalia Baumann and Quentin LoGiudice for their support with
4 the bioinformatic analyses, Julien Prados for setting up the
5 <http://genebrowser.unige.ch/humous/> website and Sergi Roig for help in figure design. We
6 thank Stéphanie Baulac and Camilla Bellone for comments on this review and Nadia Bahi-
7 Buisson and the NeuroMIG consortium for fruitful interactions and for providing gene-disorder
8 information. We thank colleagues whose work we were not able to cite due to space constraints
9 for their understanding. SC, DJ and FF laboratories take part in the NeuroMIG COST action
10 (CA16118).

11 **Fundings:** This work is supported by ERA-Net E-Rare (HETEROMICS ERARE 18-049) and
12 national funding agencies.

13 **Author contributions:** D.J., E.K., F.F. & S.C. conceived and designed the presented work.
14 D.J., E.K. & S.C. wrote the manuscript, with the help of F.F. E.K. performed the data analyses
15 and designed the figures.

16 **Competing interests:** The authors declare no competing interests.

17 **Data and materials availability:** In Fig. 5, data from mouse embryos are from (94); data
18 from human embryos and human-derived organoids are from (95, 96).

19

20

1 **FIGURE LEGENDS**

2 **Fig. 1: Key steps of corticogenesis and their susceptibility to disease.** (A) Summary of
3 embryonic corticogenesis. One neuron can be followed from its birth at the ventricular zone,
4 to its migration toward the cortical plate, and then its maturation (arrow). The role of
5 cytoskeletal processes at each of these steps is highlighted on the right. (B) Literature-based
6 summary of knowledge on the involvement of these developmental steps in specific
7 neurodevelopmental disorders. The numbers of publications were found using the text mining
8 EasyPubMed R package, which automatically identifies publications containing given the
9 combinations of keywords indicated (*e.g.* “microcephaly” and “proliferation”). These numbers
10 were then normalized for number of publications for each of the developmental steps and each
11 of the disorders. The dendrogram represents unbiased clustering of the diseases based on this
12 data. (C) Overlapping cellular processes and pleiotropic molecular mechanisms involving
13 CDK5 across development and in the adult brain. aRG, apical radial glia; bRG, basal radial
14 glia; CP, cortical plate; E, embryonic day; IN, interneuron; IPC, intermediate progenitor cell;
15 N, excitatory neuron; pcw; post-conception week; SVZ, subventricular zone; VZ, ventricular
16 zone; WM, white matter.

17

18 **Fig. 2: Cellular mechanisms at play in human cortical malformations.** Cells with abnormal
19 developmental trajectories are in pink. Reduced progenitor proliferation (sometimes associated
20 with abnormal neuron migration) leads to microcephaly. Abnormal neuron position can lead
21 to agyria and lissencephaly (reduced migration and/or progenitor misposition), polymicrogyria
22 (increased migration and/or excess of basal radial glia), cobblestone lissencephaly (excessive
23 migration), periventricular heterotopia (neurons stay close to the ventricle) or subcortical band
24 heterotopia (neurons stay below the cortex). In focal cortical dysplasia, clonal mutations affect
25 subpopulations of mature cells, which become dysmorphic (balloon cells, dysmorphic
26 neurons). Genes mentioned in this review are listed (bold: genes with published human-derived
27 brain organoid models, *: gene studied in the ferret, green: genes coding for transcription
28 factors). Pink arrowheads on MRIs highlight the malformations. MRI illustrations are from
29 (15) and (2) for polymicrogyria and focal cortical dysplasia (Reprinted from *Semin Cell Dev*
30 *Biol* vol 76, D. M. Romero, N. Bahi-Buisson & F. Francis, Genetics and mechanisms leading
31 to human cortical malformations, pages 33-75, 2018, with permission from Elsevier;
32 Definitions and classification of malformations of cortical development: practical guidelines;
33 M. Severino et al., *Brain* 2020; awaa174, with permission from Oxford University Press).

1

2 **Fig. 3: From gene mutations to diseases: levels of complexity.** (A) Levels of organization
3 during corticogenesis, from DNA (*i.e.* genes, represented by letters) to RNA and proteins (*i.e.*
4 gene expression), to cells, circuit and anatomy, and phenotype. Each circle represents a given
5 feature of that level (*e.g.* a gene, a protein, a circuit). Interactions within levels are linked
6 through complex relationships (dashed lines) to states at other levels (black lines). (B-G)
7 Examples of abnormal feature relationships across levels in disease (highlighted in pink). In
8 **B-D**, linear (monogenic), convergent (polygenic) and divergent relationships between genetic
9 defects and phenotype(s) are illustrated. (E) Hidden complexity: note that from a gene-
10 phenotype point of view, this relationship appears linear. (F) Hidden hub: different genes may
11 lead to distinct phenotypes *via* hidden feature hubs influencing several downstream pathways.
12 (G) Incomplete penetrance: not all genetic features are expressed at other levels. In A, source
13 images for human brain section and phenotype are from <https://msu.edu/~brains/brains/> and
14 ©biorender.com, respectively.

15

16 **Fig. 4: Spatiotemporal and cell type regulations of gene expression and malformations.**
17 (A) Alternative splicing of *FLNA* gene regulates its role in proliferation *versus* neuron
18 differentiation of cortical progenitors (top). Des-inhibition of the *FLNA* poison exon leads to
19 precocious neuron differentiation and periventricular heterotopia in humans (bottom, pink
20 arrowheads). (B) *DCX* mutation in human is associated with subcortical band heterotopia (pink
21 arrowheads). In mice, constitutive *Dcx* loss of function does not cause major cortical defects,
22 while acute uni/bilateral loss of function induces large subcortical heterotopia. Heterotopia are
23 highlighted in pink. KO, knock-out. (C) The numbers of publications containing the indicated
24 combinations of keywords (cell types and neurodevelopmental diseases) were identified with
25 the EasyPubMed R package as described in Fig. 1. (D) Basic connectivity rules in the mouse
26 somatosensory cortex. This connectivity appears largely conserved when neurons are
27 mispositioned, like in the “scrambled” cortex of *Reeler* mice or in case of heterotopia. ASD,
28 autism spectrum disorder; FCD, focal cortical dysplasia; HET, heterotopia; L, layer; LIS,
29 lissencephaly; MIC, microcephaly; PMG, polymicrogyria; SZ., schizophrenia. In B, source
30 image for mouse is from ©biorender.com.

31

1 **Fig. 5: Multimodal assessment of neurodevelopmental disorders.** (A) Identification of
2 developmental gene dynamics in specific cell types through single-cell RNA sequencing (top).
3 Illustration from the online resource <http://genebrowser.unige.ch/humous/> comparing data
4 from mouse, human embryos, and human derived- brain organoids (bottom). (B) Average
5 expression of genes associated with microcephaly (MIC), lissencephaly (LIS) and
6 polymicrogyria (PMG) in mouse, human embryos, and human derived- brain organoids.
7 Arrowheads show human-specific patterns. (C) Schematic representation of the probabilistic
8 distribution of affected processes in sample disorders. Pink cubes define abnormal states. (D)
9 Hypothetical patient classification through integration of multiple datasets. Here patients were
10 clustered into 2 groups based on the integration of several features, from genes to clinical
11 expression. ATAC-seq (Assay for Transposase-Accessible Chromatin with high-throughput
12 sequencing) allows to identify chromatin accessible regions. Mass-spectrometry and co-
13 immunoprecipitation allow to identified proteins based on their molecular weight and
14 interaction(s) between proteins, respectively. *In vitro* and animal model data bring
15 complementary information about the cellular effect(s) of a gene mutation. aRG, apical radial
16 glia; bRG, basal radial glia; CSF, cerebrospinal fluid; fMRI, functional magnetic resonance
17 imaging; iN, immature neurons; IPC, intermediate progenitor cells; MCD, malformations of
18 cortical development; mN mature neurons. In B, data from mouse embryos are from (94); data
19 from human embryos and human-derived organoids are from (95, 96). The 2-dimensional
20 landscapes of gene expression across time and differentiation were performed as described in
21 (94). In B, source images for mouse and human are from ©biorender.com.

22






Fig. 1: Key steps of corticogenesis and their susceptibility to disease. (A) Summary of embryonic corticogenesis. One neuron can be followed from its birth at the ventricular zone, to its migration toward the cortical plate, and then its maturation (arrow). The role of cytoskeletal processes at each of these steps is highlighted on the right. **(B)** Literature-based summary of knowledge on the involvement of these developmental steps in specific neurodevelopmental disorders. The numbers of publications were found using the text mining EasyPubMed R package, which automatically identifies publications containing given the combinations of keywords indicated (e.g. “microcephaly” and “proliferation”). These numbers were then normalized for number of publications for each of the developmental steps and each of the disorders. The dendrogram represents unbiased clustering of the diseases based on this data. **(C)** Overlapping cellular processes and pleiotropic molecular mechanisms involving CDK5 across development and in the adult brain. aRG, apical radial glia; bRG, basal radial glia; CP, cortical plate; E, embryonic day; IN, interneuron; IPC, intermediate progenitor cell; N, excitatory neuron; pcw; post-conception week; SVZ, subventricular zone; VZ, ventricular zone; WM, white matter.




Fig. 2: Cellular mechanisms at play in human cortical malformations. Cells with abnormal developmental trajectories are in pink. Reduced progenitor proliferation (sometimes associated with abnormal neuron migration) leads to microcephaly. Abnormal neuron position can lead to agyria and lissencephaly (reduced migration and/or progenitor misposition), polymicrogyria (increased migration and/or excess of basal radial glia), cobblestone lissencephaly (excessive migration), periventricular heterotopia (neurons stay close to the ventricle) or subcortical band heterotopia (neurons stay below the cortex). In focal cortical dysplasia, clonal mutations affect subpopulations of mature cells, which become dysmorphic (balloon cells, dysmorphic neurons). Genes mentioned in this review are listed (bold: genes with published human-derived brain organoid models, *: gene studied in the ferret, green: genes coding for transcription factors). Pink arrowheads on MRIs highlight the malformations. MRI illustrations are from (15) and (2) for polymicrogyria and focal cortical dysplasia (Reprinted from *Semin Cell Dev Biol* vol 76, D. M. Romero, N. Bahi-Buisson & F. Francis, Genetics and mechanisms leading to human cortical malformations, pages 33-75, 2018, with permission from Elsevier; Definitions and classification of malformations of cortical development: practical guidelines; M. Severino et al., *Brain* 2020; awaa174, with permission from Oxford University Press).




Fig. 3: From gene mutations to diseases: levels of complexity. (A) Levels of organization during corticogenesis, from DNA (*i.e.* genes, represented by letters) to RNA and proteins (*i.e.* gene expression), to cells, circuit and anatomy, and phenotype. Each circle represents a given feature of that level (*e.g.* a gene, a protein, a circuit). Interactions within levels are linked through complex relationships (dashed lines) to states at other levels (black lines). (B-G) Examples of abnormal feature relationships across levels in disease (highlighted in pink). In B-D, linear (monogenic), convergent (polygenic) and divergent relationships between genetic defects and phenotype(s) are illustrated. (E) Hidden complexity: note that from a gene-phenotype point of view, this relationship appears linear. (F) Hidden hub: different genes may lead to distinct phenotypes *via* hidden feature hubs influencing several downstream pathways. (G) Incomplete penetrance: not all genetic features are expressed at other levels. In A, source images for human brain section and phenotype are from <https://msu.edu/~brains/brains/> and ©biorender.com, respectively.




Fig. 4: Spatiotemporal and cell type regulations of gene expression and malformations. (A) Alternative splicing of *FLNA* gene regulates its role in proliferation *versus* neuron differentiation of cortical progenitors (top). Des-inhibition of the *FLNA* poison exon leads to precocious neuron differentiation and periventricular heterotopia in humans (bottom, pink arrowheads). **(B)** *DCX* mutation in human is associated with subcortical band heterotopia (pink arrowheads). In mice, constitutive *Dcx* loss of function does not cause major cortical defects, while acute uni/bilateral loss of function induces large subcortical heterotopia. Heterotopia are highlighted in pink. KO, knock-out. **(C)** The numbers of publications containing the indicated combinations of keywords (cell types and neurodevelopmental diseases) were identified with the EasyPubMed R package as described in Fig. 1. **(D)** Basic connectivity rules in the mouse somatosensory cortex. This connectivity appears largely conserved when neurons are mispositioned, like in the “scrambled” cortex of *Reeler* mice or in case of heterotopia. ASD, autism spectrum disorder; FCD, focal cortical dysplasia; HET, heterotopia; L, layer; LIS, lissencephaly; MIC, microcephaly; PMG, polymicrogyria; SZ., schizophrenia. In B, source image for mouse is from ©biorender.com.




Fig. 5: Multimodal assessment of neurodevelopmental disorders. (A) Identification of developmental gene dynamics in specific cell types through single-cell RNA sequencing (top). Illustration from the online resource <http://genebrowser.unige.ch/humous/> comparing data from mouse, human embryos, and human derived- brain organoids (bottom). (B) Average expression of genes associated with microcephaly (MIC), lissencephaly (LIS) and polymicrogyria (PMG) in mouse, human embryos, and human derived- brain organoids. Arrowheads show human-specific patterns. (C) Schematic representation of the probabilistic distribution of affected processes in sample disorders. Pink cubes define abnormal states. (D) Hypothetical patient classification through integration of multiple datasets. Here patients were clustered into 2 groups based on the integration of several features, from genes to clinical expression. ATAC-seq (Assay for Transposase-Accessible Chromatin with high-throughput sequencing) allows to identify chromatin accessible regions. Mass-spectrometry and co-immunoprecipitation allow to identified proteins based on their molecular weight and interaction(s) between proteins, respectively. *In vitro* and animal model data bring complementary information about the cellular effect(s) of a gene mutation. aRG, apical radial glia; bRG, basal radial glia; CSF, cerebrospinal fluid; fMRI, functional magnetic resonance imaging; iN, immature neurons; IPC, intermediate progenitor cells; MCD, malformations of cortical development; mN mature neurons. In B, data from mouse embryos are from (94); data from human embryos and human-derived organoids are from (95, 96). The 2-dimensional landscapes of gene expression across time and differentiation were performed as described in (94). In B, source images for mouse and human are from ©biorender.com.