



HAL
open science

Gut microbiota and vitamin status in persons with obesity: a key interplay

Lise Volland, Tiphaine Le Roy, Jean Debédats, Karine Clément

► To cite this version:

Lise Volland, Tiphaine Le Roy, Jean Debédats, Karine Clément. Gut microbiota and vitamin status in persons with obesity: a key interplay. *Obesity Reviews*, 2021, 10.1111/obr.13377 . hal-03440317

HAL Id: hal-03440317

<https://hal.sorbonne-universite.fr/hal-03440317v1>

Submitted on 22 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Gut microbiota and vitamin status in persons with obesity: a key interplay.**

2

3 **Lise Volland¹, Tiphaine Le Roy¹, Jean Debédats¹, Karine Clément^{1,2}**

4

5 **¹Sorbonne Université, Inserm, Nutrition and Obesities ; Systemic approaches (NutriOmics),**
6 **Paris, France**

7

8 **²Assistance Publique-Hôpitaux de Paris, Nutrition department, Pitié-salpêtrière hospital,**
9 **Paris France**

10

11 **Corresponding author**

12 Prof Karine Clément MD, PhD
13 Sorbonne Université/ INSERM
14 NutriOmique Research group
15 Nutrition Department, Pitié-Salpêtrière Hospital
16 91 boulevard de l'Hôpital
17 75013 Paris, France
18 Tel +33 1417 7031
19 Karine.clement@inserm.fr

20

21 **Acknowledgement**

22 The authors thank Leducq Foundation, the European commission (Join Program Initiative
23 (JPI)), the European Union's 7th Framework Program for research, technological development
24 and demonstration (HEALTH-F4-2012-305312 - METACARDIS), and the Fondation pour la
25 Recherche Médicale (FDT202106012793) for supporting the research topic addressed in this
26 review. We thank Timothy Swartz for his thorough re-reading of this article.

27 **Conflict of interest**

28 The authors have no conflict of interest.

29 **Abstract**

30

31 There are numerous factors involved in obesity progression and maintenance including
32 systemic low-grade inflammation, adipose tissue dysfunction or gut microbiota dysbiosis.
33 Recently, a growing interest has arisen for vitamins' role in obesity and related disorders, both
34 at the host and gut bacterial level. Indeed, vitamins are provided mostly by food but some,
35 from the B and K groups in particular, can be synthesized by the gut bacterial ecosystem and
36 absorbed in the colon. Knowing that vitamin deficiency can alter many important cellular
37 functions and lead to serious health issues, it is important to carefully monitor the vitamin
38 status of patients with obesity and potentially already existing comorbidities as well as
39 dysbiotic gut microbiota and thus potentially altered bacterial metabolism of vitamins. In this
40 review, we examined both murine and human studies, to assess the prevalence of sub-optimal
41 levels of several vitamins in obesity and metabolic alterations. This review also examines the
42 relationship between vitamins and the gut microbiota in terms of vitamin production and the
43 modulation of the gut bacterial ecosystem in conditions of vitamin shortage or
44 supplementation. Furthermore, some strategies to improve vitamin status of patients with
45 severe obesity are proposed within this review.

46

47 Key words: biotin, gut microbiota, micronutrients, obesity, vitamins

48

49

50

51

52

53 **Introduction**

54

55 Obesity represents a global health issue with an ever-increasing prevalence over the
56 last 50 years^{1,2}. It is expected that over 40% of the world's population will suffer from obesity
57 by 2030³. Obesity is associated with several complications such as type 2 diabetes,
58 cardiovascular diseases and cancers⁴. Triggered by multiple environmental factors, the
59 development and progression of obesity are characterized by organ structural and biological
60 alterations and systemic low-grade inflammation. Recently, the gut microbiota (GM) has
61 emerged as a contributive "organ" in the development and maintenance of obesity.

62 The GM is a complex and abundant ecosystem composed of trillions of microorganisms
63 including bacteria, viruses, archaea, fungi, and phages. The microorganisms present within the
64 gut harbor a myriad of functions, and in particular, the GM is implicated in the fermentation
65 of non-digestible food substrates. This microbial metabolism leads to the production of an
66 important variety of metabolites, which in return are known to have a broad range of
67 bioactivities, including immune modulation and intestinal barrier homeostasis. During obesity,
68 an imbalance in the GM composition, functionality, as well as a decreased bacterial richness
69 and diversity, collectively referred to as GM dysbiosis, have been reported by many studies⁵.

70 Among the numerous GM-derived metabolites (as short chain fatty acids, secondary
71 bile acid molecules, tryptophan derivatives to cite only a few), vitamins have been relatively
72 neglected in this context. Yet, gut bacteria are known to produce vitamins from the B and K
73 groups⁶. Vitamins are essential micronutrients, obtained from food and, to some extent, from
74 gut bacterial synthesis. Vitamins serve as coenzymes or cofactors for numerous processes
75 involved in host energy metabolism⁷ (Table 1). For example, B vitamins play a key role in
76 maintaining mitochondrial energy metabolism, which is altered by inadequate levels of any of

77 those vitamins⁸ (Table 1). Vitamin deficiency, due to an imbalanced diet, increased host
78 physiological requirements or altered drug-micronutrient interactions, affects host health in
79 varying ways and can lead to severe clinical consequences, ranging from fatigue to
80 dermatological, digestive, cardiovascular or neurological and depressive disorders⁹ (Table 1).
81 Dietary vitamin absorption occurs mainly in the small intestine for the dietary vitamins but it
82 is now acknowledged that the distal gut may also contribute to vitamin absorption and
83 especially bacterially-produced vitamins¹⁰. Thus, the alteration of the intestinal environment
84 and absorptive capacities potentially alters vitamins host vitamin status.

85 Despite increasing evidence of deep alterations of gut physiology as well as microbiota
86 composition and functionality observed during obesity and metabolic disorders, the
87 consequences on vitamins biosynthesis and availability have scarcely been studied. Herein,
88 we review vitamin status alterations observed in obesity and metabolic disorders and focus
89 on the production potential of vitamins by the GM as well as its modulation by both vitamin
90 supplementation and deficiency, mostly focusing on B and D vitamins. We also discuss some
91 future strategies for vitamin status management in severe obesity.

92

93 **Evidence of vitamin deficiency in obesity and related metabolic disorders in humans**

94

95 The optimal vitamin reference levels for the general population are reported by
96 national or international expert committees such as the European Food Safety Authority
97 (EFSA)¹¹ or the US Centers for Disease and Prevention through the National Health and
98 Nutrition Examination Survey (NHANES). Lately, an increasing interest for vitamin status in
99 subjects with obesity has arisen as many studies have revealed that obesity and related

100 metabolic complications are associated with deficiencies or at least sub-optimal blood
101 concentrations of many vitamins, which has been well-reviewed recently^{12,13}.

102 Briefly, a series of studies exploring different populations with metabolic alterations
103 revealed suboptimal blood level of B vitamins, including vitamin B1 (thiamine), B9 (folate), B6
104 (pyridoxine), B12 (cobalamin). A decrease of vitamin B9 has been observed in pregnant
105 women with overweight in comparison to women with a BMI ranging between 18 and 25
106 kg/m² ¹⁴. Similarly, another study on patients with severe obesity reported an important
107 decrease of vitamin B6 serum concentrations as compared to patients without obesity¹⁵.
108 Moreover, up to 75% patients with type 2 diabetes presented low blood concentration of
109 vitamin B1¹⁶. Vitamin B12 levels have also been linked with obesity with a significant negative
110 correlation reported between body mass index (BMI) and vitamin B12 blood levels in a cohort
111 of almost 1000 patients with obesity¹⁷. Moreover, the authors of this study also found a trend
112 toward decreased vitamin B12 blood concentrations in subjects with insulin resistance or
113 metabolic syndrome in comparison to participants without metabolic alterations¹⁷. In another
114 cohort of participants with overweight or obesity, 13% of participants were found to be
115 deficient in vitamin B12 ¹⁸ while in the general US population, the prevalence of vitamin B12
116 was found to be around 5% and increases with age¹⁹. Interestingly, the serum levels of vitamin
117 B12 are increased after a weight loss induced by a Low Energy Diet (LED)¹⁸. Another B vitamin,
118 vitamin B7 (biotin) has been shown to be decreased in subjects with moderate obesity²⁰ and
119 type 2 diabetes²¹. Yet, this vitamin has been neglected in the context of obesity and metabolic
120 disorders; as well as other B vitamins such as vitamins B2 (riboflavin), B3 (niacin).

121 Vitamin deficiency in metabolic disorders extends to vitamin D as well. In Geiker et al.,
122 insufficient levels of vitamin D were observed in 72% of the participants before the LED-
123 induced weight loss, and again, the serum concentration of vitamin D was increased after the

124 nutritional intervention¹⁸. Unlike B vitamins, the links between vitamin D status and obesity
125 and its complications have been investigated in a larger number of studies and were reported
126 as early as 1985²². It has been commonly accepted that vitamin D deficiency can be estimated
127 via the serum concentration of the vitamin D metabolite, 25-hydroxycholecalciferol
128 [25(OH)D], the active form of vitamin D, which integrates both the dietary intake and the
129 endogenous production. For example, analyses in a large population of older individuals
130 revealed a significant inverse association between BMI and more specifically adiposity
131 (especially in the 3rd and 4th quartile when fat percentage was divided in quartiles) and serum
132 25(OH)D concentrations²³. Moreover, several groups also showed inverse correlations
133 between 25(OH)D serum concentrations and markers of metabolic alterations such as
134 triglycerides²⁴ or fasting glucose and the HOMA-IR index, a surrogate marker of insulin
135 resistance²⁵. An observational transversal study including 73 patients with morbid obesity
136 confirmed the higher prevalence of vitamin D deficiency in comparison to the population
137 without obesity²⁴. Here, they found that 50.7% of the participants had vitamin D deficiency.
138 They also reported an increased prevalence of vitamin D deficiency associated with the
139 occurrence of metabolic syndrome in this population. This correlation was confirmed in a large
140 study of over 2 500 Chinese men and women, in which, lower 25(OH)D serum concentration
141 was associated with increased insulin resistance among the participants with overweight or
142 obesity ²⁶. Finally, a recent meta-analysis of 43 studies confirmed this inverse association
143 between serum concentration of vitamin D and risk of metabolic syndrome in the general
144 adult population²⁷. In animals, vitamin D deficiency associates with worsened obesity
145 condition, glucose intolerance and insulin resistance induced by a high-fat diet whereas no
146 such effect for animals experimenting vitamin D deficiency under standard diet were
147 observed²⁸. Thus, it seems that inadequate levels of vitamin D in the context of obesity could

148 worsen the individual metabolic health. Low levels of vitamin D are not only linked with
149 different health conditions such as loss of bone density but also diabetes, high blood pressure,
150 or multiple sclerosis. It also appears that vitamin D deficiency can impede bariatric surgery-
151 induced metabolic improvements. Interestingly, in a mouse model of vertical sleeve
152 gastrectomy, deficient levels of vitamin D were linked with decreased weight loss and glucose
153 homeostasis improvements, as well as intestinal inflammation and permeability²⁹. In addition
154 to those metabolic observations regarding vitamin D, many studies also reported strong links
155 of this vitamin with adipose tissue in terms of vitamin D storage but also modulation of
156 adipocyte differentiation or even energy metabolism and inflammation by this vitamin. These
157 aspects have been largely and well-reviewed recently³⁰.

158

159 **Alteration of vitamin status in the context of bariatric surgery**

160 Vitamin deficiency in obesity has also been investigated in subjects with severe obesity
161 undergoing bariatric surgery. For example, studies showed that the prevalence of vitamin B1
162 deficiency in patients with severe obesity before bariatric surgery is between 15.5%³¹ and
163 29%³². In the general population, vitamin B1 deficiency is associated with inadequate intake
164 and is more prevalent in underdeveloped countries whereas in industrialized countries,
165 vitamin B1 deficiency is rare and mostly related to alcoholism³³. Vitamin B9 has also been
166 found to be decreased in patients with severe obesity before the surgery at a rate of 63.2%³⁴,
167 a much higher percentage in comparison to the general population prevalence of 2% reported
168 by a recent study in the US³⁵. Importantly, these studies focused on bariatric surgery
169 candidates who follow weight-loss efforts for several months before the procedures, as
170 recommended by international guidelines³⁶. As such, these patients may have reduced
171 vitamin intake due to changes in their diet. Flancbaum and colleagues also revealed a high

172 proportion (68%) of patients presenting with a vitamin D deficiency before the surgery, which
173 was confirmed in two other cohorts with deficiency rates of 74.35%³⁷ and 76%³⁸. In
174 comparison, a study of vitamin D blood concentrations in elderly people across 11 European
175 countries reported a prevalence of vitamin D deficiency in 36% of men and 42 % of women in
176 the general population³⁹. This percentage was even lower in the US population with a
177 prevalence of vitamin D deficiency of 4.0% and 17.4% for inadequacy⁴⁰. To note, vitamin D
178 blood concentration is dependent on the season as well as geographical location, which could
179 explain such discrepancies between prevalence in different populations and studies. The two
180 latter studies investigating vitamin D blood concentrations also reported vitamin B12
181 deficiency in those patients, reaching 12.3%³⁷ and 16%³⁸ respectively.

182 Moreover, most vitamins are absorbed in the small intestine, a segment bypassed after
183 malabsorptive procedures (e.g., Roux-en-Y Gastric Bypass or biliopancreatic derivation). This
184 purposefully-induced malabsorption further deteriorates vitamin status after the surgical
185 procedure, which may not normalize despite systematic vitamin supplementation. During
186 surgical follow-up, monitoring patients' vitamin status should be reinforced, which is
187 highlighted by studies of micronutrient deficiencies after Roux-en-Y gastric bypass as well as
188 sleeve gastrectomy^{41,42}. In fact, despite medical recommendation of micronutrients and
189 especially multivitamins supplementation after the surgery, evidence of decreased
190 compliance has been reported after 10 years of follow-up increasing the risk of vitamin
191 deficiency⁴³. It is also possible that the optimization of the pre-operative vitamin status could
192 also contribute to better outcomes of the surgery. It has been suggested that vitamin D could,
193 for example, contribute to improved weight loss following bariatric surgery⁴⁴. A recent review
194 addressed this concern in terms of vitamin status and the physiological, metabolic and also
195 bacterial changes associated with bariatric surgery⁴⁵. More attention should be given to

196 vitamin status of patients before and after the bariatric surgery as suggested by recent
197 guidelines published by the British Obesity and Metabolic Surgery Society⁴⁶. The
198 aforementioned studies suggest that acting on the pre-intervention vitamin deficiency might
199 help potentialize metabolic improvement, although more clinical interventions ought to be
200 performed to confirm this hypothesis.

201

202 **Gut microbiota; a source of vitamin production**

203 The composition of the GM has now been meticulously studied over the last decade
204 thanks to major advances in sequencing technologies. Nevertheless, a recent shift towards
205 understanding the functionality of the ecosystem allows for more complete comprehension
206 of how GM modulations are involved in health and diseases. Gut bacteria produce a very large
207 number of metabolites having an impact on the host metabolism⁴⁷. The gut bacteria can
208 produce vitamin K and B vitamins⁴⁸; however the interest for the study of the relationship
209 between GM and vitamin metabolism is only recent and increasing.

210

211 The number of available bacterial genomes increased significantly with the advances
212 in sequencing technologies and metagenomic processing methods, which have allowed for
213 better characterization of GM species and their metabolic capabilities. In 2015, Magnúsdóttir
214 et al., used the PubSEED platform, which provides genome annotations across thousands of
215 genomes, to analyze the ability of human GM organisms to synthesize B vitamins⁴⁹. Thus, they
216 were able to determine that the biosynthesis of B vitamins was common among the species
217 found in the human GM. They also predicted B vitamin synthesis abilities that matched
218 published experimental data. It appears that the synthesis of vitamin B2 and vitamin B3 were
219 most commonly found among all the genomes assessed. Altogether, this study reinforces the

220 idea that the GM, as an ecosystem, is able to produce vitamins and possibly participates to
221 vitamin homeostasis of the host. Yet, this paper revealed that the production of vitamins by
222 the GM is not sufficient to cover host daily requirements. Moreover, the vitamins synthesized
223 in the gut mostly benefit the neighboring non-producing bacteria through cross-feeding
224 between organisms. To this end, bacterial genome analyses have illustrated that bacterial
225 species can be vitamin prototrophs (capable of *de novo* synthesis) or auxotrophs (dependent
226 on the uptake and salvage of vitamins) highlighting the importance of cross-feeding between
227 GM species⁵⁰.

228 Further evidence of GM mutualism for vitamins comes from co-culture *in vitro*
229 experiments showing that *Anaerobutyricum halii* is able to synthesize and provide
230 *Akkermansia muciniphila* with B12 vitamin, which is used for propionate production⁵¹. Such
231 cross-feeding mechanisms have also been observed *in vivo*, where non-producing bacteria
232 survive during limitation of vitamin intake suggesting the existence of vitamin sharing
233 between producing and non-producing bacteria⁵². Bacterial mutualism thus contributes to
234 maintaining a relative GM stability thanks to the sharing of micronutrient resources in case of
235 shortage or level modulation due to vitamin intake variability⁵². Noteworthy *Anaerobutyricum*
236 *halii* and *Akkermansia muciniphila* are both commensal bacteria associated with improved
237 metabolism^{53,54}.

238 Together with experimental work, the access to genomic annotation allowed for the
239 identification of specific vitamin producing bacterial strains. It is now known for example, that
240 species from the genera *Lactobacillus* and *Bifidobacterium* can produce vitamin B9. Indeed,
241 both *in vitro* and *in vivo* experiments demonstrated that Bifidobacterial strains are able to
242 increase the vitamin B9 concentration either in the medium or of the host^{55,56}. This ability of
243 vitamin B9 production by *Lactobacillus* and *Bifidobacterium* was assessed via the analysis of

244 genome sequences. The presence of genes and enzymes involved in different steps of the
245 vitamin B9 production pathway allowed to infer theoretical capacity of the studied strains to
246 produce this vitamin at least from a precursor molecule⁵⁷. Nevertheless, it is important to note
247 that the presence of genes involved in the production of a vitamin does not account for its
248 effective production by the bacteria. Thus, experimental validation (*in vitro* and *in vivo*)
249 remains crucial to identify potential vitamin-producing strains that could then be developed
250 as probiotics.

251 To explore the contribution of GM's vitamin synthesis, in various diseases including
252 metabolic diseases, *in vivo* experiments are needed. Germ-Free (GF) rodents are an initial
253 model used to determine the general contribution of the GM to metabolic pathways or host
254 physiology. In the absence of GM in GF mice, a systemic deficiency in different vitamins was
255 reported^{58,59}, suggesting again that those vitamins can be produced by the GM and that
256 microbial vitamin production contributes, at least partially, to vitamin homeostasis of the
257 host. Among those vitamins, earlier studies showed that GF rats developed spontaneously a
258 vitamin K deficiency leading to increased mortality of the animals⁵⁸. The side effects of vitamin
259 K deficiency were corrected by vitamin K1 supplementation and most interestingly with the
260 gut colonization of single strains of bacteria (*Escherichia coli* or *Sarcina-like micrococcus*).
261 Similar results have been found for B vitamins as rats fed a diet deprived of B vitamins stopped
262 growing after 1 to 3 weeks and died prematurely⁵⁹. Yet, in this early study, one rat behaved
263 differently and recovered from the deficiency. When its feces were added to the diet of
264 another group of rats fed the same diet, the symptoms of vitamin B deficiency disappeared in
265 those animals, suggesting that B vitamin-producing bacteria could be present in the feces
266 these rats were given.

267 Other evidence of B vitamin production by intestinal microorganisms have been
268 reported. The total output of vitamin B7 measured in feces and urine has been shown to be
269 six times greater than the intake in humans, suggesting a production by the GM^{60,61} while this
270 needs further exploration. Interestingly, vitamin B7 deficiency induced alopecia only in GF
271 animals and not in conventionally raised animals suggesting that the GM contributes to this
272 vitamin metabolism⁶². A similar phenotype was observed in vitamin B7-deficient mice treated
273 with an antibiotic, vancomycin, again suggesting a contribution of GM to vitamin B7
274 production. Moreover, those animals presented further reduction in bacterial diversity as well
275 as major increase of *Lactobacillus* genus and in particular *Lactobacillus murinus* in comparison
276 to antibiotic treated animals receiving sufficient levels of vitamin B7. Regarding other B
277 vitamins, GM depletion via antibiotic treatment resulted in a decrease of vitamin B6 levels in
278 colonic content in mice⁶³.

279 These observations all identify the GM as a major contributor to the host's vitamin
280 status. Nevertheless, the contribution of the GM's vitamin production to host vitamin
281 circulating levels remains to be experimentally established, in healthy and disease conditions.
282 Indeed, in the context of obesity, alteration in GM richness, diversity and functional anomalies
283 have repeatedly been reported^{47,64-66}. Whether these alterations are linked to deficient
284 bacterial vitamin metabolism and production need to be investigated in depth.

285

286 **Consequence of vitamin deficiencies on GM composition**

287 As raised above, the production of vitamin K by the GM has been known for several
288 decades; but the effect of its deficiency on the GM composition has been studied very
289 recently. Thus, mice fed a vitamin K deficient diet displayed reduced abundance of

290 *Lactobacillus* and enrichment in *Ruminococcus*, *Anaerostipes* genera as well as
291 *Muribaculaceae* family in comparison to animals supplemented with vitamin K⁶⁷.

292

293 A murine study investigating the effect of different micronutrient-deficient (vitamin A,
294 vitamin B9, iron, zinc, or multiple deficiencies) diet on the GM revealed that the vitamin A
295 deficiency had the higher impact⁶⁸, although vitamin A is not produced by the GM. The analysis
296 of the GM of mice fed a vitamin A deficient diet revealed an increase of *Bacteroides vulgatus*
297 and an increase of *Bacteroides dorei*. Once the deficiency was corrected via a micronutrient
298 sufficient supplementation of the diet, the initial levels of those bacteria were also restored.
299 In a human study investigating the GM children with vitamin A deficiency, changes in the
300 bacterial community composition were also reported⁶⁹. Increased abundances of *Blautia* and
301 *Faecalibacterium* genera and decreased abundances of *Bifidobacterium*, *Bacteroides* and
302 *Shigella* were observed in these children. The modulations of *Bifidobacterium* and *Bacteroides*
303 levels were even transmitted via fecal microbiota transfer of the children's feces into germ-
304 free animals. At the phylum level, a decrease of *Bacteroidetes* was observed in mice fed a
305 vitamin A-deficient diet⁷⁰. This decrease is associated with an increase of the
306 *Firmicutes/Bacteroidetes* ratio in those animals, which has been reported associated with
307 obesity and metabolic alterations⁷¹, despite discrepancies in recent studies^{72,73}. To be noted,
308 in this study, vitamin A-deficient animals presented characteristics of pre-diabetes with
309 hyperglycemia and delayed response to insulin injection, emphasizing the potential links with
310 metabolic alterations.

311

312 In the last decade, there has been an increasing number of studies investigating the
313 links between vitamin D and the GM. One of them investigated the factors explaining the most

314 the variance in alpha-diversity of the human GM, and found that the serum concentration of
315 1,25(OH)₂D was the most contributing variable, explaining 5% of the variance of the alpha-
316 diversity⁷⁴. In terms of specific compositional changes associated with vitamin D deficiency, it
317 seems that vitamin D deficiency (in models of knock-out mice for the vitamin D receptor (VDR),
318 not able to produce 1,25(OH)₂D) appeared to be associated with a decrease of the abundance
319 of the phylum *Firmicutes*⁷⁵ and the associated families *Ruminococcaceae*, *Lachnospiraceae*,
320 *Lactobacillaceae* and *Streptococcaceae*. Interestingly, in another study on VDR KO mice, a
321 significant decrease of the abundance of the genus *Lactobacillus* was observed in comparison
322 to wild-type animals⁷⁶. In a human trial on older men, a positive association was also found
323 between serum 1,25(OH)₂D levels and the abundance of the genus *Ruminococcus*⁷⁴. On the
324 contrary, vitamin D deficiency seems to be associated with an increase in *Bacteroidetes* and
325 *Proteobacteria* phyla, and more precisely *Bacteroidaceae* and *Desulfovibrionaceae* families⁷⁵.
326 This increase was again confirmed in the study from Jin and colleagues, in which an increase
327 in the genus *Bacteroides* was noticed in VDR KO animals⁷⁶. In the murine study by Zhang *et al.*,
328 investigating the effect of vitamin D deficiency on the metabolic improvements after vertical
329 sleeve gastrectomy, they also analyzed the effects on the GM composition. The authors
330 revealed that, despite known beneficial effects induced by the surgery itself, deficient levels
331 of vitamin D were associated with limited improvements of the gut dysbiosis induced by the
332 high-fat diet feeding. Indeed, the effects of the surgery, a decrease of *Firmicutes* and an
333 increase of *Bacteroidetes* phyla abundance, were weaker in the group of vitamin-D-deficient
334 animals post-surgery in comparison to the group with normal levels of vitamin D²⁹.

335 The effect of B vitamins deficiency on the GM has also been recently studied in the
336 context of metabolic diseases. In a recent study, the effects of low B6 vitamin levels on host
337 phenotype and GM of weaning rats was explored⁷⁷. The animals receiving a standard chow

338 diet with low levels of vitamin B6 displayed a decreased body weight and circulating
339 triglycerides levels compared to control animals as well as changes in the GM composition.
340 Indeed, an increase in *Lachnospiraceae* and a decrease of *Bacteroides* were observed in both
341 male and female rats.

342 In patients with obesity, an association was found between low niacin (vitamin B3)
343 intake and reduced alpha-diversity as well as levels of Bacteroidetes. These findings were
344 particularly significant in subjects with obesity and insulin-resistance but not with overt type
345 2 diabetes⁷⁸. The authors suggested that in patients with type 2 diabetes, medications such as
346 metformin known to modulate the GM^{79,80} should be taken into account to explain those
347 differences.

348 These collective findings illustrate the need to investigate the interplay between GM,
349 vitamin and host metabolism. Yet, few studies investigated the effects of vitamin deprivation
350 on the GM composition in the context of obesity and metabolic disorders. Considering the
351 links between obesity, GM dysbiosis and alteration of vitamin status as presented above, it is
352 important to study these factors to decipher the mechanisms involved.

353

354 **Effect of vitamin supplementation on GM**

355

356 Recent interest regarding the effect of vitamin supplementation on the composition of
357 the GM has led to an increase of publications on this topic. A recent review extensively
358 summarized the effects of vitamins and other dietary micronutrients such as polyphenols and
359 minerals on the composition of the GM⁸¹. Although no clear patterns of GM modulation can
360 be identified from this review due to a relatively low number of studies for some vitamins and

361 a great variability in experimental designs, it seems that vitamin supplementation can induce
362 significant changes in GM composition.

363 In the context of obesity, the establishment of multiple large cohorts in the last decade
364 has allowed the collection of important data sets of microbiome analysis, clinical and lifestyle
365 information. Data from the American Gut Project were used to assess the impact of vitamin B
366 and/or D supplementation on the GM in groups of people with normal weight (NW) or
367 overweight (OW)⁸². Despite no significant impact on alpha-diversity indexes in all groups, the
368 vitamin supplementation alone had almost no effect in NW group but had beneficial impact
369 in OW. Noteworthy, vitamin B supplementation induced a significant decrease in *Tenericutes*
370 at the phylum level and several groups at the family level (*Mogibacteriaceae*,
371 *Porphyromonadaceae*, *Coriobacteriaceae* and *Comamonadaceae*). All these bacterial families
372 have been associated with infections, inflammation and even obesity in the literature,
373 suggesting that vitamin B supplementation in the context of obesity could help decrease the
374 abundance of pathogens in the GM^{83–86}. Regarding vitamin D supplementation, effects on the
375 GM composition were observed both in the NW and OW groups. In NW group, it induced a
376 decrease of several families *Erysipelotrichaceae*, *Turicibacteraceae* and *Bifidobacteriaceae*.
377 The same effect on this last beneficial family was observed among OW participants, as well as
378 a decrease of pathogens (*Actinomycetaceae*) for this group. The effects of a combined vitamin
379 B and D supplementation was also investigated and showed a decrease of *Lachnospiraceae*
380 and *Enterobacteriaceae*, which have been shown to be associated with obesity and production
381 of endotoxins respectively^{87,88}.

382 Vitamin supplements, and in particular vitamin D, also had an important impact on the
383 functionality of the GM⁸². In NW groups, there was a decrease in genes implicated in amino-
384 acids biosynthesis and an increase of vitamin biosynthesis (related to vitamin B1 and B5) in

385 the GM of people supplemented with vitamin D. Those pathways were not impacted by
386 vitamin D in the OW group. Here again, the association of the two vitamins had specific effect
387 regarding the functional machinery of bacteria. The authors observed a decrease in the
388 potential of amino acid consumption by GM and of the degradation of polysaccharides in the
389 OW group. This paper supports the beneficial effects of vitamin D (alone or combined with
390 vitamin B) on microbial communities in a context of overweight or obesity, although effects
391 vary according to the severity of obesity.

392 Another recent trial of vitamin D supplementation over 16 weeks in overweight and
393 patients with obesity showed modulation of the GM composition⁸⁹. Despite no effect on
394 alpha-diversity, vitamin D supplementation was associated with an increase of *Lachnospira*
395 genus and a decrease of the abundance of *Blautia* genus in comparison to the group receiving
396 placebo. Moreover, the authors observed a significant correlation between serum levels of
397 25(OH)D and the abundance of *Coprococcus* genus. Yet, these studies remain quite small in
398 size as only 17 people received vitamin D supplementation and 15 received the placebo in the
399 Naderpoor study⁸⁹. In the study from Jiang and colleagues, the groups of people with
400 overweight supplemented with only vitamin D, or only vitamin B, or both vitamins comprised
401 respectively 88, 16 and 214 individuals, and supplementation was based on dietary
402 supplement intakes self-reported by patients⁸².

403 In a murine study of weaning male C57BL/6J, the effect of vitamin A supplementation
404 in condition of standard chow diet (CD) or high fat high sugar diet (HFSD) was studied⁹⁰.
405 Vitamin A was administered in the form of retinol, directly incorporated to the diet. After 9
406 weeks of follow-up, the supplementation with vitamin A (+VitA) limited HFSD-induced fat
407 mass gain but was not able to reverse metabolic alterations induced by the diet. At the GM
408 level, this study revealed that retinol supplementation prevented most of the compositional

409 changes induced by HFSD. First, vitamin A supplementation protected against the decrease of
410 alpha-diversity induced by HFSD, as HFSD+VitA animals had alpha-diversity values to the level
411 of the CD group. In CD groups, a decrease of alpha-diversity was observed in the vitamin A
412 supplemented group, to the surprise of the authors. Second, HFSD-feeding led to a decrease
413 of *Porphyromonadaceae* and *Mycoplasmataceae* and an increase of *Family XIII* in comparison
414 to CD group. All these changes were prevented by administration of vitamin A, which in
415 addition increased the abundance of *Lachnospiraceae*.

416 These studies reinforce the links between vitamins and GM as oral vitamin
417 supplementation is reported to induce modifications of the GM composition.

418

419 **Future strategies of vitamin status management in obesity**

420

421 Knowing the alteration of vitamin status in some patients with obesity, the obesity-
422 associated dysbiosis and relationships observed between GM and vitamin metabolism,
423 several strategies can be considered to optimize vitamin levels subjects with obesity.

424 Whereas obesity is often considered as a disease characterized by over-nutrition, it is
425 also well recognized that there are issues of malnutrition, explaining at least partially
426 inadequate vitamin intakes observed during obesity. Thus, a well-balanced and micronutrient-
427 sufficient diet could help prevent deficiencies in some cases. In case of deficiency or sub-
428 optimal levels, the oral vitamin supplementation also seems rational to rescue the host's
429 vitamin status. However, most vitamins are absorbed in the small intestine and only small
430 amounts of dietary intake reach the colonic bacterial ecosystem. Yet, as summarized in this
431 review, many studies showed that vitamins can modulate the distal GM composition and have
432 even been included in the definition of prebiotics by the International Scientific Association

433 for Probiotics and Prebiotics (ISAPP) in 2016. Thus, delivery tools allowing for oral vitamin
434 supplementation specifically targeting the colon might be useful. A recent control trial
435 investigated such systems to administer several vitamins (vitamin C, B2, E, D and A) in 96
436 healthy volunteers and revealed modifications of the microbial composition and metabolic
437 activity such as an increased of bacterial diversity and fecal short-chain fatty acids⁹¹. At the
438 species level, the authors observed an increase in *Bifidobacterium longum* and
439 *Anaerobutyricum hallii*, both studied for their beneficial effects in several health conditions⁹¹.
440 Similar studies in patients with severe obesity would be helpful to further investigate the
441 effects of vitamin supplementation in this context, also taking into account the potential risk
442 of over-dosing vitamin supplementation.

443 Probiotic supplementation of strains known to produce vitamins can also be a strategy
444 to rescue vitamin bacterial metabolism, altered in obesity. As reviewed by LeBlanc and
445 colleagues, it has been demonstrated that specific bacterial strains are able to produce
446 vitamins. For example, *Lactobacillus rhamnosus* GG can produce three B vitamins in culture
447 medium which are vitamin B1, B2 and B9⁹². *Bifidobacterium adolescentis* DSM 18350⁹³ has
448 been shown to increase vitamin B9 fecal levels in humans. For such a strategy, it is important
449 to identify strains capable of complete *de novo* synthesis. As proposed by Engevik and
450 colleagues, an alternative could be to rely on cross-feeding, mentioned above, and construct
451 probiotic combinations of strains depending on one another for the production of
452 intermediates involved in the vitamin synthesis⁹⁴. It would be thus of interest to investigate
453 how the host can benefit from these bacterial cross-talks. Yet, the long-term engraftment of
454 probiotics remains a challenge for long-term therapeutic perspectives.

455 Prebiotics can influence the growth and metabolism of specific gut bacteria, thus, their
456 use to promote bacterial vitamin production could also be envisaged. A recent *in silico* study

457 reported that inulin and short-chain fructo-oligosaccharides supplementation could increase
458 the production of an intermediate of the vitamin D biosynthesis pathway⁹⁵. This effect seems
459 to be mediated by the secretion of short-chain fatty acids induced by the prebiotic
460 supplementation. The authors thus suggest that prebiotic supplementation could help
461 decrease vitamin D deficiency linked to insufficiency of this intermediate in this study.

462 Appropriate *in vivo* and clinical studies should be designed to confirm the potential use
463 of these strategies in vitamin status management in severe obesity.

464

465 **Conclusion**

466 Increasing evidence of sub-optimal or even deficient blood concentrations of a number
467 of vitamins in obesity and metabolic alterations have been consistently reported. There is
468 therefore a clinical need to increase the awareness of medical care givers and patients
469 regarding vitamin status, especially for long-term follow-up and in the context of nutritional
470 or surgical interventions. The reported evidence of modulations of both bacterial vitamin
471 production and GM composition and functionality in condition of vitamin depletion and
472 supplementation reinforce the links between vitamins and GM. In this direction, the GM
473 dysbiosis observed in metabolic alterations and obesity could participate to the altered
474 vitamin host status observed in several cohorts. The restoration of a functional ecosystem via
475 a balanced diet, colon-targeted vitamin, probiotic or prebiotic supplementation of either
476 vitamin-producing single strains or combination of strains relying on cross-feeding (Figure 1)
477 can thus contribute to an improved vitamin metabolism and status. The potential effects on
478 metabolic health of such strategies should also be studied as well as the mechanisms involved
479 in these interactions between vitamins, gut microbiota and metabolic health.

480

481 **Acknowledgement**

482 The authors thank Leducq Foundation, the European commission (Join Program Initiative
483 (JPI)), the European Union's 7th Framework Program for research, technological development
484 and demonstration (HEALTH-F4-2012-305312 - METACARDIS), and the Fondation pour la
485 Recherche Médicale (FDT202106012793) for supporting the research topic addressed in this
486 review. We thank Timothy Swartz for his thorough re-reading of this article.

Table 1 – Functions, recommended levels and deficiency consequences of B and D vitamins - Adapted from *Vitamin and mineral requirements in human nutrition* ⁹⁶

Vitamin	Name	Function	Daily recommended amounts	Optimal circulating levels	Deficiency biochemical diagnosis	Deficiency consequences
Vitamin B1	Thiamin	Coenzyme functions in metabolism of carbohydrates and branched-chain amino acids	Men: 1.2 mg/Women: 1.1mg	75-195 nmol/L (Whitfield et al., 2018)	Thiamine Pyrophosphate effect (TPPE) > 25% (WHO,1999)	Beriberi, polyneuritis and neurological symptoms
Vitamin B2	Riboflavin	Coenzyme functions in numerous oxidation and reduction reactions	Men: 1.3 mg/Women: 1.0 mg	100-650 nmol/L	Erythrocyte glutathione reductase activation coefficient (EGRAC) >1.80 (Lanyau-Dominguez et al., 2020)	Growth, cheilosis, angular stomatitis, and dermatitis
Vitamin B3	Niacin	Cosubstrate/coenzyme for hydrogen transfer with numerous dehydrogenases	Men: 16 mgNEs/Women: 14 mgNEs (Niacin equivalents)	4100-69000 nmol/L	Urinary excretion of N1-methylnicotinamide (NMN) < 5.8 µmol (Redzic and Gupta, 2021)	Pellagra with diarrhoea, dermatitis, and dementia
Vitamin B5	Pantothenic acid	Constituent of coenzyme A and phosphopantetheine involved in fatty acid metabolism	Men: 5.0 mg/Women: 5.0 mg	1600-2700 nmol/L (NIH)	Serum : <1000 nmol/L (NIH)	Fatigue, sleep disturbances, impaired coordination and nausea
Vitamin B6	Pyridoxine	Coenzyme functions in metabolism of amino acids, glycogen, and sphingoid bases	Men: 1.3 mg/Women: 1.3 mg	20-200 nmol/L (NIH)	Serum: <20 nmol/L (NIH)	Nasolateral seborrhoea, glossitis and peripheral neuropathy
Vitamin B7	Biotin	Coenzyme functions in bicarbonate-dependent carboxylations	Men: 30 µg /Women: 30 µg	0.8-1.7 nmol/L	Serum: <0.8 nmol/L	Fatigue, depression, nausea, dermatitis, and muscular pains
Vitamin B9	Folate	Function in producing genetic material (RNA, DNA) and amino acids	Men: 400 µg/ Women: 400 µg	> 6.8 nmol/L	< 6.8 nmol/L	Anemia, digestive and neurological disorders
Vitamin B12	Cobalamin	Functions in the development, myelination and function of the central nervous system	Men: 2.4 µg/Women: 2.4 µg	0,185-0,600 pmol/L	Serum: < 0,18 nmol/L	Anemia, fatigue, weakness
Vitamin D	Calciferol	Maintenance of normal blood level of calcium and phosphate	Men: 10 µg/Women: 10 µg	100-200 nmol/L	Serum: <75 nmol/L	Fatigue, bone pain, muscle weakness, depression




Figure 1 – **Intended strategies for host vitamin status improvement in obesity and metabolic alterations.** Improvement of vitamin metabolism can be envisioned via nutritional intervention with micronutrient sufficient and balanced diet, but also oral vitamin supplementation, for example via colon targeted system to deliver the vitamin directly to the gut microbial ecosystem and finally via probiotic administration of either single

strains known to synthesize a specific vitamin or a bacterial community capable of cross-feeding to synthesize both vitamins and other metabolites such as short-chain fatty acids (SCFA) - *Created with BioRender.com*

References:

1. NCD Risk Factor Collaboration (NCD-RisC). Trends in adult body-mass index in 200 countries from 1975 to 2014: a pooled analysis of 1698 population-based measurement studies with 19.2 million participants. *Lancet Lond Engl.* 2016;387(10026):1377-1396. doi:10.1016/S0140-6736(16)30054-X
2. NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128.9 million children, adolescents, and adults. *Lancet Lond Engl.* 2017;390(10113):2627-2642. doi:10.1016/S0140-6736(17)32129-3
3. Kelly T, Yang W, Chen C-S, Reynolds K, He J. Global burden of obesity in 2005 and projections to 2030. *Int J Obes.* 2008;32(9):1431-1437. doi:10.1038/ijo.2008.102
4. Segula D. Complications of obesity in adults: A short review of the literature. *Malawi Med J.* 2014;26(1):20-24.
5. Crovesy L, Masterson D, Rosado EL. Profile of the gut microbiota of adults with obesity: a systematic review. *Eur J Clin Nutr.* 2020;74(9):1251-1262. doi:10.1038/s41430-020-0607-6
6. Valdes AM, Walter J, Segal E, Spector TD. Role of the gut microbiota in nutrition and health. *BMJ.* 2018;361:k2179. doi:10.1136/bmj.k2179
7. Huskisson E, Maggini S, Ruf M. The Role of Vitamins and Minerals in Energy Metabolism and Well-Being. *J Int Med Res.* 2007;35(3):277-289. doi:10.1177/147323000703500301
8. Depeint F, Bruce WR, Shangari N, Mehta R, O'Brien PJ. Mitochondrial function and toxicity: role of B vitamins on the one-carbon transfer pathways. *Chem Biol Interact.* 2006;163(1-2):113-132. doi:10.1016/j.cbi.2006.05.010
9. Porter K, Hoey L, Hughes CF, Ward M, McNulty H. Causes, Consequences and Public Health Implications of Low B-Vitamin Status in Ageing. *Nutrients.* 2016;8(11). doi:10.3390/nu8110725
10. Said HM. Intestinal absorption of water-soluble vitamins in health and disease. *Biochem J.* 2011;437(3):357-372. doi:10.1042/BJ20110326
11. European Food Safety Authority, ed. *Tolerable Upper Intake Levels for Vitamins and Minerals.* European Food Safety Authority; 2006.
12. Astrup A, Bügel S. Overfed but undernourished: recognizing nutritional inadequacies/deficiencies in patients with overweight or obesity. *Int J Obes.* 2019;43(2):219-232. doi:10.1038/s41366-018-0143-9
13. Thomas-Valdés S, Tostes M das GV, Anunciação PC, da Silva BP, Sant'Ana HMP. Association between vitamin deficiency and metabolic disorders related to obesity. *Crit Rev Food Sci Nutr.* 2017;57(15):3332-3343. doi:10.1080/10408398.2015.1117413
14. Santacruz A, Collado MC, García-Valdés L, et al. Gut microbiota composition is associated with body weight, weight gain and biochemical parameters in pregnant women. *Br J Nutr.* 2010;104(1):83-92. doi:10.1017/S0007114510000176

15. Aasheim ET, Hofsø D, Hjelmesæth J, Birkeland KI, Bøhmer T. Vitamin status in morbidly obese patients: a cross-sectional study. *Am J Clin Nutr.* 2008;87(2):362-369. doi:10.1093/ajcn/87.2.362
16. Thornalley PJ, Babaei-Jadidi R, Al Ali H, et al. High prevalence of low plasma thiamine concentration in diabetes linked to a marker of vascular disease. *Diabetologia.* 2007;50(10):2164-2170. doi:10.1007/s00125-007-0771-4
17. Baltaci D, Kutlucan A, Turker Y, et al. Association of vitamin B12 with obesity, overweight, insulin resistance and metabolic syndrome, and body fat composition; primary care-based study. *Med Glas Off Publ Med Assoc Zenica-Doboj Cant Bosnia Herzeg.* 2013;10(2):203-210.
18. Geiker NRW, Veller M, Kjoelbaek L, et al. Effect of low energy diet for eight weeks to adults with overweight or obesity on folate, retinol, vitamin B12, D and E status and the degree of inflammation: a post hoc analysis of a randomized intervention trial. *Nutr Metab.* 2018;15. doi:10.1186/s12986-018-0263-1
19. Allen LH. How common is vitamin B-12 deficiency? *Am J Clin Nutr.* 2009;89(2):693S-696S. doi:10.3945/ajcn.2008.26947A
20. Järvinen E, Ismail K, Muniandy M, et al. Biotin-dependent functions in adiposity: a study of monozygotic twin pairs. *Int J Obes.* 2016;40(5):788-795. doi:10.1038/ijo.2015.237
21. Maebashi M, Makino Y, Furukawa Y, Ohinata K, Kimura S, Sato T. Therapeutic Evaluation of the Effect of Biotin on Hyperglycemia in Patients with Non-Insulin Dependent Diabetes Mellitus. *J Clin Biochem Nutr.* 1993;14:211-218. doi:10.3164/jcbtn.14.211
22. Liel Y, Ulmer E, Shary J, Hollis BW, Bell NH. Low circulating vitamin D in obesity. *Calcif Tissue Int.* 1988;43(4):199-201. doi:10.1007/BF02555135
23. Oliai Araghi S, van Dijk SC, Ham AC, et al. BMI and body fat mass is inversely associated with vitamin D levels in older individuals. *J Nutr Health Aging.* 2015;19(10):980-985. doi:10.1007/s12603-015-0657-y
24. Botella-Carretero JJ, Alvarez-Blasco F, Villafruela JJ, Balsa JA, Vázquez C, Escobar-Morreale HF. Vitamin D deficiency is associated with the metabolic syndrome in morbid obesity. *Clin Nutr Edinb Scotl.* 2007;26(5):573-580. doi:10.1016/j.clnu.2007.05.009
25. Devaraj S, Jialal G, Cook T, Siegel D, Jialal I. Low Vitamin D Levels in Northern American Adults with the Metabolic Syndrome. *Horm Metab Res.* 2011;43(1):72-74. doi:10.1055/s-0030-1268485
26. Lu L, Yu Z, Pan A, et al. Plasma 25-Hydroxyvitamin D Concentration and Metabolic Syndrome Among Middle-Aged and Elderly Chinese Individuals. *Diabetes Care.* 2009;32(7):1278-1283. doi:10.2337/dc09-0209
27. Hajhashemy Z, Shahdadian F, Moslemi E, Mirenayat FS, Saneei P. Serum vitamin D levels in relation to metabolic syndrome: A systematic review and dose–response meta-analysis of epidemiologic studies. *Obes Rev.* 2021;22(7):e13223. doi:10.1111/obr.13223

28. Su D, Nie Y, Zhu A, et al. Vitamin D Signaling through Induction of Paneth Cell Defensins Maintains Gut Microbiota and Improves Metabolic Disorders and Hepatic Steatosis in Animal Models. *Front Physiol.* 2016;7:498. doi:10.3389/fphys.2016.00498
29. Zhang J, Feng M, Pan L, et al. Effects of vitamin D deficiency on the improvement of metabolic disorders in obese mice after vertical sleeve gastrectomy. *Sci Rep.* 2021;11(1):6036. doi:10.1038/s41598-021-85531-9
30. Park CY, Han SN. The Role of Vitamin D in Adipose Tissue Biology: Adipocyte Differentiation, Energy Metabolism, and Inflammation. *J Lipid Atheroscler.* 2021;10(2):130. doi:10.12997/jla.2021.10.2.130
31. Carrodeguas L, Kaidar-Person O, Szomstein S, Antozzi P, Rosenthal R. Preoperative thiamine deficiency in obese population undergoing laparoscopic bariatric surgery. *Surg Obes Relat Dis Off J Am Soc Bariatr Surg.* 2005;1(6):517-522; discussion 522. doi:10.1016/j.soard.2005.08.003
32. Flancbaum L, Belsley S, Drake V, Colarusso T, Tayler E. Preoperative nutritional status of patients undergoing Roux-en-Y gastric bypass for morbid obesity. *J Gastrointest Surg Off J Soc Surg Aliment Tract.* 2006;10(7):1033-1037. doi:10.1016/j.gassur.2006.03.004
33. Fattal-Valevski A. Thiamine (Vitamin B1). *J Evid-Based Complement Altern Med.* 2011;16(1):12-20. doi:10.1177/1533210110392941
34. Krzizek E-C, Brix JM, Herz CT, et al. Prevalence of Micronutrient Deficiency in Patients with Morbid Obesity Before Bariatric Surgery. *Obes Surg.* 2018;28(3):643-648. doi:10.1007/s11695-017-2902-4
35. Diaz K, Na Z, Gupta S, et al. Prevalence of Folic Acid Deficiency and Cost Effectiveness of Folic Acid Testing: A Single Center Experience. *Blood.* 2018;132(Supplement 1):4878-4878. doi:10.1182/blood-2018-99-111607
36. Mechanick JL, Apovian C, Brethauer S, et al. Clinical practice guidelines for the perioperative nutrition, metabolic, and nonsurgical support of patients undergoing bariatric procedures - 2019 update: cosponsored by American Association of Clinical Endocrinologists/American College of Endocrinology, The Obesity Society, American Society for Metabolic & Bariatric Surgery, Obesity Medicine Association, and American Society of Anesthesiologists. *Surg Obes Relat Dis Off J Am Soc Bariatr Surg.* 2020;16(2):175-247. doi:10.1016/j.soard.2019.10.025
37. Arias PM, Domeniconi EA, García M, Esquivel CM, Martínez Lascano F, Foscarini JM. Micronutrient Deficiencies After Roux-en-Y Gastric Bypass: Long-Term Results. *Obes Surg.* 2020;30(1):169-173. doi:10.1007/s11695-019-04167-x
38. Al-Mutawa A, Anderson AK, AlSabah S, Al-Mutawa M. Nutritional Status of Bariatric Surgery Candidates. *Nutrients.* 2018;10(1):67. doi:10.3390/nu10010067
39. van der Wielen RP, Löwik MR, van den Berg H, et al. Serum vitamin D concentrations among elderly people in Europe. *Lancet Lond Engl.* 1995;346(8969):207-210. doi:10.1016/s0140-6736(95)91266-5

40. Orces C, Lorenzo C, Guarneros JE. The Prevalence and Determinants of Vitamin D Inadequacy among U.S. Older Adults: National Health and Nutrition Examination Survey 2007-2014. *Cureus*. 11(8):e5300. doi:10.7759/cureus.5300
41. Verger EO, Aron-Wisnewsky J, Dao MC, et al. Micronutrient and Protein Deficiencies After Gastric Bypass and Sleeve Gastrectomy: a 1-year Follow-up. *Obes Surg*. 2016;26(4):785-796. doi:10.1007/s11695-015-1803-7
42. Ha J, Kwon Y, Kwon J-W, et al. Micronutrient status in bariatric surgery patients receiving postoperative supplementation per guidelines: Insights from a systematic review and meta-analysis of longitudinal studies. *Obes Rev*. 2021;22(7):e13249. doi:10.1111/obr.13249
43. Mehaffey JH, Mehaffey RL, Mullen MG, et al. Nutrient Deficiency 10 Years Following Roux-en-Y Gastric Bypass: Who's Responsible? *Obes Surg*. 2017;27(5):1131-1136. doi:10.1007/s11695-016-2364-0
44. Slusher AL, McAllister MJ, Huang C-J. A therapeutic role for vitamin D on obesity-associated inflammation and weight-loss intervention. *Inflamm Res*. 2015;64(8):565-575. doi:10.1007/s00011-015-0847-4
45. Aron-Wisnewsky J, Clément K. A place for vitamin supplementation and functional food in bariatric surgery? *Curr Opin Clin Nutr Metab Care*. 2019;22(6):442-448. doi:10.1097/MCO.0000000000000602
46. O'Kane M, Parretti HM, Pinkney J, et al. British Obesity and Metabolic Surgery Society Guidelines on perioperative and postoperative biochemical monitoring and micronutrient replacement for patients undergoing bariatric surgery—2020 update. *Obes Rev*. 2020;21(11):e13087. doi:10.1111/obr.13087
47. Agus A, Clément K, Sokol H. Gut microbiota-derived metabolites as central regulators in metabolic disorders. *Gut*. 2021;70(6):1174-1182. doi:10.1136/gutjnl-2020-323071
48. Hill MJ. Intestinal flora and endogenous vitamin synthesis. *Eur J Cancer Prev*. 1997;6:S43-S45. doi:10.1097/00008469-199703001-00009
49. Magnúsdóttir S, Ravcheev D, de Crécy-Lagard V, Thiele I. Systematic genome assessment of B-vitamin biosynthesis suggests co-operation among gut microbes. *Front Genet*. 2015;6:148. doi:10.3389/fgene.2015.00148
50. Rodionov DA, Arzamasov AA, Khoroshkin MS, et al. Micronutrient Requirements and Sharing Capabilities of the Human Gut Microbiome. *Front Microbiol*. 2019;10. doi:10.3389/fmicb.2019.01316
51. Belzer C, Chia LW, Aalvink S, et al. Microbial Metabolic Networks at the Mucus Layer Lead to Diet-Independent Butyrate and Vitamin B12 Production by Intestinal Symbionts. *mBio*. 2017;8(5). doi:10.1128/mBio.00770-17
52. Sharma V, Rodionov DA, Leyn SA, et al. B-Vitamin Sharing Promotes Stability of Gut Microbial Communities. *Front Microbiol*. 2019;10. doi:10.3389/fmicb.2019.01485

53. Cani PD, de Vos WM. Next-Generation Beneficial Microbes: The Case of Akkermansia muciniphila. *Front Microbiol.* 2017;8. doi:10.3389/fmicb.2017.01765
54. Udayappan S, Manneras-Holm L, Chaplin-Scott A, et al. Oral treatment with Eubacterium hallii improves insulin sensitivity in db/db mice. *Npj Biofilms Microbiomes.* 2016;2(1):1-10. doi:10.1038/npjbiofilms.2016.9
55. Pompei A, Cordisco L, Amaretti A, Zanoni S, Matteuzzi D, Rossi M. Folate Production by Bifidobacteria as a Potential Probiotic Property. *Appl Environ Microbiol.* 2007;73(1):179-185. doi:10.1128/AEM.01763-06
56. Pompei A, Cordisco L, Amaretti A, et al. Administration of Folate-Producing Bifidobacteria Enhances Folate Status in Wistar Rats. *J Nutr.* 2007;137(12):2742-2746. doi:10.1093/jn/137.12.2742
57. Rossi M, Amaretti A, Raimondi S. Folate production by probiotic bacteria. *Nutrients.* 2011;3(1):118-134. doi:10.3390/nu3010118
58. Gustafsson BE, Daft FS, McDaniel EG, Smith JC, Fitzgerald RJ. Effects of Vitamin K-Active Compounds and Intestinal Microorganisms in Vitamin K-Deficient Germfree Rats. *J Nutr.* 1962;78(4):461-468. doi:10.1093/jn/78.4.461
59. Fridericia LS, Freudenthal P, Gudjonsson S, Johansen G, Schoubye N. Refection, a Transmissible Change in the Intestinal Content, enabling Rats to grow and thrive without Vitamin B in the Food. *Epidemiology & Infection.* Published online 1927. doi:10.1017/S002217240003182X
60. Bender D. *Nutritional Biochemistry of the Vitamins.*; 2003. doi:10.1017/CBO9780511615191
61. Oppel TW. Studies of Biotin Metabolism in Man. Part I. The Excretion of Biotin in, Human Urine. *Am J Med Sci.* 1942;204(6):856-875.
62. Hayashi A, Mikami Y, Miyamoto K, et al. Intestinal Dysbiosis and Biotin Deprivation Induce Alopecia through Overgrowth of Lactobacillus murinus in Mice. *Cell Rep.* 2017;20(7):1513-1524. doi:10.1016/j.celrep.2017.07.057
63. Miki T, Goto R, Fujimoto M, Okada N, Hardt W-D. The Bactericidal Lectin RegIII β Prolongs Gut Colonization and Enteropathy in the Streptomycin Mouse Model for Salmonella Diarrhea. *Cell Host Microbe.* 2017;21(2):195-207. doi:10.1016/j.chom.2016.12.008
64. Le Chatelier E, Nielsen T, Qin J, et al. Richness of human gut microbiome correlates with metabolic markers. *Nature.* 2013;500(7464):541-546. doi:10.1038/nature12506
65. Cotillard A, Kennedy SP, Kong LC, et al. Dietary intervention impact on gut microbial gene richness. *Nature.* 2013;500(7464):585-588. doi:10.1038/nature12480
66. Aron-Wisniewsky J, Prifti E, Belda E, et al. Major microbiota dysbiosis in severe obesity: fate after bariatric surgery. *Gut.* 2019;68(1):70-82. doi:10.1136/gutjnl-2018-316103

67. Ellis JL, Karl JP, Oliverio AM, et al. Dietary vitamin K is remodeled by gut microbiota and influences community composition. *Gut Microbes*. 2021;13(1):1-16. doi:10.1080/19490976.2021.1887721
68. Hibberd MC, Wu M, Rodionov DA, et al. The effects of micronutrient deficiencies on bacterial species from the human gut microbiota. *Sci Transl Med*. 2017;9(390). doi:10.1126/scitranslmed.aal4069
69. Feng D, Chen B, Zeng B, et al. Faecal microbiota from children with vitamin A deficiency impairs colonic barrier function in germ-free mice: the possible role of alterative bile acid metabolites. *Nutrition*. Published online April 20, 2021:111274. doi:10.1016/j.nut.2021.111274
70. Tian Y, Nichols RG, Cai J, Patterson AD, Cantorna MT. Vitamin A deficiency in mice alters host and gut microbial metabolism leading to altered energy homeostasis. *J Nutr Biochem*. 2018;54:28-34. doi:10.1016/j.jnutbio.2017.10.011
71. Ley RE, Turnbaugh PJ, Klein S, Gordon JI. Microbial ecology: human gut microbes associated with obesity. *Nature*. 2006;444(7122):1022-1023. doi:10.1038/4441022a
72. Jumpertz R, Le DS, Turnbaugh PJ, et al. Energy-balance studies reveal associations between gut microbes, caloric load, and nutrient absorption in humans¹²³. *Am J Clin Nutr*. 2011;94(1):58-65. doi:10.3945/ajcn.110.010132
73. Duncan SH, Lobley GE, Holtrop G, et al. Human colonic microbiota associated with diet, obesity and weight loss. *Int J Obes*. 2008;32(11):1720-1724. doi:10.1038/ijo.2008.155
74. Thomas RL, Jiang L, Adams JS, et al. Vitamin D metabolites and the gut microbiome in older men. *Nat Commun*. 2020;11(1):5997. doi:10.1038/s41467-020-19793-8
75. Ooi JH, Li Y, Rogers CJ, Cantorna MT. Vitamin D Regulates the Gut Microbiome and Protects Mice from Dextran Sodium Sulfate–Induced Colitis¹²³. *J Nutr*. 2013;143(10):1679-1686. doi:10.3945/jn.113.180794
76. Jin D, Wu S, Zhang Y-G, et al. Lack of Vitamin D Receptor Causes Dysbiosis and Changes the Functions of the Murine Intestinal Microbiome. *Clin Ther*. 2015;37(5):996-1009.e7. doi:10.1016/j.clinthera.2015.04.004
77. Mayengbam S, Chleilat F, Reimer RA. Dietary Vitamin B6 Deficiency Impairs Gut Microbiota and Host and Microbial Metabolites in Rats. *Biomedicines*. 2020;8(11). doi:10.3390/biomedicines8110469
78. Fangmann D, Theismann E-M, Türk K, et al. Targeted Microbiome Intervention by Microencapsulated Delayed-Release Niacin Beneficially Affects Insulin Sensitivity in Humans. *Diabetes Care*. 2018;41(3):398-405. doi:10.2337/dc17-1967
79. Forslund K, Hildebrand F, Nielsen T, et al. Disentangling type 2 diabetes and metformin treatment signatures in the human gut microbiota. *Nature*. 2015;528(7581):262-266. doi:10.1038/nature15766
80. Sun L, Xie C, Wang G, et al. Gut microbiota and intestinal FXR mediate the clinical benefits of metformin. *Nat Med*. Published online November 5, 2018:1. doi:10.1038/s41591-018-0222-4

81. Yang Q, Liang Q, Balakrishnan B, Belobrajdic DP, Feng Q-J, Zhang W. Role of Dietary Nutrients in the Modulation of Gut Microbiota: A Narrative Review. *Nutrients*. 2020;12(2). doi:10.3390/nu12020381
82. Jiang S, Zhu Q, Mai M, Yang W, Du G. Vitamin B and vitamin D as modulators of gut microbiota in overweight individuals. *Int J Food Sci Nutr*. 2020;71(8):1001-1009. doi:10.1080/09637486.2020.1748580
83. Almuzara M, Cittadini R, Estraviz ML, Ellis A, Vay C. First report of *Comamonas kerstersii* causing urinary tract infection. *New Microbes New Infect*. 2018;24:4-7. doi:10.1016/j.nmni.2018.03.003
84. Giannelli V, Di Gregorio V, Iebba V, et al. Microbiota and the gut-liver axis: Bacterial translocation, inflammation and infection in cirrhosis. *World J Gastroenterol WJG*. 2014;20(45):16795-16810. doi:10.3748/wjg.v20.i45.16795
85. Clavel T, Duck W, Charrier C, Wenning M, Elson C, Haller D. *Enterorhabdus caecimuris* sp. nov., a member of the family Coriobacteriaceae isolated from a mouse model of spontaneous colitis, and emended description of the genus *Enterorhabdus* Clavel et al. 2009. *Int J Syst Evol Microbiol*. 2010;60(Pt 7):1527-1531. doi:10.1099/ijs.0.015016-0
86. Wu Y, Chi X, Zhang Q, Chen F, Deng X. Characterization of the salivary microbiome in people with obesity. *PeerJ*. 2018;6:e4458. doi:10.7717/peerj.4458
87. Kameyama K, Itoh K. Intestinal colonization by a Lachnospiraceae bacterium contributes to the development of diabetes in obese mice. *Microbes Environ*. 2014;29(4):427-430. doi:10.1264/jsme2.ME14054
88. Xie J, Peters BM, Li B, et al. Clinical features and antimicrobial resistance profiles of important Enterobacteriaceae pathogens in Guangzhou representative of Southern China, 2001-2015. *Microb Pathog*. 2017;107:206-211. doi:10.1016/j.micpath.2017.03.038
89. Naderpoor N, Mousa A, Fernanda Gomez Arango L, Barrett HL, Dekker Nitert M, de Courten B. Effect of Vitamin D Supplementation on Faecal Microbiota: A Randomised Clinical Trial. *Nutrients*. 2019;11(12). doi:10.3390/nu11122888
90. Biyong EF, Alfors S, Dumetz F, et al. Dietary vitamin A supplementation prevents early obesogenic diet-induced microbiota, neuronal and cognitive alterations. *Int J Obes*. Published online November 22, 2020:1-11. doi:10.1038/s41366-020-00723-z
91. Pham VT, Fehlbauer S, Seifert N, et al. Effects of colon-targeted vitamins on the composition and metabolic activity of the human gut microbiome— a pilot study. *Gut Microbes*. 2021;13(1):1-20. doi:10.1080/19490976.2021.1875774
92. LeBlanc JG, Chain F, Martín R, Bermúdez-Humarán LG, Courau S, Langella P. Beneficial effects on host energy metabolism of short-chain fatty acids and vitamins produced by commensal and probiotic bacteria. *Microb Cell Factories*. 2017;16(1):79. doi:10.1186/s12934-017-0691-z
93. Strozzi GP, Mogna L. Quantification of Folic Acid in Human Feces After Administration of Bifidobacterium Probiotic Strains: *J Clin Gastroenterol*. 2008;42:S179-S184. doi:10.1097/MCG.0b013e31818087d8

94. Engevik MA, Morra CN, Röth D, et al. Microbial Metabolic Capacity for Intestinal Folate Production and Modulation of Host Folate Receptors. *Front Microbiol.* 2019;10:2305. doi:10.3389/fmicb.2019.02305
95. Gokhale S, Bhaduri A. Provitamin D 3 modulation through prebiotics supplementation: simulation based assessment. *Sci Rep.* 2019;9(1):19267. doi:10.1038/s41598-019-55699-2
96. World Health Organization, Food and Agriculture Organization of the United Nations, eds. *Vitamin and Mineral Requirements in Human Nutrition*. 2nd ed. World Health Organization ; FAO; 2004.