



**HAL**  
open science

## **Prenatal Assessment of Atypical Adrenal Glands**

Emmanuelle Lesieur, Aurélie Noire, Paul Maurice, Catherine Garel,  
Jean-marie Jouannic, Katia Chaumoitre, Florence Bretelle, Jean Baptiste  
Haumonte, Guillaume Gorincour, Edwin Quarello

► **To cite this version:**

Emmanuelle Lesieur, Aurélie Noire, Paul Maurice, Catherine Garel, Jean-marie Jouannic, et al.. Prenatal Assessment of Atypical Adrenal Glands: A Systematic Approach for Diagnosis. *Journal of Ultrasound in Medicine*, 2021, 40 (8), pp.1719-1728. 10.1002/jum.15536 . hal-03461871

**HAL Id: hal-03461871**

**<https://hal.sorbonne-universite.fr/hal-03461871>**

Submitted on 1 Dec 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Prenatal assessment of atypical adrenal glands: a systematic approach for diagnosis**

Journal:	<i>Journal of Ultrasound in Medicine</i>
Manuscript ID	JUM-2020-06-0934.R1
Wiley - Manuscript type:	Case Series
Date Submitted by the Author:	n/a
Complete List of Authors:	Lesieur, Emmanuelle; Screening and diagnosis Unit Noire, Aurelie; Cabinet de Gynécologie et Obstétrique Maurice, Paul; APHP, Service de Médecine Fœtale, Hôpital Armand Trousseau Garel, Catherine; APHP, Department of pediatric imaging, Hôpital d'Enfants Armand-Trousseau Jouannic, Jean-Marie; APHP, Service de Médecine Fœtale, Hôpital Armand Trousseau Chaumoitre, Katia; APHM, : Service d'Imagerie Médicale, AP-HM Hôpital Nord Bretelle, Florence; APHM, Department of Gynaecology and Obstetrics, Gynepole Haumonte, Jean-Baptiste; Screening and Diagnosis Unit, Obstetrics and Gynecology Gorincour, Guillaume; Image2 Center Quarello, Edwin; Unite dechographies obstetricales et de diagnostic antenatal, Hopital Saint Joseph
Section:	Obstetric (Detailed Fetal Anatomy)
Keywords:	Prenatal, ultrasound, adrenal gland, MRI, counseling
Abstract:	

1 **TITLE**2 **Prenatal assessment of atypical adrenal glands: a systematic approach for diagnosis**

3

4

5

6 **AUTHORS**

7 **Emmanuelle Lesieur** <sup>(1)</sup>, **Aurélie Noire** <sup>(2)</sup>, **Paul Maurice** <sup>(3)</sup>, **Catherine Garel** <sup>(4)</sup>, **Jean-Marie Jouannic** <sup>(3)</sup>,  
8 **Katia Chaumoitre** <sup>(5)</sup>, **Florence Bretelle** <sup>(6)</sup>, **Jean Baptiste Haumonte** <sup>(1)</sup>, **Guillaume Gorincour** <sup>(7)</sup>, **Edwin**  
9 **Quarello** <sup>(1,7)</sup>

10

11 <sup>(1)</sup>: Screening and Diagnosis Unit, Obstetrics and Gynecology Department, St Joseph Hospital, 26 Bd de  
12 Louvain, Marseille Cedex 08, France.

13 <sup>(2)</sup>: Cabinet de Gynécologie et Obstétrique, 7 Av Gambetta, 83400 Hyères, France

14 <sup>(3)</sup>: Service de Médecine Fœtale, Hôpital Armand Trousseau, APHP. Sorbonne Université, 26 avenue A Netter,  
15 75012 Paris, France.

16 <sup>(4)</sup>: Department of pediatric imaging, Hôpital d'Enfants Armand-Trousseau APHP. Sorbonne Université, 26  
17 avenue A Netter, 75012 Paris France.

18 <sup>(5)</sup>: Service d'Imagerie Médicale, AP-HM Hôpital Nord, Chemin des Bourrely, 13015 Marseille, France.

19 <sup>(6)</sup>: Department of Gynaecology and Obstetrics, Gynepole, AP-HM, Assistance Publique-Hôpitaux de  
20 Marseille, Hôpital Nord, Chemin des Bourrely, 13015 Marseille AMU, Aix-Marseille Université, France.

21 <sup>(7)</sup>: IMAGE<sup>2</sup> Center, 6 rue Rocca, Marseille 13008, France

22

23 ***Correspondence :***

24 Dr Edwin QUARELLO

25 Image 2 Center

26 6 rue Rocca

27 Marseille 13008

28 France

29 [e.quarello@me.com](mailto:e.quarello@me.com)

30

31

32

33

34

35

## SUMMARY

The aim of this study was to identify specific unusual prenatal ultrasonographic patterns of the adrenal gland and to propose a systematic approach for diagnosis. Six foetuses with unusual aspects of one or both adrenal glands, detected during routine prenatal ultrasound screening, were evaluated. Prenatal and postnatal management are described. A checklist of ultrasound features was created in order to perform a detailed analysis of adrenal lesions and guide prenatal management; this includes: time of appearance, location, growth, vascularization, structure, and the presence of findings suggestive of malignancy.

Prenatal – ultrasound – adrenal gland – MRI – counseling

## INTRODUCTION

With the widespread use of foetal ultrasound (US), identification of atypical suprarenal masses has become more common. Yet, optimal prenatal diagnosis and prognosis remain challenging.<sup>1</sup> Identifying characteristic imaging findings can help achieve the most relevant diagnosis.<sup>2</sup> Evaluation of the adrenal glands is not included in standard guidelines. Atypical presentation often results from an increased size and / or abnormal appearance (cystic, enlarged, pseudo-tumoural).

Although sufficient epidemiological knowledge of these atypical suprarenal entities is still lacking, five diagnoses stand out according to their location and characteristics: neuroblastoma (NB), congenital adrenal hyperplasia (CAH), adrenal haemorrhage (AH), adrenocortical macrocyst (AM), and adrenal cyst (AC). The main diagnostic issue is the exclusion of malignant neuroblastoma.<sup>3-5</sup> The principal differential diagnosis is subdiaphragmatic extralobar pulmonary sequestration (SEPS).<sup>6</sup> Other less frequent causes can, however, be considered such as enteric duplication cyst, some renal cystic diseases, kidney duplication with upper pole moiety dilatation, and focal renal dysplasia.<sup>7-9</sup>

The definitive diagnosis of an atypical suprarenal entity using US is challenging, especially since the complications specific to each diagnosis imply very different prognoses and postnatal care.<sup>10</sup> Although in most cases the diagnosis cannot be given prenatally, the distinction between specific US features provides a clearer picture to support parental counselling. Based on a series of six cases of atypical adrenal glands, the aim of this study was to propose a systematic US approach in order to aid diagnosis and improve perinatal care and counselling.

**65 PATIENTS AND METHODS**

66 We retrospectively reviewed the US descriptions of cases with an atypical aspect of one or both adrenals  
67 detected during routine prenatal US screening in three different prenatal diagnostic centres, in Marseille and  
68 Paris, France over a five-year period. In accordance with our country's regulations, the retrospective use of  
69 descriptive data does not require specific authorization in France (IRB or equivalent). Patients, all of whom  
70 are informed during US examinations of the possible use of their data for scientific purposes (informed  
71 consent), may at any time indicate their refusal.

72 During the period 2015-2020, cases were investigated in order to characterise different atypical adrenal  
73 entities. Six cases were encountered, leading to the characterization of six different entities. This analysis  
74 focused only on prenatal US description, and all recorded foetal US images were reviewed. The prenatal and  
75 postnatal management of each case is described. All US examinations were performed by expert clinicians. If  
76 magnetic resonance imaging (MRI) was performed, a precise evaluation of the lesion was carried out. Based  
77 on the data, a checklist of US features was then created in order to establish detailed analysis of the adrenal  
78 lesion and guide prenatal management.

79

80

81

82

83

84

85

86

87

88

89

90

91

**92 RESULTS**

93 Six unusual foetuses, detected during routine prenatal US screening, were assessed. In the first case, congenital  
94 hyperplasia of the adrenal gland was diagnosed antenatally followed by postnatal management. In the second  
95 case, an atypical adrenal gland was described in a patient with Beckwith-Wiedemann syndrome. The third  
96 illustrates the prenatal diagnosis of neuroblastoma together with postnatal management. In the fourth and fifth  
97 cases, a complex and a simple haemorrhagic cyst were diagnosed antenatally respectively. The last case  
98 involved extralobar pulmonary sequestration, a differential diagnosis that was established faced with an  
99 atypical adrenal mass. The description of these cases is presented in Table 1

For Review Only

## DISCUSSION

Based on US foetal screening, the adrenal gland is one of the anatomical landmarks necessary to accurately define the abdominal perimeter in the mid and late trimesters of pregnancy. The normal appearance of adrenal glands changes with gestational age. In the first trimester, adrenal glands appear as hypoechoic triangles above the kidneys and measure about half the height of the kidney (*Figure 1*). They then gradually lengthen and take an inverted V shape above the kidney, to measure more than a third of the length of the kidney. In the third trimester, the appearance of corticomedullary differentiation is observed with a hypoechoic cortex and hyperechoic medulla (*Figure 1*).<sup>11</sup>

When an atypical adrenal gland is identified, six possible diagnoses should be initially considered, even if other, less frequent, causes are not excluded, as mentioned above (*Figure 2*).<sup>12</sup>

First, the bilateral or unilateral presentation should be considered, followed by its sonographic appearance.

When bilaterally increased in size and associated with normal sonographic appearance, this may indicate adrenal hypertrophy which can be encountered in some cases of gestational diabetes, intra-uterine growth retardation, and foetal inflammatory syndrome, especially in cases of premature rupture of the membranes.<sup>13,14</sup>

In bilateral cases with abnormal sonographic appearance, congenital adrenal hyperplasia (CAH) should be suspected (*Table 1 - Case 1, Figure 3*). This autosomal recessive disorder may be indicated by the association with two US features: a typical aspect of “cerebriform” adrenal glands, associated with virilization of the genitalia. Supported by a family history, these two US features favour the diagnosis of CAH. The prognosis is excellent with appropriate treatment.<sup>15,16</sup> Second, the presence of bilateral cysts with mixed content, possibly

partially haemorrhagic, can raise suspicion of an adrenocortical macrocyst in the setting of Beckwith-Wiedemann syndrome. Other US features should be investigated, such as foetal macrosomia with polyhydramnios, macroglossia, and omphalocele (*Table 1 – Case 2, Figure 4*).<sup>17-19</sup>

When unilateral with an abnormal US appearance, neuroblastoma is the most frequent congenital malignant tumour of the adrenal (90%). It is most often on the right side (60%), but can be bilateral. Appearance is variable, and cystic in 50% of cases (*Table 1 – Case 3, Figure 5*). The contours of neuroblastoma are regular, and the presence of thick walls, vegetations or partitions can be suggestive. The prognosis is variable and depends on its appearance, time at onset and possible secondary locations.<sup>20,21</sup> Overall, a pure cystic form has

127 a good prognosis. Early detection is associated with a good outcome in more than 90% of cases. A few cases  
128 with secondary localization<sup>22</sup> and placental dissemination, with a high risk of foetal hydrops and maternal  
129 preeclampsia, have been described, most often with a poor prognosis.<sup>23,24</sup>

130 An adrenal haemorrhage presents as a rounded or oval, echogenic, heterogeneous and avascular mass of the  
131 adrenal area with a changing sonographic appearance, which may help diagnosis. Its size gradually decreases  
132 during pregnancy follow-up. Bilateral localization can help diagnosis, as well as specific MRI features (**Table**  
133 **1 – Case 4, Figure 6**). A foetal adrenal haemorrhage can regress and disappear spontaneously during the  
134 prenatal period. Adrenal insufficiency is found very rarely, even in cases of bilateral adrenal hemorrhage.<sup>25</sup>

135 Isolated adrenal cysts (**Table 1 – Case 5, Figure 7**) can occur unilaterally. Differential diagnoses of  
136 neuroblastoma (cystic form) and adrenocortical macrocyst (Beckwith-Wiedemann syndrome) should be kept  
137 in mind. Spontaneous involution is most likely. An absence of malignancy must be confirmed at birth.<sup>26,27</sup>  
138 The differential diagnosis for neuroblastoma is difficult during the prenatal period.<sup>28</sup>

139 Extralobar pulmonary sequestration is also generally diagnosed early in pregnancy, more commonly in the  
140 mid trimester (**Table 1 – Case 6, Figure 8**). It is defined as a portion of lung tissue that is totally  
141 discontinuous from the tracheobronchial tree and usually has its own pleural covering with an anomalous  
142 systemic blood supply.<sup>6</sup> It can have a variable sonographic appearance, and is more frequently left-sided.

143 The US diagnosis is based on the demonstration of a feeding systemic artery that can be identified on colour  
144 Doppler imaging. Most often, the artery is derived from the terminal portion of the descending thoracic aorta  
145 or the upper part of the abdominal aorta.<sup>29</sup>

146 All specific prenatal features and prognostic factors of these three entities are summarized in Table 2.

147 MRI can be a useful adjunct in the prenatal diagnosis of unusual suprarenal entities. Indeed, MRI confirms  
148 US diagnosis, complements US in equivocal diagnoses, and may lead to detection of additional findings.<sup>30</sup>

149 Regarding term at delivery, although some authors have suggested that the increase in size of the adrenal gland  
150 may be related to preterm delivery,<sup>31</sup> especially when associated with polyhydramnios, others believe that  
151 there is no relationship between size of the adrenal gland and gestational age at delivery.<sup>32</sup>

152 The treatment of such entities has not yet been determined, and recommendations have changed in recent  
153 years. Initially, surgery was favoured in all cases<sup>33,34</sup>, then a wait-and-see attitude, based on close prenatal and



154 postnatal US follow-up, was promoted.<sup>4,35</sup> Overall, patients prenatally diagnosed with extralobar pulmonary  
155 sequestration have an excellent prognosis and a wait-and-see attitude, followed by possible postnatal  
156 embolization and/or resection, is recommended.<sup>36</sup> As far as adrenal haemorrhage is concerned, therapeutic  
157 abstention is required. Regarding neuroblastomas, surgical excision is performed immediately for tissue /  
158 mixed forms that have increased in volume. For prenatal cystic and stable forms, a wait-and-see attitude could  
159 be proposed initially, to be re-evaluated on a case-by-case basis.

160 Appropriate prenatal assessment and close sonographic monitoring may avoid surgery in cases of benign  
161 masses such as adrenal haemorrhage or spontaneously regressing neuroblastomas.

## 163 CONCLUSION

164 Prenatal US diagnosis of atypical adrenal glands requires a detailed analysis of the lesion, since the prognosis  
165 and morbidity vary according to final diagnoses. Knowledge and distinction of relevant US signs are necessary  
166 for optimal management. The analysis of atypical adrenal glands underlines the complexity of foetal US and  
167 the gap between prenatally suspected conditions and final postnatal diagnosis, which may be potentially very  
168 different. Optimization of US analysis may improve perinatal care and prenatal counselling.

181 **BIBLIOGRAPHY**

- 182 1. Rubenstein SC, Benacerraf BR, Retik AB, Mandell J. Fetal suprarenal masses: sonographic appearance  
 183 and differential diagnosis. *Ultrasound Obstet Gynecol Off J Int Soc Ultrasound Obstet Gynecol.*  
 184 1995;5(3):164-167. doi:10.1046/j.1469-0705.1995.05030164.x
- 185 2. Sargar KM, Khanna G, Hulett Bowling R. Imaging of Nonmalignant Adrenal Lesions in Children.  
 186 *Radiogr Rev Publ Radiol Soc N Am Inc.* 2017;37(6):1648-1664. doi:10.1148/rg.2017170043
- 187 3. Maki E, Oh K, Rogers S, Sohaey R. Imaging and Differential Diagnosis of Suprarenal Masses in the  
 188 Fetus. *J Ultrasound Med.* 2014;33(5):895-904. doi:10.7863/ultra.33.5.895
- 189 4. Sauvat F, Sarnacki S, Brisse H, et al. Outcome of suprarenal localized masses diagnosed during the  
 190 perinatal period: a retrospective multicenter study. *Cancer.* 2002;94(9):2474-2480.  
 191 doi:10.1002/cncr.10502
- 192 5. Schwärzler P, Bernard JP, Senat MV, Ville Y. Prenatal diagnosis of fetal adrenal masses: differentiation  
 193 between hemorrhage and solid tumor by color Doppler sonography. *Ultrasound Obstet Gynecol Off J*  
 194 *Int Soc Ultrasound Obstet Gynecol.* 1999;13(5):351-355. doi:10.1046/j.1469-0705.1999.13050351.x
- 195 6. Kalenahalli KV, Garg N, Goolahally LN, Reddy SP, Iyengar J. Infradiaphragmatic extralobar pulmonary  
 196 sequestration: masquerading as suprarenal mass. *J Clin Neonatol.* 2013;2(3):146-148. doi:10.4103/2249-  
 197 4847.120009
- 198 7. Brame M, Masel J, Homsy Y. Antenatal detection and management of suprarenal masses. *Urology.*  
 199 1999;54(6):1097. doi:10.1016/s0090-4295(99)00425-2
- 200 8. McNamara A, Levine D. Intraabdominal Fetal Echogenic Masses: A Practical Guide to Diagnosis and  
 201 Management. *RadioGraphics.* 2005;25(3):633-645. doi:10.1148/rg.253045124
- 202 9. Nakazawa N, Okazaki T, Miyano T. Prenatal detection of isolated gastric duplication cyst. *Pediatr Surg*  
 203 *Int.* 2005;21(10):831-834. doi:10.1007/s00383-005-1517-3
- 204 10. Yao W, Li K, Xiao X, Zheng S, Chen L. Neonatal suprarenal mass: differential diagnosis and treatment.  
 205 *J Cancer Res Clin Oncol.* 2013;139(2):281-286. doi:10.1007/s00432-012-1316-x
- 206 11. Matter D, Sick H, Warter P, Koritke JG. [Echographic study of the normal adrenal gland during  
 207 development and in the adult]. *Morphol Med.* 1983;3(1):19-25.
- 208 12. Daneman A, Baunin C, Lobo E, et al. Disappearing suprarenal masses in fetuses and infants. *Pediatr*  
 209 *Radiol.* 1997;27(8):675-681.
- 210 13. Hetkamps T, Hammer K, Möllers M, et al. Fetal Adrenal Gland Size in Gestational Diabetes Mellitus.  
 211 *Journal of perinatal medicine.* doi:10.1515/jpm-2019-0146
- 212 14. Heese S, Hammer K, Möllers M et al. Adrenal Gland Size in Growth Restricted Fetuses. *Journal of*  
 213 *perinatal medicine.* doi:10.1515/jpm-2017-0339
- 214 15. Saada J, Grebille A-G, Aubry M-C, Rafii A, Dumez Y, Benachi A. Sonography in prenatal diagnosis of  
 215 congenital adrenal hyperplasia. *Prenat Diagn.* 2004;24(8):627-630. doi:10.1002/pd.948
- 216 16. Sonographic diagnosis of fetal adrenal hyperplasia: utility for prenatal corticotherapy. Accessed March  
 217 29, 2020. <https://obgyn-onlinelibrary-wiley-com.lama.univ-amu.fr/doi/epdf/10.1002/pd.1219>
- 218 17. MacFarland SP, Mostoufi-Moab S, Zellely K, et al. Management of adrenal masses in patients with  
 219 Beckwith-Wiedemann syndrome. *Pediatr Blood Cancer.* 2017;64(8). doi:10.1002/pbc.26432

- 220 18. Wieacker P, Wilhelm C, Greiner P, Schillinger H. Prenatal diagnosis of Wiedemann-Beckwith  
221 syndrome. *J Perinat Med*. 1989;17(5):351-355. doi:10.1515/jpme.1989.17.5.351
- 222 19. Merrot T, Walz J, Anastasescu R, Chaumoître K, D'Ercole C. Prenatally detected cystic adrenal mass  
223 associated with Beckwith-Wiedemann syndrome. *Fetal Diagn Ther*. 2004;19(6):465-469.  
224 doi:10.1159/000080155
- 225 20. Menager N, Quarello E, Capelle M, et al. Prenatal diagnosis of atypical adrenal neuroblastoma with  
226 pulmonary metastases is possible: Impact on the assessment of prenatal prognosis. *Diagn Interv Imaging*.  
227 2012;93(7-8):632-634. doi:10.1016/j.diii.2012.03.005
- 228 21. Heling KS, Chaoui R, Hartung J, Kirchmair F, Bollmann R. Prenatal diagnosis of congenital  
229 neuroblastoma. Analysis of 4 cases and review of the literature. *Fetal Diagn Ther*. 1999;14(1):47-52.  
230 doi:10.1159/000020888
- 231 22. Liyanage IS, Katoch D. Ultrasonic prenatal diagnosis of liver metastases from adrenal neuroblastoma. *J*  
232 *Clin Ultrasound JCU*. 1992;20(6):401-403. doi:10.1002/jcu.1870200607
- 233 23. Le Bouar G, Lassel L, Loeuillet-Olivo L, d'Herve D, Poulain P. [Antenatal metastatic neuroblastoma:  
234 prognostic criteria. A case report]. *J Gynecol Obstet Biol Reprod (Paris)*. 2003;32(8 Pt 1):740-744.
- 235 24. Zhou S-S, Zhou Y-M, Li D, Chen N-N. Preeclampsia and future cardiovascular risk: A point of view  
236 from the clearance of plasma vasoactive amines. *Hypertens Pregnancy*. 2016;35(1):1-14.  
237 doi:10.3109/10641955.2015.1115062
- 238 25. Wang L, Zhan W, Wu D, Ni X, Luo H, Yang Y. Clinical value of serial ultrasonography in the dynamic  
239 observation of foetal cystic adrenal lesions. *Prenat Diagn*. 2018;38(11):829-834. doi:10.1002/pd.5335
- 240 26. Aslan H, Ozseker B, Gul A. Prenatal sonographic and magnetic resonance imaging diagnosis of cystic  
241 neuroblastoma. *Ultrasound Obstet Gynecol Off J Int Soc Ultrasound Obstet Gynecol*. 2004;24(6):693-  
242 694. doi:10.1002/uog.1733
- 243 27. Tait DL, Williams J, Sandstad J, Lucci JA. Benign adrenal cyst presenting in a pregnant patient. *Am J*  
244 *Perinatol*. 1997;14(8):461-464. doi:10.1055/s-2007-994180
- 245 28. Marcenaro L, Campos Varas JM, Vuletin Solis JF, et al [Differential Diagnosis and Minimally Invasive  
246 Surgery of an Antenatal Adrenal Mass]. *Revista chilena de pediatria*. doi:10.32641/rchped.v90i3.823
- 247 29. Obeidat N, Sallout B, ALAAli W. Isolated subdiaphragmatic extralobar pulmonary sequestration:  
248 masquerading as suprarenal mass with spontaneous resolution. *Clin Exp Obstet Gynecol*.  
249 2016;43(3):457-459.
- 250 30. Flanagan SM, Rubesova E, Jaramillo D, Barth RA. Fetal suprarenal masses--assessing the  
251 complementary role of magnetic resonance and ultrasound for diagnosis. *Pediatr Radiol*.  
252 2016;46(2):246-254. doi:10.1007/s00247-015-3470-1
- 253 31. Santipap M, Phupong V. Combination of Three-Dimensional Ultrasound Measurement of Foetal Adrenal  
254 Gland Enlargement and Placental Alpha microglobulin-1 for the Prediction of the Timing of Delivery  
255 Within Seven Days in Women With Threatened Preterm Labour and Preterm Labour. *Journal of*  
256 *obstetrics and gynaecology: the journal of the Institute of Obstetrics and Gynaecology*.  
257 doi:10.1080/01443615.2018.1446422
- 258 32. Lackova E, Cunderlik A, Ticha L, Gabor M. Fetal Adrenal Gland Enlargement - Prenatal and Postnatal  
259 Management. *Neuro endocrinology letters*. Published November 2017. Accessed April 5, 2020.

- 260 33. Moon S-B, Shin H-B, Seo J-M, Lee S-K. Clinical features and surgical outcome of a suprarenal mass  
261 detected before birth. *Pediatr Surg Int.* 2010;26(3):241-246. doi:10.1007/s00383-009-2531-7
- 262 34. Lin JN, Lin GJ, Hung IJ, Hsueh C. Prenatally detected tumor mass in the adrenal gland. *J Pediatr Surg.*  
263 1999;34(11):1620-1623. doi:10.1016/s0022-3468(99)90629-2
- 264 35. de Luca JL, Rousseau T, Durand C, Sagot P, Sapin E. Diagnostic and therapeutic dilemma with large  
265 prenatally detected cystic adrenal masses. *Fetal Diagn Ther.* 2002;17(1):11-16. doi:10.1159/000047998
- 266 36. Schwab ME, Braun HJ, Padilla BE. Imaging modalities and management of prenatally diagnosed  
267 suprarenal masses: an updated literature review and the experience at a high volume Fetal Treatment  
268 Center. *J Matern-Fetal Neonatal Med Off J Eur Assoc Perinat Med Fed Asia Ocean Perinat Soc Int Soc*  
269 *Perinat Obstet.* Published online January 26, 2020:1-8. doi:10.1080/14767058.2020.1716719

For Review Only

	<b>DIAGNOSIS</b>	<b>Congenital adrenal hyperplasia</b>	<b>Adrenocortical macrocyst Beckwith Wiedemann Syndrome</b>	<b>Solid neuroblastoma</b>	<b>Adrenal haemorrhage</b>	<b>Isolated adrenal cyst</b>	<b>Left extralobar pulmonary sequestration</b>	
<b>PRENATAL MANAGEMENT</b>	PATIENT'S AGE (years)	37	31	26	32	26	29	
	TRIMESTER AND WG AT DISCOVERY (EGA)	Mid trimester 24	Mid trimester 25	Late trimester 35	Late trimester 34	Late trimester 32	Mid trimester 24	
	LOCATION	Bilateral	Bilateral	Unilateral left	Unilateral right	Unilateral left	Unilateral left	
	FETAL GENDER	Male	Male	Male	Male	Female	Female	
	ANTENATAL ULTRASOUND FEATURES	<ul style="list-style-type: none"> <li>Size (mm)</li> <li>Structure</li> <li>Vascularized</li> <li>Systemic vessel</li> <li>Associated findings</li> </ul>	Increased Cerebriform aspect  No No Penis increased in size Family history of CAH	50 x 40 Multicystic with mixed content Heterogeneous No No No macroglossia	22 x 32 Hyperechoic Homogeneous Yes No No	38 x 29 Altered content Heterogeneous No No No	10 x 8 Hypoechoic Homogeneous No No No	30 x 20 Mixed component Heterogeneous Yes Yes (from the aorta) No
	FETAL MRI	<ul style="list-style-type: none"> <li>When (WG)</li> <li>Location</li> <li>Structure</li> <li>Associated findings</li> </ul>	NR NR	28 Bilateral  Cystic and heterogeneous formation  Macroglossia	37 Unilateral left Prerenal Retrogastric Well limited, Heterogeneous, No hemorrhagic signs  No	35 Unilateral right  Cystic and heterogeneous formation  No	NR NR  NR  NR	29 and 33 Thoraco abdominal junction  Plurilocular cystic formation  a systemic posterior vessel arising from the aorta
	PRENATAL EVOLUTION	Stable	Increase	Increase	Decrease	Stable	Increase	
	PROPOSED PRENATAL DIAGNOSIS	Congenital adrenal hyperplasia	Adrenocortical macrocyst Beckwith Wiedemann Syndrome	Solid neuroblastoma, Adrenal haemorrhage	Adrenal haemorrhage, Cystic neuroblastoma	Isolated adrenal cyst	Left extralobar pulmonary sequestration	
<b>POSTNATAL MANAGEMENT</b>	COMPLEMENTARY EXAMS	<ul style="list-style-type: none"> <li>Urinary catecholamine</li> <li>Androgenic marquors</li> <li>Postnatal ultrasound / MRI</li> <li>Lesion confirmed</li> <li>Associated elements</li> <li>Histological findings post-surgery</li> </ul>	Negative 21-H deficiency  Yes No  -	NR NR  NR NR  NR	Positive Normal  Yes Liver metastases (present at 1 month) Malignant cells	Negative Normal  Yes No  NR	NR NR  Yes No  No malignant cells	Negative Normal  Yes No  Lung cells – no malignant cells
	EVOLUTION	Normal aspect at one week and one month with treatment	Termination of pregnancy Pathology: fetal macrosomia, macroglossia, bilateral nephromegaly, bilateral adrenal masses with haemorrhagic lesion on the left. No malignant cells.	Spontaneous regression, then stabilization	Spontaneous regression	Stabilization	Stabilization	
	THERAPEUTIC PROTOCOL	Hydrocortisone and fludrocortisone therapy	NR	Close monitoring, then, Surgery	Close monitoring	Monitoring (spontaneous regression)	Close monitoring	

272 Table 1: Observations of fetuses with atypical adrenal gland detected during routine prenatal ultrasound. (EGA: Estimated Gestational Age ; NR: Not Realize)

		BILATERAL LOCATION			UNILATERAL LOCATION			DIFFERENTIAL DIAGNOSIS
		<i>Abnormal appearance</i>		<i>Normal appearance</i>	<i>Abnormal appearance</i>			<i>Abnormal appearance</i>
		<b>Congenital Adrenal Hyperplasia</b>	<b>Adrenocortical macrocyst (Beckwith Wiedemann Syndrome)</b>	<b>GDM FGR FIS</b>	<b>Neuroblastoma</b>	<b>Adrenal haemorrhage</b>	<b>Adrenal Cyst</b>	<b>Extralobar pulmonary sequestration</b>
<i>Characteristic ultrasound signs</i>	<b>Trimester at discovery</b>	Mid trimester Late trimester	Mid trimester Late trimester	Mid trimester Late trimester	Late trimester	Late trimester	Mid trimester Late trimester	Mid trimester Late trimester
	<b>Preference Location</b>	Bilateral	Bilateral ++ (possible unilateral)	Bilateral	Right	No preference	No preference	Left
	<b>Prenatal Evolution</b>							
	Decrease	-	-	+	+	+++	+++	+
	Stable	+++	+	+++	++	++	++	++
	Increase	-	+++	+	+++	+	+	+++
<b>Vascularization</b>								
Vascularized mass	No	No	No	Yes	No	No	Yes	
Systemic artery	No	No	No	Yes	No	No	Yes	
<b>Description of the lesion</b>								
Well limited Structure	Yes Cerebriform pattern	No Cystic, Heterogeneous	Yes Increase in size without abnormal aspect	Yes Homogeneous (solid form) or Heterogeneous (cystic form)	Yes Heterogeneous	Yes Homogeneous	No Homogeneous (nodular form) or Heterogeneous (cystic-nodular form)	
<b>Elements of malignity</b> <i>Partition, vegetation</i>	No	No	No	Yes	No	No	No	
<i>Prognostic Elements</i>	<b>Fetal complications</b>	No	Organomegaly, Macroglossia, Omphalocele Polyhydramnios	No	Possible metastases (liver ++, lungs, skull, bone marrow, placenta)	No	No	Mass effect (fattening diaphragmatic cupolas +/- pleural effusion)
	<b>Prognosis</b>	Excellent with treatment	Dependent on associated malformations	Excellent	Good (90% cure if early detection)	Excellent	Excellent	Good


273

274

Table 2: Check list of characteristic ultrasound signs according to more frequent diagnosis of suprarenal mass and prognostic elements associated.

275 (GDM: Gestational Diabetes Mellitus, FGR: fetal growth restriction, FIS: Fetal inflammatory syndrome

For Review Only




276


277 *Figure 1: Normal appearance of adrenal glands (arrow) according to the trimester of pregnancy. (a) in the first trimester, in parasagittal view it appears as*  
278 *hypoechoic triangle measuring grossly half the height of the kidney. (b) Then, from the second trimester onward, in axial view it takes an oval structure, (c)*  
279 *with appearance of a hypoechoic cortex and a hyperechoic medullary during late trimester ultrasound.*



For Review Only




*Figure 2: Diagnostic orientation of an atypical adrenal gland according to its type of location and appearance*




285

286 *Figure 3: Congenital adrenal hyperplasia at 24 estimated gestational age, with a typical "cerebriform aspect" (a, arrow), associated with a virilization of*  
287 *the genitalia (b) in a male fetus.*




288

289 *Figure 4: Ultrasound at 25 estimated gestational age in a male fetus showing bilateral partially hemorrhagic cysts on both adrenal glands. This typical*  
 290 *ultrasound appearance, on the left (a) and on the right (b) associated with macroglossia (c) was consistent with adrenocortical macrocyst in the setting of*  
 291 *Beckwith – Wiedemann syndrome. Fetal MRI was helpful, confirming the atypical appearance of the adrenals, here in frontal (d) and transverse view (e) on*  
 292 *BFTE sequences. (LAG: Left Adrenal Gland; LK: Left Kidney; RAG: Right Adrenal Gland)*




293

294 *Figure 5: A prenatal well limited, heterogeneous (a) and vascularized (b) mass was diagnosed as a solid neuroblastoma at 35 estimated gestational age in a*  
295 *male fetus. Coronal (c) and sagittal (d) T2-weighted MR images show slightly hyperintense mass of the left adrenal compartment. (e) Apparent diffusion*  
296 *coefficient map shows hypointensity mass that confirming the malignant hypothesis.*



297

298 Figure 6: A heterogeneous and avascular mass of the right adrenal gland, with a changing sonographic pattern through follow-up, on a axial (a1), parasagittal  
 299 (a2) and frontal views (b) is the typical aspect of an adrenal hemorrhage, at 34 weeks estimated gestational age, in a male fetus. Parasagittal (c) and  
 300 transverse (d) T2- HASTE weighted images show hematic sediment confirming the hemorrhagic composition of the mass.




301

302

303

*Figure 7: A well limited, isolated, hypoechoic and homogeneous cyst located above the left kidney in axial (a) and parasagittal (b) views at 32 weeks estimated gestational age in a female fetus, consistent with an adrenal cyst.*



304  
305 *Figure 8: An extralobar pulmonary sequestration in a female fetus, represented by a heterogeneous mass, with mixed component, under the left diaphragm (a)*  
306 *vascularized by a multiple feeding arteries identified on Color Doppler Imaging (b) at 24 weeks estimated gestational age. (c) Coronal SSFSE T2w confirms*  
307 *the left lower lobe mass with systemic feeding vessels (arrow) coursing into the mass.*  
308



For Review Only

**TITLE****Prenatal assessment of atypical adrenal glands: a systematic approach for diagnosis****AUTHORS****Emmanuelle Lesieur** <sup>(1)</sup>, **Aurélie Noire** <sup>(2)</sup>, **Paul Maurice** <sup>(3)</sup>, **Catherine Garel** <sup>(4)</sup>, **Jean-Marie Jouannic** <sup>(3)</sup>, **Katia Chaumoitre** <sup>(5)</sup>, **Florence Bretelle** <sup>(6)</sup>, **Jean Baptiste Haumonte** <sup>(1)</sup>, **Guillaume Gorincour** <sup>(7)</sup>, **Edwin Quarello** <sup>(1,7)</sup>

<sup>(1)</sup>: Screening and Diagnosis Unit, Obstetrics and Gynecology Department, St Joseph Hospital, 26 Bd de Louvain, Marseille Cedex 08, France.

<sup>(2)</sup>: Cabinet de Gynécologie et Obstétrique, 7 Av Gambetta, 83400 Hyères, France

<sup>(3)</sup>: Service de Médecine Fœtale, Hôpital Armand Trousseau, APHP. Sorbonne Université, 26 avenue A Netter, 75012 Paris, France.

<sup>(4)</sup>: Department of pediatric imaging, Hôpital d'Enfants Armand-Trousseau APHP. Sorbonne Université, 26 avenue A Netter, 75012 Paris France.

<sup>(5)</sup>: Service d'Imagerie Médicale, AP-HM Hôpital Nord, Chemin des Bourrely, 13015 Marseille, France.

<sup>(6)</sup>: Department of Gynaecology and Obstetrics, Gynepole, AP-HM, Assistance Publique-Hôpitaux de Marseille, Hôpital Nord, Chemin des Bourrely, 13015 Marseille AMU, Aix-Marseille Université, France.

<sup>(7)</sup>: IMAGE<sup>2</sup> Center, 6 rue Rocca, Marseille 13008, France

***Correspondence :***

Dr Edwin QUARELLO

Image 2 Center

6 rue Rocca

Marseille 13008

France

[e.quarello@me.com](mailto:e.quarello@me.com)

## 36 SUMMARY

37 The aim of this study was to identify specific unusual prenatal ultrasonographic patterns of the adrenal gland  
38 and to propose a systematic approach for diagnosis. Six foetuses with unusual aspects of one or both adrenal  
39 glands, detected during routine prenatal ultrasound screening, were evaluated. Prenatal and postnatal  
40 management are described. A checklist of ultrasound features was created in order to perform a detailed  
41 analysis of adrenal lesions and guide prenatal management; this includes: time of appearance, location,  
42 growth, vascularization, structure, and the presence of findings suggestive of malignancy.

43 Prenatal – ultrasound – adrenal gland – MRI – counseling

## 44 INTRODUCTION

45 With the widespread use of foetal ultrasound (US), identification of atypical suprarenal masses has become  
46 more common. Yet, optimal prenatal diagnosis and prognosis remain challenging.<sup>1</sup> Identifying characteristic  
47 imaging findings can help achieve the most relevant diagnosis.<sup>2</sup> Evaluation of the adrenal glands is not  
48 included in standard guidelines. Atypical presentation often results from an increased size and / or abnormal  
49 appearance (cystic, enlarged, pseudo-tumoural).

50 Although sufficient epidemiological knowledge of these atypical suprarenal entities is still lacking, five  
51 diagnoses stand out according to their location and characteristics: neuroblastoma (NB), congenital adrenal  
52 hyperplasia (CAH), adrenal haemorrhage (AH), adrenocortical macrocyst (AM), and adrenal cyst (AC). The  
53 main diagnostic issue is the exclusion of malignant neuroblastoma.<sup>3-5</sup> The principal differential diagnosis is  
54 subdiaphragmatic extralobar pulmonary sequestration (SEPS).<sup>6</sup> Other less frequent causes can, however, be  
55 considered such as enteric duplication cyst, some renal cystic diseases, kidney duplication with upper pole  
56 moiety dilatation, and focal renal dysplasia.<sup>7-9</sup>

57 The definitive diagnosis of an atypical suprarenal entity using US is challenging, especially since the  
58 complications specific to each diagnosis imply very different prognoses and postnatal care.<sup>10</sup> Although in  
59 most cases the diagnosis cannot be given prenatally, the distinction between specific US features provides  
60 a clearer picture to support parental counselling. Based on a series of six cases of atypical adrenal glands, the  
61 aim of this study was to propose a systematic US approach in order to aid diagnosis and improve perinatal  
62 care and counselling.

**PATIENTS AND METHODS**

We retrospectively reviewed the US descriptions of cases with an atypical aspect of one or both adrenals detected during routine prenatal US screening in three different prenatal diagnostic centres, in Marseille and Paris, France over a five-year period. In accordance with our country's regulations, the retrospective use of descriptive data does not require specific authorization in France (IRB or equivalent). Patients, all of whom are informed during US examinations of the possible use of their data for scientific purposes (informed consent), may at any time indicate their refusal.

During the period 2015-2020, cases were investigated in order to characterise different atypical adrenal entities. Six cases were encountered, leading to the characterization of six different entities. This analysis focused only on prenatal US description, and all recorded foetal US images were reviewed. The prenatal and postnatal management of each case is described. All US examinations were performed by expert clinicians. If magnetic resonance imaging (MRI) was performed, a precise evaluation of the lesion was carried out. Based on the data, a checklist of US features was then created in order to establish detailed analysis of the adrenal lesion and guide prenatal management.

**92 RESULTS**

93 Six unusual foetuses, detected during routine prenatal US screening, were assessed. In the first case, congenital  
94 hyperplasia of the adrenal gland was diagnosed antenatally followed by postnatal management. In the second  
95 case, an atypical adrenal gland was described in a patient with Beckwith-Wiedemann syndrome. The third  
96 illustrates the prenatal diagnosis of neuroblastoma together with postnatal management. In the fourth and fifth  
97 cases, a complex and a simple haemorrhagic cyst were diagnosed antenatally respectively. The last case  
98 involved extralobar pulmonary sequestration, a differential diagnosis that was established faced with an  
99 atypical adrenal mass. The description of these cases is presented in Table 1

For Review Only

## DISCUSSION

Based on US foetal screening, the adrenal gland is one of the anatomical landmarks necessary to accurately define the abdominal perimeter in the mid and late trimesters of pregnancy. The normal appearance of adrenal glands changes with gestational age. In the first trimester, adrenal glands appear as hypoechoic triangles above the kidneys and measure about half the height of the kidney (*Figure 1*). They then gradually lengthen and take an inverted V shape above the kidney, to measure more than a third of the length of the kidney. In the third trimester, the appearance of corticomedullary differentiation is observed with a hypoechoic cortex and hyperechoic medulla (*Figure 1*).<sup>11</sup>

When an atypical adrenal gland is identified, six possible diagnoses should be initially considered, even if other, less frequent, causes are not excluded, as mentioned above (*Figure 2*).<sup>12</sup>

First, the bilateral or unilateral presentation should be considered, followed by its sonographic appearance. When bilaterally increased in size and associated with normal sonographic appearance, this may indicate adrenal hypertrophy which can be encountered in some cases of gestational diabetes, intra-uterine growth retardation, and foetal inflammatory syndrome, especially in cases of premature rupture of the membranes.<sup>13,14</sup> In bilateral cases with abnormal sonographic appearance, congenital adrenal hyperplasia (CAH) should be suspected (*Table 1 - Case 1, Figure 3*). This autosomal recessive disorder may be indicated by the association with two US features: a typical aspect of “cerebriform” adrenal glands, associated with virilization of the genitalia. Supported by a family history, these two US features favour the diagnosis of CAH. The prognosis is excellent with appropriate treatment.<sup>15,16</sup> Second, the presence of bilateral cysts with mixed content, possibly partially haemorrhagic, can raise suspicion of an adrenocortical macrocyst in the setting of Beckwith-Wiedemann syndrome. Other US features should be investigated, such as foetal macrosomia with polyhydramnios, macroglossia, and omphalocele (*Table 1 – Case 2, Figure 4*).<sup>17-19</sup>

When unilateral with an abnormal US appearance, neuroblastoma is the most frequent congenital malignant tumour of the adrenal (90%). It is most often on the right side (60%), but can be bilateral. Appearance is variable, and cystic in 50% of cases (*Table 1 – Case 3, Figure 5*). The contours of neuroblastoma are regular, and the presence of thick walls, vegetations or partitions can be suggestive. The prognosis is variable and depends on its appearance, time at onset and possible secondary locations.<sup>20,21</sup> Overall, a pure cystic form has

127 a good prognosis. Early detection is associated with a good outcome in more than 90% of cases. A few cases  
128 with secondary localization<sup>22</sup> and placental dissemination, with a high risk of foetal hydrops and maternal  
129 preeclampsia, have been described, most often with a poor prognosis.<sup>23,24</sup>

130 An adrenal haemorrhage presents as a rounded or oval, echogenic, heterogeneous and avascular mass of the  
131 adrenal area with a changing sonographic appearance, which may help diagnosis. Its size gradually decreases  
132 during pregnancy follow-up. Bilateral localization can help diagnosis, as well as specific MRI features (**Table**  
133 **1 – Case 4, Figure 6**). A foetal adrenal haemorrhage can regress and disappear spontaneously during the  
134 prenatal period. Adrenal insufficiency is found very rarely, even in cases of bilateral adrenal hemorrhage.<sup>25</sup>

135 Isolated adrenal cysts (**Table 1 – Case 5, Figure 7**) can occur unilaterally. Differential diagnoses of  
136 neuroblastoma (cystic form) and adrenocortical macrocyst (Beckwith-Wiedemann syndrome) should be kept  
137 in mind. Spontaneous involution is most likely. An absence of malignancy must be confirmed at birth.<sup>26,27</sup>  
138 The differential diagnosis for neuroblastoma is difficult during the prenatal period.<sup>28</sup>

139 Extralobar pulmonary sequestration is also generally diagnosed early in pregnancy, more commonly in the  
140 mid trimester (**Table 1 – Case 6, Figure 8**). It is defined as a portion of lung tissue that is totally  
141 discontinuous from the tracheobronchial tree and usually has its own pleural covering with an anomalous  
142 systemic blood supply.<sup>6</sup> It can have a variable sonographic appearance, and is more frequently left-sided.  
143 The US diagnosis is based on the demonstration of a feeding systemic artery that can be identified on colour  
144 Doppler imaging. Most often, the artery is derived from the terminal portion of the descending thoracic aorta  
145 or the upper part of the abdominal aorta.<sup>29</sup>

146 All specific prenatal features and prognostic factors of these three entities are summarized in Table 2.

147 MRI can be a useful adjunct in the prenatal diagnosis of unusual suprarenal entities. Indeed, MRI confirms  
148 US diagnosis, complements US in equivocal diagnoses, and may lead to detection of additional findings.<sup>30</sup>

149 Regarding term at delivery, although some authors have suggested that the increase in size of the adrenal gland  
150 may be related to preterm delivery,<sup>31</sup> especially when associated with polyhydramnios, others believe that  
151 there is no relationship between size of the adrenal gland and gestational age at delivery.<sup>32</sup>

152 The treatment of such entities has not yet been determined, and recommendations have changed in recent  
153 years. Initially, surgery was favoured in all cases<sup>33,34</sup>, then a wait-and-see attitude, based on close prenatal and

154 postnatal US follow-up, was promoted.<sup>4,35</sup> Overall, patients prenatally diagnosed with extralobar pulmonary  
155 sequestration have an excellent prognosis and a wait-and-see attitude, followed by possible postnatal  
156 embolization and/or resection, is recommended.<sup>36</sup> As far as adrenal haemorrhage is concerned, therapeutic  
157 abstention is required. Regarding neuroblastomas, surgical excision is performed immediately for tissue /  
158 mixed forms that have increased in volume. For prenatal cystic and stable forms, a wait-and-see attitude could  
159 be proposed initially, to be re-evaluated on a case-by-case basis.

160 Appropriate prenatal assessment and close sonographic monitoring may avoid surgery in cases of benign  
161 masses such as adrenal haemorrhage or spontaneously regressing neuroblastomas.

## 163 **CONCLUSION**

164 Prenatal US diagnosis of atypical adrenal glands requires a detailed analysis of the lesion, since the prognosis  
165 and morbidity vary according to final diagnoses. Knowledge and distinction of relevant US signs are necessary  
166 for optimal management. The analysis of atypical adrenal glands underlines the complexity of foetal US and  
167 the gap between prenatally suspected conditions and final postnatal diagnosis, which may be potentially very  
168 different. Optimization of US analysis may improve perinatal care and prenatal counselling.



181 **BIBLIOGRAPHY**

- 182 1. Rubenstein SC, Benacerraf BR, Retik AB, Mandell J. Fetal suprarenal masses: sonographic appearance  
183 and differential diagnosis. *Ultrasound Obstet Gynecol Off J Int Soc Ultrasound Obstet Gynecol.*  
184 1995;5(3):164-167. doi:10.1046/j.1469-0705.1995.05030164.x
- 185 2. Sargar KM, Khanna G, Hulett Bowling R. Imaging of Nonmalignant Adrenal Lesions in Children.  
186 *Radiogr Rev Publ Radiol Soc N Am Inc.* 2017;37(6):1648-1664. doi:10.1148/rg.2017170043
- 187 3. Maki E, Oh K, Rogers S, Sohaey R. Imaging and Differential Diagnosis of Suprarenal Masses in the  
188 Fetus. *J Ultrasound Med.* 2014;33(5):895-904. doi:10.7863/ultra.33.5.895
- 189 4. Sauvat F, Sarnacki S, Brisse H, et al. Outcome of suprarenal localized masses diagnosed during the  
190 perinatal period: a retrospective multicenter study. *Cancer.* 2002;94(9):2474-2480.  
191 doi:10.1002/cncr.10502
- 192 5. Schwärzler P, Bernard JP, Senat MV, Ville Y. Prenatal diagnosis of fetal adrenal masses: differentiation  
193 between hemorrhage and solid tumor by color Doppler sonography. *Ultrasound Obstet Gynecol Off J*  
194 *Int Soc Ultrasound Obstet Gynecol.* 1999;13(5):351-355. doi:10.1046/j.1469-0705.1999.13050351.x
- 195 6. Kalenahalli KV, Garg N, Goolahally LN, Reddy SP, Iyengar J. Infradiaphragmatic extralobar pulmonary  
196 sequestration: masquerading as suprarenal mass. *J Clin Neonatol.* 2013;2(3):146-148. doi:10.4103/2249-  
197 4847.120009
- 198 7. Brame M, Masel J, Homsy Y. Antenatal detection and management of suprarenal masses. *Urology.*  
199 1999;54(6):1097. doi:10.1016/s0090-4295(99)00425-2
- 200 8. McNamara A, Levine D. Intraabdominal Fetal Echogenic Masses: A Practical Guide to Diagnosis and  
201 Management. *RadioGraphics.* 2005;25(3):633-645. doi:10.1148/rg.253045124
- 202 9. Nakazawa N, Okazaki T, Miyano T. Prenatal detection of isolated gastric duplication cyst. *Pediatr Surg*  
203 *Int.* 2005;21(10):831-834. doi:10.1007/s00383-005-1517-3
- 204 10. Yao W, Li K, Xiao X, Zheng S, Chen L. Neonatal suprarenal mass: differential diagnosis and treatment.  
205 *J Cancer Res Clin Oncol.* 2013;139(2):281-286. doi:10.1007/s00432-012-1316-x
- 206 11. Matter D, Sick H, Warter P, Koritke JG. [Echographic study of the normal adrenal gland during  
207 development and in the adult]. *Morphol Med.* 1983;3(1):19-25.
- 208 12. Daneman A, Baunin C, Lobo E, et al. Disappearing suprarenal masses in fetuses and infants. *Pediatr*  
209 *Radiol.* 1997;27(8):675-681.
- 210 13. Hetkamps T, Hammer K, Möllers M, et al. Fetal Adrenal Gland Size in Gestational Diabetes Mellitus.  
211 *Journal of perinatal medicine.* doi:10.1515/jpm-2019-0146
- 212 14. Heese S, Hammer K, Möllers M et al. Adrenal Gland Size in Growth Restricted Fetuses. *Journal of*  
213 *perinatal medicine.* doi:10.1515/jpm-2017-0339
- 214 15. Saada J, Grebille A-G, Aubry M-C, Rafii A, Dumez Y, Benachi A. Sonography in prenatal diagnosis of  
215 congenital adrenal hyperplasia. *Prenat Diagn.* 2004;24(8):627-630. doi:10.1002/pd.948
- 216 16. Sonographic diagnosis of fetal adrenal hyperplasia: utility for prenatal corticotherapy. Accessed March  
217 29, 2020. <https://obgyn-onlinelibrary-wiley-com.lama.univ-amu.fr/doi/epdf/10.1002/pd.1219>
- 218 17. MacFarland SP, Mostoufi-Moab S, Zellely K, et al. Management of adrenal masses in patients with  
219 Beckwith-Wiedemann syndrome. *Pediatr Blood Cancer.* 2017;64(8). doi:10.1002/pbc.26432

- 220 18. Wieacker P, Wilhelm C, Greiner P, Schillinger H. Prenatal diagnosis of Wiedemann-Beckwith  
221 syndrome. *J Perinat Med*. 1989;17(5):351-355. doi:10.1515/jpme.1989.17.5.351
- 222 19. Merrot T, Walz J, Anastasescu R, Chaumoître K, D'Ercole C. Prenatally detected cystic adrenal mass  
223 associated with Beckwith-Wiedemann syndrome. *Fetal Diagn Ther*. 2004;19(6):465-469.  
224 doi:10.1159/000080155
- 225 20. Menager N, Quarello E, Capelle M, et al. Prenatal diagnosis of atypical adrenal neuroblastoma with  
226 pulmonary metastases is possible: Impact on the assessment of prenatal prognosis. *Diagn Interv Imaging*.  
227 2012;93(7-8):632-634. doi:10.1016/j.diii.2012.03.005
- 228 21. Heling KS, Chaoui R, Hartung J, Kirchmair F, Bollmann R. Prenatal diagnosis of congenital  
229 neuroblastoma. Analysis of 4 cases and review of the literature. *Fetal Diagn Ther*. 1999;14(1):47-52.  
230 doi:10.1159/000020888
- 231 22. Liyanage IS, Katoch D. Ultrasonic prenatal diagnosis of liver metastases from adrenal neuroblastoma. *J*  
232 *Clin Ultrasound JCU*. 1992;20(6):401-403. doi:10.1002/jcu.1870200607
- 233 23. Le Bouar G, Lassel L, Loeuillet-Olivo L, d'Herve D, Poulain P. [Antenatal metastatic neuroblastoma:  
234 prognostic criteria. A case report]. *J Gynecol Obstet Biol Reprod (Paris)*. 2003;32(8 Pt 1):740-744.
- 235 24. Zhou S-S, Zhou Y-M, Li D, Chen N-N. Preeclampsia and future cardiovascular risk: A point of view  
236 from the clearance of plasma vasoactive amines. *Hypertens Pregnancy*. 2016;35(1):1-14.  
237 doi:10.3109/10641955.2015.1115062
- 238 25. Wang L, Zhan W, Wu D, Ni X, Luo H, Yang Y. Clinical value of serial ultrasonography in the dynamic  
239 observation of foetal cystic adrenal lesions. *Prenat Diagn*. 2018;38(11):829-834. doi:10.1002/pd.5335
- 240 26. Aslan H, Ozseker B, Gul A. Prenatal sonographic and magnetic resonance imaging diagnosis of cystic  
241 neuroblastoma. *Ultrasound Obstet Gynecol Off J Int Soc Ultrasound Obstet Gynecol*. 2004;24(6):693-  
242 694. doi:10.1002/uog.1733
- 243 27. Tait DL, Williams J, Sandstad J, Lucci JA. Benign adrenal cyst presenting in a pregnant patient. *Am J*  
244 *Perinatol*. 1997;14(8):461-464. doi:10.1055/s-2007-994180
- 245 28. Marcenaro L, Campos Varas JM, Vuletin Solis JF, et al [Differential Diagnosis and Minimally Invasive  
246 Surgery of an Antenatal Adrenal Mass]. *Revista chilena de pediatria*. doi:10.32641/rchped.v90i3.823
- 247 29. Obeidat N, Sallout B, ALAAli W. Isolated subdiaphragmatic extralobar pulmonary sequestration:  
248 masquerading as suprarenal mass with spontaneous resolution. *Clin Exp Obstet Gynecol*.  
249 2016;43(3):457-459.
- 250 30. Flanagan SM, Rubesova E, Jaramillo D, Barth RA. Fetal suprarenal masses--assessing the  
251 complementary role of magnetic resonance and ultrasound for diagnosis. *Pediatr Radiol*.  
252 2016;46(2):246-254. doi:10.1007/s00247-015-3470-1
- 253 31. Santipap M, Phupong V. Combination of Three-Dimensional Ultrasound Measurement of Foetal Adrenal  
254 Gland Enlargement and Placental Alpha microglobulin-1 for the Prediction of the Timing of Delivery  
255 Within Seven Days in Women With Threatened Preterm Labour and Preterm Labour. *Journal of*  
256 *obstetrics and gynaecology: the journal of the Institute of Obstetrics and Gynaecology*.  
257 doi:10.1080/01443615.2018.1446422
- 258 32. Lackova E, Cunderlik A, Ticha L, Gabor M. Fetal Adrenal Gland Enlargement - Prenatal and Postnatal  
259 Management. *Neuro endocrinology letters*. Published November 2017. Accessed April 5, 2020.

- 260 33. Moon S-B, Shin H-B, Seo J-M, Lee S-K. Clinical features and surgical outcome of a suprarenal mass  
261 detected before birth. *Pediatr Surg Int.* 2010;26(3):241-246. doi:10.1007/s00383-009-2531-7
- 262 34. Lin JN, Lin GJ, Hung IJ, Hsueh C. Prenatally detected tumor mass in the adrenal gland. *J Pediatr Surg.*  
263 1999;34(11):1620-1623. doi:10.1016/s0022-3468(99)90629-2
- 264 35. de Luca JL, Rousseau T, Durand C, Sagot P, Sapin E. Diagnostic and therapeutic dilemma with large  
265 prenatally detected cystic adrenal masses. *Fetal Diagn Ther.* 2002;17(1):11-16. doi:10.1159/000047998
- 266 36. Schwab ME, Braun HJ, Padilla BE. Imaging modalities and management of prenatally diagnosed  
267 suprarenal masses: an updated literature review and the experience at a high volume Fetal Treatment  
268 Center. *J Matern-Fetal Neonatal Med Off J Eur Assoc Perinat Med Fed Asia Ocean Perinat Soc Int Soc*  
269 *Perinat Obstet.* Published online January 26, 2020:1-8. doi:10.1080/14767058.2020.1716719

For Review Only

	<b>DIAGNOSIS</b>	<b>Congenital adrenal hyperplasia</b>	<b>Adrenocortical macrocyst Beckwith Wiedemann Syndrome</b>	<b>Solid neuroblastoma</b>	<b>Adrenal haemorrhage</b>	<b>Isolated adrenal cyst</b>	<b>Left extralobar pulmonary sequestration</b>	
<b>PRENATAL MANAGEMENT</b>	PATIENT'S AGE (years)	37	31	26	32	26	29	
	TRIMESTER AND WG AT DISCOVERY (EGA)	Mid trimester 24	Mid trimester 25	Late trimester 35	Late trimester 34	Late trimester 32	Mid trimester 24	
	LOCATION	Bilateral	Bilateral	Unilateral left	Unilateral right	Unilateral left	Unilateral left	
	FETAL GENDER	Male	Male	Male	Male	Female	Female	
	ANTENATAL ULTRASOUND FEATURES	<ul style="list-style-type: none"> <li>▪ Size (mm)</li> <li>▪ Structure</li> <li>▪ Vascularized</li> <li>▪ Systemic vessel</li> <li>▪ Associated findings</li> </ul>	Increased Cerebriform aspect	50 x 40 Multicystic with mixed content Heterogeneous	22 x 32 Hyperechoic Homogeneous Yes No No	38 x 29 Altered content Heterogeneous No No No	10 x 8 Hypoechoic Homogeneous No No No	30 x 20 Mixed component Heterogeneous Yes Yes (from the aorta) No
	FETAL MRI	<ul style="list-style-type: none"> <li>▪ When (WG)</li> <li>▪ Location</li> <li>▪ Structure</li> <li>▪ Associated findings</li> </ul>	NR NR	28 Bilateral	37 Unilateral left Prerenal Retrogastric Well limited, Heterogeneous, No hemorrhagic signs	35 Unilateral right	NR NR	29 and 33 Thoraco abdominal junction
	PRENATAL EVOLUTION	Stable	Increase	Increase	Increase	Decrease	Stable	Increase
PROPOSED PRENATAL DIAGNOSIS	Congenital adrenal hyperplasia	Adrenocortical macrocyst Beckwith Wiedemann Syndrome	Solid neuroblastoma, Adrenal haemorrhage	Adrenal haemorrhage, Cystic neuroblastoma	Isolated adrenal cyst	Left extralobar pulmonary sequestration		
<b>POSTNATAL MANAGEMENT</b>	COMPLEMENTARY EXAMS	<ul style="list-style-type: none"> <li>▪ Urinary catecholamine</li> <li>▪ Androgenic marquors</li> <li>▪ Postnatal ultrasound / MRI</li> </ul>	Negative 21-H deficiency	NR NR	Positive Normal	Negative Normal	Negative Normal	
		<ul style="list-style-type: none"> <li>▪ Lesion confirmed</li> <li>▪ Associated elements</li> </ul>	Yes No	NR NR	Yes Liver metastases (present at 1 month) Malignant cells	Yes No	Yes No	
		<ul style="list-style-type: none"> <li>▪ Histological findings post-surgery</li> </ul>	-	NR		NR	No malignant cells	Lung cells – no malignant cells
	EVOLUTION	Normal aspect at one week and one month with treatment	Termination of pregnancy Pathology: fetal macrosomia, macroglossia, bilateral nephromegaly, bilateral adrenal masses with haemorrhagic lesion on the left. No malignant cells.	Spontaneous regression, then stabilization	Spontaneous regression	Stabilization	Stabilization	
	THERAPEUTIC PROTOCOL	Hydrocortisone and fludrocortisone therapy	NR	Close monitoring, then, Surgery	Close monitoring	Monitoring (spontaneous regression)	Close monitoring	

272 Table 1: Observations of fetuses with atypical adrenal gland detected during routine prenatal ultrasound. (EGA: Estimated Gestational Age ; NR: Not Realize)

		BILATERAL LOCATION			UNILATERAL LOCATION			DIFFERENTIAL DIAGNOSIS
		<i>Abnormal appearance</i>		<i>Normal appearance</i>	<i>Abnormal appearance</i>			<i>Abnormal appearance</i>
		<b>Congenital Adrenal Hyperplasia</b>	<b>Adrenocortical macrocyst (Beckwith Wiedemann Syndrome)</b>	<b>GDM FGR FIS</b>	<b>Neuroblastoma</b>	<b>Adrenal haemorrhage</b>	<b>Adrenal Cyst</b>	<b>Extralobar pulmonary sequestration</b>
<i>Characteristic ultrasound signs</i>	<b>Trimester at discovery</b>	Mid trimester Late trimester	Mid trimester Late trimester	Mid trimester Late trimester	Late trimester	Late trimester	Mid trimester Late trimester	Mid trimester Late trimester
	<b>Preference Location</b>	Bilateral	Bilateral ++ (possible unilateral)	Bilateral	Right	No preference	No preference	Left
	<b>Prenatal Evolution</b>							
	Decrease	-	-	+	+	+++	+++	+
	Stable	+++	+	+++	++	++	++	++
	Increase	-	+++	+	+++	+	+	+++
<b>Vascularization</b>								
Vascularized mass	No	No	No	Yes	No	No	Yes	
Systemic artery	No	No	No	Yes	No	No	Yes	
<b>Description of the lesion</b>								
Well limited Structure	Yes Cerebriform pattern	No Cystic, Heterogeneous	Yes Increase in size without abnormal aspect	Yes Homogeneous (solid form) or Heterogeneous (cystic form)	Yes Heterogeneous	Yes Homogeneous	No Homogeneous (nodular form) or Heterogeneous (cystic-nodular form)	
<b>Elements of malignity</b> <i>Partition, vegetation</i>	No	No	No	Yes	No	No	No	
<i>Prognostic Elements</i>	<b>Fetal complications</b>	No	Organomegaly, Macroglossia, Omphalocele Polyhydramnios	No	Possible metastases (liver ++, lungs, skull, bone marrow, placenta)	No	No	Mass effect (fattening diaphragmatic cupolas +/- pleural effusion)
	<b>Prognosis</b>	Excellent with treatment	Dependent on associated malformations	Excellent	Good (90% cure if early detection)	Excellent	Excellent	Good


273

274

Table 2: Check list of characteristic ultrasound signs according to more frequent diagnosis of suprarenal mass and prognostic elements associated.

275 (GDM: Gestational Diabetes Mellitus, FGR: fetal growth restriction, FIS: Fetal inflammatory syndrome)

For Review Only



276


277 *Figure 1: Normal appearance of adrenal glands (arrow) according to the trimester of pregnancy. (a) in the first trimester, in parasagittal view it appears as*  
278 *hypoechoic triangle measuring grossly half the height of the kidney. (b) Then, from the second trimester onward, in axial view it takes an oval structure, (c)*  
279 *with appearance of a hypoechoic cortex and a hyperechoic medullary during late trimester ultrasound.*

For Review Only



281


282



283


284

*Figure 2: Diagnostic orientation of an atypical adrenal gland according to its type of location and appearance*




285

286 *Figure 3: Congenital adrenal hyperplasia at 24 estimated gestational age, with a typical "cerebriform aspect" (a, arrow), associated with a virilization of*  
287 *the genitalia (b) in a male fetus.*




288

289 *Figure 4: Ultrasound at 25 estimated gestational age in a male fetus showing bilateral partially hemorrhagic cysts on both adrenal glands. This typical*  
 290 *ultrasound appearance, on the left (a) and on the right (b) associated with macroglossia (c) was consistent with adrenocortical macrocyst in the setting of*  
 291 *Beckwith – Wiedemann syndrome. Fetal MRI was helpful, confirming the atypical appearance of the adrenals, here in frontal (d) and transverse view (e) on*  
 292 *BFTE sequences. (LAG: Left Adrenal Gland; LK: Left Kidney; RAG: Right Adrenal Gland)*



293


294 *Figure 5: A prenatal well limited, heterogeneous (a) and vascularized (b) mass was diagnosed as a solid neuroblastoma at 35 estimated gestational age in a*  
 295 *male fetus. Coronal (c) and sagittal (d) T2-weighted MR images show slightly hyperintense mass of the left adrenal compartment. (e) Apparent diffusion*  
 296 *coefficient map shows hypointensity mass that confirming the malignant hypothesis.*



297

298 *Figure 6: A heterogeneous and avascular mass of the right adrenal gland, with a changing sonographic pattern through follow-up, on a axial (a1), parasagittal*  
 299 *(a2) and frontal views (b) is the typical aspect of an adrenal hemorrhage, at 34 weeks estimated gestational age, in a male fetus. Parasagittal (c) and*  
 300 *transverse (d) T2- HASTE weighted images show hematic sediment confirming the hemorrhagic composition of the mass.*

For Review Only




302

303


304

*Figure 7: A well limited, isolated, hypoechogenic and homogeneous cyst located above the left kidney in axial (a) and parasagittal (b) views at 32 weeks estimated gestational age in a female fetus, consistent with an adrenal cyst.*




*Figure 8: An extralobar pulmonary sequestration in a female fetus, represented by a heterogeneous mass, with mixed component, under the left diaphragm (a) vascularized by a multiple feeding arteries identified on Color Doppler Imaging (b) at 24 weeks estimated gestational age. (c) Coronal SSFSE T2w confirms the left lower lobe mass with systemic feeding vessels (arrow) coursing into the mass.*






Normal appearance of adrenal glands (arrow) according to the trimester of pregnancy. (a) in the first trimester, in parasagittal view it appears as hypoechoic triangle measuring grossly half the height of the kidney. (b) Then, from the second trimester onward, in axial view it takes an oval structure, (c) with appearance of a hypoechoic cortex and a hyperechoic medullary during late trimester ultrasound.

581x201mm (72 x 72 DPI)




Diagnostic orientation of an atypical adrenal gland according to its type of location and appearance




Congenital adrenal hyperplasia at 24 estimated gestational age, with a typical "cerebriform aspect" (a, arrow), associated with a virilization of the genitalia (b) in a male fetus.

490x210mm (72 x 72 DPI)




Ultrasound at 25 estimated gestational age in a male fetus showing bilateral partially hemorrhagic cysts on both adrenal glands. This typical ultrasound appearance, on the left(a) and on the right (b) associated with macroglossia (c) was consistent with adrenocortical macrocyst in the setting of Beckwith – Wiedemann syndrome. Fetal MRI was helpful, confirming the atypical appearance of the adrenals, here in frontal (d) and transverse view (e) on BFTE sequences. (LAG: Left Adrenal Gland; LK: Left Kidney; RAG: Right Adrenal Gland)

553x352mm (72 x 72 DPI)




A prenatal well limited, heterogeneous (a) and vascularized (b) mass was diagnosed as a solid neuroblastoma at 35 estimated gestational age in a male fetus. Coronal (c) and sagittal (d) T2-weighted MR images show slightly hyperintense mass of the left adrenal compartment. (e) Apparent diffusion coefficient map shows hypointensity mass that confirming the malignant hypothesis.

570x320mm (72 x 72 DPI)




A heterogeneous and avascular mass of the right adrenal gland, with a changing sonographic pattern through follow-up, on a axial (a1), parasagittal (a2) and frontal views (b) is the typical aspect of an adrenal hemorrhage, at 34 weeks estimated gestational age, in a male fetus. Parasagittal (c) and transverse (d) T2-HASTE weighted images show hematic sediment confirming the hemorrhagic composition of the mass.

507x290mm (72 x 72 DPI)



A well limited, isolated, hypoechogenic and homogeneous cyst located above the left kidney in axial (a) and parasagittal (b) views at 32 weeks estimated gestational age in a female fetus, consistent with an adrenal cyst.

433x182mm (72 x 72 DPI)



An extralobar pulmonary sequestration in a female fetus, represented by a heterogeneous mass, with mixed component, under the left diaphragm (a) vascularized by a multiple feeding arteries identified on Color Doppler Imaging (b) at 24 weeks estimated gestational age. (c) Coronal SSFSE T2w confirms the left lower lobe mass with systemic feeding vessels (arrow) coursing into the mass.

639x258mm (72 x 72 DPI)